

Hardy's inequality in Orlicz-Sobolev spaces of variable exponent

Yoshihiro MIZUTA, Eiichi NAKAI,
Takao OHNO and Tetsu SHIMOMURA

(Received February 4, 2010)

Abstract. Our aim in this paper is to deal with a norm version of Hardy's inequality for Orlicz-Sobolev functions with $|\nabla u| \in L^{p(\cdot)} \log L^{p(\cdot)q(\cdot)}(\Omega)$ for an open set $\Omega \subset \mathbf{R}^n$. Here $p(\cdot)$ and $q(\cdot)$ are variable exponents satisfying log-Hölder and loglog-Hölder conditions, respectively. We are also concerned with the case when p attains the value 1 in some parts of the domain is included in the results.

Key words: variable exponent, Lebesgue space, Hardy's inequality, Sobolev embeddings.

1. Introduction and statement of results

In recent years, the generalized Lebesgue spaces have attracted more and more attention, in connection with the study of elasticity, fluid mechanics and differential equations with $p(\cdot)$ -growth; see for example Kováčik-Rákosník [17], Musielak [24], Orlicz [25] and Růžička [26].

Let \mathbf{R}^n denote the n -dimensional Euclidean space. In this paper, following Cruz-Uribe and Fiorenza [1], we consider variable exponents $p(\cdot)$ and $q(\cdot)$ are continuous functions on \mathbf{R}^n satisfying:

$$(p1) \quad 1 \leq p^- = \inf_{x \in \mathbf{R}^n} p(x) \leq \sup_{x \in \mathbf{R}^n} p(x) = p^+ < \infty;$$

$$(p2) \quad |p(x) - p(y)| \leq \frac{C_{\log}}{\log(e + 1/|x - y|)} \quad \text{whenever } x \in \mathbf{R}^n \text{ and } y \in \mathbf{R}^n;$$

$$(p3) \quad |p(x) - p(y)| \leq \frac{C_{\log}}{\log(e + |x|)} \quad \text{whenever } |y| \geq |x|/2;$$

$$(q1) \quad -\infty < q^- = \inf_{x \in \mathbf{R}^n} q(x) \leq \sup_{x \in \mathbf{R}^n} q(x) = q^+ < \infty;$$

$$(q2) \quad |q(x) - q(y)| \leq \frac{C_{\log \log}}{\log(e + \log(e + 1/|x - y|))} \quad \text{whenever } x \in \mathbf{R}^n \text{ and } y \in \mathbf{R}^n.$$

2000 Mathematics Subject Classification : Primary 46E30, 42B25.

The second author was supported by Grant-in-Aid for Scientific Research (C), No. 20540167, Japan Society for the Promotion of Science.

Condition (p3) implies that $p_\infty = \lim_{|x| \rightarrow \infty} p(x)$ exists and

$$|p(x) - p_\infty| \leq \frac{C_{\log}}{\log(e + |x|)} \quad \text{for all } x \in \mathbf{R}^n. \quad (1.1)$$

Set

$$\Phi_{p(\cdot), q(\cdot)}(x, t) = (t(\log(c_0 + t))^{q(x)})^{p(x)};$$

here, we assume the existence of $c_0 > e$ such that

(Φ_1) $\Phi_{p(\cdot), q(\cdot)}(x, \cdot)$ is convex on $[0, \infty)$ for every $x \in \mathbf{R}^n$.

We note by a computation of the second derivative of $\Phi(x, \cdot)$ that if there is a positive constant C_0 such that

$$C_0(p(x) - 1) + p(x)q(x) \geq 0, \quad (1.2)$$

then condition (Φ_1) holds. For example, if $p^- > 1$, then (1.2) is satisfied with $C_0 \geq -p^-q^-/(p^- - 1)$; if $p^- = 1$ and $q^- \geq 0$, then (1.2) is also satisfied with every $C_0 \geq 0$. For later use it is convenient to see from (Φ_1) that

(Φ_2) $t^{-1}\Phi_{p(\cdot), q(\cdot)}(x, t)$ is nondecreasing on $(0, \infty)$ for fixed $x \in \mathbf{R}^n$.

Let Ω be an open set in \mathbf{R}^n . Define the $\Phi_{p(\cdot), q(\cdot)}(\Omega)$ norm by

$$\|f\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} = \inf \left\{ \lambda > 0 : \int_{\Omega} \Phi_{p(\cdot), q(\cdot)} \left(x, \frac{|f(x)|}{\lambda} \right) dx \leq 1 \right\}$$

and denote by $\Phi_{p(\cdot), q(\cdot)}(\Omega)$ the space of all measurable functions f on Ω with $\|f\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} < \infty$.

We define the variable exponent Sobolev–Orlicz space by

$$W^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega) = \{u \in \Phi_{p(\cdot), q(\cdot)}(\Omega) : |\nabla u| \in \Phi_{p(\cdot), q(\cdot)}(\Omega)\}.$$

The norm $\|u\|_{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega) = \|u\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} + \|\nabla u\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)}$ makes $W^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$ a Banach space. Further we denote the closure of $C_0^\infty(\Omega)$ in $W^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$ by $W_0^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$, which is extended to be 0 outside Ω . In case $q \equiv 0$, $\Phi_{p(\cdot), q(\cdot)}(\Omega)$ and $W^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$ are denote by $L^{p(\cdot)}(\Omega)$ and $W^{1, p(\cdot)}(\Omega)$ for simplicity. For fundamental properties of these spaces, see, for example, Kováčik and Rákosník [17].

We denote by $B(x, r)$ the open ball centered at x of radius r . For a measurable set E , we denote by $|E|$ the Lebesgue measure of E .

Recently Hästö [11, Theorem 3.2] proved the following:

Theorem A *Let $\Omega \neq \mathbf{R}^n$ be an open set. Suppose $1 < p^- \leq p^+ < \infty$. Assume that Ω satisfies the measure density condition, that is, there exists a constant $k > 0$ such that*

$$|B(z, r) \cap \Omega^c| \geq k|B(z, r)| \quad (1.3)$$

for every $z \in \partial\Omega$ and $r > 0$ (see [8]). Then there exist positive constants C and b_0 such that the inequality

$$\|\delta^{b-1}u\|_{L^{p(\cdot)}(\Omega)} \leq C\|\delta^b|\nabla u|\|_{L^{p(\cdot)}(\Omega)}$$

holds for all $u \in W_0^{1,p(\cdot)}(\Omega)$ and all $0 \leq b < b_0$, where $\delta(x) = \text{dist}(x, \partial\Omega)$ and the constants C, b_0 depend on p^+, p^-, n, k .

This gives an Harjulehto, Hästö and Koskenoja [10, Theorem 3.3] in the case when Ω is bounded. In the constant exponent case, Theorem A is given by Hajlasz [7, Theorem 1]; for related results, see also Edmunds and Evans [4], Kufner and L. E. Persson [18], Wannebo [30], Kinnunen and Martio [15].

Our aim in this paper is to give Hardy's inequality for $W_0^{1,\Phi_{p(\cdot),q(\cdot)}}(\Omega)$, as an extension of Theorem A. For $0 \leq a \leq 1$, set

$$1/p_a^\sharp(x) = 1/p(x) - a/n$$

and

$$\Phi_{p_a^\sharp(\cdot),q(\cdot)}(x, t) = (t(\log(c_0 + t))^{q(x)})^{p_a^\sharp(x)}.$$

Note that

$$\Phi_{p_a^\sharp(\cdot),q(\cdot)}(x, t) = (\Phi_{p(\cdot),q(\cdot)}(x, t))^{p_a^\sharp(x)/p(x)}$$

and $\Phi_{p_a^\sharp(\cdot),q(\cdot)}$ also satisfies (Φ_1) , since $p_a^\sharp(x)/p(x) \geq 1$.

Theorem 1.1 *Let $\Omega \neq \mathbf{R}^n$ be an open set satisfying (1.3). Suppose $1 < p^- \leq p^+ < \infty$, $0 < A \leq 1$ and $0 < A < n/p^+$. Then there exist $C > 0$ and*

$0 < b_0 < 1$ depending on A such that

$$\|\delta^{a+b-1}u\|_{\Phi_{p_a^\sharp(\cdot), q(\cdot)}(\Omega)} \leq C\|\delta^b|\nabla u|\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)}$$

for all $u \in W_0^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$, $0 \leq a \leq A$ and $0 \leq b \leq b_0$.

This theorem extends Theorem A given by Hästö [11], whose crucial idea is a partition norm on $L^{p(\cdot)}(\mathbf{R}^n)$. We give a straightforward and simple proof of Theorem 1.1 in a quite different manner. In fact, we apply Poincaré's inequality and the boundedness of maximal functions, following the idea by Hedberg [13].

If $p^- > n$, then we do not need the measure density condition to derive Poincaré's inequality. In fact we will show the following result, which gives an extension of Harjulehto, Hästö and Koskenoja [10, Theorem 3.5].

Theorem 1.2 *Let $\Omega \neq \mathbf{R}^n$ be an open set. Suppose $n < p^- \leq p^+ < \infty$ and $0 < A < n/p^+$. Then there exist $C > 0$ and $0 < b_0 < 1$ depending on A such that*

$$\|\delta^{a+b-1}u\|_{\Phi_{p_a^\sharp(\cdot), q(\cdot)}(\Omega)} \leq C\|\delta^b|\nabla u|\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)}$$

for all $u \in W_0^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$, $0 \leq a \leq A$ and $0 \leq b \leq b_0$.

Finally we are concerned with the case $p^- \geq 1$:

Theorem 1.3 *Let $\Omega \neq \mathbf{R}^n$ be an open set satisfying (1.3). $0 < A < \min\{1, n/p^+\}$ and $\gamma > 1$. Let*

$$L(t) = (\log(c_0 + t + 1/t))^{-\gamma}.$$

Then there exists a constant $C > 0$ depending on A and γ such that

$$\int_{\Omega} \Phi_{p_a^\sharp(\cdot), q(\cdot)}(x, \delta(x)^{a-1}|u(x)|) L(\delta(x)^{a-1}|u(x)|) dx \leq C$$

for all $u \in W_0^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$ with $\|\nabla u\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} \leq 1$ and $0 \leq a \leq A$.

Theorem 1.3 is not always valid when $\gamma = 1$ (see Remark 4.3). The case $a = 1$, that is, Sobolev inequality for $W_0^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$, is given in [12, Theorem 1.1], which was an extension of the results by Harjulehto and Hästö

[9, Proposition 4.2(1)] and Hästö [11, Theorem 3.4].

For further related results, see Samko [27], [28], Kokilashvili and Samko [16], Diening and Samko [3], Humberto and Samko [14] and Futamura, Mizuta and Shimomura [6].

2. Proof of Theorem 1.1.

Throughout this paper, let C denote various constants independent of the variables in question and $C(a, b, \dots)$ be a constant that depends on a, b, \dots .

Denote by $W_{loc}^{1,1}(\mathbf{R}^n)$ the class of all functions u such that $\varphi u \in W^{1,1}(\mathbf{R}^n)$ for every $\varphi \in C_0^\infty(\mathbf{R}^n)$.

First let us begin with the following lemma (see the proof of Hajlasz [7, Proposition 1]).

Lemma 2.1 *Let $\Omega \neq \mathbf{R}^n$ be an open set satisfying (1.3). Then there exists a constant $C = C(n, k) > 0$ such that*

$$|u(x)| \leq C \int_{B(x, 2\delta(x))} |x - y|^{1-n} |\nabla u(y)| dy$$

for almost every $x \in \Omega$, whenever $u \in W_{loc}^{1,1}(\mathbf{R}^n)$ and $u = 0$ outside Ω .

Proof. Let $u \in W_{loc}^{1,1}(\mathbf{R}^n)$ and $u = 0$ outside Ω . By (1.3), we obtain

$$\begin{aligned} |u(x)| &= \frac{1}{|B(x, 2\delta(x)) \cap \Omega^c|} \int_{B(x, 2\delta(x)) \cap \Omega^c} |u(x) - u(y)| dy \\ &\leq \frac{C}{|B(x, 2\delta(x))|} \int_{B(x, 2\delta(x))} |u(x) - u(y)| dy \\ &\leq C \int_{B(x, 2\delta(x))} |x - y|^{1-n} |\nabla u(y)| dy \end{aligned}$$

for almost every $x \in \Omega$. □

For a locally integrable function f on \mathbf{R}^n , we consider the maximal function

$$Mf(x) = \sup_{r>0} \frac{1}{|B(x, r)|} \int_{B(x, r)} |f(y)| dy.$$

We know the following result concerning the boundedness of maximal functions in $\Phi_{p(\cdot),q(\cdot)}(\Omega)$.

Lemma 2.2 ([5, Proposition 2.2]) *Suppose $p^- > 1$. Then there exists a constant $C > 0$ such that*

$$\|Mf\|_{\Phi_{p(\cdot),q(\cdot)}(\mathbf{R}^n)} \leq C\|f\|_{\Phi_{p(\cdot),q(\cdot)}(\mathbf{R}^n)}$$

for all $f \in \Phi_{p(\cdot),q(\cdot)}(\mathbf{R}^n)$.

Lemma 2.3 *Let $\Omega \neq \mathbf{R}^n$ be an open set satisfying (1.3). Suppose $1 < p^- \leq p^+ < \infty$. Then there exist $C > 0$ and $0 < b_0 < 1$ such that*

$$\|\delta^{b-1}u\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)} \leq C\|\delta^b|\nabla u|\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)}$$

for all $u \in W_0^{1,\Phi_{p(\cdot),q(\cdot)}}(\Omega)$ and $0 \leq b \leq b_0$.

Proof. Since $\Omega \neq \mathbf{R}^n$, without loss of generality, we may assume that the origin is on the boundary $\partial\Omega$, that is,

$$0 \in \partial\Omega. \quad (2.1)$$

We first treat $u \in C_0^\infty(\Omega)$. Applying Lemma 2.1 to $\delta^b u$, we have

$$\delta(x)^b|u(x)| \leq C \int_{B(x,2\delta(x))} |x-y|^{1-n} \{b\delta(y)^{b-1}|u(y)| + \delta(y)^b|\nabla u(y)|\} dy, \quad (2.2)$$

so that

$$\delta(x)^{b-1}|u(x)| \leq CbM(\delta^{b-1}|u|)(x) + CM(\delta^b|\nabla u|)(x).$$

In view of Lemma 2.2, we find

$$\|\delta^{b-1}u\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)} \leq Cb\|\delta^{b-1}u\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)} + C\|\delta^b|\nabla u|\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)},$$

which gives

$$(1 - Cb)\|\delta^{b-1}u\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)} \leq C\|\delta^b|\nabla u|\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)}.$$

Now it suffices to take b_0 such that $1 - Cb_0 > 0$.

We next treat $u \in W_0^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$ with compact support. Then we can find a sequence $\varphi_j \in C_0^\infty(\Omega)$ such that $\varphi_j \rightarrow u$ in $W_0^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$. Suppose $u = 0$ outside $B(0, R)$. Then we may assume that $\varphi_j = 0$ outside $B(0, 2R)$. By the above discussions we have

$$\|\delta^{b-1} \varphi_j\|_{\Phi_{p(\cdot), q(\cdot)}}(\Omega) \leq C \|\delta^b |\nabla \varphi_j|\|_{\Phi_{p(\cdot), q(\cdot)}}(\Omega).$$

Since $\|\delta^b |\nabla(\varphi_j - u)|\|_{\Phi_{p(\cdot), q(\cdot)}}(\Omega)$ tends to zero as $j \rightarrow \infty$, we obtain

$$\|\delta^{b-1} u\|_{\Phi_{p(\cdot), q(\cdot)}}(\Omega) \leq C \|\delta^b |\nabla u|\|_{\Phi_{p(\cdot), q(\cdot)}}(\Omega).$$

Finally we treat a general $u \in W_0^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$. For $N > 1$ we consider a continuous function h_N on $[0, \infty)$ and N^* such that $0 \leq h_N \leq 1$ on $[0, \infty)$, $h_N = 0$ on $[0, N]$, $h_N = 0$ on $[N^*, \infty]$, $h_N(t) \leq t^{-1}$ for $t \in [N, N^*]$ and

$$\int_N^{N^*} h_N(t) dt = 1;$$

the existence of N^* is assured since $\int_N^\infty t^{-1} dt = \infty$. Set

$$H_N(x) = 1 - \int_0^{|x|} h_N(t) dt$$

for $x \in \mathbf{R}^n$. Then we know as above that

$$\begin{aligned} & \|\delta^{b-1}(H_N u)\|_{\Phi_{p(\cdot), q(\cdot)}}(\mathbf{R}^n) \\ & \leq C \|\delta^b |\nabla(H_N u)|\|_{\Phi_{p(\cdot), q(\cdot)}}(\mathbf{R}^n) \\ & \leq C \|\delta^b |\nabla H_N| u\|_{\Phi_{p(\cdot), q(\cdot)}}(\mathbf{R}^n) + C \|\delta^b H_N |\nabla u|\|_{\Phi_{p(\cdot), q(\cdot)}}(\mathbf{R}^n) \\ & \leq C \| |x|^b |\nabla H_N| u\|_{\Phi_{p(\cdot), q(\cdot)}}(\mathbf{R}^n) + C \|\delta^b |\nabla u|\|_{\Phi_{p(\cdot), q(\cdot)}}(\mathbf{R}^n). \end{aligned}$$

Since $|\nabla H_N(x)| \leq h_N(|x|) \leq |x|^{-1}$ for $x \in \Omega$, $\| |x|^b |\nabla H_N| u\|_{\Phi_{p(\cdot), q(\cdot)}}(\mathbf{R}^n)$ tends to 0 as $N \rightarrow \infty$, so that

$$\begin{aligned} \|\delta^{b-1} u\|_{\Phi_{p(\cdot), q(\cdot)}}(\mathbf{R}^n) & \leq \liminf_{N \rightarrow \infty} \|\delta^{b-1}(H_N u)\|_{\Phi_{p(\cdot), q(\cdot)}}(\mathbf{R}^n) \\ & \leq C \|\delta^b |\nabla u|\|_{\Phi_{p(\cdot), q(\cdot)}}(\mathbf{R}^n), \end{aligned}$$

which completes the proof. \square

Note that

$$\begin{aligned} & \delta(x)^{a-1} \int_{B(x, 2\delta(x))} |x-y|^{1-n} f(y) dy \\ & \leq 2^{1-a} \int_{B(x, 2\delta(x))} |x-y|^{a-n} f(y) dy \end{aligned} \quad (2.3)$$

for $x \in \Omega$, $0 \leq a \leq 1$ and nonnegative measurable function f on \mathbf{R}^n . To give its estimate, we prepare the following result.

Lemma 2.4 *Let $0 < A < n/p^+$. Then there exists a constant $C > 0$ depending on A such that*

$$I \equiv \int_{B(x, 2|x|) \setminus B(x, r)} |x-y|^{a-n} f(y) dy \leq Cr^{a-n/p(x)} (\log(c_0 + r^{-1}))^{-q(x)}$$

for all $x \in \mathbf{R}^n$, $r > 0$, $0 \leq a \leq A$ and $f \geq 0$ with $\|f\|_{\Phi_{p(\cdot), q(\cdot)}(\mathbf{R}^n)} \leq 1$.

This follows from [12, Lemmas 3.1 and 3.2]. Actually, since the estimate of the integral outside $B(x, 2|x|)$ given in the proof of [12, Lemma 3.1] is not needed here, we use [12, Lemma 3.1] if r is large, and both of [12, Lemmas 3.1 and 3.2] if r is small.

Next consider

$$J \equiv \delta(x)^{a-1} \int_{B(x, 2\delta(x))} |x-y|^{1-n} f(y) dy$$

for $x \in \Omega$ and $f \geq 0$. When $a = 0$, we find

$$J \leq CMf(x).$$

Lemma 2.5 *Let $\Omega \neq \mathbf{R}^n$ be an open set. If $0 < A \leq 1$ and $0 < A < n/p^+$, then there exists a constant $C > 0$ depending on A such that*

$$J \leq C\{Mf(x)\}^{1-ap(x)/n} \{\log(c_0 + Mf(x))\}^{-ap(x)q(x)/n}$$

for all $x \in \Omega$, $0 \leq a \leq A$ and $f \geq 0$ with $\|f\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} \leq 1$.

Proof. We have only to consider the case $a > 0$. Let $f \geq 0$ with

$$\|f\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} \leq 1.$$

First suppose $\{Mf(x)\}^{-p(x)/n} \{\log(c_0 + Mf(x))\}^{-p(x)q(x)/n} \leq \delta(x)$; we set $f = 0$ outside Ω as before. For $0 < r < \delta(x)$, we have by (2.1) and Lemma 2.4

$$\begin{aligned} J &\leq C \left\{ \delta(x)^{a-1} r Mf(x) + \int_{B(x, 2\delta(x)) \setminus B(x, r)} |x-y|^{a-n} f(y) dy \right\} \\ &\leq C \{ r^a Mf(x) + r^{a-n/p(x)} (\log(c_0 + 1/r))^{-q(x)} \}. \end{aligned}$$

Now, letting $r = \{Mf(x)\}^{-p(x)/n} \{\log(c_0 + Mf(x))\}^{-p(x)q(x)/n} (\leq \delta(x))$, we have

$$J \leq C \{Mf(x)\}^{1-ap(x)/n} \{\log(c_0 + Mf(x))\}^{-ap(x)q(x)/n}.$$

Next suppose $\{Mf(x)\}^{-p(x)/n} \{\log(c_0 + Mf(x))\}^{-p(x)q(x)/n} > \delta(x)$. Then we have

$$\begin{aligned} J &\leq C \delta(x)^a Mf(x) \\ &\leq C \{Mf(x)\}^{1-ap(x)/n} \{\log(c_0 + Mf(x))\}^{-ap(x)q(x)/n}, \end{aligned}$$

which proves the case $a > 0$. □

Now we are ready to prove Theorem 1.1.

Proof of Theorem 1.1. Let $u \in W_0^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$ with $\|\delta^b |\nabla u|\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} \leq 1$. We have by (2.2)

$$\delta(x)^{a+b-1} |u(x)| \leq C \delta(x)^{a-1} \int_{B(x, 2\delta(x))} |x-y|^{1-n} f(y) dy, \quad (2.4)$$

where $f(y) = b\delta(y)^{b-1} |u(y)| + \delta(y)^b |\nabla u(y)|$. Here note from Lemma 2.3 that

$$\|f\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} \leq C.$$

Let $0 < A \leq 1$, $0 < A < n/p^+$ and $0 \leq a \leq A$. Then we obtain by Lemma 2.5

$$\delta(x)^{a+b-1} |u(x)| \leq C Mf(x)^{p(x)/p_a^+(x)} (\log(c_0 + Mf(x)))^{-ap(x)q(x)/n}.$$

By Lemmas 2.2 and 2.3, we have

$$\begin{aligned}
\|\delta^{a+b-1}u\|_{\Phi_{p_a^\sharp(\cdot), q(\cdot)}(\Omega)} &\leq C\|Mf\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} \leq C\|f\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} \\
&\leq Cb\|\delta^{b-1}u\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} + C\|\delta^b|\nabla u|\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} \\
&\leq C\|\delta^b|\nabla u|\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)},
\end{aligned}$$

which proves the theorem. \square

Remark 2.6 Let Ω be a bounded domain in \mathbf{R}^n satisfying (1.3), $n \leq p^- \leq p^+ < \infty$ and $1 - 1/p^- > q^+$. Then there exist $C_1, C_2 > 0$ and $0 < b_0 < 1$ such that

$$\int_{\Omega} \exp(C_1(\delta(x)^{a+b-1}u(x))^{p(x)/(p(x)-p(x)q(x)-1)})dx \leq C_2$$

for all $u \in W_0^{1, \Phi_{p(\cdot), q(\cdot)}}(\Omega)$ with $\|\delta^b|\nabla u|\|_{\Phi_{p(\cdot), q(\cdot)}(\Omega)} \leq 1$, $n/p^- \leq a \leq 1$ and $0 \leq b \leq b_0$.

In fact, (2.4) gives

$$\delta(x)^{a+b-1}|u(x)| \leq C \int_{B(x, 2\delta(x))} |x-y|^{a-n} f(y) dy,$$

where $f(y) = b\delta(y)^{b-1}|u(y)| + \delta(y)^b|\nabla u(y)|$ and $n/p^- \leq a \leq 1$. As in the proof of Theorem 1.1 and [22, Lemma 4.4], we obtain

$$\delta(x)^{a+b-1}|u(x)| \leq C(\log(c_0 + Mf(x)))^{(p(x)-p(x)q(x)-1)/p(x)},$$

so that

$$\exp((C^{-1}\delta(x)^{a+b-1}|u(x)|)^{p(x)/(p(x)-p(x)q(x)-1)}) \leq c_0 + Mf(x).$$

By integration we establish the required inequality.

3. Proof of Theorem 1.2

For a proof of Theorem 1.2, we prepare the following lemma instead of Lemma 2.3. We write $f \sim g$ if there exists a constant C so that $C^{-1}g \leq f \leq Cg$.

Lemma 3.1 *Let $\Omega \neq \mathbf{R}^n$ be an open set. Suppose $n < p^- \leq p^+ < \infty$. Then there exist $C > 0$ and $0 < b_0 < 1$ such that*

$$\|\delta^{b-1}u\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)} \leq C\|\delta^b|\nabla u|\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)}$$

for all $u \in W_0^{1,\Phi_{p(\cdot),q(\cdot)}}(\Omega)$ and $0 \leq b \leq b_0$.

Proof. Let $u \in W_0^{1,\Phi_{p(\cdot),q(\cdot)}}(\Omega)$. Suppose $n < p^- \leq p^+ < \infty$. Since $u \in W_{loc}^{1,s}(\mathbf{R}^n)$ with $n < s < p^-$, u is (Hölder) continuous in Ω and

$$|u(x)| \leq C \left(\delta(x)^{s-n} \int_{B(x,2\delta(x))} |\nabla u(y)|^s dy \right)^{1/s}$$

for every $x \in \Omega$; for this, see also [7, Proposition 1], [15, (3.1)] and [23, Theorem 1]. This implies that

$$[\delta(x)^{-1}|u(x)|]^s \leq C\delta(x)^{-n} \int_{B(x,2\delta(x))} |\nabla u(y)|^s dy. \quad (3.1)$$

Applying (3.1) to $\delta^b u$ as in Lemma 2.3, we have

$$[\delta(x)^{b-1}|u(x)|]^s \leq C\delta(x)^{-n} \int_{B(x,2\delta(x))} [b\delta(y)^{b-1}|u(y)| + \delta(y)^b|\nabla u(y)|]^s dy, \quad (3.2)$$

so that

$$[\delta(x)^{b-1}|u(x)|]^s \leq C\{b^s M([\delta^{b-1}u]^s)(x) + M([\delta^b|\nabla u|^s])(x)\}.$$

In view of Lemma 2.2, we find

$$\begin{aligned} & \|[\delta^{b-1}u]^s\|_{\Phi_{p(\cdot)/s,sq(\cdot)}(\Omega)} \\ & \leq C\{b^s \|[\delta^{b-1}u]^s\|_{\Phi_{p(\cdot)/s,sq(\cdot)}(\Omega)} + \|[\delta^b|\nabla u|^s]\|_{\Phi_{p(\cdot)/s,sq(\cdot)}(\Omega)}\}, \end{aligned}$$

which gives

$$(1 - Cb^s) \|[\delta^{b-1}u]^s\|_{\Phi_{p(\cdot)/s,sq(\cdot)}(\Omega)} \leq C \|[\delta^b|\nabla u|^s]\|_{\Phi_{p(\cdot)/s,sq(\cdot)}(\Omega)}.$$

Since $\|f^s\|_{\Phi_{p(\cdot)/s,sq(\cdot)}(\Omega)} \sim \|f\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)}$, we obtain

$$(1 - Cb^s)\|\delta^{b-1}u\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)} \leq C\|\delta^b|\nabla u|\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)}.$$

Now it suffices to take b_0 such that $1 - Cb_0^s > 0$. \square

Proof of Theorem 1.2. Let $u \in W_0^{1,\Phi_{p(\cdot),q(\cdot)}}(\Omega)$. Suppose $n < s < p^- \leq p^+ < \infty$ and

$$\|\delta^b|\nabla u|\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)} \leq 1.$$

For $0 \leq a \leq A < n/p^+$, we have by (3.2)

$$\begin{aligned} [\delta(x)^{a+b-1}|u(x)|]^s &\leq C\delta(x)^{as-n} \int_{B(x,2\delta(x))} f(y)dy \\ &\leq C\delta(x)^{as-As} \int_{B(x,2\delta(x))} |x-y|^{As-n} f(y)dy, \end{aligned}$$

where $f = [b\delta(y)^{b-1}|u(y)| + \delta(y)^b|\nabla u(y)|]^s$. Here note from Lemma 3.1 that

$$\|f\|_{\Phi_{p(\cdot)/s,sq(\cdot)}(\Omega)} \leq C.$$

Hence, as in the proof of Lemma 2.5 with a , $p(x)$ and $q(x)$ replaced by as , $p(x)/s$ and $sq(x)$, respectively, we obtain

$$\delta(x)^{a+b-1}|u(x)| \leq CMf(x)^{p(x)/(sp_a^\sharp(x))}(\log(c_0 + Mf(x)))^{-ap(x)q(x)/n}.$$

By Lemmas 2.2 and 3.1, we have

$$\begin{aligned} &\|\delta^{a+b-1}u\|_{\Phi_{p_a^\sharp(\cdot),q(\cdot)}(\Omega)} \\ &\leq C\|Mf\|_{\Phi_{p(\cdot)/s,sq(\cdot)}(\Omega)} \leq C\|f\|_{\Phi_{p(\cdot)/s,sq(\cdot)}(\Omega)} \\ &\leq C\{b^s\|\delta^{b-1}u\|_{\Phi_{p(\cdot)/s,sq(\cdot)}(\Omega)} + \|[\delta^b|\nabla u|]^s\|_{\Phi_{p(\cdot)/s,sq(\cdot)}(\Omega)}\} \\ &\leq C\{b^s\|\delta^{b-1}u\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)} + \|\delta^b|\nabla u|\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)}\} \\ &\leq C\|\delta^b|\nabla u|\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)}, \end{aligned}$$

which proves the theorem. \square

4. Proof of Theorem 1.3

For a proof of Theorem 1.3, we prepare the following results.

Lemma 4.1 ([12, Lemmas 2.1–2.3]) *Let f be a nonnegative measurable function on Ω such that $\|f\|_{\Phi_{p(\cdot),q(\cdot)}(\Omega)} \leq 1$. Then there exists a constant $C > 0$ such that*

$$\{Mf(x)\}^{p(x)} \leq C\{Mg(x)(\log(c_0 + Mg(x)))^{-p(x)q(x)} + C(1 + |x|)^{-n}\},$$

and

$$\Phi_{p(\cdot),q(\cdot)}(x, Mf(x)) \leq C\{Mg(x) + (1 + |x|)^{-n}\},$$

where $g(y) = \Phi_{p(\cdot),q(\cdot)}(y, f(y))$.

The next lemma is proved along the same lines as in Stein [29, Chapter 1]; see also [20, Lemma 2.5].

Lemma 4.2 *Suppose $1 < \gamma \leq 2$. Then there exists a constant $C > 0$ such that*

$$\int_{\Omega} Mg(x)(\log(c_0 + Mg(x) + Mg(x)^{-1}))^{-\gamma} dx \leq C\|g\|_{L^1(\Omega)}$$

for all $g \in L^1(\Omega)$.

Proof. For $1 < \gamma \leq 2$, we see that $t(\log(\gamma + t + 1/t))^{-\gamma}$ is increasing on $(0, \infty)$ and

$$t(\log(c_0 + t + 1/t))^{-\gamma} \leq C(\gamma)t(\log(\gamma + t + 1/t))^{-\gamma}.$$

Hence

$$\begin{aligned} & \int_{\Omega} Mg(x)(\log(c_0 + Mg(x) + Mg(x)^{-1}))^{-\gamma} dx \\ & \leq C(\gamma) \int_{\Omega} Mg(x)(\log(\gamma + Mg(x) + Mg(x)^{-1}))^{-\gamma} dx \\ & = C(\gamma) \int_0^{\infty} \lambda(t) d(t(\log(\gamma + t + t^{-1}))^{-\gamma}), \end{aligned}$$

where $\lambda(t) = |\{x \in \Omega : Mg(x) > t\}|$. Here we note from [29, Theorem 1, Chapter 1] that

$$\lambda(t) \leq Ct^{-1} \int_{\{x \in \Omega : |g(x)| > t/2\}} |g(x)| \, dx$$

for $t > 0$. Now we obtain by Fubini's Theorem

$$\begin{aligned} & \int_{\Omega} Mg(x)(\log(c_0 + Mg(x) + Mg(x)^{-1}))^{-\gamma} dx \\ & \leq C \int_{\Omega} |g(x)| \left\{ \int_0^{2|g(x)|} t^{-1} d(t(\log(\gamma + t + t^{-1}))^{-\gamma}) \right\} dx \\ & \leq C \int_{\Omega} |g(x)| dx, \end{aligned}$$

as required. \square

Proof of Theorem 1.3. We may assume that $1 < \gamma \leq 2$. By Lemmas 2.1 and 2.5 we have

$$\begin{aligned} & \delta(x)^{a-1} |u(x)| \\ & \leq C \{M(|\nabla u|)(x)\}^{1-ap(x)/n} \{\log(c_0 + M(|\nabla u|)(x))\}^{-ap(x)q(x)/n} \end{aligned}$$

for $x \in \Omega$ and $0 \leq a \leq A (< \min\{1, n/p^+\})$. Hence Lemma 4.1 gives

$$\begin{aligned} & \Phi_{p_a^\sharp(\cdot), q(\cdot)}(x, \delta(x)^{a-1} |u(x)|) (\log(c_0 + \delta(x)^{a-1} |u(x)| + \delta(x)^{1-a} |u(x)|^{-1}))^{-\gamma} \\ & \leq C \Phi_{p(\cdot), q(\cdot)}(x, M(|\nabla u|)(x)) (\log(c_0 + M(|\nabla u|)(x) + M(|\nabla u|)(x)^{-1}))^{-\gamma} \\ & \leq C \{Mg(x)(\log(c_0 + Mg(x) + Mg(x)^{-1}))^{-\gamma} \\ & \quad + (1 + |x|)^{-n} (\log(c_0 + |x|))^{-\gamma}\}, \end{aligned}$$

where $g(y) = \Phi_{p(\cdot), q(\cdot)}(y, |\nabla u(y)|)$. Thus we obtain the required conclusion with the aid of Lemma 4.2. \square

Remark 4.3 Theorem 1.3 is not valid when $(p^- =) p^+ = 1$ and $\gamma = 1$.

To show this when $n = 1$, first consider the function

$$v(x) = (\log(\log(4/x)))^{-\beta}$$

for $\beta > 0$. Then $|\nabla v(x)| = \beta x^{-1}(\log(4/x))^{-1}(\log(\log(4/x)))^{-\beta-1}$ and

$$\int_0^1 |\nabla v(x)| dx < \infty.$$

On the other hand, if $0 < a < 1$ and $1/p_a^\sharp(x) = 1 - a$, then

$$\begin{aligned} & \int_0^{1/2} (x^{a-1}|v(x)|)^{p_a^\sharp(x)} (\log(c_0 + x^{a-1}|v(x)|))^{-1} dx \\ & \geq C \int_0^{1/2} x^{-1} (\log(\log(4/x)))^{-\beta/(1-a)} (\log(1/x))^{-1} dx = \infty \end{aligned}$$

when $-\beta/(1-a)+1 \geq 0$, that is, $0 < \beta \leq 1-a$. Now, letting $\varphi \in C^\infty((0, 1))$ such that $\varphi(x) = 1$ for $x \in (0, 1/2)$ and $\lim_{x \rightarrow 1-0} \varphi(x) = 0$, one may consider $u(x) = \varphi(x)v(x)$ for $0 < x < 1$.

References

- [1] Cruz-Urbe D. and Fiorenza A., *L log L results for the maximal operator in variable L^p spaces*. Trans. Amer. Math. Soc. **361** (2009), 2631–2647.
- [2] Diening L., *Maximal functions on generalized $L^{p(\cdot)}$ spaces*. Math. Inequal. Appl. **7**(2) (2004), 245–253.
- [3] Diening L. and Samko S., *Hardy inequality in variable exponent Lebesgue spaces*. Fract. Calc. Appl. Anal. **10**(1) (2007), 1–19.
- [4] Edmunds D. E. and Evans W. D., *Hardy operators, function spaces and embeddings*, Springer, 2004.
- [5] Maeda F.-Y., Mizuta Y. and Ohno T., *Approximate identities and Young type inequalities in variable Lebesgue-Orlicz spaces $L^{p(\cdot)}(\log L)^{q(\cdot)}$* , Ann. Acad. Sci. Fenn. Math. **35** (2010), 405–420.
- [6] Futamura T., Mizuta Y. and Shimomura T., *Integrability of maximal functions and Riesz potentials in Orlicz spaces of variable exponent*. J. Math. Anal. Appl. **366** (2010), 391–417.
- [7] Hajlasz P., *Pointwise Hardy inequalities*. Proc. Amer. Math. Soc. **127**(2) (1999), 417–423.
- [8] Hajlasz P., Koskela P. and Tuominen H., *Sobolev embeddings, extensions and measure density condition*. J. Funct. Anal. **254**(5) (2008), 1217–1234.
- [9] Harjulehto P. and Hästö P., *Sobolev inequalities for variable exponents at-*

- taining the values 1 and n . Publ. Mat. **52** (2008), 347–363.
- [10] Harjulehto P., Hästö P. and Koskenoja M., *Hardy's inequality in variable exponent Sobolev space*. Georgian Math. J. **12**(3) (2005), 431–442.
 - [11] Hästö P., *Local-to-global results in variable exponent spaces*. Math. Res. Lett. **16**(2) (2009), 263–278.
 - [12] Hästö P., Mizuta Y., Ohno T. and Shimomura T., *Sobolev inequalities for Orlicz spaces of two variable exponents*. Glasgow Math. J. **52** (2010), 227–240.
 - [13] Hedberg L. I., *On certain convolution inequalities*. Proc. Amer. Math. Soc. **36** (1972), 505–510.
 - [14] Humberto R. and Samko S., *Hardy type inequality in variable Lebesgue spaces*. Ann. Acad. Sci. Fenn. Math. **34** (2009), 279–289.
 - [15] Kinnunen J. and Martio O., *Hardy's inequalities for Sobolev functions*. Math. Res. Lett. **4**(4) (1997), 489–500.
 - [16] Kokilashvili V. and Samko, S. *Maximal and fractional operators in weighted $L^{p(x)}$ spaces*. Rev. Mat. Iberoamericana **20**(2) (2004), 493–515.
 - [17] Kováčik O. and Rákosník J., *On spaces $L^{p(x)}$ and $W^{k,p(x)}$* . Czechoslovak Math. J. **41** (1991), 592–618.
 - [18] Kufner A. and Persson L. E., *Weighted inequalities of Hardy type*, World Scientific Publishing Co., Inc., River Edge, NJ, 2003.
 - [19] Mizuta Y., *Potential theory in Euclidean spaces*, Gakkōtoshō, Tokyo, 1996.
 - [20] Mizuta Y., Ohno T. and Shimomura T., *Integrability of maximal functions for generalized Lebesgue spaces with variable exponent*. Math. Nachr. **281**(3) (2008), 386–395.
 - [21] Mizuta Y., Ohno T. and Shimomura T., *Sobolev's inequalities and vanishing integrability for Riesz potentials of functions in the generalized Lebesgue space $L^{p(\cdot)}(\log L)^{q(\cdot)}$* . J. Math. Anal. Appl. **345** (2008), 70–85.
 - [22] Mizuta Y., Ohno T. and Shimomura T., *Sobolev embeddings for Riesz potential spaces of variable exponents near 1 and Sobolev's exponent*. Bull. Sci. Math. **134**(1) (2010), 12–36.
 - [23] Mizuta Y. and Shimomura T., *Continuity and differentiability for weighted Sobolev spaces*. Proc. Amer. Math. Soc. **130** (2002), 2985–2994.
 - [24] Musielak J., *Orlicz spaces and modular spaces*. Lecture Notes Math. **1034**, Springer, 1983.
 - [25] Orlicz W., *Über konjugierte Exponentenfolgen*. Studia Math. **3** (1931), 200–211.
 - [26] Růžička M., *Electrorheological fluids: modeling and mathematical theory*. Lecture Notes in Math. **1748**, Springer, 2000.
 - [27] Samko S., *Hardy inequality in the generalized Lebesgue spaces*. Fract. Calc.

- Appl. Anal. **6**(4) (2003), 355–362.
- [28] Samko S., *Hardy-Littlewood-Stein-Weiss inequality in the Lebesgue spaces with variable exponent*. Fract. Calc. Appl. Anal. **6**(4) (2003), 421–440.
- [29] Stein E. M., *Singular integrals and differentiability properties of functions*, Princeton Univ. Press, Princeton, 1970.
- [30] Wannebo A., *Hardy inequalities*. Proc. Amer. Math. Soc. **109**(1) (1990), 85–95.

Y. Mizuta
Department of Mathematics
Graduate School of Science
Hiroshima University
Higashi-Hiroshima 739-8521, Japan
E-mail: yomizuta@hiroshima-u.ac.jp

E. Nakai
Department of Mathematics
Ibaraki University
Mito, Ibaraki 310-8512, Japan
E-mail: enakai@mx.ibaraki.ac.jp

T. Ohno
Faculty of Education and Welfare Science
Oita University
Dannoharu Oita-city 870-1192, Japan
E-mail: t-ohno@oita-u.ac.jp

T. Shimomura
Department of Mathematics
Graduate School of Education
Hiroshima University
Higashi-Hiroshima 739-8524, Japan
E-mail: tshimo@hiroshima-u.ac.jp