

## Research Article

# On Spectrum of the Laplacian in a Circle Perforated along the Boundary: Application to a Friedrichs-Type Inequality

**G. A. Chechkin,<sup>1,2</sup> Yu. O. Koroleva,<sup>1,3</sup> L.-E. Persson,<sup>2,3</sup> and P. Wall<sup>3</sup>**

<sup>1</sup> Department of Differential Equations, Faculty of Mechanics and Mathematics, Moscow Lomonosov State University, Moscow 119991, Russia

<sup>2</sup> Narvik University College, Postboks 385, 8505 Narvik, Norway

<sup>3</sup> Department of Engineering Science and Mathematics, Luleå University of Technology, 971 87 Luleå, Sweden

Correspondence should be addressed to Yu. O. Koroleva, yulia.koroleva@ltu.se

Received 24 May 2011; Accepted 30 August 2011

Academic Editor: Mayer Humi

Copyright © 2011 G. A. Chechkin et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In this paper, we construct and verify the asymptotic expansion for the spectrum of a boundary-value problem in a unit circle periodically perforated along the boundary. It is assumed that the size of perforation and the distance to the boundary of the circle are of the same smallness. As an application of the obtained results, the asymptotic behavior of the best constant in a Friedrichs-type inequality is investigated.

## 1. Introduction

We study a two-dimensional eigenvalue problem for the Laplace operator in a unit circle periodically perforated along the boundary. It is assumed that the size of perforation and the distance to the boundary of the circle are of the same smallness. The asymptotic behavior of the spectrum of the considered boundary-value problem is investigated in this paper. We construct and verify the asymptotic expansion for the eigenvalues with respect to the small parameter describing the microinhomogeneous structure of the domain. A similar problem was considered in [1] for the case of perforation located along the plane part of the boundary. The case studied in this paper is much more complicated since the eigenvalues of multiplicity more than one can appear. The technique for asymptotic analysis of such kind of problem can be found, for example, in [2, 3].

The obtained results are used for asymptotic expansion of the best constant in a Friedrichs-type inequality for functions from the space  $H^1$ , vanishing on the boundary of the perforation and satisfying homogeneous Neuman condition on the boundary of the circle. Analogous questions concerning the asymptotic behavior of the best constant in Friedrichs-type inequality in domains having microinhomogeneous structure in a neighborhood of the boundary were studied in [1, 4–11]. In the remaining part of this introduction, we will give a short description of some of the most important results in these papers to put the results obtained in this paper into a more general frame.

In paper [4], the authors proved a Friedrichs-type inequality for functions, having zero trace on the small periodically alternating pieces of the boundary of a two-dimensional domain. The total measure of the set, where the function vanishes, tends to zero. It turns out that for this case the constant in the Friedrichs-type inequality is bounded. Moreover, the precise asymptotics of the constant in the derived Friedrichs-type inequality is described as the small parameter characterizing the microinhomogeneous structure of the boundary, tends to zero.

Paper [5] is devoted to the asymptotic analysis of functions depending on the small parameter, which characterizes the microinhomogeneous structure of the domain where the functions are defined. The authors considered a boundary-value problem in a two-dimensional domain perforated nonperiodically along the boundary in the case when the diameter of circles and the distance between them have the same order. In particular, it was proved that the Dirichlet problem is the limit for the original problem. Moreover, some numerical simulations were used to illustrate the results. As an application, a Friedrichs-type inequality was derived for functions vanishing on the boundary of the cavities. It was proved that the constant in the obtained inequality is close to the constant in the inequality for functions from  $H^1$ . The three-dimensional case of the same problem is considered in [8].

In paper [9], the author considered a three-dimensional domain, which is aperiodically perforated along the boundary in the case when the diameter of the holes and the distance between them have the same order. A Friedrichs-type inequality was derived for functions from the space  $H^1$  vanishing on the boundaries of cavities. In particular, it was shown that the constant in the derived inequality tends to the constant of the classical inequality for functions from  $H^1$  when the small parameter describing the size of perforation tends to zero.

Paper [1] (see also [7]) deals with the construction of the asymptotic expansion for the first eigenvalue of a boundary-value problem for the Laplacian in a perforated domain. This asymptotics gives an asymptotic expansion for the best constant in a corresponding Friedrichs-type inequality.

Paper [11], is devoted to the Friedrichs-type inequality, where the domain is periodically and rarely perforated along the boundary. It is assumed that the functions satisfy homogeneous Neumann boundary conditions on the outer boundary and that they vanish on the perforation. In particular, it is proved that the best constant in the inequality converges to the best constant in a Friedrichs-type inequality as the size of the perforation goes to zero much faster than the period of perforation. The limit Friedrichs-type inequality is valid for functions in the Sobolev space  $H^1$ .

Some generalizations of Friedrichs-type inequalities are Hardy-type inequalities. There exist several books devoted to this topic, see [12–16]. The first attempts to generalize the classical results concerning Hardy-type inequalities in fixed domains to domains with microinhomogeneous structure one can find in [6, 10].

Paper [6] deals with a three-dimensional weighted Hardy-type inequality in the case when the domain  $\Omega$  is bounded and has nontrivial microstructure. It is assumed that the small holes are distributed periodically along the boundary. The main result is the validity of a weighted Hardy-type inequality for the class of functions from the Sobolev space  $H^1$  having zero trace on the small holes under the assumption that a weight function decreases to zero in a neighborhood of the microinhomogeneity on the boundary.

In paper [10], the author derived a new two-dimensional weighted Hardy-type inequality in a rectangle for the class of functions from the Sobolev space  $H^1$  vanishing on small alternating pieces of the boundary. The dependence of the best constant in the derived inequality on the small parameter describing the size of microinhomogeneity was established.

This paper is organized as follows: in Section 2 we give all necessary definitions and state the spectral problem. Section 3 is devoted to the construction of the leading terms of asymptotic expansion, while the complete expansions for the simple and multiple eigenvalues are constructed in Sections 4 and 5, respectively. The verification of the constructed asymptotics is given in Section 6. Finally, in Section 7, the obtained results are applied to describe the asymptotic behavior for the best constant in a Friederichs-type inequality considered in a perforated domain.

## 2. Preliminaries

Consider a unit circle  $\Omega$  centered at the origin. We introduce the polar system of coordinates  $(\theta, r)$  in  $\Omega$ . Introduce a small parameter  $\varepsilon = 2/N$ ,  $N \gg 1$ , and consider the open set  $B_\varepsilon$  which is the union of small sets periodically distributed along the boundary. Each of these small sets can be obtained from the neighboring one by rotation about the origin through the angle  $\varepsilon\pi$ . Finally, we define  $\Omega_\varepsilon = \Omega \setminus \overline{B_\varepsilon}$  and  $\partial B_\varepsilon = \Gamma_\varepsilon$ , see Figure 1. Let us describe the geometry of  $B_\varepsilon$  in details. Consider the semi-strip:

$$\Pi = \left\{ \xi : -\frac{\pi}{2} < \xi_1 < \frac{\pi}{2}, \xi_2 > 0 \right\}, \quad \Gamma := \left\{ \xi : -\frac{\pi}{2} < \xi_1 < \frac{\pi}{2}, \xi_2 = 0 \right\}. \quad (2.1)$$


Let  $B$  be an arbitrary two-dimensional open domain with a smooth boundary that is symmetric the vertical axis and lies in a disk of a fixed radius  $a < 1$  centered at the point  $(0, 1)$ , see Figure 2. Let  $B_a$  be the union of the  $\pi$ -integer translations of  $B$  along the axis  $\xi_1$ . Then we define  $B_\varepsilon$  as the image of  $B_a$  under the mapping  $\theta = \varepsilon\xi_1, r = 1 - \varepsilon\xi_2$ .

Consider the following spectral problem:


$$\begin{aligned} -\Delta u_\varepsilon &= \lambda_\varepsilon u_\varepsilon \quad \text{in } \Omega_\varepsilon, \\ u_\varepsilon &= 0 \quad \text{on } \Gamma_\varepsilon, \\ \frac{\partial u_\varepsilon}{\partial r} &= 0 \quad \text{on } \partial\Omega. \end{aligned} \quad (2.2)$$

The problem,

$$\begin{aligned} -\Delta u_0 &= \lambda_0 u_0 \quad \text{in } \Omega, \\ u_0 &= 0 \quad \text{on } \partial\Omega, \end{aligned} \quad (2.3)$$



**Figure 1:** Perforated circle.



**Figure 2:** Cell of periodicity.

is the limit one for (2.2). This fact can be established analogously as in [17, 18], by using the same technique.

*Remark 2.1.* In particular, it can be proved that the number of eigenvalues (bearing in mind the multiplicities) of the original problem converging to the eigenvalue of the limit (homogenized) problem is equal to the multiplicity of the mentioned eigenvalue of the limit problem (for the method of proof see, e.g., [19]).

*Remark 2.2.* The limit spectral problem (2.3) is studied very well. In particular, if the eigenvalue  $\lambda_0$  is simple, then the corresponding eigenfrequency  $k_0 = \sqrt{\lambda_0}$  of (2.3) is the zero-point of the Bessel-function  $\mathcal{J}_0$ , and the corresponding eigenfunction has the form  $\mathcal{J}_0(k_0 r)$ . One can find the definition of Bessel-functions, for example, in [20, Section 4.7].

The goal of this paper is to construct and verify the asymptotic expansion for the eigenvalues of (2.2). The obtained asymptotics is used for studying the behavior of the best constant in a Friedrichs-type inequality for functions belonging to the Sobolev class  $H^1(\Omega_\varepsilon, \Gamma_\varepsilon)$

(see the definition of  $H^1(\Omega_\varepsilon, \Gamma_\varepsilon)$  in Section 7). One of the main results of this paper is the following asymptotics for  $\lambda_\varepsilon$  converging to  $\lambda_0$ :

$$\lambda_\varepsilon = \lambda_0 + \sum_{i=1}^{\infty} \varepsilon^i \lambda_i, \quad (2.4)$$

where  $\lambda_i$  are some fixed constants which can be calculated according to (4.23) and (4.15) in the case of simple  $\lambda_\varepsilon$  and according to (5.10) and (4.15) when  $\lambda_\varepsilon$  is of multiplicity two. In particular,  $\lambda_1 < 0$  which implies that  $\lambda_\varepsilon < \lambda_0$ .

### 3. Construction of the Leading Terms of the Asymptotic Expansion

Suppose that  $\lambda_0$  is the simple eigenvalue for (2.3) and the corresponding eigenfunction  $u_0$  is normalized in  $L_2(\Omega)$ . Our aim is to construct the leading terms of the asymptotic expansions for  $\lambda_\varepsilon$  converging to  $\lambda_0$  as well as  $u_\varepsilon$  converging to  $u_0$ . We use the method of boundary-layer functions (see [21]) for this purpose. We are looking for eigenvalues and eigenfunctions in the following form:

$$\begin{aligned} \lambda_\varepsilon &= \lambda_0 + \varepsilon \lambda_1 + \cdots, \\ u_\varepsilon(x) &= u_0(x) + \varepsilon u_1(x) + \varepsilon \alpha_0(\theta) v(\xi) + \cdots, \end{aligned} \quad (3.1)$$

where  $\xi = (\xi_1, \xi_2)$ ,  $\xi_1 = \theta/\varepsilon$ ,  $\xi_2 = (1-r)/\varepsilon$ , and

$$\begin{aligned} u_0(x) &= \alpha_0(\theta)(1-r) + O((1-r)^2) \quad \text{as } r \rightarrow 1, \quad \alpha_0(\theta) = -\frac{\partial u_0}{\partial r} \Big|_{r=1}, \\ u_1(x) &= u_1|_{r=1} + \alpha_1(\theta)(1-r) + O((1-r)^2) \quad \text{as } r \rightarrow 1, \quad \alpha_1(\theta) = -\frac{\partial u_1}{\partial r} \Big|_{r=1}. \end{aligned} \quad (3.2)$$

Substituting the first expansion from (3.1) and the sum  $u_0 + \varepsilon u_1$  from the second expansion in (2.2) and equating terms at the same power of  $\varepsilon$ , we get the equation for  $u_1$ :

$$-\Delta_x u_1 = \lambda_0 u_1 + \lambda_1 u_0 \quad \text{in } \Omega. \quad (3.3)$$

The existence of the solution for (3.3) is given in the following proposition.

**Proposition 3.1.** *For any  $\lambda_1$ , there exists the smooth solution of (3.3) satisfying the boundary condition*

$$u_1 = -\lambda_1 \alpha_0(\theta) \left( \int_0^{2\pi} \alpha_0^2 d\theta \right)^{-1} \quad \text{on } \partial\Omega. \quad (3.4)$$

*Proof.* The existence of the smooth solution follows from the classical results on regular solutions of elliptic equations (see e.g., [22]). In order to get  $u_1$  as the unique solution, one can add the condition of mutual orthogonality:

$$\int_{\Omega} u_0 u_1 dx = 0. \quad (3.5)$$

By multiplying (3.3) by  $u_0$ , integrating (3.3) over  $\Omega$ , and twice integrating by parts the obtained equation, we find that

$$-\int_{\Omega} u_1 \Delta u_0 dx - \int_{\partial\Omega} u_1 \frac{\partial u_0}{\partial r} d\theta + \int_{\partial\Omega} u_0 \frac{\partial u_1}{\partial r} d\theta = \lambda_1 \int_{\Omega} u_0^2 dx + \lambda_0 \int_{\Omega} u_1 u_0 dx. \quad (3.6)$$

Taking into account the fact that  $u_0$  is the normalized (in  $L_2(\Omega)$ ) solution of (2.3) and since  $u_1$  satisfies (3.5), we can deduce that

$$\lambda_1 = - \int_{\partial\Omega} \frac{\partial u_0}{\partial r} u_1 d\theta = - \int_{\partial\Omega} \alpha_0(\theta) u_1 d\theta. \quad (3.7)$$

Then (3.7) leads to (3.4) and the proof is complete.  $\square$

However, the approximation  $u_0 + \varepsilon u_1$  does not satisfy the condition on  $\Gamma_\varepsilon$ . This forces us to introduce an additional term  $\alpha_0 v$  in second expansion of (3.1) to satisfy the appropriate boundary condition. We assume that the function  $v$  has exponential decay as  $\xi_2 \rightarrow \infty$  and is  $\pi$ -periodical with respect to  $\xi_1$ . Under this assumption,  $\alpha_0 v$  “almost” does not destroy (2.2) in the sense that the norm of additional contribution is small. The rigorous explanation is given in Section 6. Proceeding, we have that

$$-\Delta_x(u_0 + \varepsilon u_1 + \varepsilon \alpha_0 v + \cdots) = (\lambda_0 + \varepsilon \lambda_1 + \cdots)(u_0 + \varepsilon u_1 + \varepsilon \alpha_0 v + \cdots). \quad (3.8)$$

Taking into account (2.3) and (3.3), we see that  $v$  has to satisfy the equation

$$-\Delta_x(\alpha_0 v) = \lambda_0 \alpha_0 v. \quad (3.9)$$

Rewrite  $\Delta_x$  in polar coordinates and pass to the  $\xi$ -variables in the argument of  $v$ :

$$\begin{aligned} \Delta_x(\alpha_0 v) &= \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial}{\partial r} (\alpha_0 v) \right) + \frac{1}{r^2} \frac{\partial^2 (\alpha_0 v)}{\partial \theta^2} \\ &= \alpha_0 \frac{\partial^2 v}{\partial r^2} + \frac{\alpha_0}{r} \frac{\partial v}{\partial r} + \frac{1}{r^2} \left( v \frac{\partial^2 \alpha_0}{\partial \theta^2} + 2 \frac{\partial \alpha_0}{\partial \theta} \frac{\partial v}{\partial \theta} + \alpha_0 \frac{\partial^2 v}{\partial \theta^2} \right) \\ &= \frac{\alpha_0}{\varepsilon^2} \frac{\partial^2 v}{\partial \xi_2^2} - \frac{\alpha_0}{(\varepsilon - \varepsilon^2 \xi_2)} \frac{\partial v}{\partial \xi_2} + \frac{1}{(1 - \varepsilon \xi_2)^2} \left[ v \frac{\partial^2 \alpha_0}{\partial \theta^2} + \frac{2}{\varepsilon} \frac{\partial \alpha_0}{\partial \theta} \frac{\partial v}{\partial \xi_1} + \frac{\alpha_0}{\varepsilon^2} \frac{\partial^2 v}{\partial \xi_1^2} \right]. \end{aligned} \quad (3.10)$$

Finally, replacing formulas  $1/(\varepsilon - \varepsilon^2 \xi_2)$  and  $1/(1 - \varepsilon \xi_2)^2$  with Taylor series with respect to  $\varepsilon$ , substituting the obtained formula for  $\Delta_x(\alpha_0 v)$  in (3.9), and equating terms at  $\varepsilon^{-2}$ , we deduce that

$$\Delta_{\xi} v = 0. \quad (3.11)$$

Now we derive the boundary conditions for function  $v$ . Substituting the second series from (3.1) in boundary conditions from (2.2) and using (3.2), we have

$$\begin{aligned} 0 &= u_{\varepsilon} = u_0 + \varepsilon u_1 + \varepsilon \alpha_0 v + \cdots = \varepsilon(\alpha_0 \xi_2 + u_1|_{r=1} + \alpha_0 v) + O(\varepsilon^2), \\ 0 &= \frac{\partial u_{\varepsilon}}{\partial r} = \frac{\partial u_0}{\partial r} + \varepsilon \frac{\partial u_1}{\partial r} + \varepsilon \alpha_0 \frac{\partial v}{\partial r} + \cdots = -\alpha_0 - \varepsilon \alpha_1 - \alpha_0 \frac{\partial v}{\partial \xi_2} + \cdots, \end{aligned} \quad (3.12)$$

which implies that

$$\begin{aligned} \alpha_0 \xi_2 + u_1|_{r=1} + \alpha_0 v &= 0, \\ -\alpha_0 - \alpha_0 \frac{\partial v}{\partial \xi_2} &= 0. \end{aligned} \quad (3.13)$$

Taking into account (3.4), we derive the boundary conditions for  $v$  on  $\partial B$  and on  $\Gamma$ :

$$\begin{aligned} v &= -\xi_2 + \lambda_1 \left( \int_0^{2\pi} \alpha_0^2 d\theta \right)^{-1} \quad \text{on } \partial B, \\ \frac{\partial v}{\partial \xi_2} &= -1 \quad \text{on } \Gamma. \end{aligned} \quad (3.14)$$

Summing up (3.11) and (3.14), we get the following boundary-value problem for  $v$ :

$$\begin{aligned} \Delta_{\xi} v &= 0 \quad \Pi \setminus \overline{B}, \\ v &= -\xi_2 + \lambda_1 \left( \int_0^{2\pi} \alpha_0^2 d\theta \right)^{-1} \quad \text{on } \partial B, \\ \frac{\partial v}{\partial \xi_2} &= -1 \quad \text{on } \Gamma. \end{aligned} \quad (3.15)$$

Define the function  $Y$  as the solution of the following boundary-value problem in the cell of periodicity:

$$\begin{aligned} \Delta Y &= 0 \quad \text{in } \Pi \setminus \overline{B}, \\ Y &= 0 \quad \text{on } \partial B, \\ \frac{\partial Y}{\partial \xi_1} &= 0 \quad \text{on } \partial \Pi \setminus \Gamma, \\ \frac{\partial Y}{\partial \xi_2} &= 0 \quad \text{on } \Gamma, \\ \frac{\partial Y}{\partial \xi_2} &= 1 \quad \text{as } \xi_2 \rightarrow \infty. \end{aligned} \tag{3.16}$$

It was proved in [7] that there exists the solution of (3.16), which is even with respect to  $\xi_1$  and has the asymptotics:

$$Y(\xi) = \xi_2 + C(B) + O(e^{-\alpha \xi_2}) \quad \text{as } \xi_2 \rightarrow \infty, \tag{3.17}$$

where

$$C(B) = \int_{\Pi \setminus \overline{B}} |\nabla(Y - \xi_2)|^2 d\xi + |B| > 0, \tag{3.18}$$

and  $|B|$  is the area of the domain  $B$ .

The following lemma gives the conditions to obtain  $v$  as an exponentially decaying function as  $\xi_2 \rightarrow \infty$ .

**Lemma 3.2.** *Assume that  $F$  is  $\pi$ -periodic with respect to  $\xi_1$  function with exponential decay as  $\xi_2 \rightarrow \infty$ , and let  $v$  be a  $\pi$ -periodic solution of the boundary-value problem:*

$$\Delta v = F, \quad \xi_2 > 0; \quad v = A_1, \quad \xi \in \partial B; \quad \frac{\partial v}{\partial \xi_2} = A_2, \quad \xi \in \Gamma; \tag{3.19}$$

with finite Dirichlet integral in  $\Pi$ . Then there exists the unique weak solution, which has asymptotics  $v = C + O(e^{-\alpha \xi_2})$ ,  $\alpha > 0$ . To obtain  $v$  as a function with exponential decay as  $\xi_2 \rightarrow \infty$ , it is necessary and sufficient to have

$$\int_{\Pi \setminus B} Y F d\xi + \int_{\partial B} A_1 \frac{\partial Y}{\partial \nu_B} dS_B + \int_{\Gamma} A_2 Y d\xi_1 = 0. \tag{3.20}$$

*Proof.* The existence of the solution with asymptotics  $v = C + O(e^{-\alpha \xi_2})$  follows from the classical results on elliptic boundary-value problems in cylindric domains (see, e.g., [23] and [24, Chapters 2, 5]). Let us verify (3.20). Define  $\Pi_R = \Pi \cap \{0 < \xi_2 < R\}$  and  $\Gamma_R = \{\xi :$

$-\pi/2 < \xi_1 < \pi/2, \xi_2 = R\}$ . By multiplying the equation from (3.15) by  $Y$ , integrating it over  $\Pi_R \setminus \bar{B}$ , and using the property of  $Y$ , we get that

$$\begin{aligned}
 \int_{\Pi_R \setminus \bar{B}} F Y \, d\xi &= - \int_{\Pi_R \setminus \bar{B}} \nabla v \nabla Y \, d\xi + \int_{\Gamma_R} \frac{\partial v}{\partial \xi_2} Y \, d\xi_1 - \int_{\Gamma} \frac{\partial v}{\partial \xi_2} Y \, d\xi_1 \\
 &= \int_{\Pi_R \setminus \bar{B}} v \Delta Y \, d\xi - \int_{\Gamma_R} v \frac{\partial Y}{\partial \xi_2} \, d\xi_1 + \int_{\Gamma} v \frac{\partial Y}{\partial \xi_2} \, d\xi_1 - \int_{\partial B} v \frac{\partial Y}{\partial \nu} \, dS_B \\
 &\quad + \int_{\Gamma_R} \frac{\partial v}{\partial \xi_2} Y \, d\xi_1 - \int_{\Gamma} \frac{\partial v}{\partial \xi_2} Y \, d\xi_1 = - \int_{\Gamma_R} v \frac{\partial Y}{\partial \xi_2} \, d\xi_1 - \int_{\partial B} A_1 \frac{\partial Y}{\partial \nu} \, dS_B \\
 &\quad + \int_{\Gamma_R} \frac{\partial v}{\partial \xi_2} Y \, d\xi_1 - \int_{\Gamma} A_2 Y \, d\xi_1.
 \end{aligned} \tag{3.21}$$

Passing to the limit as  $R \rightarrow \infty$ , we obtain that

$$\int_{\Pi \setminus \bar{B}} F Y \, d\xi = -\pi C - \int_{\partial B} A_1 \frac{\partial Y}{\partial \nu} \, dS_B - \int_{\Gamma} A_2 Y \, d\xi_1. \tag{3.22}$$

This can be rewritten as

$$C = \frac{1}{\pi} \left( - \int_{\Gamma} A_2 Y \, d\xi_1 - \int_{\partial B} A_1 \frac{\partial Y}{\partial \nu} \, dS_B - \int_{\Pi \setminus \bar{B}} F Y \, d\xi \right). \tag{3.23}$$

Then  $v$  has exponential decay as  $\xi_2 \rightarrow \infty$  if and only if  $C = 0$  which is equivalent to (3.20). The proof is complete.  $\square$

In order to obtain  $v$  as function with exponential decay as  $\xi_2 \rightarrow \infty$ , one must have

$$0 = - \int_{\partial B} (-\xi_2 + K) \frac{\partial Y}{\partial \nu_B} \, dS_B + \int_{\Gamma} Y \, d\xi_1, \tag{3.24}$$

where we denote  $K = \lambda_1 \left( \int_0^{2\pi} \alpha_0^2 \, d\theta \right)^{-1}$ . However, (3.24) implies that

$$K = \left( \int_{\partial B} \xi_2 \frac{\partial Y}{\partial \nu_B} \, dS_B + \int_{\Gamma} Y \, d\xi_1 \right) \left( \int_{\partial B} \frac{\partial Y}{\partial \nu_B} \, dS_B \right)^{-1}. \tag{3.25}$$

Integrate the identities  $0 = \int_{\Pi_R \setminus \bar{B}} \Delta Y d\xi$ ,  $0 = \int_{\Pi_R \setminus \bar{B}} \xi_2 \Delta Y d\xi$ :

$$\begin{aligned} 0 &= \int_{\Pi_R \setminus \bar{B}} \Delta Y d\xi = \int_{\partial(\Pi_R \setminus \bar{B})} \frac{\partial Y}{\partial n} dS = \int_{\partial B} \frac{\partial Y}{\partial \nu_B} dS_B + \int_{\Gamma_R} \frac{\partial Y}{\partial \xi_2} d\xi_1, \\ 0 &= \int_{\Pi_R \setminus \bar{B}} (\xi_2 \Delta Y - Y \Delta \xi_2) d\xi = \int_{\partial(\Pi_R \setminus \bar{B})} \left( \xi_2 \frac{\partial Y}{\partial n} - Y \frac{\partial \xi_2}{\partial n} \right) dS \\ &= \int_{\partial B} \xi_2 \frac{\partial Y}{\partial \nu_B} dS_B + \int_{\Gamma} Y d\xi_1 + \int_{\Gamma_R} \left( \xi_2 \frac{\partial Y}{\partial \xi_2} - Y \right) d\xi_1. \end{aligned} \quad (3.26)$$

Passing to the limit as  $R \rightarrow \infty$ , we find that

$$\begin{aligned} 0 &= \int_{\partial B} \frac{\partial Y}{\partial \nu_B} dS_B + \pi, \\ 0 &= \int_{\partial B} \xi_2 \frac{\partial Y}{\partial \nu_B} dS_B + \int_{\Gamma} Y d\xi_1 - \pi C(B). \end{aligned} \quad (3.27)$$

Then (3.25) and (3.27) together with Remark 2.2 imply that

$$\lambda_1 = -C(B) \int_0^{2\pi} \alpha_0^2 d\theta = -2\pi C(B) k_0^2 (\mathcal{J}'_0)^2(k_0) < 0. \quad (3.28)$$

#### 4. Complete Expansion in the Case of the Simple Eigenvalue $\lambda_0$

Assume that  $\lambda_0$  is the simple eigenvalue of the limit problem. Now we construct the complete expansion in the following form:

$$u_\varepsilon(x) = u_\varepsilon^{\text{ex}}(x) + \chi(1-r) u_\varepsilon^{\text{in}}\left(\frac{1-r}{\varepsilon}, \frac{\theta}{\varepsilon}\right), \quad (4.1)$$

where  $\chi$  is a smooth cutoff function, which equals to one when  $1/2 < r < 1$  and zero when  $r < 1/4$ :

$$u_\varepsilon^{\text{ex}}(x) = \mathcal{J}_0(k(\varepsilon)r), \quad (4.2)$$

$$u_\varepsilon^{\text{in}}(\xi) = \sum_{i=1}^{\infty} \varepsilon^i v_i(\xi). \quad (4.3)$$

Here  $k(\varepsilon) = \sqrt{\lambda_\varepsilon} v_i(\xi)$  are  $\pi$ -periodic in  $\xi_1$  functions with exponential decay as  $\xi_2 \rightarrow \infty$ . One can easily show that (4.2) solves the equation:

$$-\Delta_x u_\varepsilon^{\text{ex}}(x) = \lambda_\varepsilon u_\varepsilon^{\text{ex}}(x) \quad (4.4)$$

if and only if  $k(\varepsilon) = \sqrt{\lambda_\varepsilon}$ .

We are looking for  $u_\varepsilon^{\text{in}}(\xi)$ , which solves the equation:

$$-\Delta_x u_\varepsilon^{\text{in}}(\xi) = \lambda_\varepsilon u_\varepsilon^{\text{in}}(\xi). \quad (4.5)$$

If (4.4) and (4.5) are satisfied, then  $u_\varepsilon$  from (4.1) is the solution of

$$-\Delta_x u_\varepsilon = \lambda_\varepsilon u_\varepsilon + F, \quad (4.6)$$

where  $F = -u_\varepsilon^{\text{in}} \Delta_x \chi - 2\nabla_x u_\varepsilon^{\text{in}} \nabla_x \chi$ . Our aim is to construct  $u_\varepsilon^{\text{in}}$  so that  $F$  will be of small order as  $\varepsilon \rightarrow 0$ . This is the reason why we need to have  $v_i$  as exponentially decaying functions.

Now we derive the formula for the Laplacian in  $\xi$ -variables:

$$\begin{aligned} \Delta_x &= \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \theta^2} \\ &= \frac{1}{\varepsilon^2} \frac{\partial^2}{\partial \xi_2^2} + \frac{1}{\varepsilon(\varepsilon \xi_2 - 1)} \frac{\partial}{\partial \xi_2} + \frac{1}{\varepsilon^2(\varepsilon \xi_2 - 1)^2} \frac{\partial^2}{\partial \xi_1^2} \\ &= \frac{1}{\varepsilon^2} \Delta_\xi + \frac{1}{\varepsilon(\varepsilon \xi_2 - 1)} \frac{\partial}{\partial \xi_2} + \frac{1}{\varepsilon^2} \left( \frac{1}{(\varepsilon \xi_2 - 1)^2} - 1 \right) \frac{\partial^2}{\partial \xi_1^2}. \end{aligned} \quad (4.7)$$

By substituting the Taylor series for the functions

$$\frac{1}{\varepsilon(\varepsilon \xi_2 - 1)}, \quad \frac{1}{\varepsilon^2} \left( \frac{1}{(\varepsilon \xi_2 - 1)^2} - 1 \right) \quad (4.8)$$

in (4.7), we get the final formula for  $\Delta_x$ :

$$\Delta_x = \frac{1}{\varepsilon^2} \Delta_\xi + \sum_{n=0}^{\infty} (n+1) \varepsilon^{n-2} \xi_2^n \frac{\partial^2}{\partial \xi_1^2} - \sum_{n=0}^{\infty} \varepsilon^{n-1} \xi_2^n \frac{\partial}{\partial \xi_2}. \quad (4.9)$$

Substituting (2.4) and (4.3) in (4.5) and taking into account (4.9), we deduce the following formula:

$$\begin{aligned} \sum_{i=1}^{\infty} \varepsilon^i \Delta_\xi v_i &= \left( \varepsilon + \varepsilon^2 \xi_2 + \cdots + \varepsilon^{n+1} \xi_2^n + \cdots \right) \sum_{i=1}^{\infty} \varepsilon^i \frac{\partial v_i}{\partial \xi_2} \\ &\quad - \left( 2\varepsilon \xi_2 + 3\varepsilon^2 \xi_2^2 + \cdots + (n+1) \varepsilon^n \xi_2^n + \cdots \right) \sum_{i=1}^{\infty} \varepsilon^i \frac{\partial^2 v_i}{\partial \xi_1^2} \\ &\quad - \left( \varepsilon^2 \lambda_0 + \varepsilon^3 \lambda_1 + \cdots + \varepsilon^{n+2} \lambda_n \right) \sum_{i=1}^{\infty} \varepsilon^i v_i. \end{aligned} \quad (4.10)$$

By equating terms of the same power of  $\varepsilon$ , we obtain that

$$\begin{aligned}
 \varepsilon^1 : \quad & \Delta_{\xi} v_1 = 0, \\
 \varepsilon^2 : \quad & \Delta_{\xi} v_2 = \frac{\partial v_1}{\partial \xi_2} - 2\xi_2 \frac{\partial^2 v_1}{\partial \xi_1^2}, \\
 & \vdots, \\
 \varepsilon^k : \quad & \Delta_{\xi} v_k = \sum_{j=1}^{k-1} \left( \xi_2^{j-1} \frac{\partial v_{k-j}}{\partial \xi_2} - (j+1)\xi_2^j \frac{\partial^2 v_{k-j}}{\partial \xi_1^2} \right) - \sum_{j=0}^{k-3} \lambda_j v_{k-j-2}, \\
 & \vdots
 \end{aligned} \tag{4.11}$$

Consider now the boundary conditions from (2.2). According to the property of  $\chi$ ,

$$u_{\varepsilon}(x) = u_{\varepsilon}^{\text{ex}}(x) + u_{\varepsilon}^{\text{in}}\left(\frac{1-r}{\varepsilon}, \frac{\theta}{\varepsilon}\right) = \mathcal{J}_0(k(\varepsilon)r) + \sum_{i=1}^{\infty} \varepsilon^i v_i(\xi), \tag{4.12}$$

in a small neighborhood of  $\partial\Omega$ . Moreover, on  $\partial\Omega$ , it yields that

$$0 = \frac{\partial u_{\varepsilon}}{\partial r} = k(\varepsilon) \mathcal{J}'_0(k(\varepsilon)) - \sum_{i=1}^{\infty} \varepsilon^{i-1} \frac{\partial v_i}{\partial \xi_2} \Big|_{\xi_2=0}. \tag{4.13}$$

We assume that the function  $k(\varepsilon)$  has asymptotics:

$$k(\varepsilon) = k_0 + \varepsilon k_1 + \dots + \varepsilon^n k_n + \dots, \tag{4.14}$$

and since  $\lambda_{\varepsilon} = k^2(\varepsilon)$ , we can derive the following formulas for  $\lambda_i$ :

$$\lambda_0 = k_0^2, \quad \lambda_1 = 2k_0 k_1, \dots, \quad \lambda_i = \sum_{j=0}^i k_j k_{i-j}. \tag{4.15}$$

Rewriting  $\mathcal{J}'_0(k(\varepsilon))$  as a Taylor series with respect to  $\varepsilon$ , we have

$$\mathcal{J}'_0(k(\varepsilon)) = \mathcal{J}'_0(k_0) + \frac{\mathcal{J}''_0(k_0)k_1\varepsilon}{1!} + \frac{(\mathcal{J}'''_0(k_0)k_1^2 + \mathcal{J}''_0(k_0)k_2)\varepsilon^2}{2!} + \dots. \tag{4.16}$$

Substituting (4.16) in (4.13), using (4.14), and equating the terms with the same powers of  $\varepsilon$ , we get the following boundary condition for  $v_i, i = 1, 2, \dots$ :

$$\frac{\partial v_i}{\partial \xi_2} = g_i(k_1, \dots, k_{i-1}) \quad \text{on } \Gamma, \tag{4.17}$$

where

$$g_1 = k_0 \mathcal{J}'_0(k_0), \quad g_2 = k_1 \mathcal{J}'_0(k_0) + k_0 k_1 \mathcal{J}''_0(k_0) \equiv 0. \quad (4.18)$$

Consider now the boundary conditions on small holes. Analogously,

$$u_\varepsilon(x) = \mathcal{J}_0(k(\varepsilon)r) + \sum_{i=1}^{\infty} \varepsilon^i v_i(\xi) = \mathcal{J}_0(k(\varepsilon)(1 - \varepsilon \xi_2)) + \sum_{i=1}^{\infty} \varepsilon^i v_i(\xi). \quad (4.19)$$

Substituting the Taylor series for  $\mathcal{J}_0(k(\varepsilon)(1 - \varepsilon \xi_2))$  with respect to  $\varepsilon$  in the last formula, using (4.14), and equating the terms with the same powers of  $\varepsilon$  in equation  $u_\varepsilon = 0$  on  $\Gamma_\varepsilon$ , we get the following boundary condition for  $v_i$ ,  $i = 1, 2, \dots$ , on  $\partial B$ :

$$v_i = -k_i \mathcal{J}'_0(k_0) + f_i(\xi_2; k_0, k_1, \dots, k_{i-1}) \quad \text{on } \partial B, \quad (4.20)$$

where  $f_i$  are polynomials of power  $i$  with respect to  $\xi_2$  with coefficients which depend on  $(k_0, k_1, \dots, k_{i-1})$ . The precise formula for  $f_i$  can be derived for each fixed  $i$ . For example, we have that

$$f_1 = k_0 \mathcal{J}'_0(k_0) \xi_2, \quad f_2 = k_1 \mathcal{J}'_0(k_0) \xi_2 - \frac{1}{2} \mathcal{J}''_0(k_0) (k_1 - k_0 \xi_2)^2. \quad (4.21)$$

The following Lemma is useful for our analysis. For the proof see for example, [3].

**Lemma 4.1.** Suppose that  $F$  and  $v$  satisfy the conditions of Lemma 3.2. (a) If  $F$  is even with respect to  $\xi_1$ , then  $v$  is even; (b) if  $F$  is odd with respect to  $\xi_1$  and  $A_1 = A_2 = 0$ , then  $v$  is odd with respect to  $\xi_1$  and decays exponentially as  $\xi_2 \rightarrow \infty$ .

**Theorem 4.2.** There exist numbers  $k_i$  and  $\pi$ -periodic in  $\xi_1$  functions  $v_i$  with finite Dirichlet integral in  $\Pi$  and exponential decay as  $\xi_2 \rightarrow \infty$ , such that these functions are solutions of the following boundary-value problems:

$$\begin{aligned} \Delta v_i &= F_i \equiv \sum_{j=1}^{i-1} \left( \xi_2^{j-1} \frac{\partial v_{i-j}}{\partial \xi_2} - (j+1) \xi_2^j \frac{\partial^2 v_{i-j}}{\partial \xi_1^2} \right) - \sum_{j=0}^{i-3} \lambda_j v_{i-j-2} \quad \text{in } \Pi \setminus \bar{B}, \\ v_i &= -k_i \mathcal{J}'_0(k_0) + f_i(\xi_2; k_0, k_1, \dots, k_{i-1}) \quad \text{on } \partial B, \\ \frac{\partial v_i}{\partial \xi_2} &= g_i(k_1, \dots, k_{i-1}) \quad \text{on } \Gamma. \end{aligned} \quad (4.22)$$

Moreover, the constants are defined by the formula:

$$k_i = -\frac{1}{\pi \mathcal{J}'_0(k_0)} \left( \int_{\Pi \setminus B} Y F_i d\xi + \int_{\partial B} f_i(\xi_2; k_0, k_1, \dots, k_{i-1}) \frac{\partial Y}{\partial \nu_B} dS_B + g_i(k_1, \dots, k_{i-1}) \int_{\Gamma} Y d\xi_1 \right) \quad (4.23)$$

In particular,

$$k_1 = -\pi C(B)k_0(\mathcal{J}')_0^2(k_0), \quad (4.24)$$

$$k_2 = \frac{k_1^2}{2k_0}. \quad (4.25)$$

*Proof.* Let  $v$  be the solution of boundary-value problem (3.15). It can be easily verified that

$$v_1 = -k_0\mathcal{J}'_0(k_0)v \quad (4.26)$$

is a solution of (4.22), (4.20), (4.17) for  $f_1, g_1$ , and  $k_1$  defined by (4.21), (4.18), and (4.24). For any  $k_2$  boundary-value problem (4.22), (4.20), (4.17) for  $v_2$  has a  $\pi$ -periodic solution with finite Dirichlet integral. By Lemma 3.2 and (3.27),  $v_2$  has exponential decay as  $\xi_2 \rightarrow \infty$  if and only if  $k_2$  is given by (4.23) for  $i = 2$ . Let us verify formula (4.25) without applying the general (4.23). It is obvious that

$$\frac{\partial^2 v_1}{\partial \xi_1^2} = -\frac{\partial^2 v_1}{\partial \xi_2^2}. \quad (4.27)$$

By using that fact one can write the boundary-value problem for  $v_2$  as

$$\begin{aligned} \Delta v_2 &= \left( \frac{\partial v_1}{\partial \xi_2} - 2\xi_2 \frac{\partial^2 v_1}{\partial \xi_1^2} \right) = \left( \frac{\partial v_1}{\partial \xi_2} + 2\xi_2 \frac{\partial^2 v_1}{\partial \xi_2^2} \right) \quad \text{in } \Pi \setminus \bar{B}, \\ v_2 &= -k_2\mathcal{J}'_0(k_0) + k_1\mathcal{J}'_0(k_0)\xi_2 - \frac{1}{2}\mathcal{J}''_0(k_0)(k_1 - k_0\xi_2)^2 \quad \text{on } \partial B, \\ \frac{\partial v_2}{\partial \xi_2} &= 0 \quad \text{on } \Gamma. \end{aligned} \quad (4.28)$$

It can be verified that the function

$$v_2 = \frac{1}{2}\xi_2^2 \frac{\partial v_1}{\partial \xi_2} \quad (4.29)$$

is  $\pi$ -periodic with finite Dirichlet integral in  $\Pi$ , has exponential decay as  $\xi_2 \rightarrow \infty$ , and satisfies problem (4.28) for  $k_2$  defined by (4.25). We can use the induction process to finalize the proof.  $\square$

Since  $k_i$  are defined by (4.23), we can calculate  $\lambda_i$  by using (4.15). Denote

$$u_{\varepsilon,N} = \mathcal{J}_0\left(\sqrt{\lambda_{\varepsilon,N}}r\right) + \chi(1-r)v_{\varepsilon,N}, \quad (4.30)$$

where  $\lambda_{\varepsilon,N}$  and  $v_{\varepsilon,N}$  are the partial sums of (2.4) and (4.3), respectively.

Theorem 4.2 implies the validity of the following useful result.

**Theorem 4.3.** *For any integer  $N > 0$ , the function  $u_{\varepsilon,N}$  is the solution of the boundary-value problem*

$$\begin{aligned} -\Delta u_{\varepsilon,N} &= \lambda_{\varepsilon,N} u_{\varepsilon,N} + F_{\varepsilon,N} \quad \text{in } \Omega_{\varepsilon}, \\ u_{\varepsilon,N} &= a_{\varepsilon,N}(\theta) \quad \text{on } \Gamma_{\varepsilon}, \\ \frac{\partial u_{\varepsilon,N}}{\partial r} &= b_{\varepsilon,N}(\theta) \quad \text{on } \partial\Omega, \end{aligned} \quad (4.31)$$

where  $\|a_{\varepsilon,N}\|_{L_2(\Gamma_{\varepsilon})} = O(\varepsilon^{N_1})$ ,  $\|b_{\varepsilon,N}\|_{L_2(\partial\Omega)} = O(\varepsilon^{N_1})$ ,  $\|F_{\varepsilon,N}\|_{L_2(\Omega_{\varepsilon})} = O(\varepsilon^{N_1})$ , and  $N_1 \rightarrow \infty$  as  $N \rightarrow \infty$ .

*Proof.* According to the definition of  $u_{\varepsilon,N}$ , we have that

$$\begin{aligned} -\Delta_x u_{\varepsilon,N} &= -\Delta_x \left( \mathcal{J}_0 \left( \sqrt{\lambda_{\varepsilon,N}} r \right) + \chi(1-r)v_{\varepsilon,N} \right) \\ &= -\Delta_x \mathcal{J}_0 \left( \sqrt{\lambda_{\varepsilon,N}} r \right) - \Delta_x \chi v_{\varepsilon,N} - 2\nabla_x \chi \nabla_x v_{\varepsilon,N} - \chi \Delta_x v_{\varepsilon,N} \\ &= \lambda_{\varepsilon,N} \mathcal{J}_0 \left( \sqrt{\lambda_{\varepsilon,N}} r \right) + \lambda_{\varepsilon,N} \chi(1-r)v_{\varepsilon,N} - \lambda_{\varepsilon,N} \chi(1-r)v_{\varepsilon,N} - \Delta_x \chi v_{\varepsilon,N} - 2\nabla_x \chi \nabla_x v_{\varepsilon,N} \\ &\quad - \chi \Delta_x v_{\varepsilon,N} \\ &= \lambda_{\varepsilon,N} u_{\varepsilon,N} + F_{\varepsilon,N}, \end{aligned} \quad (4.32)$$

where

$$F_{\varepsilon,N} = -v_{\varepsilon,N} \Delta_x \chi - \chi(\lambda_{\varepsilon,N} v_{\varepsilon,N} + \Delta_x v_{\varepsilon,N}) - 2\nabla_x \chi \nabla_x v_{\varepsilon,N} =: I_1 + I_2 + I_3. \quad (4.33)$$

Passing from  $(x_1, x_2)$  variables to polar coordinates  $(r, \theta)$ , we get that

$$\frac{\partial}{\partial x_1} = \cos \theta \frac{\partial}{\partial r} - \frac{\sin \theta}{r} \frac{\partial}{\partial \theta}, \quad \frac{\partial}{\partial x_2} = \sin \theta \frac{\partial}{\partial r} + \frac{\cos \theta}{r} \frac{\partial}{\partial \theta}, \quad (4.34)$$

$$\Delta_x = \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2}{\partial \theta^2}. \quad (4.35)$$

By using the fact that  $\lim_{x \rightarrow \infty} x e^{-\alpha x} = 0$  and due to the result of Theorem 4.2, we have that, for any  $1 \leq i \leq N$ ,

$$\varepsilon^i v_i = \varepsilon^i O\left(e^{-\alpha \xi_2}\right) = \varepsilon^N O\left(\varepsilon^{i-N} e^{-\alpha(1-r)/\varepsilon}\right) = \varepsilon^N O(\varepsilon^m) = O\left(\varepsilon^{N+m}\right), \quad (4.36)$$

where  $m$  is fixed. Hence,  $v_{\varepsilon,N} = O(\varepsilon^{N+m})$ . Similarly, taking into account (4.34) and (4.35), we can deduce that

$$\begin{aligned}\nabla_x v_{\varepsilon,N} &= O(\varepsilon^{N+m}) \left( \frac{\alpha \cos \theta}{\varepsilon} - \frac{\sin \theta}{r}, \frac{\alpha \sin \theta}{\varepsilon} + \frac{\cos \theta}{r} \right), \\ \nabla_x \chi &= (-\cos \theta \chi', -\sin \theta \chi').\end{aligned}\quad (4.37)$$

Consequently,

$$\nabla_x v_{\varepsilon,N} \nabla_x \chi = O(\varepsilon^{N+m}) O\left(\frac{1}{\varepsilon r}\right). \quad (4.38)$$

Furthermore,

$$\begin{aligned}\Delta_x v_{\varepsilon,N} &= -\frac{\alpha}{\varepsilon r} O(\varepsilon^{N+m}) + \frac{\alpha^2}{\varepsilon^2} O(\varepsilon^{N+m}) + \frac{1}{r^2} O(\varepsilon^{N+m}) = O\left(\frac{1}{\varepsilon^2 r^2}\right) O(\varepsilon^{N+m}), \\ \Delta_x \chi &= -\frac{1}{r} \chi' + \chi'' = O\left(\frac{1}{r}\right).\end{aligned}\quad (4.39)$$

According to the definition of  $\chi$ , the support of  $\nabla_x \chi$  and  $\Delta_x \chi$  is the set  $\{1/4 \leq r \leq 1/2\}$ . Summarizing, we have that

$$I_1 = O(\varepsilon^{N+m}) O\left(\frac{1}{r}\right), \quad I_2 = O(\varepsilon^{N+m}) O\left(\frac{1}{\varepsilon^2 r^2}\right), \quad I_3 = O(\varepsilon^{N+m}) O\left(\frac{1}{\varepsilon r}\right), \quad (4.40)$$

and we can derive that

$$\begin{aligned}\|F_{\varepsilon,N}\|_{L_2(\Omega_\varepsilon)}^2 &= \int_{\Omega} F_{\varepsilon,N}^2 r \, dr \, d\theta \\ &= \int_{\Omega \cap \{1/4 \leq r \leq 1\}} I_2^2 r \, dr \, d\theta + \int_{\Omega \cap \{1/4 \leq r \leq 1/2\}} \left[ (I_1 + I_3)^2 r + 2I_2(I_1 + I_3)r \right] dr \, d\theta \\ &= O(\varepsilon^{2N+2m}) O\left(\frac{1}{\varepsilon^3}\right) + O(\varepsilon^{2N+2m}) O\left(\frac{1}{\varepsilon^4}\right) = O(\varepsilon^{2N+2m}) O\left(\frac{1}{\varepsilon^4}\right).\end{aligned}\quad (4.41)$$

Therefore,

$$\|F_{\varepsilon,N}\|_{L_2(\Omega_\varepsilon)} = O(\varepsilon^{N+m-2}) = O(\varepsilon^{N_1}), \quad N_1 \longrightarrow \infty \text{ as } N \longrightarrow \infty. \quad (4.42)$$

Consider now  $u_{\varepsilon,N}$  on  $\Gamma_\varepsilon$ :

$$u_{\varepsilon,N} = \mathcal{J}_0\left(\sqrt{\lambda_{\varepsilon,N}} r\right) + v_{\varepsilon,N} = \varepsilon^{N+1} \beta_{N+1} + \varepsilon^{N+2} \beta_{N+2} + \cdots, \quad (4.43)$$

where  $\beta_j$  are the coefficients of the Taylor series of the function  $\mathcal{J}_0(\sqrt{\lambda_{\varepsilon,N}}r)$ . Hence,  $a_{\varepsilon,N} = O(\varepsilon^{N+1})$  and

$$\|a_{\varepsilon,N}\|_{L_2(\Gamma_\varepsilon)}^2 = \frac{2}{\varepsilon} \int_{\partial B_\varepsilon} a_{\varepsilon,N}^2 d\theta \sim \frac{2}{\varepsilon} 2\pi a \varepsilon O(\varepsilon^{2N+2}) = O(\varepsilon^{2N+2}), \quad (4.44)$$

which yields that  $\|a_{\varepsilon,N}\|_{L_2(\Gamma_\varepsilon)} = O(\varepsilon^{N+1}) = O(\varepsilon^{N_1})$ ,  $N_1 \rightarrow \infty$  as  $N \rightarrow \infty$ . Analogously, one can verify that  $\|b_{\varepsilon,N}\|_{L_2(\partial\Omega)} = O(\varepsilon^{N_1})$ ,  $N_1 \rightarrow \infty$  as  $N \rightarrow \infty$ . The proof is complete.  $\square$

## 5. Complete Expansion in the Case of Multiple Eigenvalue $\lambda_0$

In this section we consider the case when  $\lambda_0$  is of multiplicity two. The asymptotics of the eigenvalue were constructed in the form (2.4) and

$$u_\varepsilon(x) = u_\varepsilon^{\text{ex}}(x) + \chi(1-r)u_\varepsilon^{\text{in}}\left(\frac{1-r}{\varepsilon}, \frac{\theta}{\varepsilon}, \theta\right), \quad (5.1)$$

$$u_\varepsilon^{\text{ex}}(x) = \cos(n\theta)\mathcal{J}_n(k(\varepsilon)r), \quad (5.2)$$

$$u_\varepsilon^{\text{in}}(x) = \cos(n\theta) \sum_{i=1}^{\infty} \varepsilon^i v_i^{\text{even}}(\xi) + \sin(n\theta) \sum_{i=2}^{\infty} \varepsilon^i v_i^{\text{odd}}(\xi). \quad (5.3)$$

In this case,

$$v_i^{\text{even}} = -k_i \mathcal{J}'_n(k_0) + f_i^{(n)}(\xi_2; k_0, k_1, \dots, k_{i-1}) \quad \text{on } \partial B, \quad (5.4)$$

where  $f_i^{(n)}$  are polynomials of power  $i$  with respect to  $\xi_2$  with coefficients which depend on  $(k_0, k_1, \dots, k_{i-1})$ . Moreover,

$$\frac{\partial v_i^{\text{even}}}{\partial \xi_2} = g_i^{(n)}(k_1, \dots, k_{i-1}) \quad \text{on } \Gamma, \quad (5.5)$$

where

$$f_1^{(n)} = k_0 \mathcal{J}'_n(k_0) \xi_2, \quad f_2^{(n)} = k_1 \mathcal{J}'_0(k_0) \xi_2 - \frac{1}{2} \mathcal{J}''_0(k_0) (k_1 - k_0 \xi_2)^2, \quad (5.6)$$

$$g_1^{(n)} = k_0 \mathcal{J}'_n(k_0), \quad g_2^{(n)} = k_1 \mathcal{J}'_n(k_0) + k_0 k_1 \mathcal{J}''_n(k_0) \equiv 0,$$

$$v_i^{\text{odd}} = 0, \quad \xi \in \partial B, \quad \frac{\partial v_i^{\text{odd}}}{\partial \xi_2} = 0, \quad \xi \in \Gamma. \quad (5.7)$$

Substituting (5.3) and (2.4) in (4.5), passing to the variables  $\xi$  and  $(\theta, \rho)$ , and collecting all the terms with equal order of  $\varepsilon$ , we get two systems of equations for  $v_i^{\text{even}}$  and  $v_i^{\text{odd}}$ :

$$\begin{aligned} \Delta v_i^{\text{even}} = & \sum_{j=1}^{i-1} \left( \xi_2^{j-1} \frac{\partial v_{i-j}^{\text{even}}}{\partial \xi_2} - (j+1) \xi_2^j \frac{\partial^2 v_{i-j}^{\text{even}}}{\partial \xi_1^2} \right) - n \sum_{j=0}^{i-3} (j+1) \xi_2^j \frac{\partial v_{i-j-1}^{\text{odd}}}{\partial \xi_1} \\ & - n^2 \sum_{j=0}^{i-3} (j+1) \xi_2^j v_{i-j-2}^{\text{even}} - \sum_{j=0}^{i-3} \lambda_j v_{i-j-2}^{\text{even}} \quad \text{in } \Pi \setminus \bar{B}, \end{aligned} \quad (5.8)$$

$$\begin{aligned} \Delta v_i^{\text{odd}} = & \sum_{j=1}^{i-2} \left( \xi_2^{j-1} \frac{\partial v_{i-j}^{\text{even}}}{\partial \xi_2} - (j+1) \xi_2^j \frac{\partial^2 v_{i-j}^{\text{odd}}}{\partial \xi_1^2} \right) + n \sum_{j=0}^{i-2} (j+1) \xi_2^j \frac{\partial v_{i-j-1}^{\text{even}}}{\partial \xi_1} \\ & + n^2 \sum_{j=0}^{i-3} (j+1) \xi_2^j v_{i-j-2}^{\text{odd}} - \sum_{j=0}^{i-3} \lambda_j v_{i-j-2}^{\text{odd}} \quad \text{in } \Pi \setminus \bar{B}. \end{aligned} \quad (5.9)$$

**Theorem 5.1.** *There exist numbers  $k_i$  and  $\pi$ -periodic in  $\xi_1$  even functions  $v_i^{\text{even}}$  and odd functions  $v_i^{\text{odd}}$  with finite Dirichlet integral in  $\Pi$ , which have exponential decay as  $\xi_2 \rightarrow \infty$ , such that these functions are solutions of the boundary-value problems (5.8), (5.4), (5.5), and (5.9), (5.7), respectively. Moreover, the constants  $k_i$  are defined by the formula:*

$$k_i = -\frac{1}{\pi \mathcal{J}'_n(k_0)} \left( \int_{\Pi \setminus B} Y F_i d\xi + \int_{\partial B} f_i^{(n)}(\xi_2; k_0, k_1, \dots, k_{i-1}) \frac{\partial Y}{\partial \nu_B} dS_B + g_i^{(n)}(k_1, \dots, k_{i-1}) \int_{\Gamma} Y d\xi_1 \right), \quad (5.10)$$

*Proof.* The problems (5.8), (5.5), (5.4) for functions  $v_1^{\text{even}}, v_2^{\text{even}}$  coincide with problems (4.22), (4.17), and (4.20) (if one change  $\mathcal{J}'_0(k_0)$  by  $\mathcal{J}'_n(k_0)$  and  $f_i, g_i$  by the respective  $f_i^{(n)}, g_i^{(n)}$ ). Therefore the construction of  $v_1^{\text{even}}, v_2^{\text{even}}$  and  $k_1, k_2$  is just the same as the construction from the proof of Theorem 4.2. Due to (5.9), (5.7), the problem for  $v_2^{\text{odd}}$  is as follows:

$$\begin{aligned} \Delta v_2^{\text{odd}} &= n \xi_2 \frac{\partial v_1^{\text{even}}}{\partial \xi_1} \quad \text{in } \Pi \setminus \bar{B}, \\ v_2^{\text{odd}} &= 0 \quad \text{on } \partial B, \\ \frac{\partial v_2^{\text{odd}}}{\partial \xi_2} &= 0 \quad \text{on } \Gamma. \end{aligned} \quad (5.11)$$

The function  $v_1^{\text{even}}$  is even (due to (4.26)) and, hence, the right-hand side is odd in (5.11) and is even in (5.8). By Lemma 3.2 and Theorem 4.2, we conclude that there exists the even solution  $v_2^{\text{odd}}$  of (5.11) with exponential decay. Then we can use the iteration process to complete the proof.  $\square$

Denote

$$u_{\varepsilon, N} = \cos(n\theta) \mathcal{J}_0 \left( \sqrt{\lambda_{\varepsilon, N}} r \right) + \chi(1-r) v_{\varepsilon, N}, \quad (5.12)$$

where  $\lambda_{\varepsilon, N}$  and  $v_{\varepsilon, N}$  are the partial sums of (2.4) and (5.3), respectively.

Theorem 5.1 implies the validity of the following result.

**Theorem 5.2.** *For any integer  $N > 0$ , the function  $u_{\varepsilon,N}$  is the solution of the boundary-value problem:*

$$\begin{aligned} -\Delta u_{\varepsilon,N} &= \lambda_{\varepsilon,N} u_{\varepsilon,N} + F_{\varepsilon,N} \quad \text{in } \Omega_{\varepsilon}, \\ u_{\varepsilon,N} &= a_{\varepsilon,N}(\theta) \cos(n\theta) \quad \text{on } \Gamma_{\varepsilon}, \\ \frac{\partial u_{\varepsilon,N}}{\partial r} &= b_{\varepsilon,N}(\theta) \cos(n\theta) \quad \text{on } \partial\Omega, \end{aligned} \quad (5.13)$$

where  $\|a_{\varepsilon,N}\|_{L_2(\Gamma_{\varepsilon})} = O(\varepsilon^{N_1})$ ,  $\|b_{\varepsilon,N}\|_{L_2(\partial\Omega)} = O(\varepsilon^{N_1})$ ,  $\|F_{\varepsilon,N}\|_{L_2(\Omega_{\varepsilon})} = O(\varepsilon^{N_1})$ , and  $N_1 \rightarrow \infty$  as  $N \rightarrow \infty$ .

*Proof.* The proof is analogous to the proof of Theorem 4.3. Hence, we omit the details.  $\square$

## 6. Verification of the Asymptotics

Consider the boundary-value problem:

$$\begin{aligned} -\Delta U_{\varepsilon} &= \lambda U_{\varepsilon} + F \quad \text{in } \Omega_{\varepsilon}, \\ U_{\varepsilon} &= 0 \quad \text{on } \Gamma_{\varepsilon}, \\ \frac{\partial U_{\varepsilon}}{\partial r} &= 0 \quad \text{on } \partial\Omega, \end{aligned} \quad (6.1)$$

where  $F \in L_2(\Omega)$  and  $\lambda \neq \lambda_0$  is some fixed number.

Similarly to the techniques used in [3, 18], one can show that the boundary-value problem (6.1) has the solution  $U_{\varepsilon} \in H^1(\Omega)$  and the following representation holds:

$$U_{\varepsilon} = \frac{u_{\varepsilon}}{\lambda_{\varepsilon} - \lambda} \int_{\Omega} u_{\varepsilon} F \, dx + \tilde{U}_{\varepsilon}, \quad (6.2)$$

for  $\lambda$  close to the simple eigenvalue  $\lambda_0$  of the problem (2.3) and

$$U_{\varepsilon} = \frac{1}{\lambda_{\varepsilon} - \lambda} \sum_{i=1}^2 u_{\varepsilon}^i \int_{\Omega} u_{\varepsilon}^i F \, dx + \tilde{U}_{\varepsilon}, \quad (6.3)$$

for  $\lambda$  close to multiple eigenvalue  $\lambda_0$  of the problem (2.3). Here  $u_{\varepsilon}$  is normalized in  $L_2(\Omega)$  eigenfunctions to (2.2) and  $u_{\varepsilon}^i$  is orthonormalized in  $L_2(\Omega)$  eigenfunctions to (2.2). Moreover,

$$\|\tilde{U}_{\varepsilon}\|_{H^1} \leq C \|F\|_{L_2}, \quad (6.4)$$

where the constant  $C$  is independent on  $\varepsilon$  and  $\lambda$ . It follows from (6.2) and (6.4) that

$$\|U_\varepsilon\|_{H^1} \leq \frac{C}{\lambda_\varepsilon - \lambda} \|F\|_{L_2}. \quad (6.5)$$

Consider now the case of simple  $\lambda_0$ . Define the function:

$$U_\varepsilon^N(x) = \left(1 + \frac{1}{\varepsilon}\right) u_{\varepsilon,N}(x) - \left(1 + \frac{1}{\varepsilon}\right) a_{\varepsilon,N} + b_{\varepsilon,N} \chi(1-r)(v(\xi) + \xi_2 + C(B)), \quad (6.6)$$

where  $u_{\varepsilon,N}$  and  $v$  are the solutions of (4.31) and (3.15), respectively, and  $C(B)$  is given by (3.18). Then, by Theorem 4.3,  $U_\varepsilon^N$  is the solution of (6.1) if

$$\lambda = \lambda_{\varepsilon,N}, \quad \|F\|_{L_2} = O(\varepsilon^{N_2}), \quad N_2 \rightarrow \infty \quad \text{as } N \rightarrow \infty. \quad (6.7)$$

Taking into account (6.5), (6.7), and the fact that  $\|U_\varepsilon\|_{H^1} < \infty$ , we can conclude that for each fixed  $N$ ,

$$\lambda_\varepsilon - \lambda_{\varepsilon,N} = O(\varepsilon^{N_2}) = o(\varepsilon^N) \quad \text{as } \varepsilon \rightarrow 0. \quad (6.8)$$

Therefore the asymptotics constructed in Section 4 coincide with the expansion of  $\lambda_\varepsilon$ . For the case of multiple  $\lambda_0$ , one can use the same technique. The difference is follows: one should use (6.3) instead of (6.2) and Theorem 5.2 instead of Theorem 4.3. The asymptotics of  $\lambda_\varepsilon$  are completely verified.

## 7. Application to a Friedrichs-Type Inequality

Consider the sets  $\Omega_\varepsilon, \Gamma_\varepsilon$ , which were defined in Section 2.

**Definition 7.1.** The Sobolev class  $H^1(\Omega_\varepsilon, \Gamma_\varepsilon)$  is the class of functions from  $H^1(\Omega_\varepsilon)$  having zero trace on  $\Gamma_\varepsilon$ .

**Theorem 7.2.** Let  $u \in H^1(\Omega_\varepsilon, \Gamma_\varepsilon)$ . Then a Friedrichs-type inequality

$$\int_{\Omega_\varepsilon} u^2(x) dx \leq K_\varepsilon \int_{\Omega_\varepsilon} |\nabla u(x)|^2 dx \quad (7.1)$$

holds, where the best constant  $K_\varepsilon$  has the asymptotics

$$K_\varepsilon = \frac{1}{k_0^2} + \frac{4\pi C(B)(\mathcal{J}_0')^2(k_0)}{k_0^2} \varepsilon + o(\varepsilon), \quad (7.2)$$

as  $\varepsilon \rightarrow 0$ . Here  $k_0$  is the smallest root of the Bessel function  $\mathcal{J}_0$  and the constant  $C(B)$  is given by (3.18).

*Proof.* The geometric approach developed in [5, 9] allows us to state that there is a constant  $K > 0$  such that

$$\int_{\Omega_\varepsilon} u^2(x) dx \leq K \int_{\Omega_\varepsilon} |\nabla u(x)|^2 dx. \quad (7.3)$$

The idea and method of proof are exactly similar to the ones which were used in the mentioned papers. We are interested in the behavior of the best possible constant as  $\varepsilon \rightarrow 0$ . Clearly, the best constant  $K_\varepsilon = 1/\lambda_\varepsilon^1$ , where  $\lambda_\varepsilon^1$  is the smallest eigenvalue of the boundary-value problem (2.2) (due to the variational formulation of the smallest eigenvalue). Therefore, we can apply (2.4) and (3.28) to derive the asymptotic expansion for  $K_\varepsilon$ :

$$K_\varepsilon = (\lambda_\varepsilon^1)^{-1} = (\lambda_0^1 + \varepsilon \lambda_1^1 + o(\varepsilon))^{-1} = \frac{1}{\lambda_0^1} - \frac{2\lambda_1^1}{(\lambda_0^1)^2} \varepsilon + o(\varepsilon). \quad (7.4)$$

Since we are interested in the smallest eigenvalue  $\lambda_0^1$ , we have to choose the smallest positive root of  $\mathcal{J}_0(k_0) = 0$  as  $k_0$ , precisely,  $k_0 = 2,405$ . Then, we get, after some simple calculations and using (4.15) and (3.28),

$$K_\varepsilon = \frac{1}{k_0^2} + \frac{4\pi C(B)(\mathcal{J}'_0)^2(k_0)}{k_0^2} \varepsilon + o(\varepsilon). \quad (7.5)$$

The proof is complete. □

## Acknowledgments

This paper was completed during the stay of the second author as PostDoc at Luleå University of Technology in 2010-2011. The second author wants to express many thanks to Luleå University of Technology (Sweden) for financial support and wonderful conditions to work. The work of the first and the second authors was supported in part by RFBR (Project no. 09-01-00353) and the work of the forth author was supported by a grant from the Swedish Research Council (Project no. 621-2008-5186). The authors also thank referee for several suggestions, which have improved the final version of this paper.

## References

- [1] G. A. Chechkin, R. R. Gadył'shin, and Y. O. Koroleva, "On the eigenvalue of the Laplacian in a domain perforated along the boundary," *Doklady Mathematics*, vol. 81, no. 3, pp. 337–341, 2010, translation from *Doklady Akademii Nauk. MAIK "Nauka"*, vol. 432, no. 1, pp. 7–11, 2010.
- [2] G. A. Chechkin, "Spectrum of homogenized problem in a circle domain with many concentrated masses," in *Multi Scale Problems and Asymptotic Analysis*, vol. 24 of *GAKUTO International Series. Mathematical Sciences and Applications*, pp. 49–62, 2006.
- [3] R. R. Gadył'shin, "On the eigenvalue asymptotics for periodically clamped membrains," *St. Petersburg Mathematical Journal*, vol. 10, no. 1, pp. 1–14, 1999.
- [4] G. A. Chechkin, Y. O. Koroleva, and L.-E. Persson, "On the precise asymptotics of the constant in Friedrich's inequality for functions vanishing on the part of the boundary with microinhomogeneous structure," *Journal of Inequalities and Applications*, Article ID 34138, 13 pages, 2007.

- [5] G. A. Chechkin, Y. O. Koroleva, A. Meidell, and L.-E. Persson, "On the Friedrichs inequality in a domain perforated aperiodically along the boundary. Homogenization procedure. Asymptotics for parabolic problems," *Russian Journal of Mathematical Physics*, vol. 16, no. 1, pp. 1–16, 2009.
- [6] G. A. Chechkin, Y. O. Koroleva, L.-E. Persson, and P. Wall, "A new weighted Friedrichs-type inequality for a perforated domain with a sharp constant," *Eurasian Mathematical Journal*, vol. 2, no. 1, pp. 81–103, 2011.
- [7] G. A. Chechkin, R. R. Gadyl'shin, and Y. O. Koroleva, "On asymptotics of the first eigenvalue for Laplacian in domain perforated along the boundary," *Differential Equations*, vol. 47, no. 6, pp. 822–831, 2011.
- [8] Y. O. Koroleva, "On the Friedrichs inequality in a cube perforated periodically along the part of the boundary. Homogenization procedure," Research Report 2, Department of Mathematics, Luleå University of Technology, Luleå, Sweden, 2009.
- [9] Yu. O. Koroleva, "The Friedrichs inequality in a three-dimensional domain that is aperiodically perforated along a part of the boundary," *Russian Mathematical Surveys*, vol. 65, no. 4, pp. 788–790, 2010, translation from, *Uspekhi Matematicheskikh Nauk*, vol. 65, no. 4, pp. 199–200, 2010.
- [10] Y. O. Koroleva, "On the weighted Hardy type inequality in a fixed domain for functions vanishing on the part of the boundary," *Mathematical Inequalities and Applications*, vol. 14, no. 3, pp. 543–553, 2011.
- [11] Y. O. Koroleva, L.-E. Persson, and P. Wall, "On Friedrichs-type inequalities in domains rarely perforated along the boundary," Research Report 2, Department of Engineering Sciences and Mathematics, Luleå University of Technology, Luleå, Sweden, 2011.
- [12] V. Kokilashvili, A. Meskhi, and L.-E. Persson, *Weighted Norm Inequalities for Integral Transforms with Product Kernels*, Mathematics Research Developments Series, Nova Science Publishers, New York, NY, USA, 2010.
- [13] A. Kufner, *Weighted Sobolev Spaces*, John Wiley & Sons, New York, NY, USA, 1985.
- [14] A. Kufner, L. Maligranda, and L.-E. Persson, "The prehistory of the Hardy inequality," *American Mathematical Monthly*, vol. 113, no. 8, pp. 715–732, 2006.
- [15] A. Kufner, L. Maligranda, and L.-E. Persson, *The Hardy Inequality. About Its History and some Related Results*, Vydavetelsky Servis Publishing House, Pilsen, Czech Republic, 2007.
- [16] A. Kufner and L.-E. Persson, *Weighted Inequalities of Hardy Type*, World Scientific Publishing, River Edge, NJ, USA, 2003.
- [17] G. A. Chechkin, "On the estimation of solutions of boundary value problems in domains with concentrated masses periodically distributed along the boundary. The case of "light" masses," *Mathematical Notes*, vol. 76, no. 5-6, pp. 865–879, 2004.
- [18] G. A. Chechkin and R. R. Gadyl'shin, "On the convergence of solutions and eigenelements of a boundary value problem in a domain perforated along the boundary," *Differential Equations*, vol. 46, no. 5, pp. 667–680, 2010, translation from, *Differentsial'nye Uravneniya*, vol. 46, no. 5, pp. 665–677, 2010.
- [19] G. A. Chechkin, "Averaging of boundary value problems with singular perturbation of the boundary conditions," *Academy of Sciences Sbornik Mathematics*, vol. 79, no. 1, pp. 191–222, 1994.
- [20] N. H. Asmar, *Partial Differential Equations and Boundary Value Problems*, Prentice Hall, Upper Saddle River, NJ, USA, 2000.
- [21] M. I. Vishik and M. A. Lusternik, "Regular degeneration and the boundary layer for linear differential equations with small parameter," *Transactions of the American Mathematical Society*, vol. 35, no. 2, pp. 239–364, 1962.
- [22] S. Agmon, A. Douglis, and L. Nirenberg, "Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. II," *Communications on Pure and Applied Mathematics*, vol. 17, pp. 35–92, 1964.
- [23] S. A. Nazarov, "Junctions of singularly degenerating domains with different limit dimensions. I," *Journal of Mathematical Sciences*, vol. 80, no. 5, pp. 1989–2034, 1996, translation from, *Trudy Seminara Imeni I. G. Petrovskogo*, vol. 314, no. 18, pp. 3–78, 1995.
- [24] S. A. Nazarov and B. A. Plamenevsky, *Elliptic Problems in Domains with Piecewise Smooth Boundaries*, vol. 13 of *de Gruyter Expositions in Mathematics*, Walter de Gruyter, Berlin, Germany, 1994.