

Pointwise multipliers for functions of bounded mean oscillation

By Eiichi NAKAI and Kôzô YABUTA*

(Received Jan. 23, 1984)

(Revised April 18, 1984)

1. Introduction.

The purpose of this paper is to characterize the set of pointwise multipliers on $bmo_\phi(\mathbf{R}^n)$, which is the function space defined using the mean oscillation and a growth function ϕ .

Janson [2] has characterized pointwise multipliers on $bmo_\phi(\mathbf{T}^n)$ on the n -dimensional torus \mathbf{T}^n . We extend his result to the case of the n -dimensional Euclidean space \mathbf{R}^n .

To define $bmo_\phi(\mathbf{R}^n)$, let $I(a, r)$ be the cube $\{x \in \mathbf{R}^n; |x_i - a_i| \leq r/2, i=1, 2, \dots, n\}$ whose edges have length r and are parallel to the coordinate axes. For a cube I , we denote by $|I|$ the Lebesgue measure of I , by $M(f, I)$ or f_I the mean value of a function f on I , i.e. $|I|^{-1} \int_I f(x) dx$, and by $MO(f, I)$ the mean oscillation of f on I , i.e. $|I|^{-1} \int_I |f(x) - f_I| dx$.

We now define

$$bmo_\phi(\mathbf{R}^n) = \left\{ f \in L^1_{\text{loc}}(\mathbf{R}^n) ; \sup_{I(a, r)} \frac{MO(f, I(a, r))}{\phi(r)} < +\infty \right\},$$

where ϕ is assumed to be a positive non-decreasing function on $\mathbf{R}_+ = (0, \infty)$. Such a function is called a growth function. If two growth functions ϕ_1 and ϕ_2 are equivalent ($\phi_1 \sim \phi_2$) i.e. there is a constant $C > 0$ such that $C^{-1}\phi_1(r) \leq \phi_2(r) \leq C\phi_1(r)$, then $bmo_{\phi_1}(\mathbf{R}^n) = bmo_{\phi_2}(\mathbf{R}^n)$.

A function g on \mathbf{R}^n is called a pointwise multiplier on $bmo_\phi(\mathbf{R}^n)$, if the pointwise multiplication fg belongs to $bmo_\phi(\mathbf{R}^n)$ for all f belonging to $bmo_\phi(\mathbf{R}^n)$.

Janson's characterization is the following. If ϕ is a growth function and $\phi(r)/r$ is almost decreasing, then a function g is a pointwise multiplier on $bmo_\phi(\mathbf{T}^n)$ if and only if g belongs to $bmo_\phi(\mathbf{T}^n) \cap L^\infty(\mathbf{T}^n)$ where $\phi(r) = \phi(r) / \int_r^1 \phi(t) t^{-1} dt$. (A positive function $h(t)$ is said to be almost decreasing if there

* The second named author was partly supported by Grant-in-Aid for Co-operative Research (No. 57340004), Ministry of Education.

is a constant A such that $h(t) \leq Ah(t')$ if $t \geq t'$.)

However the case of \mathbf{R}^n is more complicated, and we must introduce a new function space similar to bmo_ϕ as follows. Let $w(x, r)$ be a positive function on $\mathbf{R}^n \times \mathbf{R}_+$. We define

$$bmo_w(\mathbf{R}^n) = \left\{ f \in L^1_{\text{loc}}(\mathbf{R}^n) ; \|f\|_{BMO_w} = \sup_{I(a,r)} \frac{MO(f, I(a, r))}{w(a, r)} < +\infty \right\}.$$

With a growth function $\phi(r)$, we always associate the function $w_\phi(x, r)$, defined by

$$w_\phi(x, r) = \phi(r) / \left(\left| \int_r^1 \phi(t) \frac{dt}{t} \right| + \int_1^{2+|x|} \phi(t) \frac{dt}{t} \right).$$

Then our main result is the following.

THEOREM 1. *Suppose $\phi(r)/r$ is almost decreasing. Then a function g is a pointwise multiplier on $bmo_\phi(\mathbf{R}^n)$ if and only if g belongs to $bmo_{w_\phi}(\mathbf{R}^n) \cap L^\infty(\mathbf{R}^n)$.*

We consider $bmo_\phi(\mathbf{R}^n)$ with the norm

$$\|f\|_{bmo_\phi} = |M(f, I(0, 1))| + \sup_{I(a,r)} \frac{MO(f, I(a, r))}{\phi(r)}.$$

Usually (see Janson [2]), $bmo_\phi(\mathbf{R}^n)$ is denoted by $BMO_\phi(\mathbf{R}^n)$ equipped with the seminorm

$$\|f\|_{BMO_\phi} = \sup_{I(a,r)} \frac{MO(f, I(a, r))}{\phi(r)}.$$

Then $BMO_\phi(\mathbf{R}^n)$ modulo constants is a Banach space, but $bmo_\phi(\mathbf{R}^n)$ is itself a Banach space modulo null-functions. To consider pointwise multipliers, the space $bmo_\phi(\mathbf{R}^n)$ is a more suitable one than $BMO_\phi(\mathbf{R}^n)$.

If we consider subspaces $bmo_\phi(\mathbf{R}^n) \cap L^p(\mathbf{R}^n)$, we obtain a similar result as follows.

THEOREM 2. *Suppose $\phi(r)/r$ is almost decreasing.*

(i) *Let $1 \leq p < \infty$. Then a function g is a pointwise multiplier from $bmo_\phi(\mathbf{R}^n) \cap L^p(\mathbf{R}^n)$ to $bmo_\phi(\mathbf{R}^n)$ if and only if $g \in bmo_\phi(\mathbf{R}^n) \cap L^\infty(\mathbf{R}^n)$, where $\psi(r) = \phi(r) / \int_{\min(1,r)}^2 \phi(t) t^{-1} dt$.*

(ii) *A function g is a pointwise multiplier from $bmo_\phi(\mathbf{R}^n) \cap L^\infty(\mathbf{R}^n)$ to $bmo_\phi(\mathbf{R}^n)$ if and only if $g \in bmo_\phi(\mathbf{R}^n) \cap L^\infty(\mathbf{R}^n)$.*

In these cases, g is a pointwise multiplier from $bmo_\phi(\mathbf{R}^n) \cap L^p(\mathbf{R}^n)$ into itself ($1 \leq p \leq \infty$).

If we define the Banach space $UBM-BMO_\phi(\mathbf{R}^n)$ by

$$\{f \in L^1_{\text{loc}}(\mathbf{R}^n) ; \|f\|_{UBM-BMO_\phi} = \|f\|_{BMO_\phi} + \sup_{a \in \mathbf{R}^n} M(f, I(a, 1)) < +\infty\},$$

then we have the following theorem similar to the torus case.

THEOREM 3. *Suppose $\phi(r)/r$ is almost decreasing. Then a function g is a pointwise multiplier from $UBM-BMO_\phi(\mathbf{R}^n)$ to $bmo_\phi(\mathbf{R}^n)$ if and only if $g \in bmo_\phi(\mathbf{R}^n) \cap L^\infty(\mathbf{R}^n)$, where ϕ is as in Theorem 2. In this case, g is a pointwise multiplier on $UBM-BMO_\phi(\mathbf{R}^n)$.*

It is known that $UBM-BMO_1(\mathbf{R}^n)$ is the dual space of the local Hardy space $h^1(\mathbf{R}^n)$, introduced by D. Goldberg [1]. Hence by duality we have, as in the torus case [2], the following:

COROLLARY 4. *A function g is a pointwise multiplier from $h^1(\mathbf{R}^n)$ to itself, if and only if $g \in bmo_\phi(\mathbf{R}^n) \cap L^\infty(\mathbf{R}^n)$, where $\phi(r) = 1 / \int_{\min(1,r)}^2 t^{-1} dt$.*

Our Theorem 1 answers a problem, which is implicitly stated in Johnson [3]. Stegenga [5] also treated the one dimensional torus case with $\phi \equiv 1$, and applied it to the boundedness problem of Toeplitz operators on the Hardy space $H^1(\mathbf{T})$. Applications of this paper will be treated in future.

Sections 2 and 3 are for the preliminaries and lemmas. In section 4 we give the proofs of Theorems 1, 2 and 3, and in section 5 we give some sufficient conditions for pointwise multipliers, and examples. The letter C will always denote a constant and does not necessarily denote the same one.

We note that the almost-decreasingness of $\phi(t)/t$ combined with the non-decreasingness of $\phi(t)$ implies that $\phi(t)$ is equivalent to a nondecreasing concave function. We have learned this from J. Peetre.

We would like to express our thanks to the referee. He gave us a proof of Lemma 3.4 simpler than ours, valid for $1 \leq p \leq \infty$, by which we could improve the case $p = \infty$ in Theorem 2.

2. Preliminaries.

First, we state some simple lemmas without proofs. (See for example Spanne [4].) We write

$$\rho(f, r) = \sup_{a \in \mathbf{R}^n, t \leq r} MO(f, I(a, t)).$$

LEMMA 2.1. $MO(f, I) \leq 2 \inf_c |I|^{-1} \int_I |f(x) - c| dx.$

LEMMA 2.2. *If $|F(x) - F(y)| \leq C|x - y|$, then $MO(F(f), I) \leq 2CMO(f, I)$.*

LEMMA 2.3. *Suppose that $I(a', r') \subset I(a, r)$. Then*

$$|M(f, I(a', r')) - M(f, I(a, r))| \leq C \int_{r'}^{2r} \frac{\rho(f, t)}{t} dt.$$

In the sequel, we always assume that $\phi(t)$ denotes a positive non-decreasing function and that $\phi(t)/t$ is almost decreasing. For each ϕ , we define strictly

positive functions $\Phi^*(r)$ and $\Phi_*(r)$:

$$\Phi^*(r) = \begin{cases} \int_1^r \phi(t)/t \, dt & (2 \leq r) \\ \int_1^2 \phi(t)/t \, dt & (0 < r < 2), \end{cases}$$

$$\Phi_*(r) = \begin{cases} \int_r^2 \phi(t)/t \, dt & (0 < r \leq 1) \\ \int_1^2 \phi(t)/t \, dt & (1 < r). \end{cases}$$

Then, by a slight modification of the proof of the theorem 2 (a) in Spanne [4, p. 601], we see that $\Phi^*(r)$ and $\Phi_*(r)$ belong to $bmo_\phi(\mathbf{R}_+)$. One can easily see that $f(|x|) \in bmo_\phi(\mathbf{R}^n)$ if $f(r) \in bmo_\phi(\mathbf{R}_+)$. Hence we have:

LEMMA 2.4. $\Phi^*(|x|), \Phi_*(|x|) \in bmo_\phi(\mathbf{R}^n)$.

Next we state some other properties of the functions $\Phi^*(r)$ and $\Phi_*(r)$.

LEMMA 2.5. (i) For any $k > 0$, there exists a constant $C_k > 0$ such that

$$C_k^{-1} \Phi^*(kr) \leq \Phi^*(r) \leq C_k \Phi^*(r/k) \quad \text{for all } r > 0.$$

(ii) For any $k > 0$, there is a constant $C_k > 0$ such that

$$C_k^{-1} \Phi_*(kr) \geq \Phi_*(r) \geq C_k \Phi_*(r/k) \quad \text{for all } r > 0.$$

(iii) There is a constant $C > 0$, depending only on the dimension n , such that

$$r^{-n} \int_0^r \Phi^*(t) t^{n-1} dt \geq C \Phi^*(r/2) \quad \text{for all } r > 0.$$

(iv) There is a constant $C > 0$ such that

$$r^{-1} \int_0^r \frac{dt}{\Phi^*(t)} \leq C \frac{\phi(r)}{\Phi^*(r)} \quad \text{for all } r \geq 2.$$

PROOF. (i) Since $\Phi^*(r)$ is non decreasing, it is clear for $0 < k \leq 1$. So we assume $k > 1$. If $r \leq kr \leq 2$, then $\Phi^*(kr) = \Phi^*(r)$. If $r \leq 2 \leq kr$, then $\Phi^*(kr) \leq \Phi^*(2k) \leq C_k \Phi^*(2) = C_k \Phi^*(r)$. And if $2 \leq r \leq kr$, then, since $\phi(t)/t$ is almost decreasing, we get

$$\begin{aligned} \Phi^*(kr) &= \int_1^{kr} \phi(t) \frac{dt}{t} = \int_{1/k}^r \phi(kt) \frac{dt}{t} \leq \int_{1/k}^r A k \phi(t) \frac{dt}{t} \\ &\leq C_k \Phi^*(r). \end{aligned}$$

Therefore we get $\Phi^*(kr) \leq C_k \Phi^*(r)$ for all $r > 0$. And hence $\Phi^*(r) \leq C_k \Phi^*(r/k)$ for all $r > 0$.

(ii) In a way similar to the case (i) we get (ii).

(iii) Since $\Phi^*(t)$ is non-decreasing, we have

$$\begin{aligned} r^{-n} \int_0^r \Phi^*(t) t^{n-1} dt &\geq r^{-n} \int_{r/2}^r \Phi^*(t) t^{n-1} dt \\ &\geq n^{-1} (1 - 2^{-n}) \Phi^*(r/2). \end{aligned}$$

(iv) Since

$$\Phi^*(t) \geq \max \{ \phi(1) \log t, \phi(1) \log 2 \} > \frac{1}{4} \phi(1) \log (e^2 + t),$$

and

$$\frac{1}{\log (e^2 + t)} \leq \frac{2}{\log (e^2 + t)} \left(1 - \frac{1}{\log (e^2 + t)} \right) = 2 \frac{d}{dt} \left(\frac{e^2 + t}{\log (e^2 + t)} \right),$$

we have

$$\begin{aligned} \frac{1}{r} \int_0^r \frac{dt}{\Phi^*(t)} &\leq \frac{8}{\phi(1)r} \left[\frac{e^2 + t}{\log (e^2 + t)} \right]_0^r < \frac{8}{\phi(1)r} \frac{e^2 + r}{\log (e^2 + r)} \\ &< C / \log r, \qquad \text{as } r \geq 2. \end{aligned}$$

Hence we have the desired inequality, since

$$\phi(r) \log r = \int_1^r \phi(t) \frac{dt}{t} \geq \int_1^r \phi(t) \frac{dt}{t} = \Phi^*(r) \quad \text{as } r \geq 2.$$

q. e. d.

REMARK 2.1. By this lemma there is a constant $C > 0$ such that

$$\begin{aligned} (2.1) \quad C^{-1} (\Phi_*(r) + \Phi^*(r) + \Phi^*(|x|)) &\leq \left| \int_r^1 \phi(t) \frac{dt}{t} \right| + \int_1^{2^{+1}x_1} \phi(t) \frac{dt}{t} \\ &\leq C (\Phi_*(r) + \Phi^*(r) + \Phi^*(|x|)). \end{aligned}$$

Finally in this section, we note one more fact (Spanne [4, p. 601]).

LEMMA 2.6. If $\int_0^1 \phi(t) t^{-1} dt < +\infty$, then

$$\omega(f, r) = \text{ess sup}_{|x-y| \leq r} |f(x) - f(y)| \leq C \int_0^r \phi(t) \frac{dt}{t} \|f\|_{BMO_\phi},$$

for any $f \in bmo_\phi(\mathbf{R}^n)$.

3. Lemmas.

To prove the theorems, we show a few lemmas in this section.

LEMMA 3.1. There is a constant $C > 0$ such that

$$|M(f, I(a, r))| \leq C \|f\|_{bmo_\phi} (\Phi_*(r) + \Phi^*(r) + \Phi^*(|a|))$$

for any $f \in bmo_\phi(\mathbf{R}^n)$ and for any cube $I(a, r)$.

PROOF. Case 1: $r \geq 1$, $|a| \geq r$. Since $I(a, r)$, $I(0, 1) \subset I(0, r+2|a|)$, by Lemma 2.3 and Lemma 2.5 (i), we have

$$\begin{aligned} & |M(f, I(a, r)) - M(f, I(0, 1))| \\ & \leq |M(f, I(a, r)) - M(f, I(0, r+2|a|))| + |M(f, I(0, 1)) - M(f, I(0, r+2|a|))| \\ & \leq C \int_r^{2(r+2|a|)} \rho(f, t) \frac{dt}{t} + C \int_1^{2(r+2|a|)} \rho(f, t) \frac{dt}{t} \\ & \leq 2C \int_1^{6|a|} \rho(f, t) \frac{dt}{t} \leq 2C \|f\|_{BMO_\phi} \int_1^{6|a|} \phi(t) \frac{dt}{t} \\ & = 2C \|f\|_{BMO_\phi} \Phi^*(6|a|) \leq C' \|f\|_{BMO_\phi} \Phi^*(|a|). \end{aligned}$$

Case 2: $1 \leq r$, $|a| \leq r$. Since $I(a, r)$, $I(0, 1) \subset I(0, r+2|a|)$, in a way similar to the case 1, we have

$$|M(f, I(a, r)) - M(f, I(0, 1))| \leq C \|f\|_{BMO_\phi} \Phi^*(r).$$

Case 3: $r \leq 1$, $1 \leq |a|$. Since $I(a, r)$, $I(0, 1) \subset I(0, r+2|a|)$, by Lemma 2.3 and Lemma 2.5 (i), we have

$$\begin{aligned} & |M(f, I(a, r)) - M(f, I(0, 1))| \\ & \leq C \int_r^{2(r+2|a|)} \rho(f, t) \frac{dt}{t} + C \int_1^{2(r+2|a|)} \rho(f, t) \frac{dt}{t} \\ & \leq C \int_r^1 \rho(f, t) \frac{dt}{t} + 2C \int_1^{6|a|} \rho(f, t) \frac{dt}{t} \\ & \leq C \|f\|_{BMO_\phi} \int_r^1 \phi(t) \frac{dt}{t} + 2C \|f\|_{BMO_\phi} \int_1^{6|a|} \phi(t) \frac{dt}{t} \\ & \leq C' \|f\|_{BMO_\phi} (\Phi_*(r) + \Phi^*(|a|)). \end{aligned}$$

Case 4: $r \leq 1$, $|a| \leq 1$. Since $I(a, r)$, $I(0, 1) \subset I(0, 3)$, by Lemma 2.3 we get

$$\begin{aligned} & |M(f, I(a, r)) - M(f, I(0, 1))| \\ & \leq |M(f, I(a, r)) - M(f, I(0, 3))| + |M(f, I(0, 1)) - M(f, I(0, 3))| \\ & \leq C \int_r^6 \rho(f, t) \frac{dt}{t} + C \int_1^6 \rho(f, t) \frac{dt}{t} \leq 2C \|f\|_{BMO_\phi} \int_r^6 \phi(t) \frac{dt}{t} \\ & \leq C' \|f\|_{BMO_\phi} \Phi_*(r). \end{aligned}$$

Summing up the above cases, we obtain

$$|M(f, I(a, r))| \leq |M(f, I(0, 1))| + C \|f\|_{BMO_\phi} (\Phi_*(r) + \Phi^*(r) + \Phi^*(|a|))$$

$$\leq C' \|f\|_{bmo_\phi} (\Phi_*(r) + \Phi^*(r) + \Phi^*(|a|)).$$

q. e. d.

REMARK 3.1. The estimate in Lemma 3.1 is sharp. In fact, consider the functions $\Phi^*(|x|)$ and $\Phi_*(|x-a|)$. Then $\|\Phi^*(|x|)\|_{bmo_\phi}, \|\Phi_*(|x-a|)\|_{bmo_\phi} < C$, independently of $a \in \mathbf{R}^n$, since $\Phi_*(r) \leq \phi(2) \max(\log(2/r), \log 2)$. If $4|a| \leq r$, using $\{x; |x| \leq r/4\} \subset I(a, r)$ we get

$$\begin{aligned} M(\Phi^*(|x|), I(a, r)) &\geq |I(a, r)|^{-1} \int_{|x| \leq r/4} \Phi^*(|x|) dx \\ &= C_1 r^{-n} \int_0^{r/4} \Phi^*(t) t^{n-1} dt \\ &\geq C_2 \Phi^*(r/8) \geq C_3 \Phi^*(r) \geq C_3 \Phi^*(|a|), \end{aligned}$$

by using Lemma 2.5 (i) and (iii). If $r < 4|a|$, by Lemma 2.5 (i)

$$\begin{aligned} M(\Phi^*(|x|), I(a, r)) &\geq C_1 M\left(\Phi^*(|x|), I\left(a, \frac{r}{4n}\right)\right) \geq C_2 \Phi^*(|a|) \\ &\geq C_2 \Phi^*\left(\frac{r}{4}\right) \geq C_3 \Phi^*(r), \end{aligned}$$

since $\Phi^*(|x|) \geq C_4 \Phi^*(|a|)$ on $I(a, r/(4n))$ by Lemma 2.5 (i). Next we consider $\Phi_*(|x-a|)$. Since $\Phi_*(r)$ is non-increasing and $\{x; |x-a| < r/2\} \subset I(a, r)$, we have

$$\begin{aligned} M(\Phi_*(|x-a|), I(a, r)) &\geq |I(a, r)|^{-1} \int_{|x-a| < r/2} \Phi_*(|x-a|) dx \\ &\geq C \Phi_*\left(\frac{r}{2}\right) \geq C' \Phi_*(r) \end{aligned}$$

by using Lemma 2.5 (ii).

LEMMA 3.2. Suppose $1 \leq p \leq \infty$. There is a constant $C > 0$ such that

$$|M(f, I(a, r))| \leq C (\|f\|_{BMO_\phi} + \|f\|_{L^p}) \Phi_*(r)$$

for any $f \in bmo_\phi(\mathbf{R}^n) \cap L^p(\mathbf{R}^n)$, and for any cube $I(a, r)$.

PROOF. If $1 \leq r$, we have by Hölder's inequality

$$|M(f, I(a, r))| \leq \left(|I|^{-1} \int_I |f(x)|^p dx \right)^{1/p} \leq \|f\|_{L^p} \leq C \Phi_*(r) \|f\|_{L^p}.$$

If $0 < r < 1$, since $I(a, r) \subset I(a, 1)$, by Lemma 2.3 we have

$$\begin{aligned} |M(f, I(a, r))| &\leq |M(f, I(a, r)) - M(f, I(a, 1))| + |M(f, I(a, 1))| \\ &\leq C \int_r^2 \rho(f, t) \frac{dt}{t} + |M(f, I(a, 1))| \\ &\leq C \|f\|_{BMO_\phi} \Phi_*(r) + \|f\|_{L^p} \end{aligned}$$

$$\leq C'(\|f\|_{BMO_\phi} + \|f\|_{L^p})\Phi_*(r).$$

q. e. d.

REMARK 3.2. Let $f(x) = \Phi_*(|x-a|) - \Phi_*(1)$. Then, since $\Phi_*(r) \leq \phi(2) \times \max(\log 2/r, \log 2)$, $\|f\|_{L^p} \leq C_p$ ($1 \leq p < \infty$), $\|f\|_{BMO_\phi} \leq C$, independently of a . As in Remark 3.1, we have $M(f, I(a, r)) \geq C\Phi_*(r)$ for $r \leq 1$.

LEMMA 3.3. Suppose $f \in bmo_\phi(\mathbf{R}^n)$ and $g \in L^\infty(\mathbf{R}^n)$. Then, fg belongs to $bmo_\phi(\mathbf{R}^n)$ if and only if

$$F(f, g) = \sup_{I(a, r)} |M(f, I(a, r))| MO(g, I(a, r)) / \phi(r) < +\infty.$$

In this case,

$$F(f, g) \leq \|fg\|_{BMO_\phi} + 2\|g\|_\infty \|f\|_{BMO_\phi}.$$

PROOF. For any cube $I = I(a, r)$, by elementary calculation (see for example Stegenga [5, p. 582]), we have

$$|MO(fg, I) - |f_I| MO(g, I)| \leq 2\|g\|_\infty MO(f, I),$$

and therefore

$$\left| \frac{MO(fg, I)}{\phi(r)} - \frac{|f_I| MO(g, I)}{\phi(r)} \right| \leq 2\|g\|_\infty \|f\|_{BMO_\phi}.$$

This implies the assertion by the definition of $bmo_\phi(\mathbf{R}^n)$.

q. e. d.

LEMMA 3.4. Suppose $1 \leq p \leq \infty$. If g is a pointwise multiplier from $bmo_\phi(\mathbf{R}^n) \cap L^p(\mathbf{R}^n)$ to $bmo_\phi(\mathbf{R}^n)$, then it follows that $g \in L^\infty(\mathbf{R}^n)$.

PROOF. First of all, since $bmo_\phi(\mathbf{R}^n) \cap L^p(\mathbf{R}^n)$ is a Banach space, equipped with the norm $\|f\|_{BMO_\phi} + \|f\|_{L^p}$, and $bmo_\phi(\mathbf{R}^n)$ is also a Banach space, we have by the closed graph theorem that

$$\|gf\|_{bmo_\phi} \leq C(\|f\|_{BMO_\phi} + \|f\|_{L^p})$$

for all $f \in bmo_\phi(\mathbf{R}^n) \cap L^p(\mathbf{R}^n)$.

For any cube $I = I(a, r)$ with $r < 1$, we define a function $h \in bmo_\phi(\mathbf{R}^n) \cap L^p(\mathbf{R}^n)$ as follows:

$$h(x) = \begin{cases} 0 & r \leq |x-a| \\ \exp(i\Phi_*(|x-a|)) - \exp(i\Phi_*(r)) & |x-a| < r. \end{cases}$$

Then, by Lemma 2.2, we get $\|h\|_{BMO_\phi} \leq C_0 \|\Phi_*(|x|)\|_{BMO_\phi}$. And, since $\text{supp } h \subset I(a, 2)$ and $|h(x)| \leq 2$, we have $\|h\|_{L^p} \leq C_p$. Hence $\|gh\|_{bmo_\phi} \leq C(\|h\|_{BMO_\phi} + \|h\|_{L^p}) \leq C_1$, independently of I . This gives

$$(3.1) \quad MO(gh, I(a, 4r)) \leq C_1 \phi(4r).$$

Let C_2 and C_3 be constants such that $\log C_2 = \pi/\phi(1)$, $1 < C_2 C_3 < C_2$, and let $L_r =$

$\{x; r/C_2 \leq |x-a| \leq r/(C_2C_3)\}$. If $x \in L_r$, then, since $\phi(r)/r$ is almost decreasing, we have

$$\begin{aligned} (\phi(r)/(AC_2C_3)) \log C_2C_3 &\leq \phi(r/(C_2C_3)) \log C_2C_3 \leq \int_{r/(C_2C_3)}^r \phi(t) \frac{dt}{t} \\ &\leq \Phi_*(|x-a|) - \Phi_*(r) \leq \int_{r/C_2}^r \phi(t) \frac{dt}{t} \leq \phi(1) \log C_2 = \pi. \end{aligned}$$

So, the inequality $|e^{i\theta} - 1| \geq 2\theta/\pi$ ($0 \leq \theta \leq \pi$) implies that $|h(x)| \geq C_4\phi(r)$ for $x \in L_r$. Let $\sigma = M(gh, I(a, 4r))$. Then we have, by considering the support of h ,

$$\begin{aligned} MO(gh, I(a, 4r))|I(a, 4r)| &= \int_{I(a, 4r)} |gh(x) - \sigma| dx \\ &\geq \int_{L_r} |gh(x) - \sigma| dx + \int_{I(a, 4r) \setminus I(a, 2r)} |\sigma| dx \\ &\geq \int_{L_r} (|gh(x) - \sigma| + |\sigma|) dx \geq \int_{L_r} |gh(x)| dx \\ &\geq C_4\phi(r) \int_{L_r} |g(x)| dx, \end{aligned}$$

and so

$$(3.2) \quad |L_r|^{-1} \int_{L_r} |g(x)| dx \leq C_5 MO(gh, I(a, 4r)) / \phi(r)$$

From (3.1) and (3.2) it follows that

$$|L_r|^{-1} \int_{L_r} |g(x)| dx \leq C_6.$$

Letting r tend to zero, we have

$$|g(a)| \leq C_6 \quad \text{a. e.}$$

q. e. d.

4. Proofs of the theorems.

PROOF OF THEOREM 1. Suppose that g is a pointwise multiplier on $bmo_\phi(\mathbf{R}^n)$. Then $g \in L^\infty$ by Lemma 3.4. Since $gf \in bmo_\phi(\mathbf{R}^n)$ for all $f \in bmo_\phi(\mathbf{R}^n)$, by Lemma 3.3 and the closed graph theorem we have

$$(4.1) \quad \sup_{I(a,r)} \frac{|f_I| MO(g, I)}{\phi(r)} < C \|f\|_{bmo_\phi}.$$

Hence, taking $f(x) = \Phi^*(|x|)$ or $\Phi_*(|x-a|)$, we have by Remark 3.1

$$(4.2) \quad \sup_{I(a,r)} (\Phi_*(r) + \Phi^*(r) + \Phi^*(|a|)) MO(g, I) / \phi(r) < +\infty,$$

and hence

$$\sup_{I(a,r)} \frac{MO(g, I(a, r))}{w_\phi(a, r)} < +\infty,$$

by using Remark 2.1. Consequently $g \in bmo_{w_\phi}(\mathbf{R}^n) \cap L^\infty(\mathbf{R}^n)$.

Conversely, suppose $g \in bmo_{w_\phi}(\mathbf{R}^n) \cap L^\infty(\mathbf{R}^n)$. For any $I=I(a, r)$ and any $f \in bmo_\phi(\mathbf{R}^n)$, by Lemma 3.1 we get

$$\begin{aligned} \frac{|f_I| MO(g, I)}{\phi(r)} &\leq C \|f\|_{bmo_\phi} (\Phi_*(r) + \Phi^*(r) + \Phi^*(|a|)) MO(g, I) / \phi(r) \\ &\leq C' \|f\|_{bmo_\phi} \frac{MO(g, I)}{w_\phi(a, r)} \leq C' \|f\|_{bmo_\phi} \|g\|_{BMO_{w_\phi}}. \end{aligned}$$

Therefore $fg \in bmo_\phi(\mathbf{R}^n)$ by Lemma 3.3, which shows that g is a pointwise multiplier on $bmo_\phi(\mathbf{R}^n)$. This proves Theorem 1.

PROOF OF THEOREM 2. (i) Case $1 \leq p < \infty$. Suppose that g is a pointwise multiplier from $bmo_\phi(\mathbf{R}^n) \cap L^p(\mathbf{R}^n)$ to $bmo_\phi(\mathbf{R}^n)$. Then g is bounded by Lemma 3.4. Hence by Lemma 3.3

$$\sup_{I(a,r)} \frac{|f_I| MO(g, I)}{\phi(r)} \leq C (\|f\|_{BMO_\phi} + \|f\|_{L^p}).$$

Taking $f(x) = \Phi_*(|x-a|) - \Phi_*(1)$, we have by Remark 3.2

$$\sup_{r \leq 1, a \in \mathbf{R}^n} \Phi_*(r) MO(g, I) / \phi(r) < +\infty.$$

According to $g \in L^\infty(\mathbf{R}^n)$, $MO(g, I) \leq 2\|g\|_\infty$. Since $\Phi_*(r)$ is constant and $\phi(r) \geq \phi(1)$ for $r \geq 1$, we have

$$\sup_{r > 1, a \in \mathbf{R}^n} \Phi_*(r) MO(g, I) / \phi(r) < +\infty.$$

Thus $g \in bmo_\phi(\mathbf{R}^n) \cap L^\infty(\mathbf{R}^n)$. Sufficiency can be proved in the same way as in Theorem 1, using Lemma 3.2 in place of Lemma 3.1. (ii) Case $p = \infty$. (Necessity) Since $1 \in bmo_\phi(\mathbf{R}^n) \cap L^\infty(\mathbf{R}^n)$, g must belong to $bmo_\phi(\mathbf{R}^n)$. By Lemma 3.4, g is bounded. (Sufficiency) We have, for any cube I ,

$$|f_I| MO(g, I) / \phi(r) \leq \|f\|_\infty MO(g, I) / \phi(r) \leq \|f\|_\infty \|g\|_{BMO_\phi}.$$

So, since g is bounded, by Lemma 3.3 we have $fg \in bmo_\phi(\mathbf{R}^n)$. This completes the proof.

PROOF OF THEOREM 3. (Necessity) Clearly we have $bmo_\phi(\mathbf{R}^n) \cap L^2(\mathbf{R}^n) \subset UBM-BMO_\phi(\mathbf{R}^n)$. Hence by Theorem 2 we have the desired conclusion. (Sufficiency) For all $r \geq 1$ and all $a \in \mathbf{R}^n$, we get

$$(4.3) \quad |M(|f|, I(a, r))| \leq 2^n \left[\sup_{b \in \mathbf{R}^n} |M(f, I(b, 1))| + \sup_{b \in \mathbf{R}^n} MO(f, I(b, 1)) \right].$$

(To show this, let j be the smallest integer satisfying $r \leq 2^j$ and take the cube

$I(a, 2^j)$, and then divide it into non-overlapping 2^{jn} cubes with side length 1. Then by the definition we get the above inequality.) Hence we get $\sup_{r \geq 1, a \in \mathbf{R}^n} |M(f, I(a, r))|/\phi(r) \leq C\|f\|_{UBM-BMO_\phi}$. As in Case 4 in Lemma 3.1 we get $|M(f, I(a, r))| \leq C\|f\|_{UBM-BMO_\phi} \Phi_*(r)$. Therefore, since g is bounded, we have $gf \in bmo_\phi(\mathbf{R}^n)$ by Lemma 3.3. q. e. d.

REMARK 4.1. By (4.3), one can easily show that $\|f\|_{UBM-BMO_\phi}$ is equivalent to

$$(4.4) \quad \sup_{0 < r \leq 1, a \in \mathbf{R}^n} |MO(f, I(a, r))|/\phi(r) + \sup_{r \geq 1, a \in \mathbf{R}^n} |M(|f|, I(a, r))|/\phi(r).$$

For the case $\phi(t) \equiv 1$, Goldberg [1, Corollary 1] introduced $UBM-BMO_1(\mathbf{R}^n)$, using (4.4), by the symbol bmo , and showed that it is the dual of the local Hardy space $h^1(\mathbf{R}^n)$.

5. Some sufficient conditions and examples.

As consequences of our theorems, we give some sufficient conditions for pointwise multipliers, corresponding to those in the torus case, Stegenga [5, Corollary 2.8] and Janson [2, p. 196].

PROPOSITION 5.1. *Suppose g satisfies the following conditions:*

(5.1) *There is a constant $M_1 > 0$ such that*

$$|g(x+y) - g(x)| \leq M_1 \phi(|y|) / [\Phi_*(|y|) + (1 - \text{sgn } \phi(0+)) \Phi^*(|x|)]$$

for all $x, y \in \mathbf{R}^n$ with $|y| \leq 1$, where $\phi(0+) = \lim_{r \downarrow 0} \phi(r)$.

(5.2) *There are constants $M_2 > 0$ and $B \in \mathbf{C}$ such that*

$$|g(x) - B| \leq M_2 / \Phi^*(|x|) \quad \text{for all } x \in \mathbf{R}^n.$$

Then, g is a pointwise multiplier on $bmo_\phi(\mathbf{R}^n)$.

PROOF. We omit the detailed proof. One has only to treat the four cases; $\{r \leq 1/\sqrt{n}, \phi(0+) = 0\}$, $\{r \leq 1/\sqrt{n}, \phi(0+) > 0\}$, $\{1/\sqrt{n} < r \leq |a|/\sqrt{n}\}$, and $\{r \geq \max(1, |a|/\sqrt{n})\}$.

As a consequence we have the following corollary, whose proof we omit.

COROLLARY 5.2. *If $g = g_1/g_2$ satisfies the following conditions:*

(5.3) *g_1 is bounded and there is a $C_1 > 0$ such that $|g_1(x) - g_1(y)| \leq C_1|x - y|$, $x, y \in \mathbf{R}^n$;*

(5.4) *There are $C_2, C_3 > 0$ such that $|g_2(x)| \geq C_2\Phi^*(|x|)$ and $|g_2(x) - g_2(y)| \leq C_3|x - y|$, $x, y \in \mathbf{R}^n$.*

Then, g is a pointwise multiplier on $bmo_\phi(\mathbf{R}^n)$.

For pointwise multipliers from $bmo_\phi \cap L^p$ to bmo_ϕ , we have:

PROPOSITION 5.3. *If g is bounded and satisfies*

$$(5.5) \quad |g(x+y) - g(x)| \leq C\phi(|y|)/\Phi_*(|y|), \quad x, y \in \mathbf{R}^n, |y| < 1,$$

then g is a pointwise multiplier from $bmo_\phi(\mathbf{R}^n) \cap L^p(\mathbf{R}^n)$ to $bmo_\phi(\mathbf{R}^n)$, ($1 \leq p \leq \infty$).

EXAMPLES. By Corollary 5.2

$$\frac{1}{\Phi^*(|x|)}, \quad \frac{\sin|x|}{\Phi^*(|x|)}, \quad \frac{1}{1+|x|}, \quad \frac{\sin\Phi^*(|x|)}{1+|x|}$$

are pointwise multipliers on $bmo_\phi(\mathbf{R}^n)$. And, for any ϕ , for which $\phi(t)/(t\Phi_*(t))$ is almost decreasing, put $\Psi_*(r) = \int_r^2 \phi(t)/(t\Phi_*(t)) dt$ for $0 < r \leq 1$ and $= \int_1^2 \phi(t)/(t\Phi_*(t)) dt$ for $1 < r$. Then, $\sin\Psi_*(|x|)/\Phi^*(|x|)$ is a pointwise multiplier on $bmo_\phi(\mathbf{R}^n)$. This gives a pointwise multiplier, which is not continuous, as in [2, p. 196].

References

- [1] D. Goldberg, A local version of real Hardy spaces, *Duke Math. J.*, **46** (1979), 27-42.
- [2] S. Janson, On functions with conditions on the mean oscillation, *Ark. Mat.*, **14** (1976), 189-196.
- [3] R. Johnson, Behaviour of BMO under certain functional operations, preprint.
- [4] S. Spanne, Some function spaces defined using the mean oscillation over cubes, *Ann. Scuola Norm. Sup. Pisa*, **19** (1965), 593-608.
- [5] D. A. Stegenga, Bounded Toeplitz operators on H^1 and applications of duality between H^1 and the functions of bounded mean oscillation, *Amer. J. Math.*, **98** (1976), 573-589.

Eiichi NAKAI

Yuki Daiichi Senior High School
1076 Yuki, Yuki-shi
Ibaraki-ken 307
Japan

Kôzô YABUTA

Department of Mathematics
Ibaraki University
Mito, Ibaraki 310
Japan