

GROWTH OF SOLUTIONS OF AN n -th ORDER LINEAR DIFFERENTIAL EQUATION WITH ENTIRE COEFFICIENTS

BENHARRAT BELAÏDI AND SAADA HAMOUDA

Abstract

We consider a differential equation $f^{(n)} + A_{n-1}(z)f^{(n-1)} + \cdots + A_1(z)f' + A_0(z)f = 0$, where $A_0(z), \dots, A_{n-1}(z)$ are entire functions with $A_0(z) \not\equiv 0$. Suppose that there exist a positive number μ , and a sequence $(z_j)_{j \in \mathbb{N}}$ with $\lim_{j \rightarrow +\infty} z_j = \infty$, and also two real numbers α, β ($0 \leq \beta < \alpha$) such that $|A_0(z_j)| \geq e^{\alpha|z_j|^\mu}$ and $|A_k(z_j)| \leq e^{\beta|z_j|^\mu}$ as $j \rightarrow +\infty$ ($k = 1, \dots, n-1$). We prove that all solutions $f \not\equiv 0$ of this equation are of infinite order. This result is a generalization of one theorem of Gundersen ([3], p. 418).

1. Introduction

For $n \geq 2$, we consider a linear differential equation

$$(1.1) \quad f^{(n)} + A_{n-1}(z)f^{(n-1)} + \cdots + A_1(z)f' + A_0(z)f = 0,$$

where $A_0(z), \dots, A_{n-1}(z)$ are entire functions with $A_0(z) \not\equiv 0$. Let $\rho(f)$ denote the order of an entire function f , that is,

$$\rho(f) = \overline{\lim}_{r \rightarrow +\infty} \frac{\log T(r, f)}{\log r} = \overline{\lim}_{r \rightarrow +\infty} \frac{\log \log M(r, f)}{\log r},$$

where $T(r, f)$ is the Nevanlinna characteristic function of f (see [6]), and $M(r, f) = \max_{|z|=r} |f(z)|$.

It is well-known that all solutions of (1.1) are entire functions, and if some of the coefficients of (1.1) are transcendental, then (1.1) has at least one solution with order $\rho(f) = +\infty$.

The question which arises is: What conditions on $A_0(z), \dots, A_{n-1}(z)$ will guarantee that every solution $f \not\equiv 0$ of (1.1) has infinite order?

In this paper we prove three results concerning this question. According to [5], [7, pp. 199–209], [8, pp. 106–108], [9, pp. 65–67], we know that if

2000 *Mathematics Subject Classification*: Primary 34M10; Secondary 30D35.

Keywords: Linear differential equations, entire functions, finite order of growth.

Received October 10, 2001; revised May 1, 2002.

$A_0(z), \dots, A_{n-1}(z)$ are polynomials with $A_0(z) \not\equiv 0$, then every solution f of (1.1) is an entire function with finite rational order.

In the study of the differential equation

$$(1.2) \quad f'' + A(z)f' + B(z)f = 0,$$

where $A(z)$ and $B(z) \not\equiv 0$ are entire functions, Gundersen proved the following result:

THEOREM 1.1 ([3, p. 418]). *Let $A(z)$ and $B(z) \not\equiv 0$ be entire functions, and let α, β, θ_1 and θ_2 be real numbers with $\alpha > 0, \beta > 0$ and $\theta_1 < \theta_2$. If*

$$(1.3) \quad |B(z)| \geq \exp\{(1 + o(1))\alpha|z|^\beta\}$$

and

$$(1.4) \quad |A(z)| \leq \exp\{o(1)|z|^\beta\}$$

as $z \rightarrow \infty$ with $\theta_1 \leq \arg z \leq \theta_2$, then every solution $f \not\equiv 0$ of (1.2) has infinite order.

Remark. Theorem 1.1 was recently extended for an n -th order linear differential equation (see [1]).

In the same paper, Gundersen also proved the following:

THEOREM 1.2 ([3, p. 417]). *Let $A(z)$ and $B(z)$ be entire functions such that either (i) $\rho(A) < \rho(B)$ or (ii) A is a polynomial and B is transcendental. Then every solution $f \not\equiv 0$ of (1.2) has infinite order.*

2. Statement and proof of results

In this paper we prove the following results:

THEOREM 2.1. *Suppose that there exist a positive number μ , and a sequence of points $(z_j)_{j \in \mathbb{N}}$ with $\lim_{j \rightarrow +\infty} z_j = \infty$, and two real numbers α, β ($0 \leq \beta < \alpha$) such that*

$$(2.1) \quad |A_0(z_j)| \geq e^{\alpha|z_j|^\mu}$$

and

$$(2.2) \quad |A_k(z_j)| \leq e^{\beta|z_j|^\mu} \quad (k = 1, \dots, n-1),$$

as $j \rightarrow +\infty$. Then every solution $f \not\equiv 0$ of (1.1) has infinite order.

From Theorem 2.1, we deduce the following two results:

COROLLARY 2.2. *Suppose that*

$$(2.3) \quad \max\{\rho(A_k) : k = 1, \dots, n-1\} < \rho(A_0).$$

Then every solution $f \not\equiv 0$ of (1.1) has infinite order.

COROLLARY 2.3. *Suppose that $A_1(z), \dots, A_{n-1}(z)$ are polynomials and $A_0(z)$ is transcendental. Then every solution $f \not\equiv 0$ of (1.1) has infinite order.*

Corollary 2.2 and Corollary 2.3 were proved by Z.-x. Chen and S.-a. Gao in [2]. In this paper, we give another proof.

In the proof of Theorem 2.1, we need the following lemma:

LEMMA 2.4 ([4, p. 89]). *Let w be a transcendental entire function of finite order ρ . Let $\Gamma = \{(k_1, j_1), (k_2, j_2), \dots, (k_m, j_m)\}$ denote a finite set of distinct pairs of integers satisfying $k_i > j_i \geq 0$ for $i = 1, \dots, m$, and let $\varepsilon > 0$ be a given constant. Then there exists a set $E \subset [0, 2\pi)$ with linear measure zero such that, if $\psi_0 \in [0, 2\pi) - E$, then there is a constant $R_0 = R_0(\psi_0) > 1$ such that for all z satisfying $\arg z = \psi_0$ with $|z| \geq R_0$ and for all $(k, j) \in \Gamma$, the following estimate hold:*

$$\left| \frac{w^{(k)}(z)}{w^{(j)}(z)} \right| \leq |z|^{(k-j)(\rho-1+\varepsilon)}.$$

Proof of Theorem 2.1. Suppose that $f \not\equiv 0$ is a solution of (1.1) with $\rho(f) < +\infty$. We can write (1.1) as

$$(2.4) \quad \frac{1}{A_0(z)} \frac{f^{(n)}}{f} + \sum_{k=1}^{n-1} \frac{A_k(z)}{A_0(z)} \frac{f^{(k)}}{f} = -1.$$

Then by Lemma 2.4, there exists a set $E \subset [0, 2\pi)$ with linear measure zero such that, if $\psi_0 \in [0, 2\pi) - E$, then there is a constant $R_0 = R_0(\psi_0) > 1$ such that for all z_j satisfying $\arg z_j = \psi_0$ with $|z_j| \geq R_0$ and for all $k = 1, 2, \dots, n$, we have

$$(2.5) \quad \left| \frac{f^{(k)}(z_j)}{f(z_j)} \right| \leq |z_j|^{kc} \quad (k = 1, \dots, n, \quad c = \rho - 1 + \varepsilon).$$

Then by (2.1), (2.2) and (2.5), we obtain that

$$(2.6) \quad \left| \frac{A_k(z_j)}{A_0(z_j)} \right| \left| \frac{f^{(k)}(z_j)}{f(z_j)} \right| \leq \frac{1}{e^{(\alpha-\beta)|z_j|^\mu}} |z_j|^{kc} \quad (k = 1, \dots, n-1, \quad c = \rho - 1 + \varepsilon).$$

Since

$$\lim_{j \rightarrow +\infty} \frac{1}{e^{(\alpha-\beta)|z_j|^\mu}} |z_j|^{kc} = 0 \quad (k = 1, \dots, n-1, \quad c = \rho - 1 + \varepsilon),$$

we see

$$\lim_{j \rightarrow +\infty} \left| \frac{A_k(z_j)}{A_0(z_j)} \right| \left| \frac{f^{(k)}(z_j)}{f(z_j)} \right| = 0 \quad (k = 1, \dots, n-1).$$

We have also from (2.1) and (2.5) that

$$(2.7) \quad \left| \frac{1}{A_0(z_j)} \right| \left| \frac{f^{(n)}(z_j)}{f(z_j)} \right| \leq \frac{1}{e^{\alpha|z_j|^\mu}} |z_j|^{nc} \quad (c = \rho - 1 + \varepsilon),$$

which implies

$$(2.8) \quad \lim_{j \rightarrow +\infty} \left| \frac{1}{A_0(z_j)} \right| \left| \frac{f^{(n)}(z_j)}{f(z_j)} \right| = 0.$$

Letting $j \rightarrow +\infty$ in the relation

$$\frac{1}{A_0(z_j)} \frac{f^{(n)}(z_j)}{f(z_j)} + \sum_{k=1}^{n-1} \frac{A_k(z_j)}{A_0(z_j)} \frac{f^{(k)}(z_j)}{f(z_j)} = -1,$$

we get a contradiction. Thus every solution $f \not\equiv 0$ of (1.1) has infinite order.

Next, we give two examples that illustrates Theorem 2.1.

Example 1. Consider the differential equation

$$(2.9) \quad f'' - f' - e^{2z}f = 0.$$

In this equation, for $z = z_j = j \rightarrow \infty$, we have

$$\begin{aligned} |A_0(z_j)| &= |-e^{2z_j}| = e^{2j} \geq e^{2|z_j|}, \\ |A_1(z_j)| &= 1 \leq e^{0|z_j|}. \end{aligned}$$

It is easy to see that the conditions (2.1) and (2.2) in Theorem 2.1 are satisfied. The two linearly independent functions $f_1(z) = e^{e^z}$ and $f_2(z) = e^{-e^z}$ are solutions of (2.9) with $\rho(f_1) = \rho(f_2) = +\infty$.

Example 2. Consider the differential equation

$$(2.10) \quad f''' - (3 + 6e^z)f'' + (2 + 6e^z + 11e^{2z})f' - 6e^{3z}f = 0.$$

In this equation, for $z = z_j = (1 + i)j \rightarrow \infty$, we have

$$\begin{aligned} |A_0(z_j)| &= |-6e^{3z_j}| = 6e^{3j} > e^{3j} = e^{3(\sqrt{2}/2)|z_j|}, \\ |A_1(z_j)| &= |2 + 6e^{z_j} + 11e^{2z_j}| \leq 19e^{2j} < e^{(5/2)j} = e^{5(\sqrt{2}/4)|z_j|} \end{aligned}$$

and

$$|A_2(z_j)| = |-(3 + 6e^{z_j})| \leq 9e^j < e^{(5/2)j} = e^{5(\sqrt{2}/4)|z_j|}.$$

Hence the conditions (2.1) and (2.2) of Theorem 2.1 are verified. The three linearly independent functions $f_1(z) = e^{e^z}$, $f_2(z) = e^{2e^z}$ and $f_3(z) = e^{3e^z}$ are solutions of (2.10) with $\rho(f_1) = \rho(f_2) = \rho(f_3) = +\infty$.

We now give a generalization of Example 2.

Example 3. Consider the differential equation

$$(2.11) \quad f^{(n)} + P_{n-1}(e^z)f^{(n-1)} + \cdots + P_1(e^z)f' + \beta e^{\alpha z}f = 0,$$

where $\alpha \in \mathbf{R}$ with $\alpha > 0$, $\beta \in \mathbf{C}^*$, and $P_1, \dots, P_{n-1}$ are polynomials such that $\max_{1 \leq k \leq n-1} \deg(P_k) < \alpha$. It follows from Theorem 2.1 that every solution $f \not\equiv 0$ of (2.11) has infinite order.

Proof of Corollary 2.2. Let $\max_{1 \leq k \leq n} \{\rho(A_k)\} = b < \rho(A_0) = a$. Then for a given ε , $0 < \varepsilon < (a - b)/2$, we have

$$(2.12) \quad |A_k(z)| < e^{|z|^{b+\varepsilon}} \quad (k = 1, \dots, n-1)$$

and

$$(2.13) \quad |A_0(z)| > e^{|z|^{a-\varepsilon}}$$

for sufficiently large $|z|$. Then by (2.13), there exists a number $\alpha > 1$ such that

$$(2.14) \quad |A_0(z)| > e^{\alpha|z|^{b+\varepsilon}}$$

for sufficiently large $|z|$. By making use of (2.12), (2.14) and Theorem 2.1, we get our result.

Proof of Corollary 2.3. As in the proof of Corollary 2.2, we obtain immediately Corollary 2.3.

Acknowledgement. The authors would like to thank the referee for his/her helpful remarks and suggestions.

REFERENCES

- [1] B. BELAÏDI AND S. HAMOUDA, Orders of solutions of an n -th order linear differential equation with entire coefficients, *Electron. J. Differential Equations*, 2001 (2001), No. 61, 5 pp.
- [2] Z.-X. CHEN AND S.-A. GAO, The complex oscillation theory of certain non-homogeneous linear differential equations with transcendental entire coefficients, *J. Math. Anal. Appl.*, **179** (1993), 403–416.
- [3] G. G. GUNDERSEN, Finite order solutions of second order linear differential equations, *Trans. Amer. Math. Soc.*, **305** (1988), 415–429.
- [4] G. G. GUNDERSEN, Estimates for the logarithmic derivative of a meromorphic function, plus similar estimates, *J. London Math. Soc.* (2), **37** (1988), 88–104.
- [5] G. G. GUNDERSEN, E. M. STEINBART AND S. WANG, The possible orders of solutions of linear differential equations with polynomial coefficients, *Trans. Amer. Math. Soc.*, **350** (1998), 1225–1247.
- [6] W. K. HAYMAN, *Meromorphic Functions*, Oxford Mathematical Monographs, Clarendon Press, Oxford, 1964.
- [7] G. JANK UND L. VOLKMANN, *Einführung in die Theorie der ganzen und meromorphen Funktionen mit Anwendungen auf Differentialgleichungen*, UTB für Wissenschaft: Grosse Reihe, Birkhäuser Verlag, Basel, 1985.

- [8] G. VALIRON, Lectures on the General Theory of Integral Functions, translated by E. F. Collingwood, Chelsea Publishing, New York, 1949.
- [9] H. WITTICH, Neuere Untersuchungen über eindeutige analytische Funktionen, Zweite, korrigierte Auflage, Ergebnisse der Mathematik und ihrer Grenzgebiete 8, Springer-Verlag, Berlin, 1968.

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MOSTAGANEM
B. P 227 MOSTAGANEM-ALGERIA
e-mail: belaidi.benharrat@caramail.com
belaidi@univ-mosta.dz

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF MOSTAGANEM
B. P 227 MOSTAGANEM-ALGERIA
e-mail: Hamouda.saada@caramail.com