

## ON EMBEDDABLE 1-CONVEX SPACES

VIOREL VĂJĂITU

(Received May 31, 1999)

### 1. Introduction

Throughout this paper all complex spaces are assumed to be reduced and with countable topology.

Let  $X$  be a complex space.  $X$  is said to be *embeddable* if it can be realized as a complex analytic subset of  $\mathbb{C}^m \times \mathbb{P}^n$  for some positive integers  $m$  and  $n$ . For instance, one checks that a complex curve of bounded Zariski dimension is embeddable.

We say that  $X$  is *1-convex* if  $X$  is a modification at finitely many points of a Stein space  $Y$ , *i.e.*, there exist a compact analytic set  $S \subset X$  without isolated points and a proper holomorphic map  $\pi : X \rightarrow Y$  such that  $\pi_*(\mathcal{O}_X) = \mathcal{O}_Y$  and  $\pi$  induces an isomorphism between  $X \setminus S$  and  $Y \setminus \pi(S)$ .  $S$  is called the *exceptional set* of  $X$  and  $Y$  the *Remmert's reduction* of  $X$ . See [16] for further properties of 1-convex spaces.

A criterion of Schneider [18] says that a 1-convex space  $X$  of bounded Zariski dimension is embeddable if, and only if, there is a holomorphic line bundle  $L$  over  $X$  such that  $L|_S$  is ample.

Using this, Bănică [3] proved that a 1-convex complex surface  $X$  of bounded Zariski dimension is *embeddable* provided that  $X$  does not admit compact two dimensional irreducible components. By extending this Colţoiu ([4], [5]) showed that every connected 1-convex manifold  $X$  with 1-dimensional exceptional set is embeddable if  $\dim(X) > 3$ . This is true also for threefolds  $X$  with some exceptions when the exceptional set contains a  $\mathbb{P}^1$  ([5]).

In this short note we reconsider Colţoiu's example from another point of view. This is based on the following proposition which may be of independent interest.

**Proposition 1.** *Let  $Y \subset \mathbb{P}^n$  be a hypersurface of degree  $d$  with isolated singularities,  $\pi : M \rightarrow Y$  a resolution of singularities, and  $H \subset \mathbb{P}^n$  a hyperplane which avoids the singular locus of  $Y$  and such that  $\Gamma := H \cap Y$  is smooth. Set  $X := M \setminus \pi^{-1}(\Gamma)$ . Then for  $n \geq 4$  the following statements are equivalent:*

- (a)  $X$  is embeddable.
- (b)  $X$  is Kähler.
- (c)  $M$  is projective.

By this and an example due to Moishezon [12] (see also [6]) we obtain:

**Theorem 1.** *There exists a 1-convex threefold  $X$  with exceptional set  $\mathbb{P}^1$  such that  $X$  is not Kähler; a fortiori  $X$  is not embeddable.*

For the proof of Proposition 1 we use several short exact sequences, Bott's formula, Thom's isomorphism, and some facts on pluriharmonic functions.

Also employing recent results due to Fujiki [9] we prove (see the next section for definitions):

**Theorem 2.** *Let  $\pi : X \longrightarrow Y$  be a finite holomorphic map of complex spaces with  $X$  of bounded Zariski dimension. If  $X$  is maximal and  $Y$  is Hodge, then it holds:*

- (a)  *$Y$  compact implies  $X$  projective.*
- (b)  *$Y$  is 1-convex implies  $X$  is 1-convex and embeddable.*

REMARK 1. Note that by [23], 1-convexity is invariant under finite holomorphic surjections. However, this does not hold for embeddability.

As a consequence of Theorem 2 we improve a well-known projectivity criterion due to Grauert [10] to:

**Proposition 2.** *Let  $X$  be a compact complex space. If  $X$  is Hodge and maximal, then  $X$  is projective.*

and the embeddability result due to Th. Peternell ([17], Theorem 2.6) to:

**Proposition 3.** *Let  $X$  be a 1-convex space of bounded Zariski dimension such that  $X$  is Hodge and maximal. Then  $X$  is embeddable.*

## 2. Continuous weakly pluriharmonic functions

Let  $X$  be a complex space. As usual,  $\mathcal{P}_X$  denotes the sheaf of germs of pluriharmonic functions on  $X$ . Then the canonical map  $\mathcal{O}_X \longrightarrow \mathcal{P}_X$  given by  $f \mapsto \operatorname{Re} f$  induces a short exact sequence

$$(\star) \quad 0 \longrightarrow \mathbb{R} \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{P}_X \longrightarrow 0.$$

Consider  $\widehat{\mathcal{P}}_X :=$  the sheaf of *continuous weakly pluriharmonic functions*, i.e., for every open subset  $U$  of  $X$ ,  $\widehat{\mathcal{P}}_X(U)$  consists of those  $h \in C^0(U, \mathbb{R})$  which are pluriharmonic on  $\operatorname{Reg}(U)$ .

Clearly  $\mathcal{P}_X \subseteq \widehat{\mathcal{P}}_X$ , and if  $\widehat{\mathcal{O}}_X$  denotes the sheaf of continuous weakly holomorphic functions, we have a natural map  $\widehat{\mathcal{O}}_X \longrightarrow \widehat{\mathcal{P}}_X$  given by  $f \mapsto \operatorname{Re} f$ .

Here we prove:

**Proposition 4.** *The canonical short sequence*

$$0 \longrightarrow \mathbb{R} \longrightarrow \widehat{\mathcal{O}}_X \longrightarrow \widehat{\mathcal{P}}_X \longrightarrow 0,$$

*is exact.*

**Proof.** We check only the surjectivity of  $\widehat{\mathcal{O}}_X \longrightarrow \widehat{\mathcal{P}}_X$ . We do this in two steps.

**STEP 1.** Suppose  $X$  is normal. Let  $\pi : M \longrightarrow X$  be a resolution of singularities. Then  $\pi_* \mathcal{P}_M = \mathcal{P}_X$  by Proposition 2.1 in [9]. Now, since on a complex manifold a continuous real-valued function  $\varphi$  is pluriharmonic if and only if  $\varphi$  and  $-\varphi$  are plurisubharmonic we obtain that  $\widehat{\mathcal{P}}_X = \mathcal{P}_X$ , whence the desired surjectivity in view of  $(\star)$ .

**STEP 2.** The general case. Let  $\nu : Y \longrightarrow X$  be the normalization of  $X$ . Let  $x_o \in X$ ,  $U$  an open neighborhood of  $x_o$ , and  $h \in \widehat{\mathcal{P}}_X(U)$ . Then, by Step 1.,  $h \circ \nu \in \mathcal{P}_Y(\nu^{-1}(U))$ . By Proposition 2.3 in [9] after shrinking  $U \ni x_o$ , there is  $f \in \widehat{\mathcal{O}}_X(U)$  such that  $\operatorname{Re} f = h$ . Note that in *loc. cit.* this is done under the additional hypothesis  $h \in C^\infty(U, \mathbb{R})$ . But our case follows *mutatis mutandis*, whence the proposition.  $\square$

Recall ([7], pp. 122–126) that a complex space  $Z$  is said to be *maximal* if  $\mathcal{O}_Z = \widehat{\mathcal{O}}_Z$  and that every complex space  $X$  admits a *maximalization*  $\widehat{X}$ , i.e.,  $\widehat{X}$  is maximal and there is a holomorphic homeomorphism  $\pi : \widehat{X} \longrightarrow X$  which induces a biholomorphic map between  $\widehat{X} \setminus \pi^{-1}(M(X))$  and  $X \setminus M(X)$ , where  $M(X)$  is the non-maximal locus of  $X$ , i.e.,  $M(X) = \{x \in X; \mathcal{O}_{X,x} \neq \widehat{\mathcal{O}}_{X,x}\}$ . Clearly every normal complex space is maximal. For this reason, maximal complex spaces are also called “weakly normal”.

**Corollary 1.** *If  $X$  is maximal, then  $\mathcal{P}_X = \widehat{\mathcal{P}}_X$ .*

**Corollary 2.** *If  $X$  is normal, then every pluriharmonic function  $h$  on  $\operatorname{Reg}(X)$  extends uniquely to a pluriharmonic function on  $X$ .*

**Proof.** Since  $h$  and  $-h$  extend uniquely to plurisubharmonic functions  $\varphi$  and  $\psi$  on  $X$ , we get  $\varphi = -\psi$ . Hence  $\varphi$  is continuous, whence  $\varphi$  is pluriharmonic by Corollary 1.  $\square$

By a *d-closed, real (1, 1)-form* (in the sense of Grauert [10]) on a complex space  $X$  we mean, a *d-closed, real (1, 1)-form*  $\omega$  on  $\operatorname{Reg}(X)$  such that every point  $x \in X$  admits an open neighborhood  $U$  on which there is  $\varphi \in C^2(U, \mathbb{R})$  with  $\omega = i\partial\bar{\partial}\varphi$  on  $\operatorname{Reg}(U)$ . This  $\varphi$  is called a *local potential function* for  $\omega$ . We say that  $\omega$  is *Kähler* if the local potentials may be chosen strongly plurisubharmonic.

Alternatively, by Moishezon [14] we define a *d-closed, real (1, 1)-form* on  $X$  as a collection  $\{(U_j, \varphi_j)\}_{j \in J}$  where  $\{U_j\}_j$  is an open covering of  $X$  and  $\varphi_j \in C^2(U_j, \mathbb{R})$  are such that  $\varphi_j - \varphi_k$  is pluriharmonic. Two such collections  $\{(U_j, \varphi_j)\}_{j \in J}$  and  $\{(V_k, \psi_k)\}_{k \in K}$  define the same form if  $\varphi_j - \psi_k$  is pluriharmonic on  $U_j \cap V_k$  for all

indices  $j$  and  $k$ .

**Corollary 3.** *For a maximal complex space  $X$  the above two notions of  $d$ -closed, real  $(1, 1)$ -forms coincide in an obvious sense.*

Proof. This is immediate by Corollary 1.  $\square$

To every  $d$ -closed, real  $(1, 1)$ -form  $\omega$  on  $X$  we associate canonically an element of  $H^1(X, \widehat{\mathcal{P}}_X)$ , which in turn goes into its *de Rham* class  $[\omega] \in H^2(X, \mathbb{R})$  via the cohomology sequence from Proposition 4.

We say that  $\omega$  is *integral* if its *de Rham* class belongs to  $\text{Im}(H^2(X, \mathbb{Z}) \rightarrow H^2(X, \mathbb{R}))$ .

One has the following (see [10], proof of Satz 3)

**Lemma 1.** *If  $\omega$  is an integral form on a maximal space  $X$ , then there is a holomorphic line bundle  $L \rightarrow X$  together with a class  $C^2$ -hermitean metric on  $L$  whose Chern form is  $\omega$ . In particular, if  $\omega$  is Kähler, then  $L$  is positive.*

Let  $X$  be a complex space.  $X$  is said to be *Kähler* if  $X$  has a Kähler form (in the sense of Grauert). We say that  $X$  is *Hodge* if it admits a Kähler form which is integral.

**Proposition 5.** *Let  $\pi : Y \rightarrow X$  be a finite holomorphic map of complex spaces such that  $X$  is Hodge. Then  $Y$  is Hodge. In particular, the maximalization  $\tilde{X}$  and the normalization  $X^*$  of  $X$  are Hodge, too.*

Proof. Let  $\{(U_j, \psi_j)\}_j$ ,  $U_j \subseteq X$ , defines a Kähler form  $\omega$  on  $X$ . Let  $V_j \subseteq U_j$  such that  $\{V_j\}_j$  is also a covering of  $X$ . Then by [22] for every  $\delta \in C^0(X, \mathbb{R})$ ,  $\delta > 0$ , there exists  $\psi \in C^\infty(Y, \mathbb{R})$ ,  $0 < \psi < \delta$ , such that  $\sigma_j := \psi_j \circ \pi + \psi$  are strongly plurisubharmonic on  $W_j := \pi^{-1}(V_j)$  for all  $j$ ; hence  $\{(W_j, \sigma_j)\}_j$  defines a Kähler form  $\pi^*\omega$  on  $Y$ . Of course  $\pi^*\omega$  depends on  $\delta$  and  $\psi$ , but this is irrelevant for our discussion. Moreover, in view of a canonical commutative diagram and Proposition 4, if  $\omega$  is integral, then  $\pi^*\omega$  is integral too.  $\square$

Now Lemma 1 and the criteria of Grauert [10] and Schneider [18] give Theorem 2.

**REMARK 2.** There is a compact, normal, two dimensional complex space  $X$  with only one singularity such that  $\text{Reg}(X)$  is Kähler, and  $X$  is *not* Kähler. (This follows from [14] and [10].)

### 3. Proof of proposition 1

The only nontrivial implication is (b)  $\Rightarrow$  (c) which we now consider. First we state:

CLAIM. The restriction map  $H^1(M, \mathcal{P}_M) \longrightarrow H^1(X, \mathcal{P}_M)$  is surjective.

The proof of this will be done in several steps.

STEP 1. *For every abelian group  $G$  we have  $H^1(\Gamma, G) = 0$ .*

Indeed, by a theorem of Siu [19], as  $Y \setminus \Gamma$  is a Stein subspace of  $\mathbb{P}^n \setminus \Gamma$ , it admits a Stein open neighborhood  $D$ ; thus  $\mathbb{P}^n \setminus \Gamma = D \cup (\mathbb{P}^n \setminus Y)$  is a union of two Stein open subsets. On the other hand, if an  $n$ -dimensional complex manifold  $\Omega$  is a union of  $q$  Stein open subsets, then  $H_c^i(\Omega, G) = 0$  for  $i \leq n - q$ . The assertion follows easily.

STEP 2.  $H^2(Y, \mathcal{O}_Y) = 0$ .

For this, we let  $\mathcal{I}_Y$  be the coherent ideal sheaf of  $Y$  in  $\mathbb{P}^n$ . Then  $\mathcal{I}_Y \simeq \mathcal{O}(-[Y])$ , where  $[Y]$  denotes the canonical line bundle associated to the divisor defined by  $Y$ .

Now Bott's formula gives the vanishing of  $H^i(\mathbb{P}^n, \mathcal{O}(k))$  for integers  $i, k$  with  $1 \leq i < n$ , and by the long exact cohomology sequence associated to the short exact sequence  $0 \longrightarrow \mathcal{I}_Y \longrightarrow \mathcal{O}_{\mathbb{P}^n} \longrightarrow \mathcal{O}_Y \longrightarrow 0$ , the assertion of Step 2 results immediately.

STEP 3. *The maps  $H^1(M, \mathcal{O}) \longrightarrow H^1(X, \mathcal{O})$  and  $H^2(M, \mathcal{O}) \longrightarrow H^2(X, \mathcal{O})$  are surjective and injective respectively.*

Let  $V$  be an arbitrary open neighborhood of  $\Gamma$  in  $Y$ . Since  $Y \setminus \Gamma$  is Stein, the Mayer-Vietoris sequence for  $Y = (Y \setminus \Gamma) \cup V$  and Step 2 give that the maps  $H^1(V, \mathcal{O}) \longrightarrow H^1(V \setminus \Gamma, \mathcal{O})$  and  $H^2(V, \mathcal{O}) \longrightarrow H^2(V \setminus \Gamma, \mathcal{O})$  are surjective and injective respectively.

Assume now  $V \subset \text{Reg}(Y)$ ; hence  $\pi^{-1}(V)$  is biholomorphic to  $V$  via  $\pi$ . This and the above discussion plus the Mayer-Vietoris sequence for  $M = X \cup \pi^{-1}(V)$  completes the proof of Step 3.

STEP 4.  $H^2(M, G) \longrightarrow H^2(X, G)$  is surjective for every abelian group  $G$ .

We view  $\Gamma$  as a smooth complex hypersurface in  $M$ . The inclusion  $X \subset M$  gives rise to an exact cohomology sequence (coefficients in any abelian group  $G$ )

$$\cdots \longrightarrow H^i(M, X; G) \longrightarrow H^i(M; G) \longrightarrow H^i(X; G) \longrightarrow H^{i+1}(M, X; G) \longrightarrow \cdots$$

On the other hand since  $\Gamma$  is a non-singular complex hypersurface, a tubular neighborhood of  $\Gamma$  is diffeomorphic to a neighborhood of the 0-section of the normal bundle of  $\Gamma$  in  $M$ . This bundle being holomorphic is naturally oriented. We thus have, see [2], a Thom isomorphism:

$$H^i(M, X; G) \cong H^{i-2}(\Gamma; G),$$

whence the assertion of Step 4 using Step 1.

(•) The proof of the claim follows by diagram chasing using Steps 3 and 4 and

the next commutative diagram with exact rows:

$$\begin{array}{ccccccc} H^1(M, \mathcal{O}) & \longrightarrow & H^1(M, \mathcal{P}) & \longrightarrow & H^2(M, \mathbb{R}) & \longrightarrow & H^2(M, \mathcal{O}) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H^1(X, \mathcal{O}) & \longrightarrow & H^1(X, \mathcal{P}) & \longrightarrow & H^2(X, \mathbb{R}) & \longrightarrow & H^2(X, \mathcal{O}), \end{array}$$

(•) For the proof of the proposition we let  $\mathcal{K}_M^{1,1}$  be the sheaf of germs of real smooth  $(1, 1)$ -forms on  $M$  which are  $d$ -closed. As usual,  $\mathcal{E}_M$  represents the sheaf of germs of smooth real functions on  $M$ . The short exact sequence on  $M$ ,

$$0 \longrightarrow \mathcal{P}_M \longrightarrow \mathcal{E}_M \longrightarrow \mathcal{K}_M^{1,1} \longrightarrow 0,$$

where the last non trivial map is given by  $\varphi \mapsto \sqrt{-1}\partial\bar{\partial}\varphi$ , induces a commutative diagram with exact rows:

$$\begin{array}{ccccccc} H^0(M, \mathcal{E}_M) & \longrightarrow & H^0(M, \mathcal{K}_M^{1,1}) & \longrightarrow & H^1(M, \mathcal{P}_M) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H^0(X, \mathcal{E}_M) & \longrightarrow & H^0(X, \mathcal{K}_M^{1,1}) & \longrightarrow & H^1(X, \mathcal{P}_M) & \longrightarrow & 0. \end{array}$$

By diagram chasing and the above claim if  $\omega$  is the Kähler form of  $X$ , then there are: a smooth,  $d$ -closed, real  $(1, 1)$ -form  $\alpha$  on  $M$  and a smooth real-valued function  $\varphi$  on  $X$  such that

$$(\P) \quad \alpha|_X - \omega = \sqrt{-1}\partial\bar{\partial}\varphi.$$

Now, select  $\chi \in C^\infty(X, \mathbb{R})$  which vanishes on a neighborhood  $\Omega$  of  $\pi^{-1}(\text{Sing}(Y))$  and equals 1 outside a compact subset of  $X$ . By  $(\P)$ , the smooth  $(1, 1)$ -form  $\omega + \sqrt{-1}\partial\bar{\partial}(\chi\varphi)$  on  $X$  extends trivially to a smooth, real, and  $d$ -closed  $(1, 1)$ -form  $\widehat{\omega}$  on  $M$ .

Let  $\beta$  be the canonical Kähler form on  $\mathbb{P}^n$ . For every  $c > 0$  define a  $d$ -closed  $(1, 1)$ -form  $\widetilde{\omega}_c$  on  $M$  by setting:

$$\widetilde{\omega}_c := \widehat{\omega} + c\pi^*(\beta).$$

Clearly  $\widetilde{\omega}_c$  restricted to  $\Omega$  is positive definite for every  $c > 0$ . On the other hand, there is  $c > 0$  sufficiently large such that  $\widetilde{\omega}_c$  is positive definite near the compact set  $M \setminus \Omega$ . Thus  $M$  is Kähler. Since  $M$  is Moishezon, by [13]  $M$  is projective.  $\square$

REMARK 3. In [20] a similar version to our Proposition 1, without any smoothness assumption on  $H \cap Y$  and with the additional assumption that  $H^2(X, \mathcal{O}_X) = 0$ , is stated.

Unfortunately, the “given proof” is wrong. See Colţoiu’s pertinent comments [5] for this and many, many other fatal errors, which, to our unpleasant surprise, are used again in [21].

#### 4. Proof of theorem 1

Let  $Y \subset \mathbb{P}^4$  be a hypersurface of degree  $d > 2$  having a nondegenerate quadratic point  $y_o$  as its only singularity [12]. Let  $\sigma : V \longrightarrow \mathbb{P}^4$  be the quadratic transform with center  $y_o$ . Set  $\Sigma := \sigma^{-1}(y_o)$ ,  $W :=$  the proper transform of  $Y$  ( $W$  is a nonsingular hypersurface in  $V$ ), and  $T := \Sigma \cap W \simeq \mathbb{P}^1 \times \mathbb{P}^1$ . Let  $S$  be one of the two factors and  $\rho : T \longrightarrow S$  the corresponding projection.

If  $N$  denotes the normal bundle of  $T$  in  $W$ , the restriction of  $N$  to each of the fibres of  $\rho$  is the negative of the hyperplane bundle, so the criterion of Nakano and Fujiki applies ([8], [15]).

In other words  $W$  is obtained by blowing-up a non singular  $M$  along a rational non singular curve  $S$ . One obtains easily a holomorphic map  $\pi : M \longrightarrow Y$  which resolve the singularity  $y_o$  of  $Y$  and  $S = \pi^{-1}(y_o) \simeq \mathbb{P}^1$ .

On the other hand, by [6],  $M$  is not Kähler if  $d > 2$ . Therefore, if we choose a linear hyperplane  $H$  in  $\mathbb{P}^4$ ,  $H \not\ni y_o$ , such that  $H \cap Y$  is smooth, then by Proposition 1,  $X := M \setminus \pi^{-1}(Y \cap H)$  is the desired example.  $\square$

REMARK 4. As a counterexample for embeddability this example is due to Colţoiu [5] where by a different method he obtained that  $H^1(X, \mathcal{O}_X) = 0$  under the additional hypothesis that  $H$  intersects  $Y$  transversally.

Here we emphasize the non-Kähler property of the example.

---

#### References

- [1] A. Andreotti and H. Grauert: *Théorèmes de finitude pour la cohomologie des espaces complexes*, Bull. Soc. Math. France, **90** (1962), 193–259.
- [2] A. Andreotti and F. Norguet: *Cycles of algebraic manifolds and  $\partial\bar{\partial}$ -cohomology*, Annali Scuola Norm. Sup. Pisa, **25** (1971), 59–114.
- [3] C. Bănică: *Sur les fibres infinitésimales d’un morphisme propre d’espaces complexes*, Sémin. Norquet 1977/79, Lecture Notes in Math., **807** (1980), 122–144.
- [4] M. Colţoiu: *On the embedding of 1-convex manifolds with 1-dimensional exceptional set*, Comment. Math. Helv., **60** (1985), 458–465.
- [5] M. Colţoiu: *On 1-convex manifolds with 1-dimensional exceptional set*, Rev. roumaine Math. pures et Appl., **43** (1998), 97–104.
- [6] M. Cornalba: *Two theorems on modifications of analytic spaces*, Invent. Math., **20** (1973), 227–247.
- [7] G. Fischer: *Complex analytic geometry*, Lecture Notes in Math., 538, Springer-Verlag, 1976.
- [8] A. Fujiki and S. Nakano: *Supplement to “On the inverse of the monoidal transformation”*, Publ. Res. Inst. Math. Sci., **7** (1971–72), 637–644.

- [9] A. Fujiki: *Closures of orbits of  $\mathbb{C}$  and  $\mathbb{C}^*$  actions*, Osaka J. Math., **33** (1996), 365–385.
- [10] H. Grauert: *Über Modifikationen und exzeptionelle analytische Mengen*, Math. Ann., **146** (1962), 331–368.
- [11] H. Grauert and R. Remmert: *Plurisubharmonische Funktionen in komplexen Räumen*, Math. Z., **65** (1956), 175–194.
- [12] B.G. Moishezon: *Irreducible exceptional submanifolds of the first kind of three-dimensional complex analytic manifolds*, Soviet. Math. Dokl., **6** (1965), 402–403.
- [13] B.G. Moishezon: *On the  $n$ -dimensional compact complex varieties with  $n$  algebraically independent meromorphic functions III*, Amer. Math. Soc. Transl., **63** (1967), 51–177.
- [14] B.G. Moishezon: *On singular kählerian spaces*, Proceedings of the International conference on manifolds and relates topics in topology, (Tokyo 1973), 343–351; Tokyo, Univ. of Tokyo Press 1974.
- [15] S. Nakano: *On the inverse of the monoidal transformation*, Publ. Res. Inst. Math. Sci., **6** (1970–71), 483–502.
- [16] R. Narasimhan: *The Levi problem for complex spaces II*, Math. Ann., **146** (1962), 195–216.
- [17] Th. Peternell: *On strongly pseudoconvex Kähler manifolds*, Invent. math., **70** (1982), 157–168.
- [18] M. Schneider: *Familien negativer Vektorraumbündel und 1-konvexe Abbildungen*, Math. Seminar Univ. Hamburg, **47–48** (1978–79), 150–170.
- [19] Y.-T. Siu: *Every Stein subvariety admits a Stein neighborhood*, Invent. math., **38** (1976), 89–100.
- [20] Vo Van. Tan: *On certain non-Kählerian strongly pseudoconvex manifolds*, The Journal of Geom. Anal., **4** (1994), 233–245.
- [21] Vo Van. Tan: *On the Kählerian geometry of 1-convex threefolds*, Forum. Math., **7** (1995), 131–146.
- [22] V. Văjăitu: *Kählerianity of  $q$ -Stein spaces*, Arch. Math., **66** (1996), 250–257.
- [23] V. Văjăitu: *Invariance of cohomological  $q$ -completeness under finite holomorphic surjections*, Manuscr. math., **82** (1994), 113–124.
- [24] S.T. Yau: *Sheaf cohomology on 1-convex manifolds*, Recent developments in Several Complex Variables, Ann. of Math. Stud., **100** (1981), 429–452.

Institute of Mathematics of the Romanian Academy  
P.O. Box 1–764, RO–70700 Bucharest, Romania  
e-mail: vvajaitu@stoilow.imar.ro