

DECOMPOSITION OF REGULAR REPRESENTATIONS FOR $U(H)_\infty$

DOUG PICKRELL

Let G denote the infinite dimensional group consisting of all unitary operators which are compact perturbations of the identity (on a fixed separable Hilbert space). Kirillov showed that G has a discrete spectrum (as a compact group does). The point of this paper is to show that there are analogues of the Peter-Weyl theorem and Frobenius reciprocity for G . For the left regular representation, the only reasonable candidate for Haar measure is a Gaussian measure. The corresponding L^2 decomposition is analogous to that for a compact group. If X is a flag homogeneous space for G , then there is a unique invariant probability measure on (a completion of) X . Frobenius reciprocity holds, for our surrogate Haar measure fibers over X precisely as in finite dimensions (this is the key observation of the paper). When X is a symmetric space, each irreducible summand contains a unique invariant direction, and this direction is the L^2 limit of the corresponding (L^2 normalized) finite dimensional spherical functions.

1. Introduction. Let H be a separable complex Hilbert space, $U(H)_\infty = \{g \in U(H): g = 1 + \text{compact operator}\}$. This group is a basic example of an infinite dimensional Banach Lie group. Kirillov proved that this group is type 1 and has a discrete spectrum ([4], [6]).

Fix an orthonormal basis e_1, e_2, \dots for H . Then $U(H)_\infty$ is the closure in the operator norm topology of $U(\infty) = \bigcup_n U(n)$, where $U(n) \cong \{g \in U(H): ge_j = e_j, j > n\}$.

Relative to this basis, view $U(H) \rightarrow M$, where M is the space of matrices $(E_{ij})_{1 \leq i, j < \infty}$, and which we identify with the space of linear operators mapping H^{alg} , the algebraic span of the $\{e_j\}$, to \mathbb{C}^∞ , the space of all formal linear combinations of the $\{e_j\}$. The left action of $U(\infty)$ on $U(H)_\infty$ extends to an action of $U(\infty)$ on M .

Let ν_G denote the Gaussian measure for the linear space $\mathcal{L}_2(H)$. We recall the following facts established in [8]: (a) every ergodic invariant probability measure for the left action of $U(\infty)$ on M is a linear equivariant image of ν_G (and itself Gaussian), (b) ν_G is the weak limit of the uniform distributions on the spaces $\sqrt{n} U(n)$, and (c) up to scaling ν_G is the only $U(\infty)$ ergodic biinvariant measure on M . For these reasons it is natural to view ν_G as a kind of Haar measure for $U(H)_\infty$, relative to its

left regular action. In this paper we will exploit the existence of this Haar type measure to decompose various regular representations of $U(H)_\infty$.

Of course the first step is to decompose the representation

$$U(H)_\infty \rightarrow U(L^2(M, d\nu_G)).$$

This is done in §2, and the decomposition is analogous to the Peter-Weyl decomposition for a compact group.

In §3 we use the Peter-Weyl decomposition to decompose the regular representations for $U(H)_\infty$ on homogeneous spaces (flag manifolds) ((3.2)). The key idea can be described in terms of the simplest example. Via the basis above view $H \cong l^2 \rightarrow \mathbf{C}^\infty$. The natural projection $\pi: \mathbf{C}^\infty \setminus \{0\} \rightarrow \mathbf{P}(\mathbf{C}^\infty)$ is $U(\infty)$ equivariant, and it pushes the Gaussian measure for H to the unique $U(\infty)$ invariant probability measure on $\mathbf{P}(\mathbf{C}^\infty)$. Now it is frequently said that Gaussian measure behaves as a uniform distribution on a sphere of infinite radius. In particular we should expect

$$(1.1) \quad L^2(\mathbf{P}(\mathbf{C}^\infty)) \cong L^2(\mathbf{C}^\infty)^{U(1)},$$

where the right hand side denotes those functions invariant under the scalar action of $U(1)$. This is correct. The key ((3.8)) is to fiber the Gaussian over the invariant measure on projective space; the fiber is the Haar measure for the unitary stabilizer (in general), in this case $U(1)$. The right hand side of (1.1) is easy to understand because of the Peter-Weyl decomposition, and this leads to Frobenius reciprocity.

In §4 we consider the special case of a symmetric space, i.e. a Grassmann manifold $\text{Gr}(n, H)$. In this case the decomposition is multiplicity free. Each irreducible component contains a unique invariant direction for the isotropy group, and this direction is the L^2 limit of the corresponding (L^2 normalized) finite dimensional spherical functions.

I thank Robert Boyer for several valuable conversations.

Notation. $dm(\cdot)$ denotes Lebesgue measure, $\mathcal{P}(\cdot)$ the polynomial algebra. If π_i is a representation for G_i , then $\pi_1 \times \pi_2$ is the (outer) tensor product representation for $G_1 \times G_2$. If $G_1 = G_2$, $\pi_1 \otimes \pi_2$ is the usual tensor product representation for $G_1 = G_2$.

2. Peter-Weyl theorem. In this section it will be convenient to view ν_G as a cylinder measure (i.e. weak distribution) on $\mathcal{L}_2(H)$ (see [5] or [9]). A function on $\mathcal{L}_2(H)$ of the form $\phi(E) = \Phi(P(E))$, where P is an orthogonal projection of rank $n < \infty$ and Φ is a bounded Borel function, will be called tame; we let \mathcal{V} denote the algebra of all tame functions. If

we set

$$E(\phi) = \int \phi d\nu_G = \int_{\mathcal{R}(P)} \Phi(x) \pi^{-n} e^{-|x|^2} dm(x)$$

where ϕ is as above, then (\mathcal{V}, E) is an integration algebra. There is a natural representation of $O(\mathcal{L}_2(H))$ as automorphisms of (\mathcal{V}, E) , hence a unitary representation on $L^2(\nu_G)$, the completion of \mathcal{V} in the norm $E(\bar{\phi}\phi)$.

We view $U(H) \times U(H) \subset O(\mathcal{L}_2(H))$ by $g \times h \cdot E = g \circ E \circ h^{-1}$. Our goal in this section is to decompose the action of $U(H) \times U(H)$ (and $U(H)_\infty \times U(H)_\infty$) on $L^2(\nu_G)$. Of course there is a natural $U(\infty) \times U(\infty)$ equivariant isomorphism of $L^2(\nu_G)$, as constructed above, and $L^2(M, \nu_G)$, when we view ν_G as a probability measure on M .

Let \mathcal{T} denote the transform defined by

$$(\mathcal{T}\phi)(w) = \int \phi(L) e^{iRe(w, E)} d\nu_G(E)$$

for $\phi \in \mathcal{V}$ and $w \in \mathcal{L}_2(H)^*$. By the corollary of Theorem 6.4 of [2], \mathcal{T} extends to a $U(H) \times U(H)$ equivariant isomorphism

$$(2.1) \quad L^2(\nu_G) \cong \mathbb{C} e^{-1/4|w|^2} \otimes \sum_{j=0}^{\infty} \hat{\mathcal{P}}^j \otimes \sum_{k=0}^{\infty} \bar{\hat{\mathcal{P}}}^k$$

where $\hat{\mathcal{P}}^j$ is $(j!)^{1/2}$ times the completion of $\mathcal{P}^j(\mathcal{L}_2(H))$ in the norm it inherits from the tensor algebra.

Suppose λ is a partition, i.e. a decreasing sequence of integers $\lambda_1 \geq \lambda_2 \geq \dots$ such that $\lambda_j = 0$ for all sufficiently large j . If $\lambda_{n+1} = 0$, then we denote by $\rho_{\lambda, n}$ the representation of $U(\mathbb{C}^n)$ with signature $(\lambda_1 \geq \dots \geq \lambda_n)$, by ρ_λ the direct limit of the $\rho_{\lambda, n}$, which extends canonically from $U(\infty)$ to a representation of $U(H)$.

In the proof of (3.1) of [8] it is shown that as a representation of $U(H)_\infty \times U(H)_\infty$ (or $U(H) \times U(H)$)

$$(2.2) \quad \hat{\mathcal{P}}^j(\mathcal{L}_2(H)) = \sum \rho_\lambda^* \times \rho_\lambda$$

where the sum is over all partitions λ with $\sum \lambda_i = j$. This proves the following

(2.3) PROPOSITION. *As a representation of $U(H) \times U(H)$ or $U(H)_\infty \times U(H)_\infty$,*

$$L^2(\nu_G) = \sum \rho_\lambda^* \otimes \rho_\mu \times \rho_\lambda \otimes \rho_\mu^*$$

where the sum is over all partitions λ, μ .

By Kirillov's classification ([4]) of the irreducible representations of $U(H)_\infty$, this decomposition is analogous to the Peter-Weyl theorem for compact groups.

The physical space for $\rho_\lambda^* \times \rho_\lambda$, as a subrepresentation of (2.2), consists of matrix coefficients for ρ_λ (ρ_λ is a subrepresentation of the action of $U(H)$ on the tensor algebra of H , so this action extends naturally to an action of $GL(H)$; the matrix coefficients restrict to polynomials on $\mathcal{L}_2(H)$). Hence the physical space for $\rho_\lambda^* \otimes \rho_\mu \times \rho_\lambda \otimes \rho_\mu^*$, as a subrepresentation of the right hand side of (2.1), consists of matrix coefficients for the action of $GL(H)$ on $\mathcal{L}_2(H_\lambda, H_\mu)$ given by $g: T \rightarrow \rho_\mu(g) \circ T \circ \rho_\lambda(g)^*$ (where ρ_λ is realized on H_λ).

To describe the corresponding subspace in $L^2(\nu_G)$, one must invert the transform \mathcal{T} . Whether this can be done in a reasonably explicit manner in general, I do not know. In the case of spherical functions, there does exist a relatively simple inversion formula (see §4, especially (4.6)).

3. Frobenius reciprocity. In this section we fix a finite set of integers $0 < n_1 < n_2 < \cdots < n_l < \infty$. Let $\text{Flag}(H) \subset \text{Gr}(n_1, H) \times \cdots \times \text{Gr}(n_l, H)$ denote the set of points (flags) $\{W_i\}$ such that $W_1 \subset W_2 \subset \cdots \subset W_l$, where $\text{Gr}(n_i, H)$ denotes the set of all n_i dimensional subspaces of H . $\text{Flag}(H)$ is a homogeneous space for $U(H)$ and $U(H)_\infty$. We let $\text{Flag}(\mathbb{C}^N)$ and $\text{Flag}(\mathbb{C}^\infty)$ denote the analogous objects for $\mathbb{C}^N \cong \text{span}\{e_j: j \leq N\}$ ($N > n_l$) and $\mathbb{C}^\infty \cong \{\text{formal linear combinations of } e_j\}$. The action of $U(\infty)$ extends from $\text{Flag}(H)$ to $\text{Flag}(\mathbb{C}^\infty)$, and there are natural embeddings $GL(N) \rightarrow GL(H)$, $\text{Flag}(\mathbb{C}^N) \rightarrow \text{Flag}(\mathbb{C}^\infty)$.

Our first task is to recall why there is a unique $U(\infty)$ invariant probability measure on $\text{Flag}(\mathbb{C}^\infty)$. In the process we will develop notation which we will employ in the remainder of the paper.

A generic flag (i.e. a point in the largest cell) of $\text{Flag}(\mathbb{C}^\infty)$ can be characterized in two ways: (a) it is of the form $\{W_j\} = \{L\mathbb{C}^{n_j}\}$, where L is a lower triangular block matrix with identity matrices on the diagonal, the block sizes being $n_1, n_2 - n_1, \dots, n_l - n_{l-1}$ along the top, $n_1, \dots, n_l - n_{l-1}, \infty$ along the side; (b) each W_j is of the form $\text{graph}(z_j)$, where $z_j \in \mathcal{L}(\mathbb{C}^{n_j}, \mathbb{C}^\infty \ominus \mathbb{C}^{n_j})$. The operator L and the set $\{z_j\}$ determine one another via the relations

$$\gamma_j \alpha_j^{-1} = z_j \quad \text{where } L = \begin{pmatrix} \alpha_j & * \\ \gamma_j & * \end{pmatrix}$$

with respect to the splittings of the domain $= \mathbb{C}^{n_j} \oplus (\mathbb{C}^{n_l} \ominus \mathbb{C}^{n_j})$ and range $= \mathbb{C}^{n_j} \oplus (\mathbb{C}^\infty \ominus \mathbb{C}^{n_j})$.

Let $L^{(N)}$ denote the projection of L to $\mathcal{L}(\mathbf{C}^{n_l}, \mathbf{C}^N)$ ($L^{(N)} = Q \circ L$, where $Q: \mathbf{C}^\infty \rightarrow \mathbf{C}^N$ is the obvious projection) (similarly for z). The diagram

$$(3.1) \quad \begin{array}{ccc} L & \rightarrow & L^{(N)} \\ \downarrow & & \downarrow \\ \{z_j\} & \rightarrow & \{z_j^{(N)}\} \end{array}$$

is commutative (the cutoff is on the left, whereas the α 's act from the right). In §4 of [8] it is shown how $U(N)$ equivariance of the map $Z_j \rightarrow Z_j^{(N)}$ implies uniqueness for the $U(\infty)$ invariant probability measure on $\text{Gr}(n_j, \mathbf{C}^\infty)$. The above diagram shows the same argument applies to flags.

Conversely, the projection

$$\pi: \mathcal{L}(\mathbf{C}^{n_l}, \mathbf{C}^\infty)' \rightarrow \text{Flag}(\mathbf{C}^\infty): E \rightarrow \{E(\mathbf{C}^{n_l})\},$$

where the prime indicates we exclude those E which are singular, is $U(\infty)$ equivariant. Thus the Gaussian measure associated to the linear space $\mathcal{L}(\mathbf{C}^{n_l}, H)$ will be mapped by π to a $U(\infty)$ invariant probability measure on $\text{Flag}(\mathbf{C}^\infty)$. This proves existence.

Let μ_0 denote the unique invariant measure on $\text{Flag}(\mathbf{C}^\infty)$. Our task is to decompose $L^2(\text{Flag}(\mathbf{C}^\infty))$.

Let $K_l = \times_1^l U(\mathbf{C}^{n_j} \ominus \mathbf{C}^{n_{j-1}})$ and $K = K_l \times U(H \ominus \mathbf{C}^{n_l})$. Let $P: M \rightarrow \mathcal{L}(\mathbf{C}^{n_l}, \mathbf{C}^\infty)$ denote the obvious projection, and $\nu = P_* \nu_G$. In this section we will ultimately prove the following

(3.2) PROPOSITION. *The pullbacks*

$$L^2(\text{Flag}(\mathbf{C}^\infty)) \xrightarrow{\pi^*} L^2(\mathcal{L}(\mathbf{C}^{n_l}, \mathbf{C}^\infty), \nu)^{K_l} \xrightarrow{P_*} L^2(M, \nu_G)^K$$

are isomorphisms, where the superscripts indicate the sets of vectors invariant under the right action of K_l and K , respectively. As a representation of $U(H)_\infty$

$$L^2(\text{Flag}(\mathbf{C}^\infty)) = \sum m(\lambda, \mu) \rho_\lambda \otimes \rho_\mu^*$$

where the sum is over those partitions with $\lambda_{n_l+1} = \mu_{n_l+1} = 0$ and

$$m(\lambda, \mu) = \dim n \left((\rho_\lambda^* \otimes \rho_\mu)^K \right) = \dim n \left((\rho_{\lambda, n_l}^* \otimes \rho_{\mu, n_l})^{K_l} \right).$$

The proof of the analogue of this proposition for a compact group is trivial, because of the existence of Haar measure. Our proof will be trivial as well, once we understand how ν is fibered over μ_0 (see (3.8) below). We will prove (3.2) at the end of this section.

We will need the following computational lemma.

For $E \in \mathcal{L}(\mathbb{C}^r, \mathbb{C}^\infty)$, recall that $E^{(N)}$ is the projection of E to $\mathcal{L}(\mathbb{C}^r, \mathbb{C}^N)$.

(3.3) LEMMA. *The scalar $(\det N^{-1}E^{(N)} * E^{(N)})^{-1}$ converges to 1 and the $r \times r$ matrices $N^{-1}E^{(N)} * E^{(N)}$ and their inverses converge to the identity in $L^p(\mathcal{L}(\mathbb{C}^r, \mathbb{C}^\infty), d\nu)$ as $N \rightarrow \infty$, for all $1 \leq p < \infty$.*

Proof. First consider $N^{-1}E^{(N)} * E^{(N)}$. For the diagonal entries

$$\begin{aligned} & \int |N^{-1}(E^{(N)} * E^{(N)})_{jj} - 1|^p d\nu(E) \\ &= \int_{\mathbb{C}^N} |N^{-1}|x|^2 - 1|^p \pi^{-N} e^{-|x|^2} dm(x) \\ &= \sum_0^p \binom{p}{k} (-1)^{p-k} N^{-k} \int_0^\infty s^{N+k+1} e^{-s} ds / \int_0^\infty s^{N-1} e^{-s} ds \\ &= \sum_0^p \binom{p}{k} (-1)^{p-k} N^{-k} (N)_k \end{aligned}$$

and this tends to 0.

For off diagonal entries

$$\begin{aligned} & \int |N^{-1}(E^{(N)} * E^{(N)})_{ij}|^{2p} d\nu(E) \\ &= \int_{\mathbb{C}^N + \mathbb{C}^N} \left| N^{-1} \sum_1^N x_j \bar{y}_j \right|^{2p} \pi^{2N} e^{-|x|^2 - |y|^2} dm(x) dm(y) \\ &= N^{-2p} \int_{\substack{1 \leq i_k, j_k \leq N \\ 1 \leq k \leq p}} \sum_{l=1}^p x_{i_l} \bar{x}_{j_l} y_{i_l} \bar{y}_{j_l} \\ &= N^{-2p} \int_{\substack{1 \leq i_k \leq N \\ 1 \leq k \leq p}} \left(\int_{\mathbb{C}} |z|^2 \pi^{-1} e^{-|z|^2} dm(z) \right)^2 = N^{-2p} N^p \end{aligned}$$

which tends to zero.

We now consider

$$(3.4) \quad \int |(\det N^{-1}E^{(N)} * E^{(N)})^{-1} - 1|^{2p} d\nu(E).$$

We use the integral formula

$$\begin{aligned} & \int \phi(E^{(N)}) d\nu(E) \\ &= c \int_{(\mathbf{R}^+)^r} \left\{ \int \phi(k_1 \lambda k_2) dk_1 dk_2 \right\} \prod_{i < j} |\lambda_i^2 - \lambda_j^2|^2 \prod_1^r 2\lambda_j \lambda_j^{2(N-r)} e^{-\lambda_j^2} d\lambda_j \end{aligned}$$

where $\lambda = \text{diag}(\lambda_1, \dots, \lambda_r)$, dk_1 and dk_2 denote the unitarily invariant probability measures on $\text{Isom}(\mathbf{C}^r, \mathbf{C}^N)$ and $U(\mathbf{C}^r)$, respectively, and c is a normalization constant (see Chapter I of [1]). We now see (3.4) equals

$$\begin{aligned} & c \int \left| \prod_1^r N u_j^{-1} - 1 \right|^{2p} \prod_{i < j} (u_i - u_j)^2 \prod_1^r u_j^{N-r} e^{-u_j} du_j \\ &= \sum_0^{2p} \binom{2p}{k} (-1)^{2p-k} c N^{rk} \int \prod_{i < j} (u_i - u_j)^2 \prod_1^r u_j^{N-r-k} e^{-u_j} du_j. \end{aligned}$$

Let $s = N - r - k$. The k th integral equals

$$\begin{aligned} & c N^{rk} \int \det^2(\mathcal{L}_i^{(s)}(u_j)) \prod_1^r u_j^s e^{-u_j} du_j \\ &= c N^{rk} \det \left(\int \mathcal{L}_i^{(s)}(u) \mathcal{L}_j^{(s)}(u) u^s e^{-u} du \right) \\ &= N^{rk} \prod_1^r \left(\int \mathcal{L}_i^{(s)2} u^s e^{-u} du / \int \mathcal{L}_i^{(N-r)2} u^{N-r} e^{-u} du \right) \end{aligned}$$

where the $\mathcal{L}_i^{(s)}$ are the Laguerre polynomials. The lemma now follows from

$$\int |\mathcal{L}_i^{(s)}|^2 u^s e^{-u} du = \Gamma(s+1) \binom{s+i}{i}$$

(see Chapter 5 of [10]). □

(3.5) LEMMA. For a generic flag $\{W_j\} = \{L\mathbf{C}^{n_j}\}$ in $\text{Flag}(\mathbf{C}^\infty)$, let $g^{(N)}(L)$ be the isometry from \mathbf{C}^{n_l} to \mathbf{C}^N obtained by applying the (block) Gram-Schmidt orthonormalization process to $L^{(N)}$. Then entry by entry $N^{1/2}g_N(L)$ has a limit in probability $g(L)(\in \mathcal{L}(\mathbf{C}^{n_l}, \mathbf{C}^\infty))$. In the case $l = 1$, we actually have $L^p(\mu_0)$ convergence, for each $1 \leq p < \infty$.

REMARK. It is almost certainly the case that the limit above is $L^p(\mu_0)$ in general. However, for $l > 1$ this seems to complicate the proof immensely. The reason is essentially that the function $\{W_j\} \rightarrow L$, which is well-defined a.e. $[\mu_0]$, does not have integrable entries. It would be desirable to establish L^p convergence, because this would yield a second

proof of (3.8) below (see the remark following the proof of (3.8)). The meaning of the convergence when $l = 1$ is explored in the next section.

Proof of (3.5). Let $E \in \mathcal{L}(\mathbf{C}^{n_l}, \mathbf{C}^\infty)$, $E \cdot \mathbf{C}^{n_l} = W_j$, so that $E = LU$ where U is (block) upper triangular. Note $E^{(N)} = L^{(N)}U$. Write $E = [E_1, \dots, E_l]$, where the E_j are the columns (similarly for L , etc.).

Let $\alpha = \alpha_1(E)$. Then, as a function of E ,

$$g_1^{(N)} = L_1^{(N)} |L_1^{(N)}|^{-1} = E_1^{(N)} \alpha^{-1} \left(\alpha (E_1^{(N)*} E_1^{(N)})^{-1} \alpha^* \right)^{1/2}.$$

The entries of E_1 are in all $L^p(\nu)$, and

$$\text{tr} \left| \alpha^{-1} \left(\alpha (E_1^{(N)*} E_1^{(N)})^{-1} \alpha^* \right)^{1/2} \right|^2 = \text{tr} (E_1^{(N)*} E_1^{(N)})^{-1}.$$

By (3.3) we have L^p convergence

$$N^{1/2} g_1^{(N)} \rightarrow L_1 (\alpha_1(E) \alpha_1(E)^*)^{1/2}$$

entry by entry as $N \rightarrow \infty$ (note the existence of the limit shows the RHS is equal to a function of L , a.e. $[\nu]$).

Now suppose we have established that $g_i^{(N)}$ has a limit g_i in probability for $1 \leq i \leq j$. We have

$$(3.6) \quad g_j^{(N)} = \left(1 - \sum_{i < j} g_i^{(N)} g_i^{(N)*} \right) L_j^{(N)} \left| \left(1 - \sum_{i < j} g_i^{(N)} g_i^{(N)*} \right) L_j^{(N)} \right|^{-1}$$

(here 1 is the $N \times N$ identity matrix).

Consider the $N \times (n_j - n_{j-1})$ matrix $g_i^{(N)} g_i^{(N)*} L_j^{(N)}$. The Hilbert-Schmidt norm is dominated by

$$(3.7) \quad \left(\text{tr} (g_i^{(N)*} g_i^{(N)})^2 \right)^{1/2} \left(\text{tr} L_j^{(N)*} L_j^{(N)} \right)^{1/2}.$$

By induction the first factor is $O(N^{-1})$ in probability. On the other hand $L_j^{(N)*} L_j^{(N)}$ is the (j, j) (block) entry of $U^{-1*} E^{(N)*} E^{(N)} U^{-1}$, which is $O(N)$ by (3.3). Therefore (3.7) is $O(N^{-1/2})$ in probability. So we certainly have $(1 - \sum_{i < j} g_i^{(N)} g_i^{(N)*}) L_j^{(N)} \rightarrow L_j$ in probability, entry by entry.

Now let $\phi_N = N^{-1/2} L_j^{(N)}$, $\psi_N = -N^{-1/2} \sum_{i < j} g_i^{(N)} g_i^{(N)*} L_j^{(N)}$. We know that $\phi_N^* \phi_N \rightarrow ((UU^*)^{-1})_{jj}$ and $\psi_N^* \psi_N \rightarrow 0$ in probability. The generalized Holder inequality

$$\text{tr} |\psi_N^* \phi_N| \leq \left(\text{tr} |\psi_N^*|^2 \right)^{1/2} \left(\text{tr} |\phi_N|^2 \right)^{1/2}$$

shows that $\psi_N^* \phi_N$ and $\phi_N^* \psi_N$ tend to zero in probability as well. This implies that $|\phi_N + \psi_N|^2 \rightarrow ((UU^*)^{-1})_{jj}$, which is strictly positive. This implies

$$N^{1/2} \left| \left(1 - \sum_{i < j} g_i^{(N)} g_i^{(N)*} \right) L_j^{(N)} \right|^{-1} = |\phi_N + \psi_N|^{-1}$$

has a limit in probability. Hence (3.6), scaled by $N^{1/2}$, has a limit in probability, entry by entry. This completes the induction. \square

(3.8) PROPOSITION. *The decomposition of the Gaussian measure ν with respect to the projection $\pi: \mathcal{L}(\mathbf{C}^{n_l}, \mathbf{C}^\infty)' \rightarrow \text{Flag}(\mathbf{C}^\infty)$ is given by*

$$\int \phi d\nu = \int_{\text{Flag}(\mathbf{C}^\infty)} \int_{K_l} \phi(g(w)k) dk d\mu_0(w).$$

Proof. Assume ϕ is a bounded continuous function based on $\mathcal{L}(\mathbf{C}^{n_l}, \mathbf{C}^m)$. For $N > m$, we have

$$\begin{aligned} & \int_{\text{Isom}(\mathbf{C}^{n_l}, \mathbf{C}^N)} \phi(N^{1/2}E) d\omega_N(E) \\ &= \int_{\text{Flag}(\mathbf{C}^N)} \int_{K_l} \phi(N^{1/2}g_N(W)k) dk d\mu_{0,N}(W) \\ &= \int_{\text{Flag}(\mathbf{C}^\infty)} \int_{K_l} \phi(N^{1/2}g_N(W)k) dk d\mu_0(W) \end{aligned}$$

where $\omega_N(\mu_{0,N})$ denotes the unique invariant probability measure for $U(N)$. Take the limit as $N \rightarrow \infty$. By (2.1) of [8] the LHS converges to the LHS of (3.8). By (3.5) the RHS converges to the RHS of (3.8). This proves (3.8). \square

(3.9) REMARK. It is possible to give a more direct, but formal, argument for (3.8) as follows.

First, via direct calculation, we fiber the Gaussian on $\mathcal{L}(\mathbf{C}^{n_l}, \mathbf{C}^N)$ over $\mu_{0,N}$ on $\text{Flag}(\mathbf{C}^N)$ (we let $L = L^{(N)}$).

$$(3.10) \quad \int_{\mathcal{L}(\mathbf{C}^{n_l}, \mathbf{C}^N)} \phi d\nu(E) = c \int \phi(LU) e^{-\text{tr}|LU|^2} dm(LU).$$

Now $dm(LU) = \prod_1^l (\det|U_{jj}|^2)^{N-n_j} dm(L) dm(U)$. To separate the L and U variables in the exponential in (3.10), we (block) orthonormalize L , which amounts to multiplying L on the right by a (block) upper triangular matrix, and then we change the U variable:

$$U = L^{-1}g_N(L)V, \quad U_{jj} = (L^{-1}g_N(L))_{jj}V_{jj},$$

$$dm(U) = \prod_1^l \left(\det |(L^{-1}g_N(L))_{jj}|^2 \right)^{n_l - n_{j-1}} dm(V) \quad (n_0 = 0).$$

This implies (3.10) equals

$$\int_{\text{Flag}(\mathbb{C}^N)} \int \phi(g_N(W)V) dv_{0,N}(V) d\mu_{0,N}(W)$$

where

$$dv_{0,N}(V) = c \prod_1^l \left(\det |V_{jj}|^2 \right)^{N-n_j} e^{-\text{tr } V^* V} dm(V)$$

is a probability measure on the (block) upper triangular matrices. The formula (3.8) then formally follows from the fact that

- (i) $ce^{-\text{tr } N|V_{ij}|^2} dm(N^{1/2}V_{ij}) \rightarrow \delta_0$ as $N \rightarrow \infty$ for $1 \leq i < j \leq l$, and
- (ii) $c(\det N|V_{jj}|^2)^{N-n_j} e^{-\text{tr } N|V_{jj}|^2} dm(N^{1/2}V_{jj}) \rightarrow dk_j$, the Haar invariant probability measure on $U(\mathbb{C}^{n_j} \ominus \mathbb{C}^{n_{j-1}})$, which can be verified using the integral formulae in the proof of (3.3).

Proof of (3.2). We first consider P^* . We have $L^2(v_G) = L^2(v)yxL^2(v^\perp)$, where $v^\perp = (1 - P)_*v$. Thus

$$L^2(v_G)^{U(H \ominus \mathbb{C}^{n_l})} = L^2(v) \otimes L^2(v^\perp)^{U(H \ominus \mathbb{C}^{n_l})} = L^2(v).$$

This shows P^* induces an isomorphism

The fact π^* induces an isomorphism follows immediately from (3.8).

(2.2) implies the claims about the multiplicity. \square

4. Symmetric space. In this section we consider the special case of a Grassmannian, $\text{Gr}(n, \mathbb{C}^\infty)$. Recall that if z is the graph coordinate, the map

$$(4.1) \quad \text{Gr}(n, \mathbb{C}^\infty) \rightarrow \text{Gr}(n, \mathbb{C}^N): z \rightarrow z^{(N)},$$

which is defined almost everywhere, is $U(\mathbb{C}^N)$ equivariant. This is equivalent to saying that the pullback defines a $U(\mathbb{C}^N)$ equivariant isometric map

$$L^2(\text{Gr}(n, \mathbb{C}^N)) \rightarrow L^2(\text{Gr}(n, \mathbb{C}^\infty)).$$

We want to study how the decomposition for $\text{Gr}(n, \mathbb{C}^N)$ converges to that for $\text{Gr}(n, \mathbb{C}^\infty)$.

Because the irreducible summands of $L^2(\text{Gr}(n, \mathbb{C}^N))$ consist of algebraic functions and the projection

$$\text{Gr}(n, \mathbb{C}^{N+k}) \rightarrow \text{Gr}(n, \mathbb{C}^N)$$

defined by (4.1) is not globally continuous, it is not the case that the irreducible summands coherently embed as $N \rightarrow \infty$. Thus the convergence is somewhat subtle. It is most easily understood in terms of spherical functions.

Now $\text{Gr}(n, H) = U/K$, where $U = U(H)$, $K = U(\mathbb{C}^n) \times U(\mathbb{C}^{n^\perp})$. It is a symmetric space of rank n . The Cartan involution is given by $\theta(x) = \begin{pmatrix} \alpha & \\ & \delta \end{pmatrix} - \begin{pmatrix} \gamma & \beta \\ & \end{pmatrix}$, where $x = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ relative to $H = \mathbb{C}^n \oplus (\mathbb{C}^n)^\perp$, $x \in \mathfrak{gl}(H)$. Let \square denote the set of all operators of the form $T = \sum_1^n t_j (e_j \otimes e_{n+j}^* + e_{n+j} \otimes e_j^*)$ with $t_j \in \mathbb{R}$. This set is a maximal abelian subalgebra of $P = \{ \begin{pmatrix} x^* & \\ & x \end{pmatrix} : x \in \mathcal{L}(\mathbb{C}^{n^\perp}, \mathbb{C}^n) \}$, the real “noncompact” part of the $\theta = -1$ eigenspace.

Suppose $T \in \square$. Then

$$\exp(it) = \sum_1^n \left(\cos t_j (e_j \otimes e_j^* + e_{n+j} \otimes e_{n+j}^*) \right. \\ \left. + i \sin t_j (e_j \otimes e_{n+j}^* + e_{n+j} \otimes e_j^*) \right)$$

plus the identity on $\{e_j : 1 \leq j \leq 2n\}^\perp$. Hence generically we have $\exp(it) = \text{graph}(z)$, where $z = \sum_1^n i \tan t_j e_j \otimes e_j^*$. Note the spectrum of $(1 + z^*z)^{-1}$ is $\{u_j\}$, where $u_j = \cos^2 t_j$.

We now recall the formulae of Berezin-Karpelevic for the spherical functions (these are proven by Hoogenboom in [3]). Let $N \geq 2n$.

(4.2) LEMMA. *The spherical functions of $\text{Gr}(n, \mathbb{C}^N)$ are parameterized by partitions μ with $\mu_{n+1} = 0$. The function corresponding to μ is a multiple of the function*

$$\psi(z) = \frac{\det \{ L_{i-1+\bar{\mu}_i}^{(k)}(u_j) \}}{\det \{ u_j^{i-1} \}}$$

where u_1, \dots, u_n is the spectrum of $(1 + z^*z)^{-1}$, the $L_i^{(k)}$ are the (Legendre) orthogonal polynomials for the probability measure $(k+1)(1-x)^k dx$ on $[0, 1]$, $k = N - 2n$, and $\bar{\mu}_j = \mu_{n+1-j}$.

Using integration in polar coordinates (see Chapter I of [1]), it is easily checked that the L^2 normalized spherical function corresponding to the partition μ is given by

$$\psi_{\mu, N}(z) = \frac{\det \{ \tilde{L}_{i-1+\bar{\mu}_i}^{(k)}(u_j) \}}{\det \{ \tilde{L}_{i-1}^{(k)}(u_j) \}},$$

where \tilde{L}_i denotes the L^2 normalization of L_i .

(4.3) PROPOSITION. *For each partition μ with $\mu_{n+1} = 0$, the functions $\psi_{\mu, N}$ have a limit ψ_μ in all $L^p(\mu_0)$, $1 \leq p < \infty$. As a function of $E \in \mathcal{L}(\mathbb{C}^n, \mathbb{C}^\infty)$,*

$$\psi_\mu(E) = \frac{\det \{ \tilde{\mathcal{L}}_{i-1+\bar{\mu}_i}(v_j) \}}{\det \{ \tilde{\mathcal{L}}_{i-1}(v_j) \}}$$

where $\{v_i\}$ is the spectrum of $\alpha(E)^*\alpha(E)$ and the $\tilde{\mathcal{L}}_i$ are the L^2 normalized Laguerre polynomials.

Proof. $\{\tilde{L}_i^{(k)}(k^{-1}y)\}$ is the system of L^2 normalized orthogonal polynomials for the probability measure $(k+1)/k(1-y/k)^k dy$ on $0 \leq y \leq k$, which tends to $e^{-y} dy$ as k (or N) $\rightarrow \infty$. From this it follows easily (or one can check the well-known formulae directly) that the coefficients of $\tilde{L}_i^{(k)}(k^{-1}y)$ converge to those of $\tilde{\mathcal{L}}_i(y)$ as $k \rightarrow \infty$.

We also know that $k(1 + Z^{(N)*}Z^{(N)})^{-1} (\cong N|L_1^{(N)}|^{-2}$ in the notation of (3.5)) converges to the $n \times n$ matrix $(\alpha\alpha^*)(E)$ in all L^p . Hence we have L^p convergence

$$\sigma(ku_1^{(N)}, \dots, ku_n^{(N)}) \rightarrow \sigma(v_1, \dots, v_n)$$

for any symmetric polynomial. This implies

$$\psi_{\mu,N} = \frac{\det\{\tilde{L}_{i-1+\bar{\mu}_i}^{(k)}(ku_i^{(N)}/k)\}}{\det\{\tilde{L}_{i-1}^{(k)}(ku_i^{(N)}/k)\}} \rightarrow \psi_{\mu}$$

in all L^p , $1 \leq p < \infty$. □

By (3.2) we know that

$$\begin{aligned} L^2(\text{Gr}(n, \mathbb{C}^\infty)) &\cong \sum \rho_\lambda^* \otimes \rho_\mu \times (\rho_\lambda \otimes \rho_\mu^*)^K \\ &= \sum \rho_\lambda^* \otimes \rho_\mu \times (\rho_{\lambda,n} \otimes \rho_{\mu,n}^*)^{U(n)} \cong \sum \rho_\mu^* \otimes \rho_\mu, \end{aligned}$$

where the second and third sums are over those partitions with $(n+1)$ th term = 0.

(4.4) PROPOSITION. For each partition μ with $\mu_{n+1} = 0$, ψ_μ is, up to a multiple, the unique K invariant vector in $\rho_\mu^* \otimes \rho_\mu$.

Proof. We will use the transform \mathcal{T} of §2, which induces an equivariant isometry

$$L^2(\mathcal{L}(\mathbb{C}^n, \mathbb{C}^\infty)) \cong \mathbb{C}e^{-1/4|w|^2} \otimes \sum_0^\infty \hat{\mathcal{P}}^j \otimes \sum_0^\infty \bar{\hat{\mathcal{P}}}^k$$

where $\hat{\mathcal{P}}^j = (j!)^{1/2} \mathcal{P}^j(\mathcal{L}(\mathbb{C}^n, H))$. We must show that $\mathcal{T}\psi_\mu$ is in the one dimensional space

$$\begin{aligned} (4.5) \quad \mathbb{C}e^{-1/4|w|^2} \otimes (\rho_\mu^* \otimes \rho_\mu \times \rho_{\mu,n} \otimes \rho_{\mu,n}^*)^{K \times U(n)} \\ = \mathbb{C}e^{-1/4|w|^2} \otimes (\rho_{\mu,n}^* \times \rho_{\mu,n} \otimes \rho_{\mu,n} \times \rho_{\mu,n}^*)^{U(n) \times U(n)} \end{aligned}$$

Let $\rho = \rho_{\lambda,n}$. One vector in the space $\rho^* \times \rho$ is the character $\chi = \text{trace } \rho$ (where we have holomorphically extended ρ to a representation of $GL(n)$). Thus $|\chi|^2$ is a vector in $\rho^* \times \rho \otimes \rho \times \rho^*$. Let $\{\varepsilon_i\}$ be an orthonormal basis for a realization V of ρ . Any hermitian form on V is a multiple of a fixed $U(n)$ invariant positive form $\langle \cdot, \cdot \rangle$. A standard argument shows that this implies

$$\int_{U(n)} \langle v_1, \rho(k)^* \varepsilon_i \rangle \langle v_2, \rho(k)^* \varepsilon_j \rangle dk = \delta(i-j) \dim(V)^{-1} \langle v_1, v_2 \rangle$$

for $v_1, v_2 \in V$. Thus

$$\begin{aligned} \int_{U(n)} |\chi|^2(kw) dk &= \sum_{i,j} \int \langle \rho(w) \varepsilon_i, \rho(k)^* \varepsilon_j \rangle \langle \rho(w) \varepsilon_i, \rho(k)^* \varepsilon_j \rangle dk \\ &= \sum_j \dim(V)^{-1} \langle \rho(w^*w) \varepsilon_j, \varepsilon_j \rangle. \end{aligned}$$

Thus, by the Weyl character formula, a nonzero vector spanning the space in (4.5) is

$$\begin{aligned} \phi_\lambda(w) &= \exp\left(-\frac{1}{4}|w|^2\right) \text{tr } \rho(w^*w) \\ &= \exp\left(-\frac{1}{4}|w|^2\right) \frac{\det(v_j^{i-1+\mu_i})}{\det(v_j^{i-1})}, \end{aligned}$$

where $\{v_j\}$ is the spectrum of w^*w .

Any K invariant vector in the range of \mathcal{T} must be a linear combination of the ϕ_λ , in particular, $\psi_\mu = \sum c_\lambda \psi_\lambda$ (where a priori we only know $\lambda_{n+1} = 0$). Now asymptotically,

$$\text{tr } \rho_{\lambda,n}(\text{diag}(t_j)) \sim \langle \rho_{\lambda,n}(\text{diag}(t_j)) v_0, v_0 \rangle = \prod_1^n t_j^{\lambda_j}$$

where v_0 is a highest weight vector, we set $t = t_1 = \cdots = t_k$, $t_{k+j} = 1$, and we let $t \rightarrow \infty$. This is also the asymptotic behavior of ψ_μ (with $\mu = \lambda$ above). The theory of homogeneous chaos (Section 6.3 of [2]) shows that if f is a polynomial of degree (p, q) (in E, \bar{E}), then $\exp(\frac{1}{4}|w|^2) \mathcal{T}f$ is of the same degree. Since ψ_μ is a symmetric function in the eigenvalues of $\alpha(E)^* \alpha(E)$, it follows that $\mathcal{T}\psi_\mu$ has the same asymptotic behavior above as ϕ_μ . Thus we must have

$$(4.6) \quad \mathcal{T}\psi_\mu = c\phi_\mu$$

which proves (4.4). \square

REFERENCES

- [1] S. Helgason, *Groups and Geometric Analysis*, Academic Press, 1984.
- [2] T. Hida, *Brownian Motion*, Springer-Verlag, 1980.
- [3] B. Hoogenboom, *Spherical functions and differential operators on complex Grassmann manifolds*, Ark. Nat., **20** (1982), 69–85.
- [4] A. A. Kirillov, *Representations of an infinite dimensional unitary group*, Soviet Math. Dokl., **14** (1973), No. 5.
- [5] H. Kuo, *Gaussian Measures in Banach Spaces*, Springer-Verlag, 1975.
- [6] G. Olshanetski, *Unitary representations of the infinite dimensional groups $U(p, \infty)$, $Sp_0(p, \infty)$, $Sp(p, \infty)$ and the corresponding motion groups*, Functional Anal. Appl., **12** (1978), 185–191.
- [7] D. Pickrell, *Measures on infinite dimensional Grassmann manifolds*, to appear in J. Functional Anal.
- [8] ———, *On $U(\infty)$ invariant measures*, submitted to Duke Math. J.
- [9] I. E. Segal, *Algebraic integration theory*, Bull. Amer. Math. Soc., **71** (1965), 419–489.
- [10] G. Szego, *Orthogonal polynomials*, AMS colloquium publications, Vol. XXIII, Fourth edition, 1978.

Received February 11, 1986. Partially supported by NSF grant DMS-8506130.

YALE UNIVERSITY
NEW HAVEN, CT 06520