

MOORE SPACES AND $w\Delta$ -SPACES

R. E. HODEL

*This paper is dedicated to Professor J. H. Roberts
on the occasion of his sixty-fifth birthday.*

This paper is a study of conditions under which a $w\Delta$ -space is a Moore space. In §2 we introduce the notion of a G_δ^* -diagonal and show that every $w\Delta$ -space with a G_δ^* -diagonal is developable. In §3 we prove that every regular θ -refinable $w\Delta$ -space with a point-countable separating open cover is a Moore space. In §4 we introduce the class of α -spaces and show that a regular $w\Delta$ -space is a Moore space if and only if it is an α -space. Finally, in §5 we study a new class of spaces which generalizes both semi-stratifiable and $w\Delta$ -spaces.

1. Preliminaries. We begin with some definitions and known results which will be used throughout this paper. Unless otherwise stated no separation axioms are assumed; however regular spaces are always T_1 and paracompact spaces are always Hausdorff. The set of natural numbers will be denoted by N .

Let X be a set, \mathcal{S} a cover of X , x an element of X . The *star of x with respect to \mathcal{S}* , denoted $\text{st}(x, \mathcal{S})$, is the union of all elements of \mathcal{S} containing x . The *order of x with respect to \mathcal{S}* , denoted $\text{ord}(x, \mathcal{S})$, is the number of elements of \mathcal{S} containing x .

A space X is *developable* if there is a sequence $\mathcal{S}_1, \mathcal{S}_2, \dots$ of open covers of X such that, for each x in X , $\{\text{st}(x, \mathcal{S}_n) : n = 1, 2, \dots\}$ is a fundamental system of neighborhoods of x . Such a sequence of open covers is called a *development* for X . A regular developable space is called a *Moore space*. Bing [1] proved that every paracompact Moore space is metrizable.

According to Borges [3] a space X is a $w\Delta$ -space if there is a sequence $\mathcal{S}_1, \mathcal{S}_2, \dots$ of open covers of X such that, for each x in X , if $x_n \in \text{st}(x, \mathcal{S}_n)$ for $n = 1, 2, \dots$ then the sequence $\langle x_n \rangle$ has a cluster point. Such a sequence of open covers is called a $w\Delta$ -sequence for X . Clearly every countably compact space is a $w\Delta$ -space, and in [3] Borges proved that every developable space and every M -space is a $w\Delta$ -space. For the relationship between $w\Delta$ -spaces, strict p -spaces, and p -spaces, see [6].

A space X is *subparacompact* if every open cover of X has a σ -discrete closed refinement. Every paracompact space is subparacompact [16], and in [8] Creede proved that every semi-stratifiable space is subparacompact. For further properties of subparacompact spaces see [5], [11], and [15].

A space X is θ -refinable if for each open cover \mathcal{V} of X there is a sequence $\mathcal{S}_1, \mathcal{S}_2, \dots$ of open refinements of \mathcal{V} such that, for each x in X , there is a n in N such that $\text{ord}(x, \mathcal{S}_n)$ is finite. Such a sequence of open covers is called a θ -refinement of \mathcal{V} . In [24] Wicke and Worrell state that every subparacompact space is θ -refinable and that a countably compact T_1 space is compact if and only if it is θ -refinable.

2. Spaces with a G_δ^* -diagonal. Recall that a space X has a G_δ -diagonal if its diagonal $\Delta = \{(x, x) : x \text{ in } X\}$ is a G_δ -subset of $X \times X$. The notion of a G_δ -diagonal plays an important role in metrization theorems; see, for example, [2], [3], [7], [14], and [22].

Every semi-stratifiable Hausdorff space has a G_δ -diagonal [8]. On the other hand the space $[0, 1] \times \{0, 1\}$ with the lexicographic order is a compact perfectly normal space which fails to have a G_δ -diagonal [14].

In [7] Ceder obtained this characterization of spaces with a G_δ -diagonal.

PROPOSITION 2.1. (Ceder) *A space X has a G_δ -diagonal if and only if there is a sequence $\mathcal{S}_1, \mathcal{S}_2, \dots$ of open covers of X such that, for any two distinct points x and y of X , there is a n in N such that $y \notin \text{st}(x, \mathcal{S}_n)$.*

In light of this characterization of a G_δ -diagonal and Borges' study of spaces with a \bar{G}_δ -diagonal (see [3]), we introduce the following definition.

DEFINITION 2.2. A space X has a G_δ^* -diagonal if there is a sequence $\mathcal{S}_1, \mathcal{S}_2, \dots$ of open covers of X such that, for any two distinct points x and y of X , there is a n in N such that $y \notin \text{st}(x, \mathcal{S}_n)^-$. Such a sequence of open covers is called a G_δ^* -sequence for X .

In [13] Kullman proved that every regular θ -refinable space with a G_δ -diagonal has a \bar{G}_δ -diagonal. Since every space with a \bar{G}_δ -diagonal has a G_δ^* -diagonal, we have the following proposition.

PROPOSITION 2.3. *Every regular θ -refinable space with a G_δ -diagonal has a G_δ^* -diagonal. In particular every regular semi-stratifiable space has a G_δ^* -diagonal.*

The next result relates the G_δ^* -diagonal property to the diagonal Δ .

PROPOSITION 2.4. *Let X be a space, let $\{V_n : n = 1, 2, \dots\}$ be a*

sequence of open subsets of $X \times X$ containing Δ , and suppose that $\bigcap_{n=1}^{\infty} \bar{V}_n = \Delta$. Then X has a G_s^* -diagonal. In particular, if X is Hausdorff and $X \times X$ is perfectly normal then X has a G_s^* -diagonal.

Proof. For $n = 1, 2, \dots$ let $\mathcal{G}_n = \{G \subseteq X: G \text{ open, } G \times G \subseteq V_n\}$. Since V_n is open and contains Δ , \mathcal{G}_n covers X . To show that $\mathcal{G}_1, \mathcal{G}_2, \dots$ is a G_s^* -sequence for X , let x and y be distinct points of X . Choose n in N such that $(x, y) \notin \bar{V}_n$, and let U and W be open neighborhoods of x and y respectively such that $(U \times W) \cap V_n = \phi$. It follows that $W \cap \text{st}(x, \mathcal{G}_n) = \phi$ and so $y \notin \text{st}(x, \mathcal{G}_n)^-$.

We now prove the main result in this section.

THEOREM 2.5. *Every $w\Delta$ -space with a G_s^* -diagonal is developable.*

Proof. Let X be a space, let $\mathcal{H}_1, \mathcal{H}_2, \dots$ be a $w\Delta$ -sequence for X , and let $\mathcal{K}_1, \mathcal{K}_2, \dots$ be a G_s^* -sequence for X . For each positive integer n let

$$\mathcal{G}_n = \left\{ G: G = \left(\bigcap_{i=1}^n H_i \right) \cap \left(\bigcap_{i=1}^n K_i \right), H_i \in \mathcal{H}_i, K_i \in \mathcal{K}_i, i = 1, \dots, n \right\}.$$

It is easy to check that \mathcal{G}_{n+1} is an open refinement of \mathcal{G}_n for all n in N and that $\mathcal{G}_1, \mathcal{G}_2, \dots$ in a $w\Delta$ -sequence and a G_s^* -sequence for X .

Suppose that $\mathcal{G}_1, \mathcal{G}_2, \dots$ is not a development for X . Then there is a point x , a neighborhood W of x , and a sequence $\langle x_n \rangle$ such that for all n , $x_n \in \text{st}(x, \mathcal{G}_n)$ and $x_n \notin W$. Since $\mathcal{G}_1, \mathcal{G}_2, \dots$ is a $w\Delta$ -sequence for X , the sequence $\langle x_n \rangle$ has a cluster point p . Clearly $p \notin W$ so $p \neq x$. Since $\mathcal{G}_1, \mathcal{G}_2, \dots$ is a G_s^* -sequence for X , there is a positive integer k and a neighborhood V of p such that $V \cap \text{st}(x, \mathcal{G}_k) = \phi$. Now for $n \geq k$, $x_n \in \text{st}(x, \mathcal{G}_n) \subseteq \text{st}(x, \mathcal{G}_k)$ and so $x_n \notin V$. This contradicts the fact that p is a cluster point of $\langle x_n \rangle$. Thus $\mathcal{G}_1, \mathcal{G}_2, \dots$ is a development for X .

COROLLARY 2.6. *The following are equivalent for a regular $w\Delta$ -space X :*

- (a) X is a Moore space.
- (b) X is semi-stratifiable.
- (c) X is θ -refinable and has a G_s -diagonal.
- (d) X has a G_s^* -diagonal.

Proof. The implication (a) \Rightarrow (b) is due to Creede [8]; (b) \Rightarrow (c) follows from results by Creede [8] and Wicke and Worrell [24]; (c) \Rightarrow (d) follows from Proposition 2.3; (d) \Rightarrow (a) follows from Theorem 2.5.

REMARK 2.7. The equivalence of (a) and (b) was first proved by Creede in [8], and the equivalence of (a) and (c) is due to Siwiec [23]. It is not known if every regular $w\Delta$ -space with a G_δ -diagonal is a Moore space. For a study of p -spaces with a G_δ -diagonal, see [13].

COROLLARY 2.8. *The following are equivalent for a regular countably compact space X :*

- (a) X is metrizable.
- (b) $X \times X \times X$ is completely normal.
- (c) $X \times X$ is perfectly normal.
- (d) X has a G_δ^* -diagonal.

Proof. Clearly (a) \Rightarrow (b); (b) \Rightarrow (c) follows from a theorem due to Katětov [12]; (c) \Rightarrow (d) follows from Proposition 2.4. To prove (d) \Rightarrow (a) observe that X is a Moore space (by Corollary 2.6) and recall that every countably compact Moore space is metrizable.

3. Separating covers. In 1938 Filippov [9] proved that every paracompact M -space with a point-countable base is metrizable. Filippov's theorem was generalized by Burke and Stoltenberg in [4], and recently Burke [6] obtained another generalization as follows.

BURKE'S THEOREM. *Every regular subparacompact $w\Delta$ -space with a point-countable base is a Moore space.*

In another direction Nagata [20] proved a metrization theorem which not only generalizes Filippov's theorem but a result by Okuyama as well [22]. In order to state Nagata's theorem succinctly we use the following terminology due to Michael [17]. A cover \mathcal{V} of a set X is said to be *separating* if given distinct points x and y of X , there is a V in \mathcal{V} such that $x \in V, y \notin V$.

NAGATA'S THEOREM. *Every paracompact M -space with a point-countable separating open cover is metrizable.*

In this section we use the techniques developed by Burke, Filippov, Nagata, and Stoltenberg, together with the results in §2, to obtain a generalization of the abovementioned theorems by Burke and Nagata.

In light of the usefulness of the concept of a θ -base in the study of developable spaces (see [24]), we begin with the following definition.

DEFINITION 3.1. A θ -separating cover of a space X is a sequence $\mathcal{G}_1, \mathcal{G}_2, \dots$ of open collections such that, for any two distinct points x and y in X , there is a n in N such that

- (a) $\text{ord}(x, \mathcal{G}_n)$ is finite;
- (b) there is a G in \mathcal{G}_n such that $x \in G$ and $y \notin G$.

The relationship between a θ -separating cover and a G_δ -diagonal is given by the following two propositions.

PROPOSITION 3.2. Let X be a space with a θ -separating cover. If every closed subset of X is a G_δ then X has a G_δ -diagonal.

Proof. Let $\mathcal{G}_1, \mathcal{G}_2, \dots$ be a θ -separating cover of X . For each pair of positive integers n and k let $\mathcal{H}_{nk} = \{H: H \neq \phi, H = \bigcap_{i=1}^k G_i, G_1, \dots, G_k \text{ distinct elements of } \mathcal{G}_n\}$ and let $F_{nk} = X - \bigcup \{H: H \in \mathcal{H}_{nk}\}$. Now F_{nk} is a closed set and so $F_{nk} = \bigcap_{j=1}^\infty W_{nkj}$, where each W_{nkj} is open. For $j = 1, 2, \dots$ let $\mathcal{K}_{nkj} = \mathcal{H}_{nk} \cup \{W_{nkj}\}$. Then each \mathcal{K}_{nkj} is an open cover of X and the sequence $\{\mathcal{K}_{nkj}: n, k, j \text{ in } N\}$ exhibits the G_δ -diagonal property for X .

PROPOSITION 3.3. Every θ -refinable space with a G_δ -diagonal has a θ -separating cover.

Proof. Let X be a θ -refinable space and let $\mathcal{G}_1, \mathcal{G}_2, \dots$ be open covers of X exhibiting the G_δ -diagonal property for X . For each n in N let $\mathcal{H}_{n1}, \mathcal{H}_{n2}, \dots$ be a θ -refinement of \mathcal{G}_n . Then

$$\{\mathcal{H}_{nk}: n = 1, 2, \dots, k = 1, 2, \dots\}$$

is a θ -separating cover of X .

The following lemmas, due to Burke and Miscenko [19], play a key role in the proof of our theorem. For the sake of completeness we sketch the proof of Burke's result. (See Remark 1.9 in [6]).

LEMMA 3.4. (Burke) Let X be a regular, θ -refinable $w\Delta$ -space. Then there is a sequence $\mathcal{G}_1, \mathcal{G}_2, \dots$ of open covers of X such that for each x in X ,

- (a) $C_x = \bigcap_{n=1}^\infty \text{st}(x, \mathcal{G}_n)$ is compact;
- (b) $\{\text{st}(x, \mathcal{G}_n): n = 1, 2, \dots\}$ is a base for C_x .

Proof. Let $\mathcal{V}_1, \mathcal{V}_2, \dots$ be a $w\Delta$ -sequence for X . By induction on n construct for each positive integer n a sequence $\mathcal{W}_{n1}, \mathcal{W}_{n2}, \dots$ of open covers of X such that

- (1) for $k = 1, 2, \dots, \{\bar{W}: W \text{ in } \mathcal{W}_{nk}\}$ refines \mathcal{V}_n and $\mathcal{W}_{ij}, 1 \leq i \leq n - 1, 1 \leq j \leq n - 1$;

(2) for each x in X there is a k in N such that $\text{ord}(x, \mathcal{W}_{nk})$ is finite.

For $n = 1, 2, \dots$ let $\mathcal{G}_n = \mathcal{W}_{n1}$. Then the sequence $\mathcal{G}_1, \mathcal{G}_2, \dots$ satisfies properties (a) and (b).

LEMMA 3.5. (Miščenko) *Let \mathcal{V} be a point-countable collection of subsets of a set X and let M be a subset of X . Then there are at most countably many finite minimal covers of M by elements of \mathcal{V} .*

We now state and prove the main result in this section.

THEOREM 3.6. *Let X be a regular, θ -refinable $w\Delta$ -space with a point-countable separating open cover. Then X is a Moore space.*

Proof. We are going to show that X has a θ -separating cover and that every closed subset of X is a G_δ . It follows by Proposition 3.2 that X has a G_δ -diagonal and hence by Corollary 2.6 X is a Moore space.

Let \mathcal{V} be a point-countable separating open cover of X . We assume that $X \in \mathcal{V}$, and hence for every subset M of X there is a finite subcollection of \mathcal{V} which covers M , namely $\{X\}$. Let $\mathcal{G}_1, \mathcal{G}_2, \dots$ be open covers of X such that for each x in X ,

- (a) $C_x = \bigcap_{n=1}^{\infty} \text{st}(x, \mathcal{G}_n)$ is compact;
- (b) $\{\text{st}(x, \mathcal{G}_n) : n = 1, 2, \dots\}$ is a base for C_x .

For each n in N let $\mathcal{H}_{n1}, \mathcal{H}_{n2}, \dots$ be a θ -refinement of \mathcal{G}_n . Recall that

- (c) \mathcal{H}_{nk} refines $\mathcal{G}_n, k = 1, 2, \dots$;
- (d) for each x in X there is a k in N such that $\text{ord}(x, \mathcal{H}_{nk})$ is finite.

X has a θ -separating cover. For each pair of positive integers n and k and for each H in \mathcal{H}_{nk} let $H(n, k, 1), H(n, k, 2), \dots$ be all finite minimal covers of H by elements of \mathcal{V} , and let

$$\mathcal{H}_{nkj} = \{H \cap V : H \in \mathcal{H}_{nk}, V \in H(n, k, j)\}.$$

To show that $\{\mathcal{H}_{nkj} : n, k, j \text{ in } N\}$ is a θ -separating cover of X , let x and y be two distinct points of X . Choose V_1 in \mathcal{V} such that $x \in V_1$ and $y \notin V_1$, and let $\{V_1, \dots, V_t\}$ be a finite cover of C_x by elements of \mathcal{V} such that $x \notin V_i$ for $i = 2, \dots, t$. Now $C_x \subseteq \bigcup_{i=1}^t V_i$ and so by (b) there is a n in N such that $\text{st}(x, \mathcal{G}_n) \subseteq \bigcup_{i=1}^t V_i$. Choose k in N such that $\text{ord}(x, \mathcal{H}_{nk})$ is finite, and let H be some element of \mathcal{H}_{nk} such that $x \in H$. Since \mathcal{H}_{nk} refines \mathcal{G}_n , $H \subseteq \text{st}(x, \mathcal{G}_n)$

and so $H \subseteq \bigcup_{i=1}^t V_i$. Choose a minimal subcollection of $\{V_1, \dots, V_t\}$ which covers H and label it $H(n, k, j)$. Note that $V_1 \in H(n, k, j)$. Thus $(H \cap V_1) \in \mathcal{H}_{nkj}$, $x \in (H \cap V_1)$, and $y \notin (H \cap V_1)$. Finally, suppose H_1, \dots, H_r are all elements of \mathcal{H}_{nk} containing x . Since $H_i(n, k, j)$ is finite for $i = 1, \dots, r$ it follows that $\text{ord}(x, \mathcal{H}_{nkj})$ is finite. This completes the proof that X has a θ -separating cover.

Every closed subset of X is a G_δ . Let M be a closed subset of X . For each pair of positive integers n and k , and for each H in \mathcal{H}_{nk} such that $H \cap M \neq \emptyset$, let $H(n, k, j), j = 1, 2, \dots$ be all finite minimal covers of $H \cap M$ by elements of \mathcal{V} . By repeatedly counting a cover if necessary, we may assume that $H(n, k, j)$ exists for all j in N . For $j = 1, 2, \dots$ let $H^*(n, k, j)$ denote the union of all elements of $H(n, k, j)$, and let $W_{nkj} = \bigcup \{H \cap (\bigcap_{i=1}^j H^*(n, k, i)) : H \in \mathcal{H}_{nk}, H \cap M \neq \emptyset\}$. Clearly each W_{nkj} is open and contains M . To complete the proof that M is a G_δ it suffices to show that if $x \notin M$ then there exist n, k , and j such that $x \notin W_{nkj}$.

First suppose that $C_x \cap M = \emptyset$. Choose n in N such that $\text{st}(x, \mathcal{G}_n) \cap M = \emptyset$, and let k and j be any positive integers. Suppose $x \in W_{nkj}$. Then there is a H in \mathcal{H}_{nk} such that $x \in H$ and $H \cap M \neq \emptyset$. Now \mathcal{H}_{nk} refines \mathcal{G}_n and so $H \subseteq \text{st}(x, \mathcal{G}_n)$. Hence $\text{st}(x, \mathcal{G}_n) \cap M \neq \emptyset$ and this contradicts the choice of n .

Next suppose that $C_x \cap M \neq \emptyset$. Let $\{V_1, \dots, V_t\}$ be a finite cover of $C_x \cap M$ by elements of \mathcal{V} such that $x \notin V_r, r = 1, \dots, t$. Choose n in N such that $\text{st}(x, \mathcal{G}_n) \subseteq (\bigcup_{r=1}^t V_r) \cup (X - M)$. Let k in N be such that $\text{ord}(x, \mathcal{H}_{nk})$ is finite and let H_1, \dots, H_s be all elements of \mathcal{H}_{nk} which contain x and intersect M . For $i = 1, \dots, s$, $H_i \subseteq \text{st}(x, \mathcal{G}_n)$ and so $H_i \cap M \subseteq \bigcup_{r=1}^t V_r$. Select from $\{V_1, \dots, V_t\}$ a minimal subcollection which covers $H_i \cap M$ and label it $H_i(n, k, j_i)$. Now $x \in H_i^*(n, k, j_i)$ and so if we take $j = \max\{j_1, \dots, j_s\}$ then $x \in W_{nkj}$.

4. α -spaces. A space with a σ -closure preserving separating closed cover is called a σ^\sharp -space. This definition was introduced by Nagata and Siwiec in [21].

PROPOSITION 4.1. *Every subparacompact space with a G_δ -diagonal is a σ^\sharp -space.*

Proof. Let X be a subparacompact space and let $\mathcal{G}_1, \mathcal{G}_2, \dots$ be open covers of X exhibiting the G_δ -diagonal property for X . For each n in N let $\mathcal{F}_{n1}, \mathcal{F}_{n2}, \dots$ be a σ -discrete closed refinement of \mathcal{G}_n . Then $\{\mathcal{F}_{nk} : n = 1, 2, \dots, k = 1, 2, \dots\}$ is a σ -closure preserving

separating closed cover of X .

In [6] Burke showed that a regular $w\Delta$ -space is a Moore space if and only if it is a σ^* -space. His method of proof suggests introducing a new class of spaces which we call α -spaces. We shall show that σ^* -spaces are α -spaces and that a regular $w\Delta$ -space is a Moore space if and only if it is an α -space.

DEFINITION 4.2. A space X is an α -space if there is a function g from $N \times X$ into the topology of X such that for each x in X ,

- (a) $\bigcap_{n=1}^{\infty} g(n, x) = \{x\}$;
- (b) if $y \in g(n, x)$ then $g(n, y) \subseteq g(n, x)$.

Such a function is called an α -function for X .

PROPOSITION 4.3. *Every σ^* -space is an α -space.*

Proof. Let $\mathcal{F}_1, \mathcal{F}_2, \dots$ be a σ -closure preserving separating closed cover of a σ^* -space X . For n in N and x in X let

$$g(n, x) = X - \bigcup \{F \in \mathcal{F}_n : x \notin F\}.$$

It is easy to check that the function g is an α -function for X .

PROPOSITION 4.4. *Every space with a σ -point finite separating open cover is an α -space. In particular, every T_1 space with a σ -point finite base is an α -space.*

Proof. Let $\mathcal{G}_1, \mathcal{G}_2, \dots$ be a σ -point finite separating open cover of a space X . We may assume that $X \in \mathcal{G}_n$ for all n in N . For $n = 1, 2, \dots$ and x in X let $g(n, x) = \bigcap \{G \text{ in } \mathcal{G}_n : x \text{ in } G\}$. Then the function g is an α -function for X .

The following characterization of semi-stratifiable spaces will be useful in proving the main theorem in this section.

LEMMA 4.5. *The following are equivalent for a space X :*

- (a) X is semi-stratifiable.
- (b) There is a function g from $N \times X$ into the topology of X such that (1) for each x in X , $\bigcap_{n=1}^{\infty} g(n, x) = \{x\}$; (2) if $x \in g(n, x_n)$ for $n = 1, 2, \dots$ then the sequence $\langle x_n \rangle$ converges to x .
- (c) There is a function g from $N \times X$ into the topology of X such that (1) for each x in X and n in N , $x \in g(n, x)$; (2) if $x \in g(n, x_n)$ for $n = 1, 2, \dots$ then x is a cluster point of the sequence $\langle x_n \rangle$.

Proof. The equivalence of (a) and (b) is due to Creede [8], and

(b) \Rightarrow (c) is obvious. To complete the proof we show that (c) \Rightarrow (b). Thus, let g be a function satisfying (c), and assume that $g(n+1, x) \subseteq g(n, x)$ for all n in N and x in X .

To prove (1) of (b), first let $y \in \bigcap_{n=1}^{\infty} g(n, x)$. Then by (2) of (c), y is a cluster point of the sequence $\{x, x, \dots\}$ and so $y \in \{x\}^-$. Next let $y \in \{x\}^-$. Then $x \in g(n, y)$ for $n = 1, 2, \dots$ so by (2) of (c) it follows that x is a cluster point of the sequence $\{y, y, \dots\}$. Thus $y \in g(n, x)$ for $n = 1, 2, \dots$ and so $y \in \bigcap_{n=1}^{\infty} g(n, x)$.

To prove (2) of (b), let $x \in g(n, x_n)$, $n = 1, 2, \dots$ and suppose that the sequence $\langle x_n \rangle$ does not converge to x . Then there is a neighborhood W of x and a subsequence $\langle x_{n_k} \rangle$ of $\langle x_n \rangle$ such that $x_{n_k} \notin W$ for all k in N . Now $x \in g(n_k, x_{n_k}) \subseteq g(k, x_{n_k})$ for $k = 1, 2, \dots$ so by (2) of (c), x is a cluster point of the sequence $\langle x_{n_k} \rangle$. But this is impossible, and so we conclude that $\langle x_n \rangle$ converges to x .

THEOREM 4.6. *A regular $w\Delta$ -space is a Moore space if and only if it is an α -space.*

Proof. By Propositions 4.1 and 4.3 every Moore space is an α -space. To complete the proof let X be a regular $w\Delta$ -space which is also an α -space and let us show that X is a Moore space. By Corollary 2.6 it suffices to show that X is semi-stratifiable.

Let $\mathcal{S}_1, \mathcal{S}_2, \dots$ be a $w\Delta$ -sequence for X , let g be an α -function for X . We may assume that for x in X and n in N , $g(n+1, x) \subseteq g(n, x)$. For x in X and $n = 1, 2, \dots$ let $h(n, x) = g(n, x) \cap \text{st}(x, \mathcal{S}_n)$. We shall show that the function h satisfies (c) of Lemma 4.5.

Clearly (1) of (c) is satisfied. To check (2) let $x \in h(n, x_n)$ for $n = 1, 2, \dots$. Then for $n = 1, 2, \dots$, $x \in \text{st}(x_n, \mathcal{S}_n)$ and so $x_n \in \text{st}(x, \mathcal{S}_n)$. Thus the sequence $\langle x_n \rangle$ has a cluster point y . Suppose $y \neq x$. Now $\{y\} = \bigcap_{n=1}^{\infty} g(n, y)$ and so there is a k in N such that $x \notin g(k, y)$. Since y is a cluster point of $\langle x_n \rangle$ there is a $m \geq k$ such that $x_m \in g(k, y)$. Since g is an α -function for X , $x_m \in g(k, y)$ implies $g(k, x_m) \subseteq g(k, y)$. But $x \in h(m, x_m) \subseteq g(m, x_m) \subseteq g(k, x_m)$ and so $x \in g(k, y)$, a contradiction. Thus $x = y$ and x is a cluster point of $\langle x_n \rangle$.

COROLLARY 4.7. *Every regular $w\Delta$ -space with a σ -point finite separating open cover is a Moore space.*

COROLLARY 4.8. *Every regular countably compact space with a σ -point finite separating open cover is metrizable.*

5. A generalization of semi-stratifiable and $w\Delta$ -spaces. Let X be a space and let g be a function from $N \times X$ into the topology of

X such that for all x in X and n in N , $x \in g(n, x)$. Consider the following properties of the function g .

(A) If $x \in g(n, x_n)$ and $y_n \in g(n, x_n)$ for $n = 1, 2, \dots$ then x is a cluster point of the sequence $\langle y_n \rangle$.

(B) If $x \in g(n, x_n)$ and $y_n \in g(n, x_n)$ for $n = 1, 2, \dots$ then the sequence $\langle y_n \rangle$ has a cluster point.

(C) If $x_n \in g(n, x)$ for $n = 1, 2, \dots$ then x is a cluster point of the sequence $\langle x_n \rangle$.

(D) If $x_n \in g(n, x)$ for $n = 1, 2, \dots$ then the sequence $\langle x_n \rangle$ has a cluster point.

(E) If $x \in g(n, x_n)$ for $n = 1, 2, \dots$ then x is a cluster point of the sequence $\langle x_n \rangle$.

(F) If $x \in g(n, x_n)$ for $n = 1, 2, \dots$ then the sequence $\langle x_n \rangle$ has a cluster point.

In [10] Heath proved that developable spaces can be characterized in terms of a function g satisfying (A), and similarly $w\Delta$ -spaces can be characterized in terms of a function g satisfying (B). Clearly 1st countable spaces are characterized by (C), and (D) is precisely the definition of a q -space [18]. Finally, as proved in §4, semi-stratifiable spaces are characterized by a function g satisfying (E). These observations suggest introducing a new class of spaces, based on (F), which generalizes semi-stratifiable and $w\Delta$ -spaces.

DEFINITION 5.1. A space X is a β -space if there is a function g from $N \times X$ into the topology of X such that

(a) for all x in X and n in N , $x \in g(n, x)$;

(b) if $x \in g(n, x_n)$ for $n = 1, 2, \dots$ then the sequence $\langle x_n \rangle$ has a cluster point.

Such a function is called a β -function for X .

THEOREM 5.2. The following are equivalent for a regular space X :

(a) X is semi-stratifiable.

(b) X is a β -space with a G_s^* -diagonal.

(c) X is an α -space and a β -space.

Proof. Clearly (a) \Rightarrow (b) and (a) \Rightarrow (c). To prove (b) \Rightarrow (a) let g be a β -function for X and let $\mathcal{G}_1, \mathcal{G}_2, \dots$ be a G_s^* -sequence for X , where it is assumed that \mathcal{G}_{n+1} refines \mathcal{G}_n for all n . For x in X and n in N let $h(n, x) = g(n, x) \cap \text{st}(x, \mathcal{G}_n)$. Then h satisfies (c) of Lemma 4.5 and so X is semi-stratifiable.

To prove (c) \Rightarrow (a) let g be a β -function for X and let h be an α -function for X , where $h(n+1, x) \subseteq h(n, x)$ for all n in N and x

in X . For x in X and $n = 1, 2, \dots$ let $k(n, x) = g(n, x) \cap h(n, x)$. Then k satisfies (c) of Lemma 4.5 and so X is semi-stratifiable.

REMARK 5.3. The implication (d) \Rightarrow (a) of Corollary 2.6 and Theorem 4.6 can be proved using the above theorem together with Creede's result that every regular semi-stratifiable $w\Delta$ -space is a Moore space.

6. Summary. The relationship between some of the classes of spaces considered in this paper can be summarized in a diagram as follows.




Fig. 1

REFERENCES

1. R. H. Bing, *Metrization of topological spaces*, *Canad. J. Math.*, **3** (1951), 175-186.
2. C. J. R. Borges, *On stratifiable spaces*, *Pacific J. Math.*, **17** (1966), 1-16.
3. ———, *On metrizability of topological spaces*, *Canad. J. Math.*, **20** (1968), 795-804.
4. D. K. Burke and R. A. Stoltenberg, *A note on p -spaces and Moore spaces*, *Pacific J. Math.*, **30** (1969), 601-608.
5. D. K. Burke, *On subparacompact spaces*, *Proc. Amer. Math. Soc.*, **23** (1969), 655-663.
6. ———, *On p -spaces and $w\Delta$ -spaces*, to appear.
7. J. G. Ceder, *Some generalizations of metric spaces*, *Pacific J. Math.*, **11** (1961), 105-126.
8. G. Creede, *Concerning semi-stratifiable spaces*, *Pacific J. Math.*, **32** (1970), 47-54.
9. V. V. Filippov, *On feathered paracompacta*, *Soviet Math. Dokl.*, **9** (1968), 161-164.
10. R. W. Heath, *Arc-wise connectedness in semi-metric spaces*, *Pacific J. Math.*, **12** (1962), 1301-1319.
11. R. E. Hodel, *A note on subparacompact spaces*, *Proc. Amer. Math. Soc.*, **25** (1970), 842-845.
12. M. Katetov, *Complete normality of cartesian products*, *Fund. Math.*, **35** (1948), 271-274.
13. D. E. Kullman, *A note on developable spaces and p -spaces*, to appear in *Proc. Amer. Math. Soc.*
14. D. J. Lutzer, *A metrization theorem for linearly orderable spaces*, *Proc. Amer. Math. Soc.*, **22** (1969), 557-558.
15. L. F. McAuley, *A note on complete collectionwise normality and paracompactness*, *Proc. Amer. Math. Soc.*, **9** (1958), 796-799.
16. E. Michael, *A note on paracompact spaces*, *Proc. Amer. Math. Soc.*, **4** (1953), 831-838.

17. ———, *On Nagami's Σ -spaces and some related matters*, Proceedings of the Washington State University Conference on General Topology (1970), 13-19.
18. ———, *A note on closed maps and compact sets*, Israel J. Math., **2** (1964), 173-167.
19. A. S. Miščenko, *Spaces with point-countable bases*, Soviet Math. Dokl., **3** (1962), 855-858.
20. J. Nagata, *A note on Filipov's theorem*, Proc. Japan Acad., **45** (1969), 30-33.
21. J. Nagata and F. Siwiec, *A note on nets and metrization*, Proc. Japan Acad., **44** (1968), 623-627.
22. A. Okuyama, *On metrizability of M -spaces*, Proc. Japan Acad., **40** (1964), 176-179.
23. F. Siwiec, *Metrizability of some M -spaces*, to appear.
24. H. H. Wicke and J. M. Worrell Jr., *Characterizations of developable spaces*, Canad. J. Math., **17** (1965), 820-830.

Received June 8, 1970.

DUKE UNIVERSITY