

THE SUBSET OF PIECEWISE-LINEAR MAPPINGS IS DENSE IN THE SPACE OF K -QUASICONFORMAL MAPPINGS OF THE PLANE

BY

SIGBERT JAENISCH

Justus-Liebig-Universität, Giessen, Germany

I. Introduction

For each index n from the set \mathbb{N} of natural numbers, let \mathcal{N}_n denote the *regular net of equilateral triangles* in the complex plane \mathbb{C} , whose vertex set consists of the points $[p + (\frac{1}{2} + i\sqrt{3}/2)q]2^{-n}$ with integers p and q .

A mapping $\varphi: \mathbb{C} \rightarrow \mathbb{C}$ is called *linear*, if there are constants $a, b, c \in \mathbb{C}$ such that $\varphi(z) = az + bz^* + c$; the superscript star denotes complex conjugation. A mapping $\varphi: \mathbb{C} \rightarrow \mathbb{C}$ is said to be *piecewise-linear* with respect to the net \mathcal{N}_n , if its restrictions to the triangles of \mathcal{N}_n are linear mappings. We define the *piecewise-linearized mapping* $\varphi^{(n)}: \mathbb{C} \rightarrow \mathbb{C}$ for a mapping $\varphi: \mathbb{C} \rightarrow \mathbb{C}$ with respect to the net \mathcal{N}_n as follows: $\varphi^{(n)}$ is piecewise-linear with respect to \mathcal{N}_n , and it coincides with φ on the vertex set of \mathcal{N}_n .

The set of continuous mappings $\varphi: \mathbb{C} \rightarrow \mathbb{C}$ will be considered as a *topological space with the compact-open topology*; this induces *convergence* in the sense of uniform convergence on compact subsets. *Approximation* means convergence to a given mapping. Each continuous mapping $\varphi: \mathbb{C} \rightarrow \mathbb{C}$ is approximated by its piecewise-linearized mappings $\varphi^{(n)}$.

In the subspace of quasiconformal mappings of the plane, there is the problem: can each φ be approximated by φ_n which are piecewise-linear with respect to \mathcal{N}_n ?

METHOD OF BEURLING AND AHLFORS. *Let a quasiconformal mapping $\varphi: \mathbb{C} \rightarrow \mathbb{C}$ have maximal dilatation $K(\varphi) < \sqrt{3}$. Then, φ is approximated by the piecewise-linearized mappings $\varphi^{(n)}$; $\varphi^{(n)}$ is quasiconformal (Ahlfors [2], 768; [3], 298); $\varphi^{(n)}$ has maximal dilatation $K(\varphi^{(n)}) \leq \xi[K(\varphi)]$, where ξ is a certain function involving elliptic integrals (Agard [1], 739); for each index n , there are some φ such that $K(\varphi^{(n)}) = \xi[K(\varphi)]$ holds (Agard [1], 739); moreover, there are some φ such that $K(\varphi^{(n)}) = \xi[K(\varphi)]$ holds for all indices n ([4], 49).*

AGARD'S METHOD. Let a quasiconformal mapping $\varphi: \mathbf{C} \rightarrow \mathbf{C}$ be composed as $\varphi = \tilde{\varphi}_p \circ \dots \circ \tilde{\varphi}_1$ by quasiconformal mappings $\tilde{\varphi}_q: \mathbf{C} \rightarrow \mathbf{C}$ with maximal dilatations $K(\tilde{\varphi}_q) = [K(\varphi)]^{1/p} < \sqrt[3]{3}$. Then φ is approximated by the mappings $\varphi_n := \tilde{\varphi}_p^{(n)} \circ \dots \circ \tilde{\varphi}_1^{(n)}$; φ_n is piecewise-linear, though not with respect to \mathfrak{N}_n ; φ_n is quasiconformal; for $\varepsilon > 0$ and p sufficiently large, φ_n has maximal dilatation $K(\varphi_n) \leq [K(\varphi)]^{3,243\dots} + \varepsilon$ (Agard [1], 740); further, there are some φ such that

$$K(\varphi_n) > [K(\varphi)]^{3,243\dots}$$

holds for all indices n and all admissible p ([4], 51).

THEOREM (abridged version). Let a quasiconformal mapping $\varphi: \mathbf{C} \rightarrow \mathbf{C}$ have maximal dilatation $K(\varphi) > 1$. Then, φ can be approximated ("well" in the sense of Lehto-Virtanen [5], 194) by quasiconformal mappings $\varphi_n: \mathbf{C} \rightarrow \mathbf{C}$ with maximal dilatations $K(\varphi_n) < K(\varphi)$, which are piecewise-linear with respect to certain prescribed nets \mathfrak{N}_n of triangles.

COROLLARY. The subset of piecewise-linear mappings is dense in the space of K -quasiconformal mappings of the plane.

With regard to maximal dilatations, Agard's result is weaker than our theorem. This is caused by the use of a sufficient condition, which is an unnecessary limitation: to approximate by piecewise linearizations of the given mapping or of mappings composing it. Now, consider the following necessary, but insufficient condition.

PROPOSITION. Let a quasiconformal mapping $\varphi: \mathbf{C} \rightarrow \mathbf{C}$ be approximated by quasiconformal mappings $\varphi_n: \mathbf{C} \rightarrow \mathbf{C}$, which are piecewise-linear with respect to arbitrary nets \mathfrak{N}_n . Then, φ can be approximated by quasiconformal mappings $\psi_n: \mathbf{C} \rightarrow \mathbf{C}$ such that $\varphi_n = \psi_n^{(n)}$.

Taking the φ_n as ψ_n proves it. To apply this proposition, we take certain sufficiently smooth ψ_n with maximal dilatations $K(\psi_n) < K(\varphi)$.

2. Linear mappings of a triangle

A triangle in \mathbf{C} is given by its vertices, three non-collinear points $z_1, z_2, z_3 \in \mathbf{C}$. We represent the vertices and thus the triangle by the triple $Z := (z_1, z_2, z_3)$ in the Cartesian product space \mathbf{C}^3 . The point set of the triangle is $\mathbf{T} := \text{conv } Z$.

Let a linear mapping $\varphi: \mathbf{C} \rightarrow \mathbf{C}$ be given by $\varphi(z) := az + bz^* + c$ with constants $a, b, c \in \mathbf{C}$. Then, $\varphi^3(z', z'', z''') := (\varphi(z'), \varphi(z''), \varphi(z'''))$ defines the Cartesian product mapping $\varphi^3: \mathbf{C}^3 \rightarrow \mathbf{C}^3$. Let us introduce the notation $E := (1, 1, 1)$. φ is uniquely determined by its restriction to the vertex set of a triangle $\mathbf{T} = \text{conv } Z$. From

$$W := \varphi^3(Z) = aZ + bZ^* + cE,$$

Cramer's rule allows us to compute a, b, c for given triples Z (non-collinear) and W .

LEMMA 1. Let $Z:=(z_1, z_2, z_3)$ denote the triple of vertices of a triangle in \mathbf{C} , and let $W:=(w_1, w_2, w_3)$, $\tilde{W}:=(\tilde{w}_1, \tilde{w}_2, \tilde{w}_3)$ be triples in \mathbf{C}^3 . Then, there are uniquely determined linear mappings $\varphi, \tilde{\varphi}: \mathbf{C} \rightarrow \mathbf{C}$ such that $\varphi^3(Z) = W$ and $\tilde{\varphi}^3(Z) = \tilde{W}$. φ has the representation

$$\varphi(z) = \frac{z \cdot \det(W, Z^*, E) + z^* \cdot \det(Z, W, E) + \det(Z, Z^*, W)}{\det(Z, Z^*, E)}.$$

φ is quasiconformal if and only if $\det(W, Z^*, E) > \det(Z, W, E)$; then the complex dilatation is

$$\kappa = \frac{\det(Z, W, E)}{\det(W, Z^*, E)}.$$

If φ and $\tilde{\varphi}$ are quasiconformal, then the difference of their complex dilatations is

$$\tilde{\kappa} - \kappa = \frac{\det(W, \tilde{W}, E) \cdot \det(Z, Z^*, E)}{\det(W, Z^*, E) \cdot \det(\tilde{W}, Z^*, E)}.$$

Proof. It is trivial to check a, b, c and $\kappa = b/a$. In order to show the last result, we work with known formulas for inner “ \cdot ” and outer “ \times ” multiplication of 3-vectors:

$$\begin{aligned} &(\tilde{\kappa} - \kappa) \cdot \det(W, Z^*, E) \cdot \det(\tilde{W}, Z^*, E) \\ &= \det(Z, \tilde{W}, E) \cdot \det(W, Z^*, E) - \det(Z, W, E) \cdot \det(\tilde{W}, Z^*, E) \\ &= [\det(Z, \tilde{W}, E) \cdot W - \det(Z, W, E) \cdot \tilde{W}] \cdot (Z^* \times E) \\ &= [(E \times Z) \times (W \times \tilde{W})] \cdot (Z^* \times E) \\ &= [\det(W, \tilde{W}, E) \cdot Z - \det(W, \tilde{W}, Z) \cdot E] \cdot (Z^* \times E) \\ &= \det(W, \tilde{W}, E) \cdot \det(Z, Z^*, E). \end{aligned}$$

3. Linearization of a nearly linear mapping

Let a triangle $\mathbf{T} = \text{conv } Z$ in \mathbf{C} be given, and consider a continuous mapping $\varphi: \mathbf{T} \rightarrow \mathbf{C}$. The *linearized mapping* $\hat{\varphi}: \mathbf{C} \rightarrow \mathbf{C}$ for φ with respect to \mathbf{T} is defined as follows: $\hat{\varphi}$ is the uniquely determined linear mapping with $\hat{\varphi}^3(Z) = \varphi^3(Z)$.

If the mapping $\varphi: \mathbf{T} \rightarrow \mathbf{C}$ is quasiconformal, then its complex dilatation $\kappa(z)$ equals

$$\frac{\partial \varphi}{\partial z^*}(z) \bigg/ \frac{\partial \varphi}{\partial z}(z)$$

almost everywhere in \mathbf{T} . We want to guarantee that $\hat{\varphi}$ is quasiconformal with complex dilatation $\hat{\kappa}$ near $\kappa(z)$. Since a condition on $\kappa(z)$ alone would allow a subsequent conformal distortion of $\varphi(\mathbf{T})$ and thus of $\varphi^3(Z)$, let us try a limitation of both $\frac{\partial \varphi}{\partial z^*}(z)$ and $\frac{\partial \varphi}{\partial z}(z)$; this leads to the following lemma.

LEMMA 2. Let a triangle \mathbf{T} in \mathbf{C} have angles $\alpha_1, \alpha_2, \alpha_3$ and put $\beta := \max\{\alpha_1, \alpha_2, \alpha_3, \pi/2\}$. Consider a continuously-differentiable quasiconformal mapping $\psi: \mathbf{T} \rightarrow \mathbf{C}$, and the linearized mapping $\hat{\psi}: \mathbf{C} \rightarrow \mathbf{C}$ for ψ with respect to \mathbf{T} . Let the differential of ψ satisfy an inequality

$$|d\psi(z) - (adz + bdz^*)| \leq \varepsilon |dz|$$

with constants $a, b \in \mathbf{C}$ and a positive constant $\varepsilon < \frac{1}{2}(|a| - |b|)\sin\beta$. Then $\hat{\psi}$ is quasiconformal, and its complex dilatation $\hat{\kappa}$ satisfies the inequalities

$$\left| \hat{\kappa} - \frac{b}{a} \right| \leq \frac{|b/a| + 1}{|a/\varepsilon| \sin\beta - 1}, \quad |\hat{\kappa}| \leq \frac{|b| \sin\beta + \varepsilon}{|a| \sin\beta - \varepsilon}.$$

Proof. We are going to assume the following special conditions. In case that $\beta > \pi/2$, the maximal angle is $\alpha_1 = \beta$, the corresponding vertex is $z_1 = 0$; further, $\psi(0) = 0$. In case that $\beta = \pi/2$, the orthocenter of \mathbf{T} lies at the origin 0; further, $\psi(0) = 0$. This can be achieved by a renumbering of the vertices of \mathbf{T} , a translatory mapping of \mathbf{T} , and composition of ψ between two translatory mappings. Clearly, the values of the differentials and complex dilatations in Lemma 2 remain unchanged.

Now, the geometrical properties of \mathbf{T} imply: $|(E \times Z) \cdot Z^*| = |E \times Z| \cdot |Z| \sin\beta$; absolute values are taken componentwise: $|Z| = |(z_1, z_2, z_3)| := (|z_1|, |z_2|, |z_3|)$.

Let us define linear mappings $\varphi, \tilde{\varphi}: \mathbf{C} \rightarrow \mathbf{C}$ by $\varphi(z) := az + bz^*$ and $\tilde{\varphi} := \hat{\psi}$. Integrating the differentials $d\psi(z)$ and $d\varphi(z)$ along the segments from 0 to the vertices of \mathbf{T} , we get

$$\tilde{\varphi}^3(Z) = \hat{\psi}^3(Z) = \psi^3(Z) = \varphi^3(Z) + D;$$

here, the triple D satisfies $|D| \leq \varepsilon |Z|$ componentwise. Thus, we can estimate as follows: $|(E \times Z) \cdot D| \leq |E \times Z| \cdot |D| \leq \varepsilon |E \times Z| \cdot |Z|$. Next, we apply Lemma 1; the above formula and estimate will be used some lines further.

$$\begin{aligned} \hat{\kappa} - \kappa &= \frac{\det(aZ + bZ^*, aZ + bZ^* + D, E) \cdot \det(Z, Z^*, E)}{\det(aZ + bZ^*, Z^*, E) \cdot \det(aZ + bZ^* + D, Z^*, E)} \\ &= \frac{[a \cdot \det(Z, D, E) + b \cdot \det(Z^*, D, E)] \cdot \det(Z, Z^*, E)}{[a \cdot \det(Z, Z^*, E)] [a \cdot \det(Z, Z^*, E) + \det(D, Z^*, E)]} \\ &= \frac{(E \times Z) \cdot D + (b/a)(E \times Z^*) \cdot D}{a(E \times Z) \cdot Z^* + (Z^* \times E) \cdot D}, \end{aligned}$$

hence
$$|\hat{\kappa} - \kappa| \leq \frac{\varepsilon |E \times Z| \cdot |Z| + \varepsilon |b/a| \cdot |E \times Z| \cdot |Z|}{|a| \cdot |E \times Z| \cdot |Z| \sin\beta - \varepsilon |E \times Z| \cdot |Z|} = \frac{1 + |b/a|}{|a/\varepsilon| \sin\beta - 1};$$

and finally
$$|\hat{\kappa}| \leq |\hat{\kappa} - \kappa| + |\kappa| \leq \frac{\varepsilon + \varepsilon |b/a|}{|a| \sin\beta - \varepsilon} + \left| \frac{b}{a} \right| = \frac{|b| \sin\beta + \varepsilon}{|a| \sin\beta - \varepsilon}.$$

4. Nets of non-degenerating triangles

For each index $n \in \mathbb{N}$, let us consider a locally-finite covering \mathcal{N}_n of \mathbb{C} by triangles, such that any two different triangles of \mathcal{N}_n intersect in a common side, in a common vertex, or not at all. The \mathcal{N}_n will be called *general nets of triangles*.

The triangles of a general net can be very different in size; we want to describe the diameters of the triangles with the help of a continuous function $\delta_n: \mathbb{C} \rightarrow \mathbb{R}$.

For any triangle $T \in \mathcal{N}_n$ and any point $z \in \mathbb{C}$, put

$$T_z := (T - z) \cap (z - T) := \{w \in \mathbb{C} : z \pm w \in T\}.$$

In fact, T_z is the largest subset of $T - z$ which is symmetrical with respect to the origin 0. T_z is void for $z \notin T \in \mathcal{N}_n$. If $z \in T \in \mathcal{N}_n$, then T_z is a non-void convex set which depends on \mathcal{N}_n , depends continuously on z , but does not depend on the particular T chosen. We introduce the *symmetrical diameter* $\delta_n(z)$ for the net \mathcal{N}_n with respect to the point z as

$$\delta_n(z) := \max \{ \text{diam} [(T - z) \cap (z - T)] : T \in \mathcal{N}_n \}.$$

Let the triangle $T = \text{conv} \{z_1, z_2, z_3\}$ of the net \mathcal{N}_n have angles $\alpha_1, \alpha_2, \alpha_3$ and put $\beta := \max \{ \alpha_1, \alpha_2, \alpha_3, \pi/2 \}$. Elementary calculations lead to an upper bound and to a condition of Lipschitz type:

$$\max \{ \delta_n(z) : z \in T \} = \text{diam } T = \max \{ |z_3 - z_2|, |z_1 - z_3|, |z_2 - z_1| \},$$

$$|\delta_n(z'') - \delta_n(z')| \leq |z'' - z'| \cdot 2 \tan \frac{\beta}{2} \text{ for } z', z'' \in T.$$

Next, let us consider the shape of the triangles. We need good estimates for $\hat{\kappa}$; yet, Lemma 2 is asymptotically sharp for $\beta \uparrow \pi$, because there are examples with

$$|\hat{\kappa}| = \frac{|b| \sin \beta + \varepsilon \sin (\beta/2)}{|a| \sin \beta - \varepsilon \sin (\beta/2)}.$$

We can allow small angles, but we must prevent the triangles from being too obtuse. Therefore, we speak of a *non-degenerating sequence of nets* \mathcal{N}_n , if all angles of all triangles of all nets are bounded away from π .

This makes the functions $\delta_n: \mathbb{C} \rightarrow \mathbb{R}$ uniformly continuous, even equicontinuous. Hence, a well-known theorem ([5], 74, Hilfssatz 5.1) implies the next lemma.

LEMMA 3. *Consider a non-degenerating sequence of nets \mathcal{N}_n of triangles in \mathbb{C} . Let the symmetrical diameters $\delta_n(z)$ converge to 0 pointwise on a dense subset of \mathbb{C} . Then the δ_n converge to 0 in the compact-open topology.*

5. Approximation by smooth mappings with smaller dilatations

We are going to approximate a quasiconformal mapping $\varphi: \mathbb{C} \rightarrow \mathbb{C}$ by auxiliary mappings $\tilde{\psi}_n: \mathbb{C} \rightarrow \mathbb{C}$. We require certain smoothness properties which are to be used later.

LEMMA 4. *Let a quasiconformal mapping $\varphi: \mathbb{C} \rightarrow \mathbb{C}$ have complex dilatation $\kappa(z)$. Then, there exists a sequence of continuously-differentiable quasiconformal mappings $\tilde{\psi}_n: \mathbb{C} \rightarrow \mathbb{C}$, converging to φ in the compact-open topology, with complex dilatations $\tilde{\kappa}_n(z)$ converging to $\kappa(z)$ pointwise almost everywhere in \mathbb{C} , and satisfying the following conditions. Each $\tilde{\psi}_n$ is conformal outside some compact set; $\tilde{\psi}_n(0) = \varphi(0)$ and $\tilde{\psi}_n(1) = \varphi(1)$ holds;*

$$\frac{K(\tilde{\psi}_n) - 1}{K(\tilde{\psi}_n) + 1} \leq \left(1 - \frac{1}{n}\right) \frac{K(\varphi) - 1}{K(\varphi) + 1},$$

for each index n and for each $\varepsilon > 0$ there are constants ϱ_n, σ_n with $0 < \varrho_n, \sigma_n < \infty$ such that if two points $z', z'' \in \mathbb{C}$ fulfil either $|z'' - z'| \leq \varrho_n$ or $|z'|, |z''| \geq \sigma_n$, then $d\tilde{\psi}_n$ satisfies

$$|d\tilde{\psi}_n(z'') - d\tilde{\psi}_n(z')| \leq \varepsilon |dz|.$$

Proof. We use the standard method of defining a quasiconformal mapping implicitly by its complex dilatation. Let us start with a sequence of real-analytic functions $\kappa_n: \mathbb{C} \rightarrow \mathbb{C}$, with $\kappa_n(z)$ converging to $\kappa(z)$ pointwise almost everywhere in \mathbb{C} , and such that

$$\sup \{ |\kappa_n(z)| : |z| \leq n + \pi/2 \} \leq [K(\varphi) - 1] / [K(\varphi) + 1].$$

Further, let us put

$$\chi_n(\tau) := \begin{cases} 1 - \frac{1}{n} & \text{for } \tau \leq n \\ \left(1 - \frac{1}{n}\right) [1 - \sin^2(\tau - n)] & \text{for } n < \tau < n + \frac{\pi}{2} \\ 0 & \text{for } \tau \geq n + \frac{\pi}{2} \end{cases}$$

This defines a continuously differentiable function $\chi_n: \mathbb{R} \rightarrow \mathbb{R}$. By $\tilde{\kappa}_n(z) := \kappa_n(z)\chi_n(|z|)$, we get a continuously differentiable function $\tilde{\kappa}_n: \mathbb{C} \rightarrow \mathbb{C}$ with the required properties.

The generalized Riemann mapping theorem ([5], 204) guarantees a corresponding quasiconformal mapping $\tilde{\psi}_n: \mathbb{C} \rightarrow \mathbb{C}$ with $\tilde{\psi}_n(0) = \varphi(0)$ and $\tilde{\psi}_n(1) = \varphi(1)$. The condition for the differential $d\tilde{\psi}_n$ follows from the smoothness of $\tilde{\kappa}_n$, because $\tilde{\psi}_n$ is regular ([5], 244) in \mathbb{C} and conformal outside a compact subset of \mathbb{C} .

Finally, a well-known normality argument ([5], 218) combined with the fact that our convergence is derived from a topology, leads to the conclusion that the $\tilde{\psi}_n$ approximate φ .

6. Approximation by piecewise-linear mappings with smaller dilatations

THEOREM. *For a non-degenerating sequence of nets \mathcal{N}_n of triangles in \mathbb{C} , let the symmetrical diameters $\delta_n(z)$ converge to 0 pointwise on a dense subset of \mathbb{C} . Further, let a quasiconformal mapping $\varphi: \mathbb{C} \rightarrow \mathbb{C}$ have complex dilatation $\kappa(z)$ and maximal dilatation $K(\varphi) > 1$. Then there exist quasiconformal mappings $\varphi_n: \mathbb{C} \rightarrow \mathbb{C}$, converging to φ in the compact-open topology, with complex dilatations $\kappa_n(z)$ converging to $\kappa(z)$ pointwise almost everywhere in \mathbb{C} , such that φ_n has maximal dilatation $K(\varphi_n) < K(\varphi)$, and φ_n is piecewise-linear with respect to the net \mathcal{N}_n .*

Proof. I. In this part, we apply Lemmas 3 and 4. — Lemma 4 approximates the given mapping φ by certain mappings $\tilde{\varphi}_n$. Since $\tilde{\varphi}_n$ is a regular quasiconformal mapping, conformal outside of a certain compact subset of \mathbb{C} , the absolute value of $\partial\tilde{\varphi}_n/\partial z$ has a positive lower bound γ_n . As the sequence of the nets \mathcal{N}_n is non-degenerating, we can find a constant β_0 with $\pi/2 \leq \beta_0 < \pi$ which is an upper bound for the angles of the triangles of the nets \mathcal{N}_n . For each index n , we define the positive constant

$$\varepsilon_n := \frac{\gamma_n}{3n} \left(1 - \frac{1}{K(\varphi)} \right) \sin \beta_0.$$

Putting $\varepsilon := \varepsilon_n$ in Lemma 4, we get constants ϱ_n, σ_n such that $|z'' - z'| \leq \varrho_n$ or $|z'|, |z''| \geq \sigma_n$ implies the inequality $|d\tilde{\varphi}_n(z'') - d\tilde{\varphi}_n(z')| \leq \varepsilon_n |dz|$.

Lemma 3 ensures us that the symmetrical diameters $\delta_n(z)$ converge to 0 uniformly on compact subsets of \mathbb{C} . For each index m , and for all sufficiently large indices n , any triangle $\mathbf{T} \in \mathcal{N}_n$ will satisfy either $\text{diam } \mathbf{T} \leq \varrho_m$ or $\mathbf{T} \subset \{z: |z| \geq \sigma_m\}$. We can choose an isotonic and surjective function $\lambda: \mathbb{N} \rightarrow \mathbb{N}$ such that the pairs (m, n) with $m \leq \lambda(n)$ are admissible for the above statement.

II. Choice of the mappings φ_n . — Since for each index m , the mapping $\tilde{\varphi}_m$ is approximated by its piecewise-linearized mappings $\tilde{\varphi}_m^{(n)}$, it follows that we have $|\tilde{\varphi}_m^{(n)}(0) - \varphi(0)| = |\tilde{\varphi}_m^{(n)}(0) - \tilde{\varphi}_m(0)| \leq 2^{-m}$ and $|\tilde{\varphi}_m^{(n)}(1) - \varphi(1)| = |\tilde{\varphi}_m^{(n)}(1) - \tilde{\varphi}_m(1)| \leq 2^{-m}$ for all sufficiently large indices n . We can choose an isotonic and surjective function $\mu: \mathbb{N} \rightarrow \mathbb{N}$ such that the pairs (m, n) with $m \leq \mu(n)$ are admissible above.

Taking $\nu := \min \{\lambda, \mu\}$, we define another isotonic and surjective function $\nu: \mathbb{N} \rightarrow \mathbb{N}$. We put $m := \nu(n)$; this makes the pairs $(m, n) = (\nu(n), n)$ admissible in the sense of parts I and II. With the sequence of the mappings $\tilde{\varphi}_m$, we associate the *diluted sequence* consisting of the mappings $\psi_n := \tilde{\varphi}_{\nu(n)}$. Finally, let us define $\varphi_n := \psi_n^{(n)}$; these mappings $\varphi_n: \mathbb{C} \rightarrow \mathbb{C}$ are piecewise-linear with respect to the nets \mathcal{N}_n .

III. Application of Lemma 2. — Let us take any one of the nets \mathcal{N}_n , any one of the triangles $T \in \mathcal{N}_n$, and any one of the points $z \in T$. We define $\psi: T \rightarrow \mathbb{C}$ as the restriction of ψ_n to T ; its linearization $\hat{\psi}: \mathbb{C} \rightarrow \mathbb{C}$ with respect to T coincides with $\varphi_n = \psi_n^{(n)}$ on T . Finally, we put $a := \frac{\partial \psi_n}{\partial z}(z)$, $b := \frac{\partial \psi_n}{\partial z^*}(z)$, $\varepsilon := \varepsilon_{\nu(n)}$. Now we apply Lemma 2 and use the following estimates: $|a| \geq \gamma_{\nu(n)}$; $\sin \beta \geq \sin \beta_0$; $|b/a| \leq [1 - 1/\nu(n)][K(\varphi) - 1]/[K(\varphi) + 1]$.

$$\begin{aligned} |\kappa_n(z) - \tilde{\kappa}_{\nu(n)}| &= \left| \hat{\kappa} - \frac{b}{a} \right| \leq \frac{|b/a| + 1}{|a/\varepsilon| \sin \beta - 1} \leq \frac{[1 - 1/\nu(n)][K(\varphi) - 1]/[K(\varphi) + 1] + 1}{3\nu(n)K(\varphi)/[K(\varphi) - 1] - 1} \\ &\leq \frac{2\nu(n)K(\varphi) - [K(\varphi) - 1]}{3\nu(n)K(\varphi) - [K(\varphi) - 1]} \cdot \frac{1}{\nu(n)} \cdot \frac{K(\varphi) - 1}{K(\varphi) + 1} \leq \frac{2}{3\nu(n)} \cdot \frac{K(\varphi) - 1}{K(\varphi) + 1}, \end{aligned}$$

hence

$$|\kappa_n(z)| \leq \left| \hat{\kappa} - \frac{b}{a} \right| + \left| \frac{b}{a} \right| \leq \left[1 - \frac{1}{3\nu(n)} \right] \cdot \frac{K(\varphi) - 1}{K(\varphi) + 1};$$

and finally

$$\frac{1 + |\kappa_n(z)|}{1 - |\kappa_n(z)|} \leq \frac{6\nu(n)K(\varphi) - [K(\varphi) - 1]}{6\nu(n) + [K(\varphi) - 1]} < K(\varphi).$$

IV. Properties of the mappings φ_n . — Part III implies the quasiconformality of φ_n in each triangle $T \in \mathcal{N}_n$ and thus in \mathbb{C} . The estimates for $\kappa_n(z)$ hold in $\bigcup \{\text{int } T: T \in \mathcal{N}_n\}$, which is almost everywhere in \mathbb{C} ; since we have upper bounds depending only on $\nu(n)$ and $K(\varphi)$, we can deduce $K(\varphi_n) < K(\varphi)$. For $n \rightarrow \infty$, we find that $\kappa_n(z) - \tilde{\kappa}_{\nu(n)} \rightarrow 0$, and $\tilde{\kappa}_{\nu(n)} \rightarrow \kappa(z)$, hence $\kappa_n(z) \rightarrow \kappa(z)$ pointwise almost everywhere in \mathbb{C} . If we combine this with $\varphi_n(0) \rightarrow \varphi(0)$ and $\varphi_n(1) \rightarrow \varphi(1)$, we can conclude that the corresponding mappings $\varphi_n: \mathbb{C} \rightarrow \mathbb{C}$ approximate $\varphi: \mathbb{C} \rightarrow \mathbb{C}$ in the compact-open topology.

References

- [1]. AGARD, S., An extremal problem in quasiconformal mappings. *Duke Math. J.*, 33 (1966), 735–741.
- [2]. AHLFORS, L. V., Extension of quasiconformal mappings from two to three dimensions. *Proc. Nat. Acad. Sci. USA*, 51 (1964), 768–771.
- [3]. ——— The modular function and geometric properties of quasiconformal mappings. *Proc. Conf. Compl. Anal.* (Minneapolis, 1964), Springer, Berlin, 1965, 296–300.
- [4]. JAENISCH, S., Über die Approximierbarkeit quasikonformer Abbildungen der Ebene mittels stückweiser Linearisierung. *Mitt. Math. Sem. Giessen*, 79 (1968), 29–53.
- [5]. LEHTO, O. & VIRTANEN, K. I., Quasikonforme Abbildungen. Springer, Berlin, 1965.

Received April 2, 1968, in revised form December 18, 1968