

Research Article

Controllability Analysis of Linear Discrete Time Systems with Time Delay in State

Hong Shi,¹ Guangming Xie,^{2,3} and Wenguang Luo^{4,5}

¹ Department of Mathematics and Physics, Beijing Institute of Petrochemical Technology, Beijing 102617, China

² State Key Laboratory for Turbulence and Complex Systems, College of Engineering, Peking University, Beijing 100871, China

³ School of Electrical and Electronics Engineering, East China Jiaotong University, Nanchang 330013, China

⁴ School of Electric and Information Engineering, Guangxi University of Science and Technology, Liuzhou 545006, China

⁵ Guangxi Key Laboratory of Automobile Components and Vehicle Technology, Liuzhou 545006, China

Correspondence should be addressed to Guangming Xie, xiegm@pku.edu.cn

Received 27 August 2012; Accepted 10 October 2012

Academic Editor: Valery Y. Glizer

Copyright © 2012 Hong Shi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The controllability issues for linear discrete-time systems with delay in state are addressed. By introducing a new concept, the minimum controllability realization index (MinCRI), the characteristic of controllability is revealed. It is proved that the MinCRI of a system with state delay exists and is finite. Based on this result, a necessary and sufficient condition for the controllability of discrete-time linear systems with state delay is established.

1. Introduction

The concept of controllability of dynamical systems was first proposed by Kalman in 1960s [1]. Since then, controllability of dynamical systems has been studied by many authors in various contexts [2–11], because controllability turns out to be a fundamental concept in modern control theory and has close connections with pole assignment, structure decomposition, quadratic control, and so forth [12, 13]. On the other hand, time delay phenomena are very common in practical systems, for instance, in economic, biological, and physiological systems. Studying the time delay phenomena in control systems has become

an important topic in control theory. Chyung studied the controllability for linear time-invariant systems with constant time delay in the control functions. Simple algebraic-type necessary and sufficient criteria are established [3, 4]. However, once the time delay appears in state, the problem becomes much more complex. There are some preliminary results in [5, 6]. However, all these results are not suitable for verification and application.

In this paper, we consider the discrete-time case, the system model described as follows:

$$x(k+1) = Ax(k) + Dx(k-h) + Bu(k), \quad (1.1)$$

where $x(k) \in \mathbb{R}^n$ is the state, $u(k) \in \mathbb{R}^p$ is the input, $A, D \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times p}$ are constant matrices, and positive integer h is the length of the steps of time delay. The initial states $x(-h), x(-h+1), \dots, x(0)$ are given arbitrarily.

The controllability discussed here refers to the unconstrained controllability, or say completely controllability.

Definition 1.1 (controllability). System (1.1) is said to be (completely) controllable, if for any initial state $x(-h), x(-h+1), \dots, x(0)$ and any final state x_f , there exist a positive integer k and input $u(0), \dots, u(k-1)$ such that $x(k) = x_f$.

Definition 1.2 (controllability realization index, (CRI)). For system (1.1), if there exists a positive integer K such that for any initial state $x(-h), x(1-h), \dots, x(0)$ and any terminal state x_f , there exist input $u(0), \dots, u(K-1)$ such that $x(K) = x_f$; then one calls K the controllability realization index (CRI) of system (1.1). Obviously, if exists, such K is not unique, so one calls the smallest K among them the minimum controllability realization index (MinCRI).

In this paper, our main concern is the following. *Can the controllability of system (1.1) be realized completely in finite steps, or say can the MinCRI of the system be finite?*

Here we demonstrate that the answer to this question is *yes*. And in the case of planar systems, the exact value of the controllability realization index is obtained, and we will prove it is independent of the choices of A and D .

This paper is organized as follows. In Section 2 some concepts are introduced, which will be used in the later discussion. Section 3 contains the main results. Finally, the conclusion is provided in Section 4.

2. Preliminaries

Denote by $\mathbb{N}$ the nonnegative integer set, $\mathbb{R}$ the real number set, respectively. The matrices $A_1, \dots, A_N \in \mathbb{R}^{n \times n}$ are said to be linearly dependent in $\mathbb{R}^{n \times n}$, if there exist scalars $c_1, \dots, c_N \in \mathbb{R}$, not all zero, such that $\sum_{i=1}^N c_i A_i = 0$. In what follows, $\text{span}\{A_1, \dots, A_N\}$ will be used to denote the linear subspace constructed by the linear combinations of matrices $\{A_1, \dots, A_N\}$.

Definition 2.1 (symmetrical subspace). Given matrices $A, D \in \mathbb{R}^{n \times n}$, one defines a subset of $\mathbb{R}^{n \times n}$ as follows:

$$P[A, D] := \left\{ f(A, D) \mid f(A, D) = \sum_{m=1}^M c_m A^{i_{m,1}} D^{j_{m,1}} \dots A^{i_{m,n_m}} D^{j_{m,n_m}}, \right. \\ \left. n_m \in \mathbb{N}, i_{m,1}, j_{m,1}, \dots, i_{m,n_m}, j_{m,n_m} \in \mathbb{N}, 0 \leq i_{m,1}, j_{m,1}, \dots, i_{m,n_m}, j_{m,n_m} \leq n, \right. \\ \left. c_m \in \mathbb{R}, m = 1, \dots, M; M \in \mathbb{N} \right\}. \quad (2.1)$$

Then a one-to-one mapping $\Theta : P[A, D] \rightarrow P[A, D]$ is constructed as follows:

give any $f(A, D) \in P[A, D]$, supposing $f(A, D) = \sum_{m=1}^M c_m A^{i_{m,1}} D^{j_{m,1}} \dots A^{i_{m,n_m}} D^{j_{m,n_m}}$, one defines

$$\Theta(f(A, D)) := \sum_{m=1}^M c_m \left(A^{i_{m,1}} D^{j_{m,1}} \dots A^{i_{m,n_m}} D^{j_{m,n_m}} + D^{j_{m,n_m}} A^{i_{m,n_m}} \dots D^{j_{m,1}} A^{i_{m,1}} \right). \quad (2.2)$$

Then we get a subspace of $P[A, D]$ by

$$P_{\text{Sym}}[A, D] := \{ \Theta(f(A, D)) \mid \forall f(A, D) \in P[A, D] \}. \quad (2.3)$$

We call $P_{\text{Sym}}[A, D]$ the *symmetrical subspace* of $P[A, D]$.

By Definition 2.1, it is easy to verify that $I, A, D, AD + DA, ADA, DAD \in P_{\text{Sym}}[A, D]$, where I is the identity matrix in $\mathbb{R}^{n \times n}$.

Now, we introduce a matrix sequence $\{G_k\}_{k=0}^{\infty} \in \mathbb{R}^{n \times n}$ as follows:

$$G_k = \begin{cases} A^k, & \text{if } k = 0, 1, \dots, h, \\ G_{k-1}A + G_{k-1-h}D, & \text{if } k = h+1, h+2, \dots \end{cases} \quad (2.4)$$

Then the solution of system (1.1) can be expressed as

$$x(k+1) = \Psi_k(x(0), \dots, x(-h)) + \sum_{i=0}^k G_{k-i} B u(i), \quad k > 0, \quad (2.5)$$

where $\Psi_k(x(0), \dots, x(-h))$ is the part of the solution with zero input.

Lemma 2.2. Given the matrix sequence $\{G_k\}_{k=0}^{\infty}$ in (2.3), the following statements hold:

- (a) there are no similar terms between $G_k A$ and $G_{k-h} D$;
- (b) for any G_k , it can be expressed as

$$G_k = \sum_{\forall f(A, D) \in P_{\text{Sym}}[A, D]} f(A, D), \quad (2.6)$$

where $Q_k[A, D]$ is a subset of $\mathbb{R}^{n \times n}$ defined as follows:

$$Q_k[A, D] = \left\{ f(A, D) \mid f(A, D) = A^{i_1} D^{j_1} \cdots A^{i_k} D^{j_k}, \right. \\ \left. \forall i_1, j_1, \dots, i_k, j_k \in \mathbb{N}, \sum_{m=1}^k i_m + (h+1) \sum_{m=1}^k j_m = k \right\}; \quad (2.7)$$

(c) for all $k \in \mathbb{N}$, $G_k \in P_{\text{Sym}}[A, D]$.

Proof. Statement (a) is nearly self-evident because any term from $G_k A$ is ended by A , whereas any term from $G_{k-h} D$ is ended by D .

To prove the result of statement (b), mathematical induction is invoked.

(I) The first $h+1$ terms of $\{G_k\}_{k=0}^\infty$ are $I, A, \dots, A^h$, and it is obvious that $Q_k[A, D] = \{A^k\}$, for $k = 0, 1, \dots, h$. Thus, for $k = 0, 1, \dots, h$, (2.5) holds.

(II) Assume that, for $l = k, k+1, \dots, k+h$, G_l is expressed in form (2.5). We will prove that G_{k+h+1} can be expressed in form (2.5) as well.

First, we prove that, for any k , we have

$$Q_{k+1}[A, D] = Q_k[A, D]A \cup Q_{k-h}[A, D]D, \quad (2.8)$$

where the sets $Q_k[A, D]A$, $Q_{k-h}[A, D]D$ are defined as

$$Q_k[A, D]A = \{f(A, D)A \mid f(A, D) \in Q_k[A, D]\}, \\ Q_{k-h}[A, D]D = \{f(A, D)D \mid f(A, D) \in Q_{k-h}[A, D]\}. \quad (2.9)$$

By the assumption, we have

$$Q_k[A, D]A \\ = \{f(A, D)A \mid f(A, D) \in Q_k[A, D]\} \\ = \left\{ f(A, D)A \mid f(A, D) = A^{i_1} D^{j_1} \cdots A^{i_k} D^{j_k}, i_1, j_1, \dots, i_k, j_k \in \mathbb{N}, \right. \\ \left. \sum_{m=1}^k i_m + (h+1) \sum_{m=1}^k j_m = k \right\},$$

$$\begin{aligned}
&= \left\{ g(A, D) \mid g(A, D) = A^{i_1} D^{j_1} \cdots A^{i_{k+1}} D^{j_{k+1}}, i_1, j_1, \dots, i_k, j_k \in \mathbb{N}, \right. \\
&\quad \left. i_{k+1} = 1, j_{k+1} = 0, \sum_{m=1}^{k+1} i_m + (h+1) \sum_{m=1}^{k+1} j_m = k+1 \right\} \\
&\subseteq Q_{k+1}[A, D], \\
&Q_{k-h}[A, D]D, \\
&= \{ f(A, D)D \mid f(A, D) \in Q_{k-h}[A, D] \} \\
&= \left\{ f(A, D)D \mid f(A, D) = A^{i_1} D^{j_1} \cdots A^{i_{k-h}} D^{j_{k-h}}, i_1, j_1, \dots, i_{k-h}, j_{k-h} \in \mathbb{N}, \right. \\
&\quad \left. \sum_{m=1}^{k-h} i_m + (h+1) \sum_{m=1}^{k-h} j_m = k-h \right\} \\
&= \left\{ g(A, D) \mid g(A, D) = A^{i_1} D^{j_1} \cdots A^{i_{k+1}} D^{j_{k+1}}, i_1, j_1, \dots, i_{k-h}, j_{k-h} \in \mathbb{N}, \right. \\
&\quad \left. i_{k-h+1} = \cdots = i_{k+1} = 0, j_{k-h+1} = \cdots = j_k = 0, j_{k+1} = 1, \right. \\
&\quad \left. \sum_{m=1}^{k+1} i_m + (h+1) \sum_{m=1}^{k+1} j_m = k+1 \right\} \\
&\subseteq Q_{k+1}[A, D].
\end{aligned} \tag{2.10}$$

Thus, we have $Q_k[A, D]A \cup Q_{k-h}[A, D]D \subseteq Q_{k+1}[A, D]$.

On the contrary, given any $f(A, D) \in Q_{k+1}[A, D]$, suppose that

$$f(A, D) = A^{i_1} D^{j_1} \cdots A^{i_{k+1}} D^{j_{k+1}}, \tag{2.11}$$

where $i_1, j_1, \dots, i_{k+1}, j_{k+1} \in \mathbb{N}$, $\sum_{m=1}^{k+1} i_m + (h+1) \sum_{m=1}^{k+1} j_m = k+1$.

There are two cases: (i) the final term of $f(A, D)$ is A^{i_m} , $i_m > 0$, $1 \leq m \leq k+1$, then there exists a matrix polynomial $g(A, D)$ such that $f(A, D) = g(A, D)A$, then it is easy to verify that $g(A, D) \in Q_k[A, D]$, and it follows that $f(A, D) \in Q_k[A, D]A$; (ii) the final term of $f(A, D)$ is D^{j_m} , $j_m > 0$, $1 \leq m \leq k+1$, then there exists a matrix polynomial $g(A, D)$ such that $f(A, D) = g(A, D)D$, then it is easy to verify that $g(A, D) \in Q_{k-h}[A, D]$, and it follows that $f(A, D) \in Q_{k-h}[A, D]D$.

Thus, we have $Q_k[A, D]A \cup Q_{k-h}[A, D]D \supseteq Q_{k+1}[A, D]$. Hence, we know that (2.7) holds.

Secondly, since $G_{k+h+1} = G_{k+h}A + G_kD$, we have

$$\begin{aligned}
 G_{k+1+h} &= \sum_{f(A,D) \in Q_{k+h}[A,D]} f(A,D)A + \sum_{f(A,D) \in Q_k[A,D]} f(A,D)D \\
 &= \sum_{g(A,D) \in Q_{k+h}[A,D] \cup Q_k[A,D]} g(A,D) \\
 &= \sum_{g(A,D) \in Q_{k+h+1}[A,D]} g(A,D).
 \end{aligned} \tag{2.12}$$

Thus, (2.5) holds for $k + h + 1$. Hence, statement (b) holds.

As for statement (c), it can be deduced naturally from statement (b). Note that if $A^{i_1}D^{j_1} \dots A^{i_k}D^{j_k} \in Q_k[A, D]$, then $A^{i_1}D^{j_1} \dots A^{i_k}D^{j_k} \in Q_k[A, D]$ as well. So they must appear in G_k . By Definition 2.1, $G_k \in P_{\text{Sym}}[A, D]$. $\square$

3. Main Results

First, we consider the general case.

Theorem 3.1. *System (1.1) is controllable if and only if $\text{rank}[G_0B, G_1B, \dots, G_KB, \dots] = n$.*

Proof. From (2.4) and Definition 1.1, this theorem holds naturally. $\square$

Theorem 3.2. *For system (1.1), there exists $K \in \mathbb{N}$ such that*

$$\text{rank}[G_0B, G_1B, \dots, G_KB, \dots] = \text{rank}[G_0B, G_1B, \dots, G_KB]. \tag{3.1}$$

Proof. It is obvious that $\text{rank}[G_0B, G_1B, \dots, G_KB] = \dim(\text{span}\{G_0B, G_1B, \dots, G_KB\})$. Consider an auxiliary scalar sequence $\{d_k\}_{k=0}^\infty$, where $d_k = \dim(\text{Span}\{G_0B, G_1B, \dots, G_kB\})$. By definition, $d_k \leq d_{k+1} \leq n^2$ for any k . Hence, there exists a constant d^* such that, $\lim_{k \rightarrow \infty} d_k = d^*$. Then for $\varepsilon = 0.5$, there exists $K > 0$ such that for all $k > K$, we have $d^* - 0.5 < d_k < d^* + 0.5$. It implies that $d_k = d^*$. $\square$

Remark 3.3. From Theorem 3.2, it is clear that the controllability of the system (1.1) can be realized completely within first K steps during the evolving process, which means K is just a CRI of system (1.1), so the existence of MinCRI of linear discrete-time delay systems is ensured and the value of MinCRI is finite. Moreover, K is dependent on A, D , and h .

Now we consider the second-order case, that is, $n = 2$.

Theorem 3.4. *If $n = 2$, then $2h + 4$ is a CRI of system (1.1).*

Before giving the proof of Theorem 3.4, we first give some lemmas.

Lemma 3.5. *Given $A, D \in \mathbb{R}^{2 \times 2}$, for any $k \in \mathbb{N}$, one has $(AD)^k A, (DA)^k D, (AD)^k + (DA)^k \in \text{span}\{I, A, D, DA + AD, ADA, DAD\}$.*

Proof. According to the different cases, we formulate the proof into two cases.

(a) I, A, D are linearly dependent. Without loss of generality, suppose that

$$A = \lambda_1 I + \lambda_2 D. \quad (3.2)$$

Then we have

$$\begin{aligned} (AD)^k A &= [(\lambda_1 I + \lambda_2 D)D]^k (\lambda_1 I + \lambda_2 D) \\ (DA)^k D &= [D(\lambda_1 I + \lambda_2 D)]^k D \\ (AD)^k + (DA)^k &= [D(\lambda_1 I + \lambda_2 D)D]^k + [D(\lambda_1 I + \lambda_2 D)]^k. \end{aligned} \quad (3.3)$$

It is obvious that $(AD)^k A, (DA)^k D, (AD)^k + (DA)^k \in \text{span}\{I, D\}$.

(b) I, A, D are linearly independent. If $\text{span}\{I, A, D, DA + AD, ADA, DAD\} = \mathbb{R}^{2 \times 2}$, the proof is completed. Otherwise, suppose that $DA + AD, ADA, DAD \in \text{span}\{I, A, D\}$. We use mathematical induction.

(i) For $k = 1$, by the assumption, $(AD)^1 A, (DA)^1 D, AD + DA \in \text{span}\{I, A, D\}$.

(ii) Suppose that for k , $(AD)^k A, (DA)^k D \in \text{span}\{I, A, D\}$. Assume

$$\begin{aligned} (AD)^k A &= \lambda_1 I + \lambda_2 A + \lambda_3 D \\ (DA)^k D &= \lambda_4 I + \lambda_5 A + \lambda_6 D. \end{aligned} \quad (3.4)$$

For $k + 1$, we have

$$\begin{aligned} (AD)^{k+1} A &= A[(DA)^k D] = A(\lambda_4 I + \lambda_5 A + \lambda_6 D)A = \lambda_4 A^2 + \lambda_5 A^3 + \lambda_6 ADA \\ (DA)^{k+1} D &= D(\lambda_1 I + \lambda_2 A + \lambda_3 D)D = \lambda_1 D^2 + \lambda_2 DAD + \lambda_3 D^3 \\ (AD)^{k+1} + (DA)^{k+1} &= (AD)^{k-1} ADAD + (DA)^{k-1} DADA \\ &= (AD)^{k-1} A(\lambda_4 I + \lambda_5 A + \lambda_6 D) + (DA)^{k-1} (\lambda_4 I + \lambda_5 A + \lambda_6 D)A \\ &= \lambda_6 (AD)^k + \lambda_6 (DA)^k + (AD)^{k-1} A(\lambda_4 I + \lambda_5 A) + (DA)^{k-1} (\lambda_4 I + \lambda_5 A)A. \end{aligned} \quad (3.5)$$

It is obvious that $(AD)^{k+1} A, (DA)^{k+1} D, (AD)^{k+1} + (DA)^{k+1} \in \text{span}\{I, A, D\}$. $\square$

Lemma 3.6. Given $A, D \in \mathbb{R}^{2 \times 2}$, one has $\text{span}\{I, A, D, DA + AD, ADA, DAD\} = P_{\text{Sym}}[A, D]$.

Proof. First, it is obvious that

$$\text{span}\{I, A, D, DA + AD, ADA, DAD\} \subseteq P_{\text{Sym}}[A, D]. \quad (3.6)$$

Now we prove that $\text{span}\{I, A, D, DA + AD, ADA, DAD\} \supseteq P_{\text{Sym}}[A, D]$.

For any $f(A, D) \in P_{\text{Sym}}[A, D]$, assume that

$$f(A, D) = \sum_{m=1}^M c_m \left(A^{i_{m,1}} D^{j_{m,1}} \dots A^{i_{m,n_m}} D^{j_{m,n_m}} + D^{j_{m,n_m}} A^{i_{m,n_m}} \dots D^{j_{m,1}} A^{i_{m,1}} \right), \quad (3.7)$$

where $n_m \in \mathbb{N}$, $i_{m,1}, j_{m,1}, \dots, i_{m,n_m}, j_{m,n_m} \in \{0, 1\}$, $c_m \in \mathbb{R}$, $m = 1, \dots, M$; $M \in \mathbb{N}$.

It is easy to see that if we could prove that each

$$\begin{aligned} & A^{i_{m,1}} D^{j_{m,1}} \dots A^{i_{m,n_m}} D^{j_{m,n_m}} + D^{j_{m,n_m}} A^{i_{m,n_m}} \dots D^{j_{m,1}} A^{i_{m,1}} \\ & \in \text{span}\{I, A, D, DA + AD, ADA, DAD\}, \end{aligned} \quad (3.8)$$

then we would have proved that

$$f(A, D) \in \text{span}\{I, A, D, DA + AD, ADA, DAD\}. \quad (3.9)$$

It is easy to verify that

$$A^{i_{m,1}} D^{j_{m,1}} \dots A^{i_{m,n_m}} D^{j_{m,n_m}} + D^{j_{m,n_m}} A^{i_{m,n_m}} \dots D^{j_{m,1}} A^{i_{m,1}} \quad (3.10)$$

can only be rewritten in three different forms:

- (a) $(AD)^k A$, where $k \in \mathbb{N}$;
- (b) $(DA)^k D$, where $k \in \mathbb{N}$;
- (c) $(AD)^k + (DA)^k$, where $k \in \mathbb{N}$.

By Lemma 3.5, we know that (3.8) holds. Thus, we have

$$f(A, D) \in \text{span}\{I, A, D, DA + AD, ADA, DAD\}. \quad (3.11)$$

It follows that

$$P_{\text{Sym}}[A, D] \subseteq \text{span}\{I, A, D, DA + AD, ADA, DAD\}. \quad (3.12)$$

□

Now we are in a position to prove Theorem 3.4.

Proof of Theorem 3.4. By the definition of CRI, we only need to prove that

$$\text{span}\{G_0 B, G_1 B, \dots, G_k B, \dots\} = \text{span}\{G_0 B, G_1 B, \dots, G_{2h+3} B\}. \quad (3.13)$$

By Lemmas 2.2 and 3.6, we have

$$\text{span}\{G_0, G_1, \dots, G_k, \dots\} \subseteq \text{span}\{I, A, D, AD + DA, ADA, DAD\}. \quad (3.14)$$

Now we prove that

$$\text{span}\{G_0, G_1, \dots, G_{2h+3}\} = \text{span}\{I, A, D, AD + DA, ADA, DAD\}. \quad (3.15)$$

By the definition of G_k , we have $G_k = A^k$, $k = 0, 1, \dots, h$, $G_{h+1} = A^{h+1} + D$, $G_{h+2} = A^{h+2} + DA + AD$, $G_{h+3} = A^{h+3} + DA^2 + ADA + A^2D$. Note that $DA^2 + A^2D$ can be linearly expressed by $I, A, D, AD + DA$; then, we have

$$\text{span}\{G_0, G_1, \dots, G_{h+3}\} = \text{span}\{I, A, D, AD + DA, ADA\}. \quad (3.16)$$

Consider the term DAD ; it first appears in G_{2h+3} . It is easy to verify that all other terms in G_{2h+3} can be linearly expressed by $I, A, D, AD + DA, ADA$. Thus, we know that (3.14) holds. This implies that (3.8) holds. $\square$

Without proof, we can get the following corollaries directly.

Corollary 3.7. *Assuming $n = 2$, system (1.1) is controllable if and only if*

$$\text{rank}[B, AB, DB, (AD + DA)B, ADAB, DADB] = 2. \quad (3.17)$$

Corollary 3.8. *Assuming $n = 2$, if system (1.1) is controllable, then the controllability can be realized in $2h + 4$ steps; that is, one can select appropriate $u(0), u(1), \dots, u(2h + 3)$ such that system (1.1) can be driven from any initial state to any terminal state.*

Remark 3.9. Note that the parameter h does not appear in Corollary 3.7. This implies that the specific number of the delay steps has nothing to do with the controllability of the system.

Remark 3.10. Lemma 3.6 plays a key role in the proof of Theorem 3.4. When $n > 2$, it is difficult to build up a similar result like Lemma 3.6 for $n > 2$. Thus the CRI problem for more general case is still open.

4. Examples

In this section, we present some numerical examples to illustrate the validity of our theoretic results.

Example 4.1. Consider the system (1.1) with $n = 2$, $h = 2$, and

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \quad D = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \quad (4.1)$$

By simple calculation, we have $\text{rank}[B, AB, DB, (AD + DA)B, ADAB, DADB] = 2$. By Corollary 3.7, the system should be controllable. In fact, we have

$$x(4) = \Phi(x(0), x(-1), x(-2)) + \begin{bmatrix} I, A, A^2, D \end{bmatrix} \begin{bmatrix} u(3) \\ u(2) \\ u(1) \\ u(0) \end{bmatrix}. \quad (4.2)$$

Letting $u(2) = u(1) = 0$, we can select suitable $u(3)$ and $u(0)$ such that $x(4)$ be any state in $\mathbb{R}^2$. Thus, the system is controllable indeed.

Example 4.2. Consider the system (1.1) with $n = 2$, $h = 2$, and

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad D = A, \quad B = \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \quad (4.3)$$

By simple calculation, we have $\text{rank}[B, AB, DB, (AD + DA)B, ADAB, DADB] = 1$. By Corollary 3.7, the system should not be controllable. In fact, it is easy to see that the second element of the state $x(k)$ is not affected by any input $u(k)$. Thus, the system is not controllable indeed.

5. Conclusion

This paper discussed the controllability of linear discrete-time systems with delay in state. After introducing a new concept called MinCRI to describe the controllability feature of delay systems, we proved the existence and finiteness of MinCRI. Based on this, a necessary and sufficient condition for the controllability of linear delay systems has been derived.

Acknowledgments

The authors are grateful to the Associate Editor and the reviewers for their helpful and valuable comments and suggestions for improving this paper. This work is supported by the National Natural Science Foundation (NNSF) of China (60774089, 10972003, and 60736022). This work is supported by N2008-07 (08010702014) from Beijing Institute of Petrochemical Technology and supported by the Opening Project of Guangxi Key Laboratory of Automobile Components and Vehicle Technology(NO.2012KFZD03).

References

- [1] R. E. Kalman, "On the general theory of control systems," in *Proceedings of the 1st International Conference on Automatic Control*, vol. 1, pp. 481–492, Moscow, Russia, 1960.
- [2] R. E. Kalman, Y. C. Ho, and K. S. Narendra, "Controllability of linear dynamical systems," vol. 1, pp. 189–213, 1963.
- [3] D. H. Chyung, "On the controllability of linear systems with delay in control," *IEEE Transactions on Automatic Control*, vol. 15, no. 2, pp. 255–257, 1970.
- [4] D. H. Chyung, "Controllability of linear systems with multiple delays in control," *IEEE Transactions on Automatic Control*, vol. 15, no. 6, pp. 694–695, 1970.

- [5] K. Watanabe, "Further study of spectral controllability of systems with multiple commensurate delays in state variables," *International Journal of Control*, vol. 39, no. 3, pp. 497–505, 1984.
- [6] V. N. Phat, "Controllability of discrete-time systems with multiple delays on controls and states," *International Journal of Control*, vol. 49, no. 5, pp. 1645–1654, 1989.
- [7] V. N. Phat and T. C. Dieu, "Constrained controllability of linear discrete nonstationary systems in Banach spaces," *SIAM Journal on Control and Optimization*, vol. 30, no. 6, pp. 1311–1318, 1992.
- [8] A. V. Savkin and I. R. Petersen, "Weak robust controllability and observability of uncertain linear systems," *IEEE Transactions on Automatic Control*, vol. 44, no. 5, pp. 1037–1041, 1999.
- [9] V. Y. Glizer, "Euclidean space controllability of singularly perturbed linear systems with state delay," *Systems & Control Letters*, vol. 43, no. 3, pp. 181–191, 2001.
- [10] C. J. Wang and H. E. Liao, "Impulse observability and impulse controllability of linear time-varying singular systems," *Automatica*, vol. 37, no. 11, pp. 1867–1872, 2001.
- [11] P. Chen and H. Qin, "Controllability of linear systems in Banach spaces," *Systems & Control Letters*, vol. 45, no. 2, pp. 155–161, 2002.
- [12] T. Kailath, *Linear Systems*, Prentice-Hall, Englewood Cliffs, NJ, USA, 1980.
- [13] E. D. Sontag, *Mathematical Control Theory: Deterministic Finite-Dimensional Systems*, vol. 6, Springer, New York, NY, USA, 1990.