

## EXISTENCE OF STRONG SOLUTIONS FOR STOCHASTIC POROUS MEDIA EQUATION UNDER GENERAL MONOTONICITY CONDITIONS

BY VIOREL BARBU,<sup>1</sup> GIUSEPPE DA PRATO<sup>2</sup> AND MICHAEL RÖCKNER<sup>3</sup>

*University Al. I. Cuza and Institute of Mathematics “Octav Mayer,”  
Scuola Normale Superiore di Pisa and University of Bielefeld  
and Purdue University*

This paper addresses the existence and uniqueness of strong solutions to stochastic porous media equations  $dX - \Delta\Psi(X) dt = B(X) dW(t)$  in bounded domains of  $\mathbb{R}^d$  with Dirichlet boundary conditions. Here  $\Psi$  is a maximal monotone graph in  $\mathbb{R} \times \mathbb{R}$  (possibly multivalued) with the domain and range all of  $\mathbb{R}$ . Compared with the existing literature on stochastic porous media equations, no growth condition on  $\Psi$  is assumed and the diffusion coefficient  $\Psi$  might be multivalued and discontinuous. The latter case is encountered in stochastic models for self-organized criticality or phase transition.

**1. Introduction.** This work is concerned with existence and uniqueness of solutions to stochastic porous media equations

$$(1.1) \quad \begin{cases} dX(t) - \Delta\Psi(X(t)) dt = B(X(t)) dW(t), & \text{in } (0, T) \times \mathcal{O} := Q_T, \\ \Psi(X(t)) = 0, & \text{on } (0, T) \times \partial\mathcal{O} := \Sigma_T, \\ X(0) = x, & \text{in } \mathcal{O}, \end{cases}$$

where  $\mathcal{O}$  is an open, bounded domain of  $\mathbb{R}^d$ ,  $d \geq 1$ , with smooth boundary  $\partial\mathcal{O}$ ,  $W(t)$  is a cylindrical Wiener process on  $L^2(\mathcal{O})$ , while  $B$  is a Lipschitz continuous operator from  $H := H^{-1}(\mathcal{O})$  to the space of Hilbert–Schmidt operators on  $L^2(\mathcal{O})$ . [See hypothesis  $(H_2)$  below.]

The function  $\Psi : \mathbb{R} \rightarrow \mathbb{R}$  (or more generally the multivalued function  $\Psi : \mathbb{R} \rightarrow 2^{\mathbb{R}}$ ) is a maximal monotone graph in  $\mathbb{R} \times \mathbb{R}$ . (See the definition in Section 1.1 below.)

Existence results for (1.1) were obtained in [8] (see also [3, 4]) in the special case  $B = \sqrt{Q}$ , with  $Q$  linear nonnegative,  $\text{Tr } Q < +\infty$  and  $\Psi \in C^1(\mathbb{R})$  satisfying

---

Received March 2007; revised February 2008.

<sup>1</sup>Supported by PN-II ID.404 (2007–2010).

<sup>2</sup>Supported by the research program “Equazioni di Kolmogorov” from the Italian “Ministero della Ricerca Scientifica e Tecnologica.”

<sup>3</sup>Supported by the SFB-701 and the BIBOS-Research Center.

*AMS 2000 subject classifications.* 76S05, 60H15.

*Key words and phrases.* Stochastic porous media equation, Wiener process, convex functions, Itô’s formula.

the growth condition

$$(1.2) \quad k_3 + k_1 |s|^{r-1} \leq \Psi'(s) \leq k_2(1 + |s|^{r-1}), \quad s \in \mathbb{R},$$

where  $k_1, k_2 > 0, k_3 \in \mathbb{R}, r > 1$ .

Under these growth conditions on  $\Psi$ , (1.1) covers many important models describing the dynamics of an ideal gas in a porous medium (see, e.g., [1]) but excludes, however, other significant physical models such as plasma fast diffusion [5], which arises for  $\Psi(s) = \sqrt{s}$ , and phase transitions or dynamics of saturated underground water flows (the Richards equation). In the latter case, multivalued monotone graphs  $\Psi$  might appear as in [12]. Recently, in [15] (see also [14]), the existence results of [8] were extended to the case of monotone nonlinearities  $\Psi$  such that  $s \mapsto s\Psi(s)$  is (comparable to) a  $\Delta_2$ -regular Young function (cf. assumption (A1) in [15]) thus including the fast diffusion model. As a matter of fact, in the line of the classical work of Krylov and Rozovskii [10] the approach used in [15] is a variational one, that is, one considers the stochastic equation (1.1) in a duality setting induced by a functional triplet  $V \subset H \subset V'$  and this requires one to find appropriate spaces  $V$  and  $H$ . This was done in [15] in an elaborate way even with  $\mathcal{O}$  unbounded and with  $\Delta$  replaced by very general (not necessarily differential) operators  $L$ .

The method we use here is quite different—essentially an  $L^1$ -approach relying on weak compactness techniques in  $L^1(Q_T)$  via the Dunford–Pettis theorem which involve minimal growth assumptions on  $\Psi$ . Restricted to single-valued continuous functions  $\Psi$  the main result, Theorem 2.2 below, gives existence and uniqueness of solutions only assuming that  $\lim_{s \rightarrow +\infty} \Psi(s) = +\infty, \lim_{s \rightarrow -\infty} \Psi(s) = -\infty, \Psi$  monotonically increasing and

$$(1.3) \quad \limsup_{|s| \rightarrow +\infty} \frac{\int_0^{-s} \Psi(t) dt}{\int_0^s \Psi(t) dt} < +\infty.$$

We note that the assumptions on  $\Psi$  in [15] imply our assumptions. In this sense, under assumption  $(H_2)$  below on the noise, the results of this paper extend those in [15] in case  $L = \Delta$  if  $\mathcal{O}$  is bounded and if the coefficients do not depend on  $(t, \omega)$ . The latter two were not assumed in [15]. On the other hand, a growth condition on  $\Psi$  is imposed in [15] (cf. [15], Lemma 3.2) which is not done here. Another main point of this paper is that  $\Psi$  is no longer assumed to be continuous, it might be multivalued and with exponential growth to  $\pm\infty$  [for instance, of the form  $\exp(a|x|^p)$ ]. We note that (1.3) is not a growth condition at  $+\infty$  but a kind of symmetry condition about the behavior of  $\Psi$  at  $\pm\infty$ . If  $\Psi$  is a maximal monotone graph with potential  $j$  (i.e.,  $\Psi = \partial j$ ) then (1.3) takes the form [see hypothesis  $(H_3)$  below]

$$\limsup_{|s| \rightarrow +\infty} \frac{j(-s)}{j(s)} < +\infty.$$

Anyway this condition is automatically satisfied for even monotonically increasing functions  $\Psi$  or, for example, if a condition of the form (1.2) is satisfied. We note,

however, that because of our very general conditions on  $\Psi$  the solution of (1.1) will be pathwise only weakly continuous in  $H$ . The question of pathwise strong continuity of solutions, however, remains open. The main reason is the absence of a variational setting for problem (1.1) (see [10, 14]) in the present situation. Another major technical difficulty encountered here in the proofs is that the integration by parts formula or Itô’s formula (see Lemmas 3.1 and 3.2 below) cannot be applied directly because of the same reason. It should be said, however, that the  $L^1$  approach used here, which allows us to treat very general nonlinearities, is applicable to deterministic equations as well and seems to be new also in that context. On the other hand, the existence for the deterministic part of (1.1) is an immediate consequence of the Crandall–Liggett generation theorem for nonlinear semigroups of contractions (see [2]) which is, however, not applicable to stochastic equations.

1.1. *Notation.*  $\mathcal{O}$  is a bounded open subset of  $\mathbb{R}^d$ ,  $d \geq 1$ , with smooth boundary  $\partial\mathcal{O}$ . We set

$$Q_T = (0, T) \times \mathcal{O}, \quad \Sigma_T = (0, T) \times \partial\mathcal{O}.$$

$L^p(\mathcal{O})$ ,  $L^p(Q_T)$ ,  $p \geq 1$ , are standard  $L^p$ -function spaces and  $H_0^1(\mathcal{O})$ ,  $H^k(\mathcal{O})$  are Sobolev spaces on  $\mathcal{O}$ . By  $H := H^{-1}(\mathcal{O})$  we denote the dual of  $H_0^1(\mathcal{O})$  with the norm and the scalar product given by

$$|u|_{-1} := (A^{-1}u, u)^{1/2}, \quad \langle u, v \rangle_{-1} = (A^{-1}u, v),$$

respectively, where  $(\cdot, \cdot)$  is the pairing between  $H_0^1(\mathcal{O})$  and  $H^{-1}$  and the scalar product of  $L^2(\mathcal{O})$ . Here  $A$  denotes the Laplace operator with Dirichlet homogeneous boundary conditions, that is,

$$(1.4) \quad Au = -\Delta u, \quad u \in D(A) = H^2(\mathcal{O}) \cap H_0^1(\mathcal{O}).$$

Given a Hilbert space  $U$ , the norm of  $U$  will be denoted by  $|\cdot|_U$  and the scalar product by  $(\cdot, \cdot)_U$ . By  $C([0, T]; U)$  we shall denote the space of  $U$ -valued continuous functions on  $[0, T]$  and by  $C^w([0, T]; U)$  the space of weakly continuous functions from  $[0, T]$  to  $U$ .

Given two Hilbert spaces  $U$  and  $V$  we shall denote the space of linear continuous operators from  $U$  to  $V$  by  $L(U, V)$  and the space of Hilbert–Schmidt operators  $F : U \rightarrow V$  by  $L_{HS}(U, V)$ , with norm

$$(1.5) \quad \|F\|_{L_{HS}(U, V)} := \left( \sum_{i=1}^{\infty} |F e_i|_V^2 \right)^{1/2},$$

where  $\{e_i\}$  is an orthonormal basis in  $U$ .

If  $j : \mathbb{R} \rightarrow (-\infty, +\infty]$  is a lower semicontinuous convex function let  $\partial j : \mathbb{R} \rightarrow 2^{\mathbb{R}}$  denote the subdifferential of  $j$ , that is,

$$\partial j(y) = \{\theta \in \mathbb{R} : j(y) \leq j(z) + \theta(y - z) \forall z \in \mathbb{R}\}$$

and let  $j^*$  denote the conjugate of  $j$  (the Legendre transform of  $j$ ),

$$j^*(p) = \sup\{py - j(y) : y \in \mathbb{R}\}.$$

We recall that  $\partial j^* = (\partial j)^{-1}$  (see, e.g., [2, 6]),

$$(1.6) \quad j(y) + j^*(p) = py \quad \text{if and only if} \quad p \in \partial j(y)$$

and

$$(1.7) \quad j(u) + j^*(p) \geq pu \quad \text{for all } p, u \in \mathbb{R}.$$

Given a multivalued function  $\Phi : \mathbb{R} \rightarrow 2^{\mathbb{R}}$ , its domain is denoted by  $D(\Phi) = \{u \in \mathbb{R} : \Phi(u) \neq \emptyset\}$ .  $R(\Phi) = \{v : v \in \Phi(u), u \in D(\Phi)\}$  is its range. The function  $\Phi$  is said to be a maximal monotone graph if it is monotone, that is,

$$(y_1 - y_2)(p_1 - p_2) \geq 0 \quad \text{for all } p_i \in \Phi(y_i), i = 1, 2$$

and  $R(1 + \Phi) = \mathbb{R}$ .

Given a maximal monotone graph  $\Psi : \mathbb{R} \rightarrow 2^{\mathbb{R}}$ , there is a unique lower semicontinuous convex function  $j : \mathbb{R} \rightarrow (-\infty, +\infty]$  such that  $\Psi := \partial j$ . The function  $j$  is unique up to an additive constant and is called the *potential* of  $\Psi$ .

For the maximal monotone graph  $\Psi$  we set

$$\Psi_\lambda = \frac{1}{\lambda}(1 - (1 + \lambda\Psi)^{-1}) \in \Psi(1 + \lambda\Psi)^{-1}, \quad \lambda > 0,$$

which is called the *Yosida* approximation of  $\Psi$ . Here 1 stands for the identity function. The function  $\Psi_\lambda$  is Lipschitzian and monotonically increasing.

We set  $j_\lambda(u) = \int_0^u \Psi_\lambda(r) dr$  and recall that it is equal to the Moreau approximation of  $j$ , that is,

$$(1.8) \quad j_\lambda(u) = \min \left\{ j(v) + \frac{1}{2\lambda}|u - v|^2 : v \in \mathbb{R} \right\}.$$

We have

$$(1.9) \quad j_\lambda(u) = j((1 + \lambda\Psi)^{-1}u) + \frac{1}{2\lambda}|u - (1 + \lambda\Psi)^{-1}u|^2.$$

**2. The main result.**

2.1. *Hypotheses.*  $(H_1)$   $W(t)$  is a cylindrical Wiener process on  $L^2(\mathcal{O})$  defined by

$$(2.1) \quad W(t) = \sum_{k=1}^{\infty} \beta_k(t)e_k,$$

where  $\{\beta_k\}$  is a sequence of mutually independent Brownian motions on a filtered probability space  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$ , with right-continuous filtration and  $\{e_k\}$  is

an orthonormal basis in  $L^2(\mathcal{O})$ . To be more specific,  $\{e_k\}$  will be chosen as the normalized sequence of eigenfunctions of the operator  $A$ , hence  $e_k \in L^p(\mathcal{O})$  for all  $k \in \mathbb{N}$ ,  $p \geq 1$ .

( $H_2$ )  $B$  is Lipschitzian from  $H = H^{-1}(\mathcal{O})$  to  $L_{HS}(L^2(\mathcal{O}), D(A^\gamma))$ , where  $\gamma > d/2$ .

( $H_3$ )  $\Psi : \mathbb{R} \rightarrow 2^{\mathbb{R}}$  is a maximal monotone graph on  $\mathbb{R} \times \mathbb{R}$  such that  $0 \in \Psi(0)$ ,

$$(2.2) \quad D(\Psi) = \mathbb{R}, \quad R(\Psi) = \mathbb{R}$$

and

$$(2.3) \quad \limsup_{|s| \rightarrow +\infty} \frac{j(-s)}{j(s)} < +\infty.$$

Here  $j : \mathbb{R} \rightarrow \mathbb{R}$  is the potential of  $\Psi$ , that is,  $\partial j = \Psi$ , which under assumption (2.2) is a continuous convex function. Since  $0 \in \Psi(0)$ , by definition we have  $j(0) = \inf j$ . Hence subtracting  $j(0)$  we can take  $j$  such that  $j(0) = 0$  and  $j \geq 0$  and therefore we may assume that  $j^* \geq j^*(0) = 0$ . We recall (see, e.g., [2, 6]) that the condition  $R(\Psi) = \mathbb{R}$  is equivalent to

$$(2.4) \quad j(y) < \infty \quad \forall y \in \mathbb{R} \quad \lim_{|y| \rightarrow \infty} \frac{j(y)}{|y|} = +\infty,$$

while the condition  $D(\Psi) = \mathbb{R}$  is equivalent to

$$(2.5) \quad j^*(y) < \infty \quad \forall y \in \mathbb{R} \quad \lim_{|y| \rightarrow \infty} \frac{j^*(y)}{|y|} = +\infty.$$

In particular, hypothesis ( $H_3$ ) automatically holds if  $\Psi$  is a monotonically increasing, continuous function on  $\mathbb{R}$  satisfying condition (1.3) and

$$\lim_{s \rightarrow +\infty} \Psi(s) = +\infty, \quad \lim_{s \rightarrow -\infty} \Psi(s) = -\infty.$$

In particular, it is satisfied by functions  $\Psi$  satisfying (1.2) for  $r > 0$  or, more generally, by those satisfying assumption (A1) in [15].

We need more notation. Given a Banach space  $Z$ ,

$$C_W([0, T]; Z) = C([0, T]; L^2(\Omega, \mathcal{F}, \mathbb{P}; Z))$$

shall denote the space of all continuous adapted stochastic processes which are mean square continuous. The space

$$L^2_W([0, T]; Z) = L^2([0, T]; L^2(\Omega, \mathcal{F}, \mathbb{P}; Z))$$

is similarly defined (see, e.g., [7, 9]).

**DEFINITION 2.1.** An adapted process  $X \in C_W([0, T]; H) \cap L^1((0, T) \times \mathcal{O} \times \Omega)$ , such that  $X \in C^w([0, T], H)$ ,  $\mathbb{P}$ -a.s., is said to be a strong solution to (1.1) if

there exists a process  $\eta \in L^1((0, T) \times \mathcal{O} \times \Omega)$  such that

$$(2.6) \quad \eta(t, \xi) \in \Psi(X(t, \xi)) \quad \text{a.e. } (t, \xi) \in Q_T, \mathbb{P}\text{-a.s.},$$

$$(2.7) \quad \int_0^\bullet \eta(s) ds \in C^w([0, T]; H_0^1(\mathcal{O})),$$

$$(2.8) \quad X(t) - \Delta \int_0^t \eta(s) ds = x + \int_0^t B(X(s)) dW(s) \quad \forall t \in [0, T], \mathbb{P}\text{-a.s.},$$

$$(2.9) \quad j(X), j^*(\eta) \in L^1((0, T) \times \mathcal{O} \times \Omega).$$

[Here  $\int_0^t \eta(s) ds$  is initially defined as on  $L^1(\mathcal{O})$ -valued Bochner integral.] Of course, if  $\Psi$  is single-valued (2.6)–(2.8) reduce to

$$(2.10) \quad \int_0^\bullet \Psi(X(s)) ds \in C^w([0, T]; H_0^1(\mathcal{O}))$$

and

$$(2.11) \quad X(t) - \Delta \int_0^t \Psi(X(s)) ds = x + \int_0^t B(X(s)) dW(s) \quad \forall t \in [0, T], \mathbb{P}\text{-a.s.}$$

We note that  $X$ , as in Definition 2.1, is automatically predictable.

Theorem 2.2 below is the main result of this work.

**THEOREM 2.2.** *Under hypotheses  $(H_1)$ ,  $(H_2)$ ,  $(H_3)$ , for each  $x \in H$  there is a unique strong solution  $X = X(t, x)$  to (1.1). Moreover, the following estimate holds:*

$$(2.12) \quad \mathbb{E}|X(t, x) - X(t, y)|_{-1}^2 \leq C|x - y|_{-1}^2 \quad \text{for all } t \geq 0,$$

where  $C$  is independent of  $x, y \in H$ .

Theorem 2.2 will be proved in Section 4 via fixed-point arguments. Before, in Section 3 we shall establish the existence of solutions for the equation

$$(2.13) \quad \begin{cases} dY(t) - \Delta \Psi(Y(t)) dt = G(t) dW(t), & \text{in } Q_T, \\ \Psi(Y(t)) = 0, & \text{on } \Sigma_T, \\ Y(0) = x, & \text{in } \mathcal{O}, \end{cases}$$

where  $G : [0, T] \rightarrow L_{HS}(L^2(\mathcal{O}), D(A^\gamma))$  is a predictable process such that

$$(2.14) \quad \mathbb{E} \int_0^T \|G(t)\|_{L_{HS}(L^2(\mathcal{O}), D(A^\gamma))}^2 dt < +\infty$$

and  $\gamma > d/2$ . Here  $G dW$  is given by

$$G dW = \sum_{k=1}^\infty G e_k d\beta_k.$$

A solution of (2.13) is defined to be an adapted process  $Y$  satisfying along with  $\eta \in L^1((0, T) \times \mathcal{O} \times \Omega)$  conditions (2.6)–(2.9) where  $B(X)$  is replaced by  $G$ .

**THEOREM 2.3.** *Under hypotheses  $(H_1)$ ,  $(H_3)$ , (2.14), for each  $x \in H$  there is a unique strong solution  $Y = Y_G(t, x)$  to (2.13) in the sense of Definition 2.1. Moreover, the following estimate holds:*

$$(2.15) \quad \mathbb{E}|Y_{G_1}(t, x) - Y_{G_2}(t, y)|_{-1}^2 \leq |x - y|_{-1}^2 + \mathbb{E} \int_0^t \|G_1(s) - G_2(s)\|_{L_{HS}(L^2(\mathcal{O}), H)}^2 ds$$

*for all  $t \geq 0$*

*for all  $x, y \in H$  and  $G_1, G_2$  satisfying (2.14).*

**REMARK 2.4.** It should be noted that assumption  $(H_2)$  excludes the case where the covariance operator  $B$  is of the form  $B(X) = X$ , that is, the case of multiplicative noise.

**REMARK 2.5.** Assumption  $(H_3)$ , for example, allows monotonically increasing functions  $\Psi$  which are continuous from the right on  $\mathbb{R}$  and have a finite number of jumps  $r_1, r_2, \dots, r_N$ . However in this case one must fill the jumps by replacing the function  $\Psi$  by the maximal monotone (multivalued) graph  $\tilde{\Psi}(r) = \Psi(r)$  for  $r$  different from  $r_i$  and  $\tilde{\Psi}(r_i) = [\Psi(r_i) - \Psi(r_{i-1}) - 0]$ . Such a situation might arise in modeling underground water flows (see, e.g., [12]). In this case  $\Psi$  is the diffusivity function and (1.1) reduces to the Richards equation. It must also be said that Theorems 2.2 and 2.3 have natural extensions to equations of the form

$$(2.16) \quad dX(t) - \Delta \Psi(X(t)) dt + \Phi(X(t)) dt = B(X(t)) dW(t),$$

where  $\Phi$  is a suitable monotonically increasing and continuous function (see [15]). We note also that stochastic models of self-organized criticality lead to equations of form (1.1) where  $\Psi(r) = H(r - x_c)$  and  $H$  is the Heaviside function.

As in [15], one might consider the case where  $\Psi = \Psi(X, \omega)$ ,  $\omega \in \Omega$ , but we do not go into details here. We also note that assumption  $D(\Psi) = \mathbb{R}$  in hypothesis  $(H_3)$  excludes a situation of the following type:

$$(2.17) \quad \Psi(s) = \begin{cases} \Psi_1(s), & \text{for } s < s_0, \Psi(s_0) = (\Psi_1(s_0), +\infty), \\ \emptyset, & \text{for } s > s_0, \end{cases}$$

where  $\Psi_1$  is a continuous monotonically increasing function and  $\Psi_1(r) \rightarrow -\infty$  as  $r \rightarrow -\infty$ . In this case problem (1.1) reduces to a stochastic variational inequality and it is relevant in the description of saturation processes in infiltration. An analysis similar to that to be developed below shows that, in Definition 2.1, the function  $\eta$  is not an  $L^1$ -function but rather a bounded measure on  $\mathcal{Q}_T$  and  $\int_0^\bullet \eta(s) ds$  must be replaced by its distribution function. We plan to give details in a subsequent paper.

Another situation of interest covered by our assumptions (see also [15]) is that of logarithmic diffusion equations arising in plasma physics; for example, see [13]. In this case  $\Psi(s) = \log(\mu + |s|) \text{sign}(s)$ .

**3. Proof of Theorem 2.3.** For every  $\lambda > 0$  consider the approximating equation

$$(3.1) \quad \begin{cases} dX_\lambda(t) - \Delta(\Psi_\lambda(X_\lambda(t)) + \lambda X_\lambda(t)) dt \\ \quad = G(t) dW(t), & \text{in } (0, T) \times \mathcal{O} := Q_T, \\ \Psi_\lambda(X_\lambda(t)) + \lambda X_\lambda(t) = 0, & \text{on } (0, T) \times \partial\mathcal{O}, \\ X_\lambda(0) = x, & \text{in } \mathcal{O}, \end{cases}$$

which has a unique solution  $X_\lambda \in C_W([0, T]; H)$  such that

$$X_\lambda, \Psi_\lambda(X_\lambda) \in L^2_W(0, T; H_0^1(\mathcal{O})).$$

Indeed, setting  $y_\lambda(t) = X_\lambda(t) - W_G(t)$ , where  $W_G(t) = \int_0^t G(s) dW(s)$ , we may rewrite (3.1) as a random equation

$$(3.2) \quad \begin{cases} y'_\lambda(t) - \Delta\tilde{\Psi}_\lambda(y_\lambda(t) + W_G(t)) = 0, & \mathbb{P}\text{-a.s. in } Q_T, \\ \tilde{\Psi}_\lambda(y_\lambda(t) + W_G(t)) = 0, & \text{on } (0, T) \times \partial\mathcal{O}, \\ y_\lambda(0) = x, & \text{in } \mathcal{O}, \end{cases}$$

where  $\tilde{\Psi}_\lambda(y) = \Psi_\lambda(y) + \lambda y, \lambda > 0$ . Note that  $\tilde{\Psi}_\lambda(0) = 0$ .

For each  $\omega \in \Omega$  the operator  $\Gamma(t) : H_0^1(\mathcal{O}) \rightarrow H^{-1}(\mathcal{O})$ , defined by

$$\Gamma(t)y = -\Delta\tilde{\Psi}_\lambda(y + W_G(t)), \quad y \in H_0^1(\mathcal{O}),$$

is continuous, monotone and coercive, that is,

$$\begin{aligned} (\Gamma(t)y, y) &\geq \lambda|y + W_G(t)|_{H_0^1(\mathcal{O})}^2 - (\Gamma(t)y, W_G(t)) \\ &\geq \frac{\lambda}{2}|y|_{H_0^1(\mathcal{O})}^2 - C_\lambda|W_G(t)|_{H_0^1(\mathcal{O})}^2. \end{aligned}$$

Then, by classical existence theory for nonlinear equations (see, e.g., [11]), (3.2) has a unique solution

$$y_\lambda \in C([0, T]; L^2(\mathcal{O})) \cap L^2(0, T; H_0^1(\mathcal{O}))$$

with  $y'_\lambda \in L^2(0, T; H^{-1}(\mathcal{O}))$ . The function  $X_\lambda(t) = y_\lambda(t) + W_G(t)$  is, of course, an adapted process because the solution  $y_\lambda$  to (3.2) is a continuous function of  $W_G$  and so it satisfies the required condition.

3.1. *A priori estimates.* From now on we fix  $\omega \in \Omega$  and work with the corresponding solution  $y_\lambda$  to (3.2). We have

$$(3.3) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} |y_\lambda(t)|_{-1}^2 + (\tilde{\Psi}_\lambda(y_\lambda(t) + W_G(t)), y_\lambda(t) + W_G(t)) \\ = (\tilde{\Psi}_\lambda(y_\lambda(t) + W_G(t)), W_G(t)), \end{aligned}$$

which is equivalent to

$$(3.4) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} |y_\lambda(t)|_{-1}^2 + (\Psi_\lambda(y_\lambda(t) + W_G(t)), y_\lambda(t) + W_G(t)) \\ = -\lambda(y_\lambda(t), y_\lambda(t) + W_G(t)) + (\Psi_\lambda(y_\lambda(t) + W_G(t)), W_G(t)). \end{aligned}$$



Now set  $j_\lambda(u) = \int_0^u \Psi_\lambda(r) dr$  and let  $j_\lambda^*$  denote the conjugate of  $j_\lambda$ . Moreover, we set

$$(3.5) \quad z_\lambda = (1 + \lambda\Psi)^{-1}(y_\lambda + W_G), \quad \eta_\lambda = \Psi_\lambda(j_\lambda + W_G).$$

The aim of this subsection is to prove the following estimate: There exists a (random) constant  $C_1$  such that

$$(3.6) \quad \begin{aligned} & \frac{1}{2}|y_\lambda(t)|_{-1}^2 + \int_0^t \int_{\mathcal{O}} (j(z_\lambda) + j^*(\eta_\lambda)) d\xi ds \\ & + \frac{1}{2\lambda} \int_0^t \int_{\mathcal{O}} (y_\lambda + W_G - z_\lambda)^2 d\xi ds \leq C_1(1 + |x|_{-1}^2), \quad t \in [0, T]. \end{aligned}$$

By (1.6) we have

$$(3.7) \quad \begin{aligned} & j_\lambda^*(\Psi_\lambda(y_\lambda(t) + W_G(t))) + j_\lambda(y_\lambda(t) + W_G(t)) \\ & = \Psi_\lambda(y_\lambda(t) + W_G(t))(y_\lambda(t) + W_G(t)). \end{aligned}$$

Substituting this identity into (3.4) yields

$$(3.8) \quad \begin{aligned} & \frac{1}{2}|y_\lambda(t)|_{-1}^2 + \int_0^t \int_{\mathcal{O}} (j_\lambda(y_\lambda(s) + W_G(s)) + j_\lambda^*(\Psi_\lambda(y_\lambda(s) + W_G(s)))) d\xi ds \\ & = \frac{1}{2}|x|_{-1}^2 + \int_0^t \int_{\mathcal{O}} (\Psi_\lambda(y_\lambda(s) + W_G(s))W_G(s)) d\xi ds \\ & \quad - \lambda \int_0^t \int_{\mathcal{O}} y_\lambda(s)(y_\lambda(s) + W_G(s)) d\xi ds. \end{aligned}$$

Then, using (1.9), (3.5) and the fact that  $j_\lambda^* \geq j^*$  for all  $\lambda > 0$ , by (3.8) we see that

$$(3.9) \quad \begin{aligned} & \frac{1}{2}|y_\lambda(t)|_{-1}^2 + \int_0^t \int_{\mathcal{O}} (j(z_\lambda(s)) + j^*(\eta_\lambda(s))) d\xi ds \\ & \quad + \frac{1}{2\lambda} \int_0^t \int_{\mathcal{O}} (y_\lambda(s) + W_G(s) - z_\lambda(s))^2 d\xi ds \\ & \leq \frac{1}{2}|x|_{-1}^2 + \int_0^t \int_{\mathcal{O}} \eta_\lambda(s)W_G(s) d\xi ds \\ & \quad - \lambda \int_0^t \int_{\mathcal{O}} y_\lambda(s)(y_\lambda(s) + W_G(s)) d\xi ds. \end{aligned}$$

We now estimate the first integral from the right-hand side of (3.9) as follows:

$$(3.10) \quad \left| \int_0^t \int_{\mathcal{O}} \eta_\lambda(s)W_G(s) d\xi ds \right| \leq \delta \int_0^t \int_{\mathcal{O}} |\eta_\lambda(s)| d\xi ds,$$

where  $\delta := \sup_{s \in [0, T]} |W_G(s)|_{L^\infty(\mathcal{O})} < +\infty$ . We note that by assumption (2.14) and since  $\gamma > d/2$  it follows by Sobolev embedding that  $W_G(\cdot)$  has continuous sample paths in  $D(A^\nu) \subset L^\infty(\mathcal{O})$  and so  $\delta$  is indeed finite.

Substituting (3.10) in (3.9) yields

$$\begin{aligned} & \frac{1}{2}|y_\lambda(t)|_{-1}^2 + \int_0^t \int_{\mathcal{O}} (j(z_\lambda(s)) + j^*(\eta_\lambda(s))) d\xi ds \\ & + \frac{1}{2\lambda} \int_0^t \int_{\mathcal{O}} (y_\lambda(s) + W_G - z_\lambda(s))^2 d\xi ds \\ & \leq \frac{1}{2}|x|_{-1}^2 + \delta \int_0^t \int_{\mathcal{O}} |\eta_\lambda(s)| d\xi ds - \lambda \int_0^t \int_{\mathcal{O}} y_\lambda(s)(y_\lambda(s) + W_G(s)) d\xi ds. \end{aligned}$$

Since

$$-y_\lambda(s)(y_\lambda(s) + W_G(s)) \leq -\frac{1}{2}|y_\lambda(s)|^2 + \frac{1}{2}W_G^2(s),$$

we find

$$\begin{aligned} & \frac{1}{2}|y_\lambda(t)|_{-1}^2 + \int_0^t \int_{\mathcal{O}} (j(z_\lambda(s)) + j^*(\eta_\lambda(s))) d\xi ds \\ & + \frac{\lambda}{2} \int_0^t \int_{\mathcal{O}} |y_\lambda(s)|^2 d\xi ds \\ (3.11) \quad & + \frac{1}{2\lambda} \int_0^t \int_{\mathcal{O}} (y_\lambda(s) + W_G(s) - z_\lambda(s))^2 d\xi ds \\ & \leq \left( \frac{1}{2}|x|_{-1}^2 + \delta \int_0^t \int_{\mathcal{O}} |\eta_\lambda(s)| d\xi ds + \frac{\lambda}{2} \int_0^t \int_{\mathcal{O}} W_G^2(s) d\xi ds \right), \\ & \hspace{20em} t \in [0, T]. \end{aligned}$$

On the other hand, we recall that condition  $D(\Psi) = \mathbb{R}$  is equivalent with

$$(3.12) \quad j^* < \infty \quad \text{and} \quad \lim_{|p| \rightarrow \infty} \frac{j^*(p)}{|p|} = +\infty.$$

So, there exists  $N = N(\omega)$  such that

$$|\eta_\lambda(s)| > N \quad \Rightarrow \quad j^*(\eta_\lambda(s)) > 2C\delta|\eta_\lambda(s)|.$$

Consequently, for  $C > |Q_T|$  we have that

$$\begin{aligned} \int_0^t \int_{\mathcal{O}} |\eta_\lambda(s)| d\xi ds &= \int \int_{|\eta_\lambda(s)| > N} |\eta_\lambda(s)| d\xi ds + \int \int_{|\eta_\lambda(s)| \leq N} |\eta_\lambda(s)| d\xi ds \\ &\leq \frac{1}{2C\delta} \int_0^t \int_{\mathcal{O}} j^*(\eta_\lambda(s)) d\xi ds + NC\delta. \end{aligned}$$

Substituting this into (3.11), since  $j \geq 0$ , we obtain (3.6), which in particular implies

$$(3.13) \quad \int_0^t \int_{\mathcal{O}} (j(z_\lambda(s)) + j^*(\eta_\lambda(s))) d\xi ds \leq C_1(1 + |x|_{-1}^2)$$

and

$$(3.14) \quad \int_0^t \int_{\mathcal{O}} (y_\lambda + W_G - z_\lambda)^2 d\xi ds \leq 2\lambda C_1(1 + |x|_{-1}^2).$$

3.2. *Convergence for  $\lambda \rightarrow 0$ .* Since by (2.4) and (2.5)

$$(3.15) \quad \lim_{|u| \rightarrow \infty} j(u)/|u| = \infty, \quad \lim_{|u| \rightarrow \infty} j^*(u)/|u| = \infty,$$

we deduce from (3.13) that the sequences  $\{z_\lambda\}$  and  $\{\eta_\lambda\}$  are bounded and equi-integrable in  $L^1(Q_T)$ . Then by the Dunford–Pettis theorem the sequences  $\{z_\lambda\}$  and  $\{\eta_\lambda\}$  are weakly compact in  $L^1(Q_T)$ . Hence along a subsequence, again denoted by  $\lambda$ , we have

$$(3.16) \quad z_\lambda \rightharpoonup z, \quad \eta_\lambda \rightharpoonup \eta \quad \text{weakly in } L^1(Q_T) \text{ as } \lambda \rightarrow 0.$$

Moreover, by (3.14) we see that  $z = y + W_G$ , where

$$(3.17) \quad y_\lambda \rightharpoonup y \quad \text{weakly}^* \text{ in } L^\infty(0, T; H) \text{ and weakly in } L^1(Q_T).$$

Also note that by (3.2) we have for every  $t \in [0, T]$

$$(3.18) \quad y_\lambda(t) - \Delta \left( \int_0^t (\eta_\lambda(s) + \lambda(y_\lambda(s) + W_G(s))) ds \right) = x$$

and so the sequence  $\{\int_0^\bullet (\eta_\lambda(s) + \lambda y_\lambda(s)) ds\}$  is bounded in  $L^\infty(0, T; H_0^1(\mathcal{O}))$ . Hence, selecting a further subsequence if necessary [see (3.8)], we have

$$(3.19) \quad \lim_{\lambda \rightarrow 0} \int_0^\bullet (\eta_\lambda(s) + \lambda y_\lambda(s)) ds = \int_0^\bullet \eta(s) ds$$

weakly\* in  $L^\infty(0, T; H_0^1(\mathcal{O}))$ .

So, by (3.18) we find

$$(3.20) \quad y(t) + A \int_0^t \eta(s) ds = x \quad \text{a.e. } t \in [0, T].$$

Since

$$\int_0^\bullet \eta(s) ds \in C([0, T]; L^1(\mathcal{O})) \cap L^\infty(0, T; H_0^1(\mathcal{O})),$$

$t \mapsto \int_0^t \eta(s) ds$  is weakly continuous in  $H_0^1(\mathcal{O})$  and therefore we infer that so is  $t \mapsto A \int_0^t \eta(s) ds$  in  $H$ . Thus the function

$$(3.21) \quad \tilde{y}(t) := -A \int_0^t \eta(s) ds + x, \quad t \in [0, T],$$

is an  $H$ -valued weakly continuous version of  $y$ . Furthermore, we claim that for  $\lambda \rightarrow 0$

$$y_\lambda(t) \rightharpoonup \tilde{y}(t) \quad \text{weakly in } H \forall t \in [0, T].$$

Indeed, since  $\eta_\lambda \rightharpoonup \eta$  weakly in  $L^1(Q_T)$  and  $\lambda(y_\lambda + W_G) \rightarrow 0$  weakly in  $L^1(Q_T)$  [since it even converges strongly in  $L^2(Q_T)$  to zero by (3.11)], it follows that for every  $t \in [0, T]$

$$\int_0^t (\eta_\lambda(s) + \lambda(y_\lambda(s) + W_G(s))) ds \rightharpoonup \int_0^t \eta(s) ds \quad \text{weakly in } L^1(\mathcal{O}).$$

Hence by (3.18) and the definition of  $\tilde{\eta}$  we obtain that for every  $t \in [0, T]$

$$(-\Delta)^{-1}y_\lambda(t) \rightarrow (-\Delta)^{-1}\tilde{y}(t) \quad \text{weakly in } L^1(\mathcal{O}).$$

Since by (3.11)  $y_\lambda(t)$ ,  $\lambda > 0$ , are bounded in  $H$ , the above immediately implies the claim.

From now on we shall consider this particular version  $\tilde{y}$  of  $y$  defined in (3.21). For simplicity we denote it again by  $y$ ; so we have

$$y_\lambda(t) \rightarrow y(t) \quad \text{weakly in } H \quad \forall t \in [0, T].$$

We can also rewrite (3.21) as

$$(3.22) \quad y(t) - \Delta\eta(t) = 0 \quad \text{in } \mathcal{D}'(Q_T), \quad y(0) = x.$$

Now we are going to show that

$$(3.23) \quad \eta(t, \xi) \in \Psi(y(t, \xi) + W_G(t, \xi)) \quad \text{a.e. } (t, \xi) \in Q_T.$$

For this we shall need the following inequality:

$$(3.24) \quad \liminf_{\lambda \rightarrow 0} \int_{Q_T} y_\lambda \eta_\lambda \, d\xi \, dt \leq \int_{Q_T} y \eta \, d\xi \, dt.$$

3.2.1. *Proof of (3.24).* We first recall (1.6), which yields

$$j_\lambda(y_\lambda + W_G) + j_\lambda^*(\eta_\lambda) = (y_\lambda + W_G)\eta_\lambda \quad \text{a.e. in } Q_T$$

and so, by (1.9) and since  $j_\lambda^* \geq j^*$ , we have

$$j(y_\lambda + W_G) + j^*(\eta_\lambda) \leq (y_\lambda + W_G)\eta_\lambda \quad \text{a.e. in } Q_T,$$

which yields

$$\int_{Q_T} (j(y_\lambda + W_G) + j^*(\eta_\lambda)) \, d\xi \, dt \leq \int_{Q_T} (y_\lambda + W_G)\eta_\lambda \, d\xi \, dt.$$

Since the convex functional

$$(z, \zeta) \rightarrow \int_{Q_T} (j(z) + j^*(\zeta)) \, d\xi \, dt$$

is lower semicontinuous on  $L^1(Q_T) \times L^1(Q_T)$  (and consequently weakly lower semicontinuous on this space) we obtain that

$$(3.25) \quad \begin{aligned} & \int_{Q_T} (j(y + W_G) + j^*(\eta)) \, d\xi \, dt \\ & \leq \liminf_{\lambda \rightarrow 0} \int_{Q_T} y_\lambda \eta_\lambda \, d\xi \, dt + \int_{Q_T} W_G \eta \, d\xi \, dt. \end{aligned}$$

Furthermore, by (3.6) and again by the weak lower semicontinuity of convex integrals in  $L^1(Q_T)$ , it follows that

$$(3.26) \quad j(y + W_G), \quad j^*(\eta) \in L^1(Q_T).$$

On the other hand, since  $j(u) + j^*(p) \geq up$  for all  $u, p \in \mathbb{R}$  [see (1.7)], we have

$$(3.27) \quad (W_G + y)\eta \leq j(y + W_G) + j^*(\eta) \quad \text{a.e. in } Q_T.$$

Moreover, by assumption (2.3) we see that for every  $M > 0$  there exists  $R = R(M) \geq 0$ , such that

$$j(-y - W_G) \leq Mj(y + W_G) \quad \text{on } Q^R,$$

where

$$Q^R = \{(t, \xi) \in Q_T : |y(t, \xi) + W_G(t, \xi)| \geq R\}.$$

Since  $j(y + W_G) \in L^1(Q_T)$  we have, by continuity of  $j$ ,

$$(3.28) \quad j(-y - W_G) \leq h \quad \text{a.e. in } Q_T,$$

where  $h \in L^1(Q_T)$ . On the other hand, since  $j$  is bounded from below we have

$$(3.29) \quad j(-y - W_G) \in L^1(Q_T).$$

Taking into account that by virtue of the same inequality (1.7), besides (3.27), we have that

$$(3.30) \quad -(y + W_G)\eta \leq j(-y - W_G) + j^*(\eta) \quad \text{a.e. in } Q_T,$$

by (3.27) and (3.28) it follows that a.e. in  $Q_T$  we have

$$|(W_G + y)\eta| \leq \max\{j(y + W_G) + j^*(\eta), j(-y - W_G) + j^*(\eta)\} \in L^1(Q_T)$$

and therefore  $y\eta \in L^1(Q_T)$  as claimed [recall that  $W_G \in L^\infty(Q_T)$ ].

Now we come back to (3.4), which by integration yields

$$(3.31) \quad \frac{1}{2}(|y_\lambda(T)|_{-1}^2 - |x|_{-1}^2) + \int_{Q_T} y_\lambda \eta_\lambda d\xi dt + \lambda \int_{Q_T} y_\lambda (y_\lambda + W_G) d\xi dt = 0.$$

Taking into account that

$$(3.32) \quad y_\lambda(T) \rightarrow y(T) \quad \text{weakly in } H,$$

by (3.31) we have that

$$(3.33) \quad \liminf_{\lambda \rightarrow 0} \int_{Q_T} y_\lambda \eta_\lambda d\xi dt \leq -\frac{1}{2}(|y(T)|_{-1}^2 - |x|_{-1}^2).$$

In order to complete the proof one needs an integration by parts formula in (3.21) or (3.22), obtained by multiplying the equation by  $y$  and integrating on  $Q_T$ . Formally this is possible because  $y\eta \in L^1(Q_T)$  and  $y(t) \in H^{-1}(\mathcal{O})$  for all  $t \in [0, T]$ . But, in order to prove it rigorously, one must give sense to  $(y'(t), y(t))$ . Lemma 3.1 below answers this question positively and by (3.33) also proves (3.24).

We first note that since  $j, j^*$  are nonnegative and convex and  $j(0) = 0 = j^*(0)$ , we have for all measurable  $f : Q_T \rightarrow \mathbb{R}$  and  $\alpha \in [0, 1]$ ,

$$j(f) \in L^1(Q_T) \quad \Rightarrow \quad j(\alpha f) \in L^1(Q_T)$$

and

$$j^*(f) \in L^1(Q_T) \Rightarrow j^*(\alpha f) \in L^1(Q_T).$$

Furthermore, as in the proof of (3.28) by (2.3) we obtain that

$$j(f) \in L^1(Q_T) \Rightarrow j(-f) \in L^1(Q_T).$$

By (2.3) we see that the latter is also true for  $j^*$ , if  $f \in L^1(Q_T)$  and  $\alpha$  is small enough. Indeed by (2.3) there are  $M, R > 0$  such that

$$j(-s) \leq Mj(s), \quad \text{if } |s| \geq R,$$

hence replacing  $s$  by  $(-s)$  we get

$$\frac{1}{M}j(s) \leq j(-s), \quad \text{if } |s| \geq R.$$

Now an elementary calculation implies that for all  $p \in \mathbb{R}$

$$j^*(-p) \leq R|p| + \frac{1}{M}j^*(Mp).$$

Hence

$$j^*(-p/M) \leq \frac{R}{M}|p| + \frac{1}{M}j^*(p).$$

Therefore for  $\alpha := 1/M$  we have

$$0 \leq j^*(-\alpha f) \leq \frac{R}{M}|f| + \frac{1}{M}j^*(f) \in L^1(Q_T).$$

Hence,  $y$  and  $\eta$  constructed above fulfill all conditions in the following lemma since  $W_G \in L^\infty(Q_T)$ .

LEMMA 3.1. *Let  $y \in C^w([0, T]; H^{-1}(\mathcal{O})) \cap L^1(Q_T)$  and  $\eta \in L^1(Q_T) \cap L^\infty(0, T; H^1(\mathcal{O}))$  satisfy*

$$(3.34) \quad y(t) + A \int_0^t \eta(s) ds = x, \quad t \in [0, T].$$

*Furthermore, assume that for some  $\alpha > 0$ ,  $j(\alpha y), j^*(\alpha \eta) \in L^1(Q_T)$ . Then  $y\eta \in L^1(Q_T)$ ,*

$$(3.35) \quad \int_{Q_T} y\eta d\xi dt = -\frac{1}{2}(|y(T)|_{L^1}^2 - |x|_{L^1}^2)$$

and

$$Y_\varepsilon \Sigma_\varepsilon \rightarrow y\eta \quad \text{in } L^1(Q_T),$$

where  $Y_\varepsilon, \Sigma_\varepsilon$  are defined in (3.36) below.

PROOF. We set for  $\varepsilon > 0$

$$(3.36) \quad Y_\varepsilon = (1 + \varepsilon A)^{-m} y, \quad \Sigma_\varepsilon = (1 + \varepsilon A)^{-m} \eta,$$

where  $m \in \mathbb{N}$  is such that  $m > \max\{2, (d + 2)/4\}$ . Then

$$Y_\varepsilon \in C^w([0, T]; H_0^1(\mathcal{O}) \cap H^{2m-1}(\mathcal{O})) \subset C^w([0, T]; H_0^1(\mathcal{O}) \cap C(\overline{\mathcal{O}}))$$

and

$$\Sigma_\varepsilon \in L^1(0, T; W^{2,q}(\mathcal{O})), \quad 1 < q < \frac{d}{d-1}.$$

Hence  $Y_\varepsilon \Sigma_\varepsilon \in L^1(Q_T)$  and for  $\varepsilon \rightarrow 0$

$$(3.37) \quad \begin{cases} Y_\varepsilon(t) \rightarrow y(t), & \text{strongly in } H^{-1}(\mathcal{O}) \forall t \in [0, T], \\ Y_\varepsilon \rightarrow y, & \text{strongly in } L^1(Q_T), \\ \Sigma_\varepsilon \rightarrow \eta, & \text{strongly in } L^1(Q_T), \\ \int_0^t \Sigma_\varepsilon(s) ds \rightarrow \int_0^t \eta(s) ds, & \text{strongly in } H_0^1(\mathcal{O}) \forall t \in [0, T]. \end{cases}$$

Here we note that the last fact follows because (3.34) implies that  $\int_0^\bullet \eta(s) ds \in C^w([0, T]; H_0^1(\mathcal{O}))$ . We have also by (3.34)

$$Y_\varepsilon(t) + A \int_0^t \Sigma_\varepsilon(s) ds = (1 + \varepsilon A)^{-m} x \quad \forall t \in [0, T],$$

which implies

$$\frac{d}{dt} Y_\varepsilon(t) + A \Sigma_\varepsilon(t) = 0$$

and, taking the inner product in  $H^{-1}(\mathcal{O})$  with  $Y_\varepsilon(t)$ , we obtain

$$\frac{1}{2} \frac{d}{dt} |Y_\varepsilon(t)|_{-1}^2 + \int_{\mathcal{O}} \Sigma_\varepsilon(t) Y_\varepsilon(t) d\xi = 0 \quad \text{a.e. } t \in [0, T].$$

Hence

$$(3.38) \quad \lim_{\varepsilon \rightarrow 0} \int_{Q_T} \Sigma_\varepsilon(t) Y_\varepsilon(t) d\xi dt = -\frac{1}{2} (|y(T)|_{-1}^2 - |x|_{-1}^2)$$

and by (3.37) we may assume that for  $\varepsilon \rightarrow 0$

$$(3.39) \quad Y_\varepsilon \rightarrow y, \quad \Sigma_\varepsilon \rightarrow \eta \quad \text{a.e. in } Q_T.$$

Moreover by (1.7) we have

$$(3.40) \quad \alpha^2 \Sigma_\varepsilon Y_\varepsilon \leq j(\alpha Y_\varepsilon) + j^*(\alpha \Sigma_\varepsilon), \quad -\alpha^2 \Sigma_\varepsilon Y_\varepsilon \leq j(-\alpha Y_\varepsilon) + j^*(\alpha \Sigma_\varepsilon) \quad \text{a.e. in } Q_T.$$

Now we claim that for  $\varepsilon \rightarrow 0$

$$(3.41) \quad \begin{aligned} j(\alpha Y_\varepsilon) &\rightarrow j(\alpha y), & j^*(\alpha \Sigma_\varepsilon) &\rightarrow j^*(\alpha \eta), \\ j(-\alpha Y_\varepsilon) &\rightarrow j(-\alpha y) & \text{in } L^1(Q_T). \end{aligned}$$

By (3.39) these convergences hold a.e. in  $Q_T$ . So, in order to prove (3.41) it suffices to show that  $\{j(\alpha Y_\varepsilon)\}, \{j^*(\alpha \Sigma_\varepsilon)\}, \{j(-\alpha Y_\varepsilon)\}$  are equi-integrable on  $Q_T$  and that they are weakly compact in  $L^1(Q_T)$ . To this end let  $y \in L^1(\mathcal{O})$  and let  $Y_\varepsilon := (1 + \varepsilon A)^{-1}y$ , that is,  $Y_\varepsilon$  is the solution to the Dirichlet problem

$$(3.42) \quad \begin{cases} Y_\varepsilon - \varepsilon \Delta Y_\varepsilon = y, & \text{in } \mathcal{O}, \\ Y_\varepsilon = 0, & \text{on } \partial \mathcal{O}. \end{cases}$$

It may be represented as

$$(3.43) \quad Y_\varepsilon(\xi) = \int_{\mathcal{O}} G(\xi, \xi_1) y(\xi_1) d\xi_1 \quad \forall \xi \in \mathcal{O},$$

where  $G$  is the associated Green function. It is well known that  $\int_{\mathcal{O}} G(\xi, \xi_1) d\xi_1$  is the solution to (3.42) with  $y = 1$  so that by the maximum principle we have  $0 < \int_{\mathcal{O}} G(\xi, \xi_1) d\xi_1 \leq 1$  for all  $\xi \in \mathcal{O}$ .

We may rewrite  $Y_\varepsilon$  as

$$Y_\varepsilon(\xi) = \int_{\mathcal{O}} G(\xi, \xi_2) d\xi_2 \int_{\mathcal{O}} \tilde{G}(\xi, \xi_1) y(\xi_1) d\xi_1 \quad \forall \xi \in \mathcal{O},$$

where

$$\tilde{G}(\xi, \xi_1) = \frac{G(\xi, \xi_1)}{\int_{\mathcal{O}} G(\xi, \xi_2) d\xi_2}$$

and so  $\int_{\mathcal{O}} \tilde{G}(\xi, \xi_1) d\xi_1 = 1$  for all  $\xi \in \mathcal{O}$ .

Then, if  $j(y) \in L^1(\mathcal{O})$ ,  $j(0) = 0$  by Jensen's inequality we have

$$\begin{aligned} j(Y_\varepsilon(\xi)) &\leq \int_{\mathcal{O}} G(\xi, \xi_2) d\xi_2 \int_{\mathcal{O}} \tilde{G}(\xi, \xi_1) j(y(\xi_1)) d\xi_1 \\ &= \int_{\mathcal{O}} G(\xi, \xi_1) j(y(\xi_1)) d\xi_1 \quad \forall \xi \in \mathcal{O}. \end{aligned}$$

Hence we have shown that for any  $y \in L^1(\mathcal{O})$  with  $j(y) \in L^1(\mathcal{O})$ ,

$$j((1 + \varepsilon A)^{-1}y) \leq (1 + \varepsilon A)^{-1}j(y).$$

Iterating and using the fact that  $(1 + \varepsilon A)^{-1}$  preserves positivity we get for all  $m \in \mathbb{N}$

$$(3.44) \quad j((1 + \varepsilon A)^{-m}y) \leq (1 + \varepsilon A)^{-m}j(y) \quad \text{a.e. in } \mathcal{O}.$$

Now let  $y$  be as in the assertion of the lemma and  $Y_\varepsilon$  as in (3.36). Integrating over  $Q_T$ , since  $(1 + \varepsilon A)^{-m}$  is a contraction on  $L^1(\mathcal{O})$ , (3.44) applied to  $\alpha y$  implies

$$\int_{Q_T} j(\alpha Y_\varepsilon(\xi, t)) d\xi dt \leq \int_{Q_T} j(\alpha y(\xi_2, t)) d\xi_2 dt.$$



Taking into account that  $j(\alpha y) \in L^1(Q_T)$  we infer that  $\{j(\alpha Y_\varepsilon)\}$  is equi-integrable on  $Q_T$ . The same argument applies to  $\{j^*(\alpha \Sigma_\varepsilon)\}, \{j(-\alpha Y_\varepsilon)\}$ .

Then (3.40) implies that the sequence  $\{\Sigma_\varepsilon Y_\varepsilon\}$  is equi-integrable on  $Q_T$  and consequently by the Dunford–Pettis theorem, weakly compact in  $L^1(Q_T)$ . Since  $\{\Sigma_\varepsilon Y_\varepsilon\}$  is a.e. convergent to  $y\eta$  we infer that for  $\varepsilon \rightarrow 0$

$$(3.45) \quad \Sigma_\varepsilon Y_\varepsilon \rightarrow y\eta \quad \text{strongly in } L^1(Q_T),$$

which, combined with (3.38), implies (3.35) as desired.  $\square$

3.2.2. *Proof of (3.23).* We have

$$j(z_\lambda) - j(u) \leq \eta_\lambda(z_\lambda - u) \quad \forall u \in \mathbb{R} \text{ a.e. in } Q_T.$$

Integrating over  $Q_T$  yields

$$\int_{Q_T} j(z_\lambda) d\xi dt \leq \int_{Q_T} j(u) d\xi dt + \int_{Q_T} \eta_\lambda(z_\lambda - u) d\xi dt \quad \forall u \in L^\infty(Q_T).$$

Note that by the definition of  $\Psi_\lambda$  we have

$$z_\lambda = -\lambda\eta_\lambda + y_\lambda + W_G.$$

Therefore, since  $z = y + W_G$ , by (3.24) and Fatou’s lemma we can let  $\lambda \rightarrow 0$  to obtain

$$\int_{Q_T} j(z) d\xi dt - \int_{Q_T} j(u) d\xi dt \leq \int_{Q_T} \eta(z - u) d\xi dt \quad \forall u \in L^\infty(Q_T).$$

Now by Lusin’s theorem for each  $\epsilon > 0$  there is a compact subset  $Q_\epsilon \subset Q_T$  such that  $(d\xi \otimes dt)(Q_T \setminus Q_\epsilon) \leq \epsilon$  and  $y, \eta$  are continuous on  $Q_\epsilon$ . Let  $(t_0, x_0)$  be a Lebesgue point for  $y, \eta$  and  $y\eta$  and let  $B_r$  be the ball of center  $(t_0, x_0)$  and radius  $r$ . We take

$$u(t, \xi) = \begin{cases} z(t, \xi), & \text{if } (t, \xi) \in Q_\epsilon \cap B_r^c, \\ v, & \text{if } (t, \xi) \in (Q_\epsilon \cap B_r) \cup (Q_T \setminus Q_\epsilon). \end{cases}$$

Here  $v$  is arbitrary in  $\mathbb{R}$ . Since  $u$  is bounded we can substitute into the above inequality to get

$$\int_{B_r \cap Q_\epsilon} (j(z) - j(v) - \eta(z - v)) d\xi dt \leq \int_{Q_T \setminus Q_\epsilon} (\eta(z - v) + j(v) - j(z)) d\xi dt.$$

Letting  $\epsilon \rightarrow 0$  we obtain that

$$\int_{B_r} (j(z) - j(v) - \eta(z - v)) d\xi dt \leq 0 \quad \forall v \in \mathbb{R}, r > 0.$$

This yields

$$j(z(t_0, x_0)) \leq j(v) + \eta(t_0, x_0)(z(t_0, x_0) - v) \quad \forall v \in \mathbb{R}$$

and therefore  $\eta(t_0, x_0) \in \partial j(z(t_0, x_0)) = \Psi(z(t_0, x_0))$ . Since almost all points of  $Q_T$  are Lebesgue points we get (3.23) as claimed.

3.3. *Completion of proof of Theorem 2.3.* Let us first summarize what we have proved for the pair  $(y, \eta) \in L^1(Q_T) \times L^1(Q_T)$ . We have

$$\begin{aligned}
 &y \in C^w([0, T]; H), \quad \int_0^\bullet \eta(s) ds \in C^w([0, T]; H_0^1(\mathcal{O})), \\
 &\eta(t, \xi) \in \Psi(y(t, \xi)) \quad \text{for a.e. } (t, \xi) \in Q^T, \\
 &y(t) + A \int_0^t \eta(s) ds = x, \quad t \in [0, T], \\
 &j(\alpha y), j^*(\alpha y) \in L^1(Q_T) \quad \text{for some } \alpha \in (0, 1].
 \end{aligned}$$

We claim that  $(y, \eta)$  is the only such pair. Indeed, if  $(\tilde{y}, \tilde{\eta})$  is another, then

$$j\left(\frac{\alpha}{2}(y - \tilde{y})\right) \leq \frac{1}{2}j(\alpha y) + \frac{1}{2}j(-\alpha \tilde{y})$$

and

$$j^*\left(\frac{\alpha}{2}(y - \tilde{y})\right) \leq \frac{1}{2}j^*(\alpha y) + \frac{1}{2}j^*(-\alpha \tilde{y}).$$

But as we have seen before Lemma 3.1 the right-hand sides are in  $L^1(Q_T)$ . Hence  $y - \tilde{y}, \eta - \tilde{\eta}$  fulfill all conditions of Lemma 3.1 and adopting the notation from there we have for  $\varepsilon > 0$

$$\begin{aligned}
 Y_\varepsilon(t) - \tilde{Y}_\varepsilon(t) &= \Delta \int_0^t (\Sigma_\varepsilon(s) - \tilde{\Sigma}_\varepsilon(s)) ds \\
 &= \int_0^t \Delta(\Sigma_\varepsilon(s) - \tilde{\Sigma}_\varepsilon(s)) ds, \quad t \in [0, T].
 \end{aligned}$$

Differentiating and subsequently taking the inner product in  $H$  with  $Y_\varepsilon(t) - \tilde{Y}_\varepsilon(t)$  and integrating again we arrive at

$$\begin{aligned}
 &\frac{1}{2} |(1 + \varepsilon A)^{-m}(Y_\varepsilon(t) - \tilde{Y}_\varepsilon(t))|_{-1}^2 \\
 &= \int_0^t \int_{\mathcal{O}} (Y_\varepsilon(s) - \tilde{Y}_\varepsilon(s))(\Sigma_\varepsilon(s) - \tilde{\Sigma}_\varepsilon(s)) d\xi ds \\
 &= \int_0^t \int_{\mathcal{O}} (1 + \varepsilon A)^{-m}(y(s) - \tilde{y}(s))(1 + \varepsilon A)^{-m}(\eta(s) - \tilde{\eta}(s)) d\xi ds, \\
 & \hspace{20em} t \in [0, T].
 \end{aligned}$$

Letting  $\varepsilon \rightarrow 0$  and applying Lemma 3.1 we obtain that for  $t \in [0, T]$

$$\frac{1}{2} |y(t) - \tilde{y}(t)|_{-1}^2 = \int_0^t \int_{\mathcal{O}} (y(s) - \tilde{y}(s))(\eta(s) - \tilde{\eta}(s)) d\xi ds \leq 0$$

by the monotonicity of  $\Psi$ .

Now let us consider the  $\omega$ -dependence of  $y$  and  $\eta$ . By (3.21), (3.23) we know that  $y = y(t, \xi, \omega)$  is the solution to the equation

$$(3.46) \quad \begin{cases} y'(t) - \Delta \Psi(y(t) + W_G(t)(\omega)) = 0, & \text{a.e. } t \in [0, T], \\ y(0) = x \end{cases}$$

and as seen earlier for  $\eta = \eta(t, \xi, \omega)$  as in (3.16)

$$(3.47) \quad \begin{aligned} y &\in C^w([0, T]; H) \cap L^1(Q_T), & \eta &\in L^1(Q_T), \\ \int_0^\bullet \eta(s) ds &\in C^w([0, T]; H_0^1(\mathcal{O})) \end{aligned}$$

and

$$(3.48) \quad \eta(t, \xi, \omega) \in \Psi(y(t, \xi, \omega)) + W_G(t, \xi, \omega) \quad \text{a.e. } (t, \xi, \omega) \in Q_T \times \Omega.$$

By the above uniqueness of  $(y, \eta)$ , it follows that for any sequence  $\lambda \rightarrow \infty$  we have  $\mathbb{P}$ -a.s.

$$\begin{aligned} y_\lambda(t) &\rightarrow y(t) && \text{weakly in } H = H^{-1}(\mathcal{O}) \forall t \in [0, T], \\ y_\lambda &\rightarrow y && \text{weakly in } L^1(Q_T), \\ \int_0^t \eta_\lambda(s) ds &\rightarrow \int_0^t \eta(s) ds && \text{weakly in } L^1(\mathcal{O}) \forall t \in [0, T] \\ &&& \text{and weakly in } H_0^1(\mathcal{O}) \text{ a.e. } t \in [0, T], \\ \eta_\lambda &\rightarrow \eta && \text{weakly in } L^1(Q_T). \end{aligned}$$

Therefore  $y$  and  $\eta$  are strong  $L^1(Q_T)$ -limits of a sequence of convex combinations of  $y_\lambda, \eta_\lambda$ , respectively, and since  $y_\lambda$  and  $\eta_\lambda$  are predictable processes, it follows that so are  $y$  and  $\eta$ . In particular, this means that  $Y(t) = y(t) + W_G(t)$  is an  $H$ -valued weakly continuous adapted process and that the following equation is satisfied:

$$(3.49) \quad Y(t) - \Delta \int_0^t \eta(s) ds = x + \int_0^t G(s) dW(s), \quad t \in [0, T].$$

Equivalently

$$(3.50) \quad \begin{cases} dY(t) - \Delta \Psi(Y(t)) dt = G(t) dW(t), \\ Y(0) = x. \end{cases}$$

In order to prove that  $Y$  is a solution of (3.50) in the sense of Definition 2.1 with  $G(t)$  replacing  $B(X(t))$  and to prove uniqueness and some energy estimates for solutions to (3.50) we need an Itô's formula-type result. As in the case of Lemma 3.1 the difficulty is that the integral

$$\int_{Q_T} \Psi(Y) Y d\xi dt$$

might not be (in general) well defined, taking into account that  $\Psi(Y), Y \in L^1(Q_T)$  only. We have, however, the following.

LEMMA 3.2. *Let  $Y$  be an  $H$ -valued weakly continuous adapted process satisfying (3.49). Then the following equality holds:*

$$\begin{aligned}
 \frac{1}{2}|Y(t)|_{-1}^2 &= \frac{1}{2}|x|_{-1}^2 - \int_0^t \int_{\mathcal{O}} \eta(s)Y(s) d\xi ds \\
 (3.51) \qquad &+ \int_0^t \langle Y(s), G(s) dW(s) \rangle_{-1} \\
 &+ \frac{1}{2} \int_0^t \|G(s)\|_{L_{HS}(L^2(\mathcal{O}),H)}^2 ds, \qquad \mathbb{P}\text{-a.s.}
 \end{aligned}$$

Furthermore,  $Y \in C_W([0, T]; H) \cap L^1((0, T) \times \mathcal{O} \times \Omega)$ , and  $\eta \in L^1((0, T) \times \mathcal{O} \times \Omega)$  and all conditions (2.6)–(2.9) are satisfied.

PROOF. By Lemma 3.1 we have that  $Y\eta \in L^1(Q_T)$ . Next we introduce the sequences (see the proof of Lemma 3.1) for  $m \in \mathbb{N}$

$$Y_\varepsilon = (1 + \varepsilon A)^{-m} Y, \qquad \Sigma_\varepsilon = (1 + \varepsilon A)^{-m} \eta.$$

For large enough  $m$  we can apply Itô’s formula to the problem

$$\begin{cases} dY_\varepsilon(t) + A\Sigma_\varepsilon(t) = (1 + \varepsilon A)^{-m} G dW(t), \\ Y_\varepsilon(0) = (1 + \varepsilon A)^{-m} x = x_\varepsilon. \end{cases}$$

We have

$$\begin{aligned}
 \frac{1}{2}|Y_\varepsilon(t)|_{-1}^2 &= \frac{1}{2}|x_\varepsilon|_{-1}^2 - \int_0^t \int_{\mathcal{O}} \Sigma_\varepsilon(s)Y_\varepsilon(s) d\xi ds \\
 (3.53) \qquad &+ \int_0^t \langle Y_\varepsilon(s), G_\varepsilon(s) dW(s) \rangle_{-1} \\
 &+ \frac{1}{2} \int_0^t \|G_\varepsilon(s)\|_{L_{HS}(L^2(\mathcal{O}),H)}^2 ds, \qquad t \in [0, T],
 \end{aligned}$$

where  $G_\varepsilon = (1 + \varepsilon A)^{-m} G$ . Letting  $\varepsilon \rightarrow 0$  [since  $W_G \in L^\infty(Q_T)$ ] we get by (3.45)

$$\int_{Q_T} Y_\varepsilon \Sigma_\varepsilon d\xi ds \rightarrow \int_{Q_T} Y \eta d\xi ds, \qquad \mathbb{P}\text{-a.s.}$$

Furthermore

$$Y_\varepsilon(t) \rightarrow Y(t) \qquad \text{strongly in } H^{-1}(\mathcal{O}) \forall t \in [0, T],$$

which by virtue of (3.53) yields (3.51), if we prove first that for  $t \in [0, T]$

$$(3.54) \qquad \mathbb{P} - \lim_{\varepsilon \rightarrow 0} \int_0^t \langle Y_\varepsilon(s), G_\varepsilon(s) dW(s) \rangle = \int_0^t \langle Y(s), G(s) dW(s) \rangle.$$

We shall even show that this convergence in probability is locally uniform in  $t$ . We have by a standard consequence of the Burkholder–Davis–Gundy inequality for

$p = 1$  (see, e.g., [14], Corollary D-0.2) that for  $\bar{Y}_\varepsilon := (1 + \varepsilon A)^{-2m} Y$  and  $\delta_1, \delta_2 > 0$

$$(3.55) \quad \mathbb{P} \left[ \sup_{t \in [0, T]} \left| \int_0^t \langle Y(s), G(s) dW(s) \rangle - \int_0^t \langle Y_\varepsilon(s), G_\varepsilon(s) dW(s) \rangle \right| \geq \delta_1 \right] \leq \frac{3\delta_2}{\delta_1} + \mathbb{P} \left[ \int_0^T \|G(s)\|_{L_{HS}(L^2(\mathcal{O}), H)}^2 |Y(s) - \bar{Y}_\varepsilon(s)|_{-1}^2 ds \geq \delta_2 \right].$$

Since  $Y \in C^w([0, T]; H)$ ,  $\mathbb{P}$ -a.s. and  $(1 + \varepsilon A)^{-1}$  is a contraction on  $H$  we have

$$\sup_{s \in [0, T]} |Y(s) - \bar{Y}_\varepsilon(s)|_{-1} \leq 2 \sup_{s \in [0, T]} |Y(s)|_{-1}^2, \quad \mathbb{P}\text{-a.s.}$$

Hence by (2.14) the second term on the right-hand side of (3.55) converges to zero as  $\varepsilon \rightarrow 0$ . Taking subsequently  $\delta_2 \rightarrow 0$ , (3.55) implies (3.54). We emphasize that, since the left-hand side of (3.51) is not continuous  $\mathbb{P}$ -a.s. (though all terms on the right-hand side are), the  $\mathbb{P}$ -zero set of  $\omega \in \Omega$  for which (3.51) does not hold might depend on  $t$ .

Next we want to prove that

$$(3.56) \quad \mathbb{E} \left[ \sup_{t \in [0, T]} |Y(t)|_{-1}^2 \right] < \infty.$$

To this end first note that by (3.48) and (1.6) we have

$$(3.57) \quad \eta(s)Y(s) = j(Y(s)) + j^*(\eta(s)) \geq 0,$$

hence (3.51) implies that for every  $t \in [0, T]$

$$(3.58) \quad |Y(t)|_{-1}^2 \leq |x|_{-1}^2 + N_t + \int_0^t \|G(s)\|_{L_{HS}(L^2(\mathcal{O}), H)}^2 ds, \quad \mathbb{P}\text{-a.s.},$$

where

$$N_t := \int_0^t \langle Y(s), G(s) dW(s) \rangle_{-1}, \quad t \geq 0,$$

is a continuous local martingale such that

$$\langle N \rangle_t = 2 \int_0^t |G^*(s)Y(s)|_{L^2(\mathcal{O})}^2 ds, \quad t \geq 0,$$

where  $G^*(s)$  is the adjoint of  $G(s) : L^2(\mathcal{O}) \rightarrow H$ . We shall prove that

$$(3.59) \quad \mathbb{E} \left[ \sup_{t \in [0, T]} |N_t| \right] < +\infty.$$

By the Burkholder–Davis–Gundy inequality for  $p = 1$  applied to the stopping times

$$\tau_N := \inf\{t \geq 0 : |N_t| \geq N\} \wedge T, \quad N \in \mathbb{N},$$

we obtain

$$\begin{aligned}
 & \mathbb{E} \left[ \sup_{t \in [0, \tau_N]} |N_t| \right] \\
 (3.60) \quad & \leq 3E \left[ \sup_{s \in [0, \tau_N]} |Y(s)|_{-1} \left( 4 \int_0^{\tau_N} \|G(s)\|_{L_{HS}(L^2(\mathcal{O}); H)}^2 ds \right)^{1/2} \right] \\
 & \leq 6C \left( \mathbb{E} \left[ \sup_{s \in [0, \tau_N]} |Y(s)|_{-1}^2 \right] \right)^{1/2},
 \end{aligned}$$

where

$$C := \left( \mathbb{E} \left[ \int_0^T \|G(s)\|_{L_{HS}(L^2(\mathcal{O}); H)}^2 ds \right] \right)^{1/2} < \infty.$$

Since  $Y \in C^w([0, T]; H)$ , we know that  $s \mapsto |Y(s)|_{-1}^2$  is lower semicontinuous. Therefore by (3.58)

$$\begin{aligned}
 \sup_{s \in [0, \tau_N]} |Y(s)|_{-1}^2 &= \sup_{s \in [0, \tau_N] \cap \mathbb{Q}} |Y(s)|_{-1}^2 \\
 &\leq |x|_{-1}^2 + \sup_{s \in [0, \tau_N]} |N_s| + \int_0^T \|G(s)\|_{L_{HS}(L^2(\mathcal{O}); H)}^2 ds, \quad \mathbb{P}\text{-a.s.}
 \end{aligned}$$

So (3.60) implies that for all  $N \in \mathbb{N}$

$$\left( \mathbb{E} \left[ \sup_{t \in [0, \tau_N]} |N_t| \right] \right)^2 \leq 36C^2 \left[ |x|_{-1}^2 + \mathbb{E} \left[ \sup_{s \in [0, \tau_N]} |N_s| \right] + C^2 \right],$$

which entails that

$$\sup_{N \in \mathbb{N}} \mathbb{E} \left[ \sup_{t \in [0, \tau_N]} |N_t| \right] < \infty.$$

By monotone convergence this implies (3.59), since  $N_t$  has continuous sample paths and  $\tau_N \uparrow T$  as  $N \rightarrow \infty$ . Now (3.58) implies that (4.11) holds.

By (3.59), (3.57) and (3.51) it follows that

$$(3.61) \quad \eta Y \in L^1((0, T) \times \mathcal{O} \times \Omega).$$

Hence by (3.57)

$$j(Y), j^*(\eta) \in L^1((0, T) \times \mathcal{O} \times \Omega)$$

and therefore

$$Y, \eta \in L^1((0, T) \times \mathcal{O} \times \Omega).$$

Taking the expectation in (3.51) we see that  $t \mapsto \mathbb{E}[|Y(t)|_{-1}^2]$  is continuous. Since  $Y \in C^w([0, T]; H)$ ,  $\mathbb{P}$ -a.s., (3.61) then also implies that  $Y \in C_w([0, T]; H)$ . This in turn, together with (3.49), also implies that (2.7) holds.  $\square$

Now we come back to the proof of Theorem 2.3. We first note that Lemma 3.2 also implies the uniqueness of the solution  $Y$  and estimate (2.15). Indeed by (3.55) and the monotonicity of  $\Psi$  we have for  $YG_i, i = 1, 2$ , the estimate (2.15). This concludes the proof of Theorem 2.3.

**4. Proof of Theorem 2.2.** Consider the space

$$(4.1) \quad \mathcal{K} = \left\{ X \in C_W([0, T]; H) \cap L^1((0, T) \times \mathcal{O} \times \Omega) : X \text{ predictable,} \right. \\ \left. \sup_{t \in [0, T]} \mathbb{E}[e^{-2\alpha t} |X(t)|_{-1}^2] \leq M_1^2, \mathbb{E} \int_{Q_T} j(X(s)) d\xi ds \leq M_2 \right\},$$

where  $\alpha > 0, M_1 > 0$  and  $M_2 > 0$  will be specified later.

The space  $\mathcal{K}$  is endowed with the metric induced by the norm

$$\|X\|_\alpha = \left( \sup_{t \in [0, T]} \mathbb{E}[e^{-2\alpha t} |X(t)|_{-1}^2] \right)^{1/2}.$$

Note that  $\mathcal{K}$  is closed in the norm  $\|\cdot\|_\alpha$ . Indeed, if  $X_n \rightarrow X$  in  $\|\cdot\|_\alpha$  then since

$$\mathbb{E} \int_{Q_T} j(X_n(s)) d\xi ds \leq M_2 \quad \forall n \in \mathbb{N},$$

(3.15) implies that

$$X_n \rightarrow X \quad \text{in } L^1((0, T) \times \mathcal{O} \times \Omega)$$

and by Fatou's lemma we get

$$\mathbb{E} \int_{Q_T} j(X(s)) d\xi ds \leq M_2$$

as claimed. Now consider the mapping  $\Gamma : \mathcal{K} \rightarrow C_W([0, T]; H) \cap L^1((0, T) \times \mathcal{O} \times \Omega)$  defined by

$$(4.2) \quad Y = \Gamma(X),$$

where  $Y \in C_W([0, T]; H) \cap L^1((0, T) \times \mathcal{O} \times \Omega)$  is the solution in the sense of Definition 2.1 of the problem

$$(4.3) \quad \begin{cases} dY(t) - \Delta \Psi(Y(t)) dt = B(X(t)) dW(t), & \text{in } Q_T, \\ \Psi(Y(t)) = 0, & \text{on } \Sigma_T, \\ Y(0) = x, & \text{in } \mathcal{O}. \end{cases}$$

We shall prove that for  $\alpha, M_1, M_2$  suitably chosen,  $\Gamma$  maps  $\mathcal{K}$  into itself and it is a contraction in the norm  $\|\cdot\|_\alpha$ .

Indeed by (3.51) and (1.6) for any solution  $Y$  to (4.3) we have that

$$\begin{aligned} & \frac{1}{2}|Y(t)|_{-1}^2 + \int_0^t \int_{\mathcal{O}} (j(Y(s)) + j^*(\eta(s))) d\xi ds \\ &= \int_0^t \langle Y(s), B(X(s)) dW(s) \rangle_{-1} \\ & \quad + \frac{1}{2} \int_0^t \|B(X(s))\|_{LHS(L^2(\mathcal{O}),H)}^2 ds + \frac{1}{2}|x|_{-1}^2, \quad t \in [0, T], \mathbb{P}\text{-a.s.} \end{aligned}$$

By hypothesis  $(H_2)$  we have

$$\begin{aligned} & \frac{1}{2} \sup_{t \in [0, T]} \mathbb{E}[e^{-2\alpha t} |Y(t)|_{-1}^2] + e^{-2\alpha t} \mathbb{E} \int_0^t \int_{\mathcal{O}} (j(Y(s)) + j^*(\eta(s))) d\xi ds \\ & \leq \frac{1}{2}|x|_{-1}^2 + \frac{L^2}{2} \sup_{t \in [0, T]} \left[ e^{-2\alpha t} \int_0^t \mathbb{E}|X(s)|_{-1}^2 ds \right] \\ & \leq \frac{1}{2}|x|_{-1}^2 + \frac{L^2}{2} \sup_{t \in [0, T]} \int_0^t e^{-2\alpha(t-s)} \mathbb{E}e^{-2\alpha s} |X(s)|_{-1}^2 ds \leq \frac{1}{2}|x|_{-1}^2 + \frac{L^2 M_1^2}{4\alpha}. \end{aligned}$$

Hence

$$\sup_{t \in [0, T]} \mathbb{E}[e^{-2\alpha t} |Y(t)|_{-1}^2] \leq \frac{L^2 M_1^2}{2\alpha} + |x|_{-1}^2$$

and

$$\mathbb{E} \int_{Q_T} (j(Y(s)) + j^*(\eta(s))) d\xi \leq \left( \frac{L^2 M_1^2}{2\alpha} + |x|_{-1}^2 \right) e^{2\alpha T}.$$

Hence for  $\alpha > L^2$ ,  $M_1^2 > 2|x|_{-1}^2$  and  $M_2 \geq M_1^2 e^{2\alpha T}$  we have that  $Y \in \mathcal{K}$  and the operator  $\Gamma$  maps  $\mathcal{K}$  into itself. By a similar computation involving hypothesis  $(H_2)$  we see that for  $M_1, M_2$  and  $\alpha$  suitably chosen we have

$$(4.4) \quad \|Y_1 - Y_2\|_{\alpha} \leq \frac{C}{\sqrt{\alpha}} \|X_1 - X_2\|_{\alpha},$$

where  $Y_i = \Gamma X_i, i = 1, 2$ . Hence for a suitable  $\alpha$ ,  $\Gamma$  is a contraction and so equation  $X = \Gamma(X)$  has a unique solution in  $\Gamma$ . This completes the proof.

**Acknowledgments.** The authors are indebted to the anonymous referee for a number of useful suggestions which led to an improved presentation of this paper.

A part of this work was done during a very pleasant stay of the last named author at the Rumanian Academy of Sciences in Iasi whose warm hospitality is gratefully acknowledged.

REFERENCES

[1] ARONSON, D. G. (1986). The porous medium equation. In *Nonlinear Diffusion Problems (Montecatini Terme, 1985). Lecture Notes in Mathematics 1224* 1–46. Springer, Berlin. MR877986



- [2] BARBU, V. (1993). *Analysis and Control of Nonlinear Infinite-Dimensional Systems. Mathematics in Science and Engineering* **190**. Academic Press, Boston. [MR1195128](#)
- [3] BARBU, V. and DA PRATO, G. (2002). A phase field system perturbed by noise. *Nonlinear Anal.* **51** 1087–1099. [MR1926087](#)
- [4] BARBU, V., BOGACHEV, V. I., DA PRATO, G. and RÖCKNER, M. (2006). Weak solutions to the stochastic porous media equation via Kolmogorov equations: The degenerate case. *J. Funct. Anal.* **237** 54–75. [MR2239258](#)
- [5] BERRYMAN, J. G. and HOLLAND, C. J. (1980). Stability of the separable solution for fast diffusion. *Arch. Rational Mech. Anal.* **74** 379–388. [MR588035](#)
- [6] BRÉZIS, H. (1973). *Opérateurs maximaux monotones et semigroupes de contraction dans les espaces de Hilbert*. North-Holland, Amsterdam. [MR0348562](#)
- [7] DA PRATO, G. (2004). *Kolmogorov Equations for Stochastic PDEs*. Birkhäuser, Basel. [MR2111320](#)
- [8] DA PRATO, G., RÖCKNER, M., ROZOVSKII, B. L. and WANG, F.-Y. (2006). Strong solutions of stochastic generalized porous media equations: Existence, uniqueness, and ergodicity. *Comm. Partial Differential Equations* **31** 277–291. [MR2209754](#)
- [9] DA PRATO, G. and ZABCZYK, J. (1996). *Ergodicity for Infinite-Dimensional Systems. London Mathematical Society Lecture Note Series* **229**. Cambridge Univ. Press, Cambridge. [MR1417491](#)
- [10] KRYLOV, N. V. and ROZOVSKIĬ, B. L. (1979). Stochastic evolution equations. In *Current Problems in Mathematics* **14** 71–147, 256. Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Informatsii, Moscow. In Russian. [MR570795](#)
- [11] LIONS, J.-L. (1969). *Quelques méthodes de résolution des problèmes aux limites non linéaires*. Gauthier-Villars, Paris. [MR0259693](#)
- [12] MARINOSCHI, G. (2006). *Functional Approach to Nonlinear Models of Water Flow in Soils. Mathematical Modelling: Theory and Applications* **21**. Springer, Dordrecht. [MR2258329](#)
- [13] PEDRON, I. J., MENDES, R. S., BURATTO, T. J., MALACARNE, L. C. and LENZI, E. K. (2005). Logarithmic diffusion and porous media equations: A unified description. *Phys. Rev.* **E72** 03 1106.
- [14] PRÉVÔT, C. and RÖCKNER, M. (2007). *A Concise Course on Stochastic Partial Differential Equations. Lecture Notes in Mathematics* **1905**. Springer, Berlin. [MR2329435](#)
- [15] REN, J., RÖCKNER, M. and WANG, F.-Y. (2007). Stochastic generalized porous media and fast diffusion equations. *J. Differential Equations* **238** 118–152. [MR2334594](#)

V. BARBU  
 UNIVERSITY AL. I. CUZA  
 AND  
 INSTITUTE OF MATHEMATICS “OCTAV MAYER”  
 IASI  
 ROMANIA  
 E-MAIL: [vb41@uaic.ro](mailto:vb41@uaic.ro)

G. DA PRATO  
 SCUOLA NORMALE SUPERIORE DI PISA  
 PIAZZA DEI CAVALIERI 7  
 PISA 56126  
 ITALY  
 E-MAIL: [daprato@sns.it](mailto:daprato@sns.it)

M. RÖCKNER  
 FACULTY OF MATHEMATICS  
 UNIVERSITY OF BIELEFELD  
 GERMANY  
 AND  
 DEPARTMENT OF MATHEMATICS AND STATISTICS  
 PURDUE UNIVERSITY  
 USA