

ON THE APPROXIMATE MAXIMUM LIKELIHOOD ESTIMATION FOR DIFFUSION PROCESSES

BY JINYUAN CHANG¹ AND SONG XI CHEN^{1,2}

Peking University, and Peking University and Iowa State University

The transition density of a diffusion process does not admit an explicit expression in general, which prevents the full maximum likelihood estimation (MLE) based on discretely observed sample paths. Ait-Sahalia [*J. Finance* **54** (1999) 1361–1395; *Econometrica* **70** (2002) 223–262] proposed asymptotic expansions to the transition densities of diffusion processes, which lead to an approximate maximum likelihood estimation (AMLE) for parameters. Built on Ait-Sahalia's [*Econometrica* **70** (2002) 223–262; *Ann. Statist.* **36** (2008) 906–937] proposal and analysis on the AMLE, we establish the consistency and convergence rate of the AMLE, which reveal the roles played by the number of terms used in the asymptotic density expansions and the sampling interval between successive observations. We find conditions under which the AMLE has the same asymptotic distribution as that of the full MLE. A first order approximation to the Fisher information matrix is proposed.

1. Introduction. Continuous-time diffusion processes defined by stochastic differential equations [Karatzas and Shreve (1991), Øksendal (2000), Protter (2004)] are the basic stochastic modeling tools in the modern financial theory and applications. Diffusion models are commonly employed to describe the price dynamics of a financial asset or a portfolio of assets. An eminent application is in deriving the price of a derivative contract on an asset or a group of assets. The celebrated Black–Scholes–Merton option pricing formula [Black and Scholes (1973), Merton (1973)] was obtained by assuming that the underlying asset followed a geometric Brownian motion such that the log price process of the underlying asset followed an Ornstein–Uhlenbeck diffusion process. The widely used Vasicek (1977) and Cox, Ingersoll and Ross (1985) pricing formulas for the zero coupon bond were developed based on two specific mean-reverting diffusion processes with a constant or the square root [Feller (1952)] diffusion functions, respectively. Other pricing formulas have also been developed for assets defined by other processes; see Bakshi, Cao and Chen (1997) and Dumas, Fleming and Whaley (1998).

Received February 2011; revised August 2011.

¹Supported by Center for Statistical Science at Peking University and a National Natural Science Foundation of China Zhongdian Program Grant (No. 11131002).

²Supported by NSF Grants DMS-06-04563 and DMS-07-14978.

MSC2010 subject classifications. Primary 62M05; secondary 62F12.

Key words and phrases. Asymptotic expansion, asymptotic normality, consistency, discrete time observation, maximum likelihood estimation.

In the implementations of the aforementioned pricing formula, the parameters of the diffusion processes which describe the underlying assets dynamics have to be estimated based on empirical observations. Sundareshan (2000) gave a comprehensive survey on the financial applications of continuous-time stochastic models which were largely the diffusion processes. Fan (2005) provided an overview on nonparametric estimation for diffusion processes. Other related works include Bibby and Sørensen (1995), Wang (2002), Fan and Zhang (2003), Fan and Wang (2007), Mykland and Zhang (2009) and Aït-Sahalia, Mykland and Zhang (2011).

There are several challenges to be faced when estimating parameters of diffusion processes. One challenge is that despite being continuous-time models, the processes are only observed at discrete time points rather than observed continuously over time. The discrete observations prevent the use of the relatively straightforward likelihood expressions [Prakasa Rao (1999)] available for continuously observed diffusion processes. Another challenge is that despite the fact that the diffusion processes are Markovian, their transition densities from one time point to the next do not have finite analytic expressions, except for only a few specific processes. This means that the efficient maximum likelihood estimation (MLE) cannot be readily implemented for most of these processes.

In ground-breaking works, Aït-Sahalia (1999, 2002) established series expansions to approximate the transition densities of univariate diffusion processes. Similar expansions have been proposed for multivariate processes in Aït-Sahalia (2008). These density approximations, as advocated by Aït-Sahalia, are then employed to form approximate likelihood functions, which are maximized to obtain the approximate maximum likelihood estimators (AMLEs). Aït-Sahalia (2002, 2008) demonstrated that the approximate likelihood converges to the true likelihood as the number of terms in the series expansions goes to infinity. He also provided some results on the consistency of the AMLEs. Numerical evaluations of the transition density approximations as conducted in Aït-Sahalia (1999), Stramer and Yan (2007a, 2007b) and others, have shown good performance in the numerical approximation of the underlying transition densities. The approach has opened a very accessible route for obtaining parameter estimators for diffusion processes, and for estimating other quantities which are functions of the transition density, as commonly encountered in finance. Indeed, Aït-Sahalia and Kimmel (2007, 2010) demonstrated two such applications in stochastic volatility models and the affine term structure models, respectively. Tang and Chen (2009) provided some results on the AMLE based on the one-term expansion for the mean-reverting processes. They revealed that there was an extra leading order bias term in the AMLE due to the density approximation.

Although the above-mentioned results on the transition density approximation and the AMLE had been provided, there are some key questions that remain to be addressed. One is on the consistency of the AMLE. While Aït-Sahalia (2002, 2008) contained some results on consistency, there is more to be explored. There

are two key ingredients in Aït-Sahalia's density approximation. One is J , the number of terms used in the approximation, and the other is δ , the length of the sampling interval between successive observations. In this paper, we study explicitly the roles played by J and δ on the consistency of the AMLE, and quantify their roles on the convergence rate. Another question is under what conditions on J and δ , does the AMLE have the same asymptotic distribution as the full MLE. Here, we consider two regimes: (i) δ is fixed, and $J \rightarrow \infty$; (ii) J is fixed, but $\delta \rightarrow 0$, representing two views of asymptotics. In the case of $\delta \rightarrow 0$, it is found that $J \geq 2$ is necessary to ensure the AMLE having the same asymptotic normality as the MLE. Like the transition density, the Fisher information matrix, the quantity that defines the efficiency of the full MLE, is unknown analytically; even the underlying transition density is known. We show in this paper that an approximation to the Fisher information matrix can be obtained based on the one-term density approximation.

The paper is organized as follows. In Section 2, we outline the transition density approximations of Aït-Sahalia (1999, 2002). Some preliminary analysis is needed for studying the AMLE is presented in Section 3. Section 4 establishes the consistency and convergence rates of the AMLE. Asymptotic normality of the AMLE and its equivalence to the full MLE are addressed in Section 5. Section 6 discusses the approximation for the Fisher information matrix. Simulation results are reported in Section 7. Technical conditions and details of proofs are relegated to the Appendix.

2. Transition density approximation. Consider a univariate diffusion process $(X_t)_{t \geq 0}$ defined by a stochastic differential equation

$$(2.1) \quad dX_t = \mu(X_t; \theta) dt + \sigma(X_t; \theta) dB_t,$$

where μ and σ are, respectively, the drift and diffusion functions and B_t is the standard Brownian motion. Both the drift and diffusion functions are known except for an unknown parameter vector θ taking values in a set $\Theta \subseteq \mathbb{R}^d$.

Given a sampling interval $\delta > 0$, let $f_X(x|x_0, \delta; \theta)$ be the transition density of $X_{t+\delta}$ given $X_t = x_0$ for $(x_0, x) \in \mathcal{X} \times \mathcal{X}$, where \mathcal{X} is the domain of X_t . Despite the parametric forms of the drift and the diffusion functions that are available in (2.1), a closed-form expression for $f_X(x|x_0, \delta; \theta)$ is not generally available for most of the processes. In most cases, the density is only known to satisfy the Kolmogorov backward and forward partial differential equations. In ground-breaking works, Aït-Sahalia (1999, 2002) proposed asymptotic expansions to approximate the transition density.

The approach of Aït-Sahalia is the following. He first transformed X_t to a diffusion process with unit diffusion function by

$$(2.2) \quad Y_t = \gamma(X_t; \theta) := \int^{X_t} \frac{du}{\sigma(u; \theta)},$$

which satisfies $dY_t = \mu_Y(Y_t; \theta) dt + dB_t$, where

$$\mu_Y(y; \theta) = \frac{\mu(\gamma^{-1}(y; \theta); \theta)}{\sigma(\gamma^{-1}(y; \theta); \theta)} - \frac{1}{2} \frac{\partial \sigma}{\partial x}(\gamma^{-1}(y; \theta); \theta).$$

Let $f_Y(y|y_0, \delta; \theta)$ be the transition density of $Y_{t+\delta}$ given $Y_t = y_0$. The two density functions are related according to

$$(2.3) \quad f_X(x_t|x_{t-1}, \delta; \theta) = \sigma^{-1}(x_t; \theta) \cdot f_Y(\gamma(x_t; \theta)|\gamma(x_{t-1}; \theta), \delta; \theta).$$

To ensure convergence of the expansions, Ait-Sahalia standardized $Y_{t+\delta}$ by $Z_{t+\delta} = \delta^{-1/2}(Y_{t+\delta} - y_0)$. Let $f_Z(z|y_0, \delta; \theta)$ denote the conditional density of $Z_{t+\delta}$ given $Z_t = 0$, which is related to f_Y by

$$f_Z(z|y_0, \delta; \theta) = \delta^{1/2} f_Y(\delta^{1/2}z + y_0|y_0, \delta; \theta).$$

Let $\{H_j(z)\}_{j=1}^\infty$ be the Hermite polynomials

$$H_j(z) = \phi^{-1}(z) \frac{d^j \phi(z)}{dz^j},$$

which are orthogonal with respect to the standard normal density ϕ , namely $\int H_j(z)H_k(z)\phi(x) dx = 0$ if $j \neq k$. A formal Hermite orthogonal series expansion to the density $f_Z(z|y_0, \delta; \theta)$ is

$$(2.4) \quad f_Z^H(z|y_0, \delta; \theta) = \phi(z) \sum_{j=0}^\infty \eta_j(y_0, \delta; \theta) H_j(z),$$

where the coefficients

$$\begin{aligned} \eta_j(y_0, \delta; \theta) &= (j!)^{-1} \int H_j(z) f_Z(z|y_0, \delta; \theta) dz \\ &= (j!)^{-1} \mathbb{E}[H_j(\delta^{-1/2}(Y_{t+\delta} - y_0)) | Y_t = y_0; \theta]. \end{aligned}$$

The last conditional expectation has no analytic expression in general, although it may be simulated using the method proposed in Beskos et al. (2006). Ait-Sahalia proposed Taylor expansions for this conditional expectation with respect to the sampling interval δ based on the infinitesimal generator of Y_t . For twice continuously differentiable function g , the infinitesimal generator of Y_t is

$$(2.5) \quad \mathcal{A}_\theta g(y) = \mu_Y(y; \theta) \frac{\partial g}{\partial y} + \frac{1}{2} \frac{\partial^2 g}{\partial y^2}.$$

A K -term Taylor series expansion to $\mathbb{E}[H_j(\delta^{-1/2}(Y_{t+\delta} - y_0)) | Y_t = y_0; \theta]$ is

$$(2.6) \quad \begin{aligned} &\mathbb{E}[H_j(\delta^{-1/2}(Y_{t+\delta} - y_0)) | Y_t = y_0; \theta] \\ &= \sum_{k=0}^K \mathcal{A}_\theta^k H_j(\delta^{-1/2}(y - y_0)) \Big|_{y=y_0} \frac{\delta^k}{k!} \\ &\quad + \mathbb{E}[\mathcal{A}_\theta^{k+1} H_j(\delta^{-1/2}(Y_{t+\delta^*} - y_0)) | Y_t = y_0; \theta] \frac{\delta^{k+1}}{(k+1)!}. \end{aligned}$$

Substituting (2.6) to the orthogonal expansion (2.4) followed by gathering terms according to the powers of δ , a J -term expansion to the transition density $f_Y(y, \delta|y_0; \theta)$ is

$$f_Y^{(J)}(y|y_0, \delta; \theta) = \delta^{-1/2} \phi\left(\frac{y - y_0}{\delta^{1/2}}\right) \exp\left(\int_{y_0}^y \mu_Y(u; \theta) du\right) \sum_{j=0}^J c_j(y|y_0; \theta) \frac{\delta^j}{j!},$$

where $c_0(y|y_0; \theta) \equiv 1$ and for $j \geq 1$,

$$\begin{aligned} c_j(y|y_0; \theta) &= j(y - y_0)^{-j} \\ &\times \int_{y_0}^y (w - y_0)^{j-1} \\ &\times \left\{ \lambda_Y(w; \theta) c_{j-1}(w|y_0; \theta) + \frac{1}{2} \frac{\partial^2 c_{j-1}(w|y_0; \theta)}{\partial w^2} \right\} dw. \end{aligned}$$

Here $\lambda_Y(y; \theta) = -\{\mu_Y^2(y; \theta) + \partial \mu_Y(y; \theta) / \partial y\} / 2$.

Transforming back from y to x via (2.2) and (2.3), the J -term expansion to $f_X(x|x_0, \delta; \theta)$ is

$$\begin{aligned} f_X^{(J)}(x|x_0, \delta; \theta) &= \sigma^{-1}(x; \theta) \delta^{-1/2} \phi\left(\frac{\gamma(x; \theta) - \gamma(x_0; \theta)}{\delta^{1/2}}\right) \\ &\times \exp\left\{ \int_{x_0}^x \frac{\mu_Y(\gamma(u; \theta); \theta)}{\sigma(u; \theta)} du \right\} \sum_{j=0}^J c_j(\gamma(x; \theta) | \gamma(x_0; \theta); \theta) \frac{\delta^j}{j!}. \end{aligned}$$

Although it employs the Hermite polynomials and has the Gaussian density as the leading term as an Edgeworth expansion does, the transition density expansion is not an Edgeworth expansion. This is because the latter is for density functions of statistics admitting the central limit theorem, which differs from the current context of expanding the transition density. [Ait-Sahalia \(2002\)](#) demonstrated that as $J \rightarrow \infty$,

$$(2.7) \quad f_X^{(J)}(x|x_0, \delta; \theta) \rightarrow f_X(x|x_0, \delta; \theta)$$

uniformly with respect to $\theta \in \Theta$ and x_0 over compact subsets of \mathcal{X} . The convergence is also uniform with respect to x over subsets of \mathcal{X} depending on the property of $\sigma(x; \theta)$.

Define

$$\begin{aligned} A_1(x|x_0, \delta; \theta) &= -\log\{\sigma(x; \theta)\} - \frac{1}{2\delta} \{\gamma(x; \theta) - \gamma(x_0; \theta)\}^2, \\ A_2(x|x_0, \delta; \theta) &= \int_{x_0}^x \frac{\mu_Y(\gamma(u; \theta); \theta)}{\sigma(u; \theta)} du \end{aligned}$$

and

$$A_3(x|x_0, \delta; \theta) = \log \left\{ \sum_{j=0}^J c_j(\gamma(x; \theta) | \gamma(x_0; \theta); \theta) \delta^j / j! \right\}.$$

If $\sum_{j=0}^\infty |c_j(y|y_0, \delta; \theta)| \delta^j / j! < \infty$ on $\mathcal{Y} \times \mathcal{Y}$ with probability one, where \mathcal{Y} is the domain of Y_t , we can define $\tilde{A}_3(x|x_0, \delta; \theta) = \log\{\sum_{j=0}^\infty c_j(y|y_0; \theta) \delta^j / j!\}$. Then the result in (2.7) implies that

$$\begin{aligned} \log f_X(x|x_0, \delta; \theta) \\ (2.8) \quad &= -\log \sqrt{2\pi\delta} + A_1(x|x_0, \delta; \theta) + A_2(x|x_0, \delta; \theta) \\ &+ \tilde{A}_3(x|x_0, \delta; \theta). \end{aligned}$$

Expression (2.8) is the starting point for our analysis.

Given a set of discrete observations $\{X_{t\delta}\}_{t=1}^n$ with equal sampling length δ of the diffusion process $(X_t)_{t \geq 0}$, to simplify notations, we write X_t for $X_{t\delta}$ and hide δ in the expressions for the transition density f_X and its approximations. At the same time, we use f and $f^{(J)}$ to express f_X and $f_X^{(J)}$, respectively. Based on the J -term expansion to the true transition density, the J -term approximate log-likelihood function given in Ait-Sahalia (2002) is

$$\begin{aligned} \ell_{n,\delta}^{(J)}(\theta) &= -n \log \sqrt{2\pi\delta} + \sum_{t=1}^n A_1(X_t | X_{t-1}, \delta; \theta) \\ &+ \sum_{t=1}^n A_2(X_t | X_{t-1}, \delta; \theta) + \sum_{t=1}^n A_3(X_t | X_{t-1}, \delta; \theta). \end{aligned}$$

Let $\hat{\theta}_{n,\delta}^{(J)} = \arg \max_{\theta \in \Theta} \ell_{n,\delta}^{(J)}(\theta)$ be the approximate MLE (AMLE) and $\hat{\theta}_{n,\delta}$ be the true MLE that maximizes the full likelihood

$$\ell_{n,\delta}(\theta) = \sum_{t=1}^n \log f(X_t | X_{t-1}, \delta; \theta).$$

To keep the notation simple, we write $\hat{\theta}_n^{(J)} = \hat{\theta}_{n,\delta}^{(J)}$ and $\hat{\theta}_n = \hat{\theta}_{n,\delta}$ by suppressing δ in subscripts.

3. Preliminaries. Under regular circumstances as assumed by condition (A.2)(ii) in the Appendix, the full MLE $\hat{\theta}_n$ and the J -term approximate MLE $\hat{\theta}_n^{(J)}$ satisfy their respective likelihood score equations so that

$$(3.1) \quad \sum_{t=1}^n \nabla_\theta \log f(X_t | X_{t-1}, \delta; \hat{\theta}_n) = \sum_{t=1}^n \nabla_\theta \log f^{(J)}(X_t | X_{t-1}, \delta; \hat{\theta}_n^{(J)}) = 0.$$

Subtracting $\sum_{t=1}^n \nabla_{\theta} \log f^{(J)}(X_t|X_{t-1}, \delta; \theta_0)$ from both sides of (3.1),

$$\begin{aligned}
 & \sum_{t=1}^n \nabla_{\theta} \log f^{(J)}(X_t|X_{t-1}, \delta; \hat{\theta}_n^{(J)}) - \sum_{t=1}^n \nabla_{\theta} \log f^{(J)}(X_t|X_{t-1}, \delta; \theta_0) \\
 (3.2) \quad &= \sum_{t=1}^n \nabla_{\theta} [\tilde{A}_3(X_t|X_{t-1}, \delta; \theta_0) - A_3(X_t|X_{t-1}, \delta; \theta_0)] \\
 & \quad + \sum_{t=1}^n \nabla_{\theta} \log f(X_t|X_{t-1}, \delta; \hat{\theta}_n) - \sum_{t=1}^n \nabla_{\theta} \log f(X_t|X_{t-1}, \delta; \theta_0).
 \end{aligned}$$

Carrying out Taylor expansions on both sides of (3.2), we can get

$$\begin{aligned}
 & \frac{1}{n} \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f^{(J)}(X_t|X_{t-1}, \delta; \theta_0) \cdot (\hat{\theta}_n^{(J)} - \theta_0) \\
 & \quad + \frac{1}{2} [E_d \otimes (\hat{\theta}_n^{(J)} - \theta_0)'] \cdot \frac{1}{n} \sum_{t=1}^n \nabla_{\theta\theta\theta}^3 \log f^{(J)}(X_t|X_{t-1}, \delta; \tilde{\theta}) \cdot (\hat{\theta}_n^{(J)} - \theta_0) \\
 (3.3) \quad &= \frac{1}{n} \sum_{t=1}^n \nabla_{\theta} [\tilde{A}_3(X_t|X_{t-1}, \delta; \theta_0) - A_3(X_t|X_{t-1}, \delta; \theta_0)] \\
 & \quad + \frac{1}{n} \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f(X_t|X_{t-1}, \delta; \theta_0) \cdot (\hat{\theta}_n - \theta_0) \\
 & \quad + \frac{1}{2} [E_d \otimes (\hat{\theta}_n - \theta_0)'] \cdot \frac{1}{n} \sum_{t=1}^n \nabla_{\theta\theta\theta}^3 \log f(X_t|X_{t-1}, \delta; \bar{\theta}) \cdot (\hat{\theta}_n - \theta_0),
 \end{aligned}$$

where E_d is the $d \times d$ identity matrix, $\tilde{\theta}$ is on the joint line between $\hat{\theta}_n^{(J)}$ and θ_0 and $\bar{\theta}$ is on the joint line between $\hat{\theta}_n$ and θ_0 . Here we define

$$\nabla_{\theta\theta\theta}^3 \log f(X_t|X_{t-1}, \delta; \theta) := \begin{pmatrix} \partial^3 \log f(X_t|X_{t-1}, \delta; \theta) / \partial \theta \partial \theta' \partial \theta_1 \\ \vdots \\ \partial^3 \log f(X_t|X_{t-1}, \delta; \theta) / \partial \theta \partial \theta' \partial \theta_d \end{pmatrix},$$

which is a $d^2 \times d$ matrix, and $\nabla_{\theta\theta\theta}^3 \log f^{(J)}(X_t|X_{t-1}, \delta; \theta)$ is similarly defined. Furthermore, let

$$\begin{aligned}
 F_n(\theta_0, J, \delta) &= n^{-1} \sum_{t=1}^n \nabla_{\theta\theta}^2 [\tilde{A}_3(X_t|X_{t-1}, \delta; \theta_0) - A_3(X_t|X_{t-1}, \delta; \theta_0)], \\
 U_n(\theta_0, J, \delta) &= n^{-1} \sum_{t=1}^n \nabla_{\theta} [\tilde{A}_3(X_t|X_{t-1}, \delta; \theta_0) - A_3(X_t|X_{t-1}, \delta; \theta_0)]
 \end{aligned}$$

and

$$N_n(\theta_0, J, \delta) = n^{-1} \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f^{(J)}(X_t|X_{t-1}, \delta; \theta_0).$$

Then (3.3) can be written as

$$\begin{aligned} & N_n(\theta_0, J, \delta)(\hat{\theta}_n^{(J)} - \theta_0) + \Delta_{n1}(\hat{\theta}_n^{(J)}, \theta_0) \\ (3.4) \quad & = U_n(\theta_0, J, \delta) + [N_n(\theta_0, J, \delta) + F_n(\theta_0, J, \delta)](\hat{\theta}_n - \theta_0) \\ & \quad + \Delta_{n2}(\hat{\theta}_n, \theta_0), \end{aligned}$$

where $\Delta_{n1}(\hat{\theta}_n^{(J)}, \theta_0)$ and $\Delta_{n2}(\hat{\theta}_n, \theta_0)$ denote the remainder terms whose explicit expressions can be obtained by matching (3.3) with (3.4).

Expansion (3.4) is the starting point in our studies for the consistency and asymptotic distribution of the AMLE. Indeed, the asymptotic properties of the AMLE will be evaluated under two regimes regarding J and δ . The first one is that

$$(3.5) \quad \delta \text{ is fixed} \quad \text{but } J \rightarrow \infty,$$

which is the situation considered in Aït-Sahalia (2002). The second regime allows that

$$(3.6) \quad J \text{ is fixed,} \quad \delta \rightarrow 0 \quad \text{but } n\delta \rightarrow \infty,$$

which is more tuned with an implementation of the density approximation with a fixed number of terms.

We will first present some results which are valid for any fixed J and δ . Let $\|A\|_2 = \{\rho(A'A)\}^{1/2}$ be the spectral norm of a matrix A , where $\rho(A'A)$ denotes the largest eigen-value of $A'A$. The following proposition describes properties for the quantities that appear in (3.4).

PROPOSITION 1. *Under conditions (A.1), (A.3), (A.4), (A.6), (A.7) given in the Appendix, there exists a positive constant Δ such that for any positive integer J and $\delta \in (0, \Delta)$:*

- (a) $\mathbb{E}\{F_n(\theta_0, J, \delta)\}$, $\mathbb{E}\{U_n(\theta_0, J, \delta)\}$ and $\mathbb{E}\{N_n(\theta_0, J, \delta)\}$ exist;
- (b) $\Delta_{n1}(\hat{\theta}_n^{(J)}, \theta_0) = O_p\{\|\hat{\theta}_n^{(J)} - \theta_0\|_2^2\}$ and $\Delta_{n2}(\hat{\theta}_n, \theta_0) = O_p\{\|\hat{\theta}_n - \theta_0\|_2^2\}$ as $n \rightarrow \infty$.

Let $I(\delta) = -\mathbb{E}\nabla_{\theta\theta}^2 \log f(X_t|X_{t-1}, \delta; \theta_0)$ be the Fisher information matrix, which we assume is invertible in condition (A.5). It is expected that the expected value of $N_n(\theta_0, J, \delta)$, denoted by $N(\theta_0, J, \delta)$, will converge to $-I(\delta)$, as $J \rightarrow \infty$ for each fixed δ or J being fixed but $\delta \rightarrow 0$. The following proposition bounds the difference between $N(\theta_0, J, \delta)$ and $-I(\delta)$ for each fixed J and δ .

PROPOSITION 2. *Under conditions (A.1), (A.4), (A.6), (A.7) given in the Appendix, there exist two positive constants $\tilde{\Delta}$ and C , that are not dependent on J and δ , such that for any positive integer J and $\delta \in (0, \tilde{\Delta})$,*

$$\|N(\theta_0, J, \delta) + I(\delta)\|_2 \leq C\delta^{J+1}.$$

As $I(\delta)$ is invertible for each fixed $\delta > 0$, $N_n(\theta_0, J, \delta)$ will be invertible with probability approaching one as $J \rightarrow \infty$ for a fixed δ . However, if $\delta \rightarrow 0$, the limit of the Fisher information $I(0) := \lim_{\delta \rightarrow 0} I(\delta)$, as well as $N(\theta_0, J, 0)$, may be singular. This is the case for some Ornstein–Uhlenbeck processes as shown in Section 6. The following proposition provides another account on $N(\theta_0, J, \delta)$ and its deviation from $-I(\delta)$, as well as the convergence of $N^{-1}(\theta_0, J, \delta)U(\theta_0, J, \delta)$, where $U(\theta_0, J, \delta)$ denotes the expected value of $U_n(\theta_0, J, \delta)$ for each pair of fixed J and δ .

PROPOSITION 3. *Under conditions (A.1), (A.3)–(A.7) given in the Appendix, there exist two constants C_1, C_2 , that are not dependent on J and δ , and a constant $\underline{\Delta} > 0$ such that for any positive integer J and $\delta \in (0, \underline{\Delta})$,*

$$\|N^{-1}(\theta_0, J, \delta)I(\delta) + E_d\|_2 \leq C_1\delta^J \quad \text{and} \quad \|N^{-1}(\theta_0, J, \delta)U(\theta_0, J, \delta)\|_2 \leq C_2\delta^J.$$

The next proposition describes the convergence rate for the difference between the first derivatives of the full log-likelihood and the approximate log-likelihood.

PROPOSITION 4. *Under conditions (A.1), (A.4), (A.6), (A.7) given in the Appendix, there exist two finite positive constants $\tilde{\Delta}$ and C , not dependent on J and δ , such that for any $J, \delta \in (0, \tilde{\Delta}]$ and n ,*

$$\mathbb{E}\left\{\sup_{\theta \in \Theta} \|n^{-1} \cdot \nabla_{\theta} [\ell_{n,\delta}(\theta) - \ell_{n,\delta}^{(J)}(\theta)]\|_2\right\} \leq C\delta^{J+1}.$$

The following proposition together with Proposition 4 is needed to establish the consistency of the AMLE.

PROPOSITION 5. *Under conditions (A.1), (A.3), (A.4), (A.6), (A.7) given in the Appendix, there exists a constant $\hat{\Delta} > 0$ such that*

$$\sup_{\theta \in \Theta} \left\| \frac{1}{n} \sum_{t=1}^n \nabla_{\theta} \log f(X_t|X_{t-1}, \delta; \theta) - \mathbb{E} \nabla_{\theta} \log f(X_t|X_{t-1}, \delta; \theta) \right\|_2 \xrightarrow{P} 0$$

for (i) $\delta \in (0, \hat{\Delta}]$ being fixed, $n \rightarrow \infty$, or (ii) $n \rightarrow \infty, \delta \rightarrow 0$ but $n\delta \rightarrow \infty$.

As the full MLE $\hat{\theta}_n$ is a key bridge for the AMLE, we report in the following proposition the asymptotic normality of the MLE which covers both cases of fixed δ and diminishing δ case.

PROPOSITION 6. Under conditions (A.1)–(A.7) given in the Appendix,

$$\sqrt{n}I^{1/2}(\delta)(\hat{\theta}_n - \theta_0) \xrightarrow{d} N(0, E_d) \quad \text{as } n\delta^3 \rightarrow \infty,$$

where E_d is $d \times d$ identity matrix.

The requirement of $n\delta^3 \rightarrow \infty$ in the above proposition is to cover the case where $I(0) = \lim_{\delta \rightarrow 0} I(\delta)$ is singular, as spelled out in the proof given in the Appendix. If such case is ruled out, for instance, via the so-call Jacobsen condition [Jacobsen (2001), Sørensen (2007)], the more standard $n\delta \rightarrow \infty$ is sufficient; see also Gobet (2002) for related results.

4. Consistency. We consider in this section the consistency of the AMLE $\hat{\theta}_n^{(J)}$ and establish its convergence rate under the two asymptotic regimes given in (3.5) and (3.6), respectively. The two asymptotic regimes were also considered in Ait-Sahalia (2002, 2008). For a fixed sampling interval δ , Ait-Sahalia (2002) proved that there existed a sequence $J_n \rightarrow \infty$ such that $\hat{\theta}_n^{(J_n)} - \hat{\theta}_n \xrightarrow{P} 0$ under P_{θ_0} as $n \rightarrow \infty$, where P_{θ_0} is the underlying probability measure. Based on the consistency of $\hat{\theta}_n$, we know that the consistency of $\hat{\theta}_n^{(J_n)}$ is hold. For a fixed J , Ait-Sahalia (2008) proved that there existed a sequence $\{\delta_n\}$ vanishing to zero such that $\sqrt{n}I^{1/2}(\delta_n)(\hat{\theta}_{n,\delta_n}^{(J)} - \theta_0) = O_p(1)$.

In this paper, we will give more explicit guidelines on how to select the aforementioned sequences J_n and δ_n so that the AMLE is consistent. Our study here begins with (3.1), which together with Propositions 4 and 5 lead to the following result on the consistency of the AMLE under the two asymptotic regimes, respectively.

THEOREM 1. Under conditions (A.1)–(A.4), (A.6), (A.7) given in the Appendix, $\hat{\theta}_n^{(J)} - \theta_0 \xrightarrow{P} 0$ under either: (i) $\delta \in (0, \tilde{\Delta} \wedge \hat{\Delta}]$ being fixed, $J \rightarrow \infty$ and $n \rightarrow \infty$, or (ii) J being fixed, $n \rightarrow \infty$, $\delta \rightarrow 0$ but $n\delta \rightarrow \infty$.

By Proposition 2 and condition (A.5), multiply $N^{-1}(\theta_0, J, \delta)$ on both sides of (3.4), we have

$$\begin{aligned} & \hat{\theta}_n^{(J)} - \theta_0 \\ (4.1) \quad & = N^{-1}U_n + N^{-1}(N_n + F_n)(\hat{\theta}_n - \theta_0) - N^{-1}(N_n - N)(\hat{\theta}_n^{(J)} - \theta_0) \\ & \quad - N^{-1}\Delta_{n1}(\hat{\theta}_n^{(J)}, \theta_0) + N^{-1}\Delta_{n2}(\hat{\theta}_n, \theta_0). \end{aligned}$$

From this together with Proposition 4 and Theorem 1, we can establish the convergence rate of the AMLE.

THEOREM 2. Under conditions (A.1)–(A.7) given in the *Appendix*,

$$\hat{\theta}_n^{(J)} - \theta_0 = \begin{cases} O_p\{\delta^{J+1} + (n\delta)^{-1/2}\}, & \text{if } \delta \in (0, \tilde{\Delta} \wedge \hat{\Delta}] \text{ is fixed and } J \rightarrow \infty; \\ O_p\{\delta^J + (n\delta)^{-1/2}\}, & \text{if } J \text{ is fixed, } \delta \rightarrow 0 \text{ but } n\delta^3 \rightarrow \infty. \end{cases}$$

The above theorem reveals the impacts of the sampling interval δ and the number of terms J used in the density approximation on the convergence rate. In particular, the rate of AMLE has an extra δ^{J+1} or δ^J term in addition to the standard rate $(n\delta)^{-1/2}$ of the full MLE. This extra term is the result of the density approximation, and its particular form suggests that the sampling interval δ has to be less than 1 in order to make the AMLE $\hat{\theta}_n^{(J)}$ converge to θ_0 . It is apparent that the higher the J is, the less impact the extra term has on the AMLE $\hat{\theta}_n^{(J)}$.

5. Asymptotic distribution. In this section, we consider the asymptotic distribution of the AMLE $\hat{\theta}_n^{(J)}$. Our investigations are organized according to two asymptotic regimes: (i) δ fixed, $J \rightarrow \infty$ and (ii) J fixed, $\delta \rightarrow 0$ but $n\delta \rightarrow \infty$.

5.1. *Fixed δ , $J \rightarrow \infty$.* This is a simple case to treat. Under this setting, we note from Proposition 2 and condition (A.5) that $N^{-1}(\theta_0, J, \delta) = O(1)$ uniformly for any J . Utilizing the result in Theorem 2, expansion (4.1) becomes

$$\hat{\theta}_n^{(J)} - \theta_0 = N^{-1}U_n + (\hat{\theta}_n - \theta_0) + O_p(n^{-1/2}\delta^{J-1/2} + n^{-1}\delta^{-1} + \delta^{2J+2}).$$

Hence, note that $U_n = O_p(\delta^{J+1})$,

$$\begin{aligned} \sqrt{n}I^{1/2}(\delta)(\hat{\theta}_n^{(J)} - \theta_0) &= \sqrt{n}I^{1/2}(\delta)(\hat{\theta}_n - \theta_0) + O_p(\delta^{J-1/2} + n^{-1/2}\delta^{-1} + n^{1/2}\delta^{J+1}). \end{aligned}$$

If $n\delta^{2J+2} \rightarrow 0$, then

$$\sqrt{n}I^{1/2}(\delta)(\hat{\theta}_n^{(J)} - \theta_0) \xrightarrow{d} N(0, E_d).$$

Therefore, the AMLE has the same asymptotic distribution as the full MLE $\hat{\theta}_n$. This is attained by requesting $n\delta^{2J+2} \rightarrow 0$ in addition to $J \rightarrow \infty$. If $n\delta^{2J+2} \rightarrow c > 0$, the AMLE is still asymptotically normal but would have an inflated variance due to the contribution from the first term involving U_n . Apart from this, the asymptotic mean will no longer be zero. Hence, it is much desirable to have $n\delta^{2J+2} \rightarrow 0$. The latter condition prescribes a rule on the selection of the $J = J_n(\delta)$. By choosing an $\varepsilon > 0$ so that $\delta^{2J+2} = n^{-1-\varepsilon}$ for each pair of n and δ , then

$$J = J_n(\delta) = \frac{-1 - \varepsilon}{2 \log \delta} \log n - 1 > \frac{-1}{2 \log \delta} \log n - 1.$$

The integer truncation of the above lower bound plus one can be used as a reference value for the number of terms used in the density approximation for each given pair of (n, δ) .

TABLE 1
The least approximation term selection to guarantee the AMLE has the same asymptotic distribution as the full MLE for special sampling interval δ and sample size n

δ	$n = 500$	$n = 1,000$	$n = 2,000$	$n = 4,000$
1/252	1	1	1	1
1/52	1	1	1	1
1/12	1	1	1	1
1/4	2	2	2	2
1/2	4	4	5	5
3/4	10	12	13	14

Table 1 reports such reference values of J assigned by the above formula for a set of (n, δ) combinations commonly encountered in empirical studies. It shows that for monthly frequency or less ($\delta \leq 1/12$), one term approximation is adequate, and for $\delta = 1/4$, $J = 2$ is needed. However, there is a dramatic increase in J as the sampling length is larger than $1/4$: demanding at least four terms for $\delta = 1/2$ (half yearly) or at least ten terms for $\delta = 3/4$. The number of terms also increases for these higher δ values as n increases, although the rate of this increase is much slower than that as δ is increased. The latter may be understood that for a given δ , as n increases, the chance of having extreme values in the tails of the transition distribution increases. As the density approximation is less accurate in the tails than in the main body of the distribution, there is a need for having more terms in the density approximation.

5.2. J fixed, $\delta \rightarrow 0$ but $n\delta \rightarrow \infty$. Our starting point is the expansion (4.1). As $N_n - N = O_p\{(n\delta)^{-1/2}\}$, $N^{-1}(N_n - N) = o_p(1)$ if $n\delta^3 \rightarrow \infty$, which is also required in the asymptotic normality of the full MLE as outlined in Proposition 6. We will show in the following that $n\delta^3 \rightarrow \infty$ is also necessary to ensure AMLE sharing the same asymptotic distribution as the full MLE. It is understood that in order for $\hat{\theta}_n^{(J)}$ having the same asymptotic distribution as $\hat{\theta}_n$, it is required that

$$N^{-1}U_n, N^{-1}\Delta_{n1}(\hat{\theta}_n^{(J)}, \theta_0) \text{ and } N^{-1}\Delta_{n2}(\hat{\theta}_n, \theta_0) \quad \text{are all } o_p\{\|\hat{\theta}_n^{(J)} - \theta_0\|_2\}.$$

We will demonstrate in the following that the above requirements can be attained by $n\delta^3 \rightarrow \infty$ and $J \geq 2$. Hence, under these circumstances, $\hat{\theta}_n^{(J)}$ has the same asymptotic distribution as $\hat{\theta}_n$. Later we will demonstrate that this equivalence in the asymptotic distribution is quite unlikely for $J = 1$. Our analysis needs to expand (3.2) to the quadratic terms. To this end, let us define

$$M_n(\theta_0, J, \delta) = n^{-1} \sum_{t=1}^n \nabla_{\theta\theta\theta}^3 \log f^{(J)}(X_t|X_{t-1}, \delta; \theta_0)$$

and

$$T_n(\theta_0, J, \delta) = n^{-1} \sum_{t=1}^n \nabla_{\theta\theta\theta}^3 \log f(X_t | X_{t-1}, \delta; \theta_0).$$

By further expanding to quadratic terms, (4.1) can be written as

$$\begin{aligned} & \hat{\theta}_n^{(J)} - \theta_0 \\ &= N^{-1}U_n + N^{-1}(N_n + F_n)(\hat{\theta}_n - \theta_0) - N^{-1}(N_n - N)(\hat{\theta}_n^{(J)} - \theta_0) \\ (5.1) \quad & - \frac{1}{2}N^{-1}[E_d \otimes (\hat{\theta}_n^{(J)} - \theta_0)']M_n(\hat{\theta}_n^{(J)} - \theta_0) \\ & + \frac{1}{2}N^{-1}[E_d \otimes (\hat{\theta}_n - \theta_0)']T_n(\hat{\theta}_n - \theta_0) \\ & - N^{-1}\tilde{\Delta}_{n1}(\hat{\theta}_n^{(J)}, \theta_0) + N^{-1}\tilde{\Delta}_{n2}(\hat{\theta}_n, \theta_0), \end{aligned}$$

where $\tilde{\Delta}_{n1}(\hat{\theta}_n^{(J)}, \theta_0)$ and $\tilde{\Delta}_{n2}(\hat{\theta}_n, \theta_0)$ are remainder terms. Using the same method in the proof of Proposition 1, it can be shown that $\tilde{\Delta}_{n1}(\hat{\theta}_n^{(J)}, \theta_0) = O_p\{\|\hat{\theta}_n^{(J)} - \theta_0\|_2^3\}$ and $\tilde{\Delta}_{n2}(\hat{\theta}_n, \theta_0) = O_p\{\|\hat{\theta}_n - \theta_0\|_2^3\}$.

In order to make $\hat{\theta}_n^{(J)}$ have the same asymptotic distribution as $\hat{\theta}_n$, the two quadratic terms on the right-hand side of (5.1) have to be smaller order of $\hat{\theta}_n^{(J)} - \theta_0$ and $\hat{\theta}_n - \theta_0$, respectively, namely

$$N^{-1}[E_d \otimes (\hat{\theta}_n^{(J)} - \theta_0)']M_n(\hat{\theta}_n^{(J)} - \theta_0) = o_p\{\|\hat{\theta}_n^{(J)} - \theta_0\|_2\}$$

or equivalently

$$(5.2) \quad N^{-1}[E_d \otimes (\hat{\theta}_n^{(J)} - \theta_0)'] = o_p(1)$$

and

$$N^{-1}[E_d \otimes (\hat{\theta}_n - \theta_0)']T_n(\hat{\theta}_n - \theta_0) = o_p\{\|\hat{\theta}_n - \theta_0\|_2\}$$

or equivalently

$$(5.3) \quad n\delta^3 \rightarrow \infty$$

since $\hat{\theta}_n - \theta_0 = O_p\{(n\delta)^{-1/2}\}$ and $N^{-1} = O(\delta^{-1})$.

As $\hat{\theta}_n^{(J)} - \theta_0 = O_p\{\delta^J + (n\delta)^{-1/2}\}$, (5.2) requires that $\delta^{J-1} + n^{-1/2}\delta^{-3/2} \rightarrow 0$. Hence, in order to make $\hat{\theta}_n^{(J)}$ have the same asymptotic distribution as $\hat{\theta}_n$, it is necessary to have

$$(5.4) \quad J \geq 2 \quad \text{and} \quad n\delta^3 \rightarrow \infty.$$

Now we consider the case of $J = 1$. To ensure the remainder terms $N^{-1} \times \Delta_{n1}(\hat{\theta}_n^{(J)}, \theta_0)$ and $N^{-1}\Delta_{n2}(\hat{\theta}_n, \theta_0)$ are negligible, by a similar argument applied above for the case of $J \geq 2$, it is also necessary to assume $n\delta^3 \rightarrow \infty$. From Theorem 2, $\hat{\theta}_n^{(1)} - \theta_0 = O_p\{\delta + (n\delta)^{-1/2}\}$. To gain insight on the situation, we need to

find out the order of magnitude of the quadratic term in (5.1), namely the order of magnitude of

$$S_n = N^{-1}[E_d \otimes (\hat{\theta}_n^{(1)} - \theta_0)']M_n(\hat{\theta}_n^{(1)} - \theta_0) - N^{-1}[E_d \otimes (\hat{\theta}_n - \theta_0)']T_n(\hat{\theta}_n - \theta_0).$$

With this notation, (5.1) can be written as

$$(5.5) \quad \begin{aligned} \hat{\theta}_n^{(J)} - \theta_0 &= N^{-1}U_n + N^{-1}(N_n + F_n)(\hat{\theta}_n - \theta_0) - \frac{1}{2}S_n \\ &+ o_p\{(n\delta)^{-1/2}\} + O_p(\delta^2). \end{aligned}$$

Define an operator between two vectors A and B ,

$$A * B = [E_d \otimes A']M_n B + [E_d \otimes B']M_n A.$$

By repeated substitutions, it can be shown that

$$\begin{aligned} S_n &= \frac{1}{2}N^{-1}[(N^{-1}U_n) * (N^{-1}U_n)] + \frac{1}{2}N^{-1}[(\frac{1}{2}S_n) * (\frac{1}{2}S_n)] \\ &- N^{-1}[(N^{-1}U_n) * (\frac{1}{2}S_n)] + o_p(\delta). \end{aligned}$$

As $U_n = O_p(\delta^2)$ for $J = 1$ and $N^{-1} = O(\delta^{-1})$, it can be deduced from the above equation that $S_n = O_p(\delta)$. Hence, for $J = 1$ if we require $n\delta^3 \rightarrow \infty$, the quadratic term S_n will contribute to the leading order of $\hat{\theta}_n^{(1)} - \theta_0$. If we do not require $n\delta^3 \rightarrow \infty$, then the sum of remainder terms, $N^{-1}\tilde{\Delta}_{n1}(\hat{\theta}_n^{(J)}, \theta_0) + N^{-1}\tilde{\Delta}_{n2}(\hat{\theta}_n, \theta_0)$ will not be controlled. Hence, if $J = 1$, it is very likely that the asymptotic distribution of $\hat{\theta}_n^{(J)}$ will differ from that of $\hat{\theta}_n$ unless $U_n = 0$ with probability one. In the rare case of $U_n = 0$, it is possible for $\hat{\theta}_n^{(1)}$ and $\hat{\theta}_n$ to share the same limiting distribution.

Therefore, in order to guarantee that $\hat{\theta}_n^{(J)}$ has the same asymptotic distribution as $\hat{\theta}_n$ under $\delta \rightarrow 0$, we need to use the AMLE based on at least two-term expansions, while satisfying $n\delta^3 \rightarrow \infty$, which we will assume in the rest of this section.

Note that $\hat{\theta}_n^{(J)} - \theta_0 = O_p\{\delta^J + (n\delta)^{-1/2}\}$. Then,

$$\begin{aligned} \hat{\theta}_n^{(J)} - \theta_0 &= N^{-1}U_n + (\hat{\theta}_n - \theta_0) \\ &+ O_p(n^{-1/2}\delta^{J-3/2}) + N^{-1} \cdot O_p(\delta^{2J} + n^{-1}\delta^{-1}). \end{aligned}$$

Furthermore,

$$\begin{aligned} \sqrt{n}I^{1/2}(\delta)(\hat{\theta}_n^{(J)} - \theta_0) &= \sqrt{n}I^{-1/2}(\delta)I(\delta)N^{-1}U_n + \sqrt{n}I^{1/2}(\delta)(\hat{\theta}_n - \theta_0) + O_p(\delta^{J-3/2}) \\ &+ \sqrt{n}I^{-1/2}(\delta)I(\delta)N^{-1} \cdot O_p(\delta^{2J} + n^{-1}\delta^{-1}) \\ &= \sqrt{n}I^{1/2}(\delta)(\hat{\theta}_n - \theta_0) + O_p(\delta^{J-3/2} + n^{-1/2}\delta^{-3/2} + n^{1/2}\delta^{J+1/2}). \end{aligned}$$

Hence, for any $J \geq 2$ such that $n\delta^3 \rightarrow \infty$ and $n\delta^{2J+1} \rightarrow 0$,

$$\sqrt{n}I^{1/2}(\delta)(\hat{\theta}_n^{(J)} - \theta_0) \xrightarrow{d} N(0, E_d).$$

This result shows that, when δ vanishes to zero, in order to guarantee the AMLE has the same asymptotic distribution as full MLE, we need to pick the approximation order $J \geq 2$, while maintaining $n\delta^3 \rightarrow \infty$ and $n\delta^{2J+1} \rightarrow 0$.

The following theorem summarizes the asymptotic normality under both asymptotic regimes.

THEOREM 3. *Under conditions (A.1)–(A.7) given in the Appendix,*

$$\sqrt{n}I^{1/2}(\delta)(\hat{\theta}_n^{(J)} - \theta_0) \xrightarrow{d} N(0, E_d)$$

for: (i) $\delta \in (0, \tilde{\Delta} \wedge \dot{\Delta}]$ being fixed, $n \rightarrow \infty$, $J \rightarrow \infty$ but $n\delta^{2J+2} \rightarrow 0$ or (ii) $J \geq 2$ being fixed, $n \rightarrow \infty$, $\delta \rightarrow 0$ but $n\delta^3 \rightarrow \infty$ and $n\delta^{2J+1} \rightarrow 0$.

5.3. Asymptotic bias and variance. The remainder of this section is devoted to the consideration of the asymptotic bias and variance of the AMLE under the two asymptotic regimes. Given our analysis in the early part of this section, our consideration will be focused on the situations where the asymptotic normality of the AMLE can be assumed, namely under: (i) δ being fixed, $J \rightarrow \infty$, $n \rightarrow \infty$ but $n\delta^{2J+2} \rightarrow 0$ or (ii) $J \geq 2$ being fixed, $\delta \rightarrow 0$, $n\delta^3 \rightarrow \infty$ but $n\delta^{2J+1} \rightarrow 0$.

In the case of δ being fixed and $J \rightarrow \infty$, from (5.1) and provided $n\delta^{2J+2} \rightarrow 0$, we have

$$\begin{aligned} \hat{\theta}_n^{(J)} - \theta_0 &= N^{-1}U_n + N^{-1}(N_n + F_n)(\hat{\theta}_n - \theta_0) - N^{-1}(N_n - N)N^{-1}U_n \\ &\quad - N^{-1}(N_n - N)N^{-1}(N_n + F_n)(\hat{\theta}_n - \theta_0) \\ &\quad - \frac{1}{2}N^{-1}\{E_d \otimes [N^{-1}U_n + N^{-1}(N_n + F_n)(\hat{\theta}_n - \theta_0)]'\} \\ &\quad \times M_n[N^{-1}U_n + N^{-1}(N_n + F_n)(\hat{\theta}_n - \theta_0)] \\ &\quad + \frac{1}{2}N^{-1}[E_d \otimes (\hat{\theta}_n - \theta_0)']T_n(\hat{\theta}_n - \theta_0) + O_p(n^{-3/2}) \\ &= N^{-1}U_n + [E_d - N^{-1}(N_n - N)]N^{-1}(N_n + F_n)(\hat{\theta}_n - \theta_0) \\ &\quad + O_p(n^{-1/2}\delta^{J+1}) + O_p(n^{-3/2}). \end{aligned}$$

Then, the leading order bias of $\hat{\theta}_n^{(J)}$ is

$$(5.6) \quad B(\theta_0, J, \delta) = N^{-1}U + \mathbb{E}\{[E_d - N^{-1}(N_n - N)]N^{-1}(N_n + F_n)(\hat{\theta}_n - \theta_0)\},$$

and the leading order variance is

$$(5.7) \quad V(\theta_0, J, \delta) = N^{-1}I(\delta) \text{Var}(\hat{\theta}_n)I(\delta)N^{-1}.$$

In the case of $J \geq 2$ being fixed, $\delta \rightarrow 0$ and $n\delta^3 \rightarrow \infty$ but $n\delta^{2J+1} \rightarrow 0$, it can be shown by a similar argument to that for the fixed δ case above, the asymptotic bias and variance have the same forms as (5.6) and (5.7), respectively. Both (5.6) and (5.7) will be used to calibrate with the simulated bias and variance in the simulation study in Section 7. For $J = 1$ and $\delta \rightarrow 0$, there are difficulties in obtaining an expression for the bias of the AMLE in general due to the same dilemma in controlling the reminder terms and the quadratic term S_n as outlined in Section 5.2.

6. Approximating Fisher information matrix. We demonstrate in this section that the approximation of the transition density provides a way to approximate the Fisher information matrix. Fisher information matrix $I(\delta)$ is a key quantity associated with inference based on the full MLE. It defines the asymptotic efficiency and convergence rate. From Proposition 2, a natural candidate to approximate $I(\delta)$ is $-N(\theta_0, J, \delta)$ based on the J -term expansion. To simplify our exposition, our consideration here is focused under the following diffusion process:

$$(6.1) \quad dX_t = \mu(X_t; \eta) dt + \sigma(X_t; \xi) dB_t,$$

where $\eta = (\eta_1, \dots, \eta_{d_1})'$ and $\xi = (\xi_1, \dots, \xi_{d_2})'$ are distinct drift and diffusion parameters, respectively. The whole parameter $\theta = (\eta', \xi')'$. Here, we provide an explicit expression $N(\theta_0, 1, \delta)$ based on the one-term density expansion. Expressions for higher J values may be made via more extensive derivations.

Let μ_i, μ_{ij} and so on denote partial derivatives with respect to η_i, η_i and η_j , respectively; and σ_i and $\sigma_{x,j}$ and so on denote partial derivatives with respect to ξ_i , and x and ξ_j , respectively. By the one-term ($J = 1$) transition density approximation, derivations given in Chang and Chen (2011) show that

$$\mathbb{E}\left(\frac{\partial^2 \log f^{(1)}}{\partial \eta_i \partial \eta_j}\right) =: \delta \cdot N_{11}^{(1)} + O(\delta^2), \quad \mathbb{E}\left(\frac{\partial^2 \log f^{(1)}}{\partial \eta_i \partial \xi_j}\right) =: \delta \cdot N_{12}^{(1)} + O(\delta^2)$$

and

$$\mathbb{E}\left(\frac{\partial^2 \log f^{(1)}}{\partial \xi_i \partial \xi_j}\right) =: -2\mathbb{E}(\sigma^{-2} \sigma_i \sigma_j) + \delta \cdot N_{22}^{(1)} + O(\delta^2),$$

where

$$\begin{aligned} N_{11}^{(1)} &= \mathbb{E}\{-\sigma^{-2} \mu_i \mu_j - \mu \sigma^{-2} \mu_{ij} + \sigma^{-1} \mu_{ij} \sigma_x - \frac{1}{2} \mu_{xij}\}, \\ N_{12}^{(1)} &= \mathbb{E}\{2\mu \sigma^{-3} \mu_i \sigma_j - \sigma^{-2} \mu_i \sigma_x \sigma_j + \sigma^{-1} \mu_i \sigma_{xj}\}, \\ N_{22}^{(1)} &= \mathbb{E}\{-6\mu^2 \sigma^{-4} \sigma_i \sigma_j + 16\mu \sigma^{-3} \sigma_x \sigma_i \sigma_j + 2\mu^2 \sigma^{-3} \sigma_{ij} - 3\sigma^{-2} \mu_x \sigma_i \sigma_j \\ &\quad - \frac{19}{2} \sigma^{-2} \sigma_x^2 \sigma_i \sigma_j - \frac{9}{2} \mu \sigma^{-2} \sigma_x \sigma_{ij} - 5\mu \sigma^{-2} \sigma_{xi} \sigma_j - 5\mu \sigma^{-2} \sigma_{xj} \sigma_i \\ &\quad + \sigma^{-1} \mu_x \sigma_{ij} + 4\sigma^{-1} \sigma_{xx} \sigma_i \sigma_j + \frac{11}{2} \sigma^{-1} \sigma_x \sigma_{xi} \sigma_j + \frac{11}{2} \sigma^{-1} \sigma_x \sigma_{xj} \sigma_i \\ &\quad + \frac{3}{2} \sigma^{-1} \sigma_x^2 \sigma_{ij} + \frac{5}{2} \mu \sigma^{-1} \sigma_{xij} - \frac{3}{4} \sigma_{xx} \sigma_{ij} - \frac{5}{2} \sigma_{xi} \sigma_{xj} - \frac{3}{2} \sigma_x \sigma_{xij} \\ &\quad - \sigma_{xxi} \sigma_j - \sigma_{xxj} \sigma_i + \frac{3}{4} \sigma \sigma_{xij}\}. \end{aligned}$$

Thus

$$(6.2) \quad N(\theta_0, 1, \delta) = \begin{pmatrix} \delta \cdot N_{11}^{(1)} & \delta \cdot N_{12}^{(1)} \\ \delta \cdot N_{12}^{(1)'} & -2 \cdot \mathbb{E}(\sigma^{-2} \sigma_i \sigma_j) + \delta \cdot N_{22}^{(1)} \end{pmatrix} + O(\delta^2).$$

We learn from Proposition 2 that $-N(\theta_0, 1, \delta)$ provides a leading order approximation to $I(\delta)$ with a reminder term at the order of δ^2 . Equation (6.2) confirms

that as $\delta \rightarrow 0$, given the asymptotic normality of the full MLE $\hat{\theta}_n$ as conveyed by Proposition 6, that the convergence rate of the full MLE for the drift parameters η is $(n\delta)^{-1/2}$ whereas that for the diffusion parameters ξ is $n^{-1/2}$, faster than the drift parameter estimator. Our study confirms the results of Gobet (2002), Sørensen (2007) and Tang and Chen (2009).

In the rest of the section, we will derive the Fisher information matrix approximation for two specific diffusion processes. Both are widely employed in modeling of the interest rate dynamics.

6.1. *Vasicek model.* Consider the Vasicek (1977) model,

$$(6.3) \quad dX_t = \kappa(\alpha - X_t) dt + \sigma dB_t,$$

which is also the Ornstein–Uhlenbeck process. The conditional distribution of X_t given X_{t-1} is

$$X_t | X_{t-1} \sim N\left\{X_{t-1}e^{-\kappa\delta} + \alpha(1 - e^{-\kappa\delta}), \frac{1}{2}\sigma^2\kappa^{-1}(1 - e^{-2\kappa\delta})\right\},$$

and the stationary distribution of $\{X_t\}$ is $N(\alpha, \frac{\sigma^2}{2\kappa})$. It yields that the information matrix of $\theta = (\kappa, \alpha, \sigma)'$ is $I(\delta) = (I_{ij})_{3 \times 3}$ where

$$\begin{aligned} I_{11} &= \frac{1}{2\kappa^2} + \frac{\delta[\kappa\delta + \kappa\delta e^{2\kappa\delta} - 2e^{2\kappa\delta} + 2]}{\kappa(e^{2\kappa\delta} - 1)^2} = \frac{\delta}{2\kappa} + O(\delta^2), & I_{12} = I_{21} &= 0, \\ I_{13} = I_{31} &= \frac{(1 + 2\kappa\delta) - e^{2\kappa\delta}}{\sigma\kappa(e^{2\kappa\delta} - 1)} = -\frac{\delta}{\sigma} + O(\delta^2), \\ I_{22} &= \frac{2\kappa(e^{\kappa\delta} - 1)^2}{\sigma^2(e^{2\kappa\delta} - 1)} = \frac{\kappa^2\delta}{\sigma^2} + O(\delta^2), & I_{23} = I_{32} &= 0 \quad \text{and} \quad I_{33} = \frac{2}{\sigma^2}. \end{aligned}$$

These mean that

$$(6.4) \quad I(\delta) = \begin{pmatrix} \delta \cdot (2\kappa)^{-1} & 0 & -\delta \cdot \sigma^{-1} \\ 0 & \delta \cdot \kappa^2\sigma^{-2} & 0 \\ -\delta \cdot \sigma^{-1} & 0 & 2\sigma^{-2} \end{pmatrix} + O(\delta^2).$$

Hence $I(0) = \lim_{\delta \rightarrow 0} I(\delta)$ is singular, an issue we have raised earlier, which makes us assume that $\delta I^{-1}(\delta)$'s largest eigenvalue is bounded in condition (A.5).

Using the approximation formula in (6.2), we have

$$N(\theta, 1, \delta) = \begin{pmatrix} -\delta \cdot (2\kappa)^{-1} & 0 & \delta \cdot \sigma^{-1} \\ 0 & -\delta \cdot \kappa^2\sigma^{-2} & 0 \\ \delta \cdot \sigma^{-1} & 0 & -2\sigma^{-2} \end{pmatrix} + O(\delta^2).$$

This means the leading order term of $-N(\theta, 1, \delta)$ is identical with that of the true Fisher information matrix in (6.4).

6.2. *Cox–Ingersoll–Ross model.* Consider the Cox–Ingersoll–Ross (CIR) model [Cox, Ingersoll and Ross (1985)],

$$(6.5) \quad dX_t = \kappa(\alpha - X_t) dt + \sigma\sqrt{X_t} dB_t,$$

which is also Feller’s (1952) square root process.

Let $\theta = (\kappa, \alpha, \sigma)'$ and $c = 4\kappa\sigma^{-2}(1 - e^{-\kappa\delta})^{-1}$. The conditional distribution of cX_t given X_{t-1} is

$$cX_t|X_{t-1} \sim \chi^2_\nu(\lambda),$$

where the distribution is a noncentral χ^2 distribution with degree of freedom $\nu = 4\kappa\alpha\sigma^{-2}$ and noncentral parameter $\lambda = cX_{t-1}e^{-\kappa\delta}$. The transition density of X_t given X_{t-1} is

$$f(X_t|X_{t-1}, \delta; \theta) = \frac{c}{2}e^{-u-v} \left(\frac{v}{u}\right)^{q/2} I_q(2\sqrt{uv}),$$

where $u = cX_{t-1}e^{-\kappa\delta}/2$, $v = cX_t/2$, $q = 2\kappa\alpha/\sigma^2 - 1 \geq 0$, and I_q is the modified Bessel function of the first kind of order q . If $2\kappa\alpha > \sigma^2$, then the stationary distribution of $\{X_t\}$ is $\Gamma(\frac{2\kappa\alpha}{\sigma^2}, \frac{\sigma^2}{2\kappa})$.

Although the second partial derivations of the log transition density function can be derived after some labor that is involved with differentiating the modified Bessel function of the first kind, acquiring an expression for the Fisher information matrix is a rather hard task, largely due to the difficulty in deriving the expectations. In contrast, using the approximation formula (6.2), we can obtain the approximation for the opposite Fisher information matrix,

$$N(\theta_0, 1, \delta) = \begin{pmatrix} N_{11} & N_{12} & N_{13} \\ N_{21} & N_{22} & N_{23} \\ N_{31} & N_{32} & N_{33} \end{pmatrix} + O(\delta^2),$$

where

$$\begin{aligned} N_{11} &= \delta \cdot \frac{\alpha^2\sigma^2 - 2\kappa\alpha^2 + \alpha\sigma^2}{\sigma^4 - 2\kappa\alpha\sigma^2}, \\ N_{12} = N_{21} &= \delta \cdot \frac{4\kappa\alpha\sigma^2 - \sigma^4 - 8\kappa^2\alpha + 4\kappa\sigma^2}{2\sigma^4 - 4\kappa\alpha\sigma^2}, \\ N_{13} = N_{31} &= -\delta \cdot \frac{2\kappa\alpha^2\sigma^2 - 4\kappa^2\alpha^2 + 2\kappa\alpha\sigma^2}{\sigma^5 - 2\kappa\alpha\sigma^3}, \\ N_{22} &= \delta \cdot \frac{\kappa^2}{\sigma^2 - 2\kappa\alpha}, \\ N_{23} = N_{32} &= -\delta \cdot \frac{2\kappa^2\alpha\sigma^2 - 4\kappa^3\alpha + 2\kappa^2\sigma^2}{\sigma^5 - 2\kappa\alpha\sigma^3} \end{aligned}$$

and

$$N_{33} = \frac{-2}{\sigma^2} + \delta \cdot (24\kappa^2\alpha^2\sigma^2 - 48\kappa^3\alpha^2 + 48\kappa^2\alpha\sigma^2 - 24\kappa\alpha\sigma^4 + 36\kappa\sigma^4 + 4\sigma^5 + 9\sigma^6)(4\sigma^6 - 8\kappa\alpha\sigma^4)^{-1}.$$

Using $-N(\theta_0, 1, \delta)$, we can get the approximation of the Fisher information matrix. This approximation may be used in carrying out statistical inference on the CIR processes.

6.3. Observed Fisher information. The major application for the asymptotic normality of both the full and approximate MLEs is for statistical inference of θ , which include confidence regions and testing hypotheses for θ . For such purposes, the Fisher information $I(\delta)$ needs to be estimated. A natural candidate would be $-N_n(\hat{\theta}_n^{(J)}, J, \delta)$. Although it converges to $I(\delta)$ at the rate of $O_p\{(n\delta)^{-1/2} + \delta^J\}$ or $O_p\{(n\delta)^{-1/2} + \delta^{J+1}\}$, depending on whether δ is fixed or diminishing, $-N_n(\hat{\theta}_n^{(J)}, J, \delta)$ may not be nonnegative definite, which can hinder the acquisition of $\{-N_n(\hat{\theta}_n^{(J)}, J, \delta)\}^{1/2}$. To get around this issue, by noticing that $I(\delta)$ is the variance of the likelihood score, we consider

$$\tilde{I}_n(\theta, J, \delta) = \frac{1}{n} \sum_{t=1}^n [\nabla_{\theta} \log f^{(J)}(X_t|X_{t-1}, \delta; \theta)][\nabla_{\theta} \log f^{(J)}(X_t|X_{t-1}, \delta; \theta)]'$$

as an estimator of $I(\delta)$. The following theorem shows this by replacing $I(\delta)$ with $\tilde{I}_n(\hat{\theta}_n^{(J)}, J, \delta)$ in Theorem 3.

THEOREM 4. *Under conditions (A.1)–(A.7) given in the Appendix,*

$$\sqrt{n} \tilde{I}_n^{1/2}(\hat{\theta}_n^{(J)}, J, \delta)(\hat{\theta}_n^{(J)} - \theta_0) \xrightarrow{d} N(0, E_d)$$

for: (i) $\delta \in (0, \tilde{\Delta} \wedge \hat{\Delta}]$ being fixed, $n \rightarrow \infty$, $J \rightarrow \infty$ but $n\delta^{2J+2} \rightarrow 0$ or (ii) $J \geq 2$ being fixed, $n \rightarrow \infty$, $\delta \rightarrow 0$ but $n\delta^3 \rightarrow \infty$ and $n\delta^{2J+1} \rightarrow 0$.

Confidence regions and testing hypotheses can be readily carried out by utilizing the above results.

7. Simulation. We report results from simulation studies which are designed to confirm the theoretical findings on the AMLE as reported in the earlier sections. To allow verification with the full MLE, we considered the Vasicek and CIR diffusion models reported in the previous section as both models permit the full MLE. The two asymptotic regimes were experimented: the fixed δ and the diminishing δ with $n\delta^3 \rightarrow \infty$.

The first part of the simulation is about the case in which δ is fixed. The parameters used in the simulated Vasicek and CIR models were $\theta = (\kappa, \alpha, \sigma)'$ =

$(0.858, 0.0891, 0.0468)'$ and $\theta = (\kappa, \alpha, \sigma)' = (0.892, 0.09, 0.1817)'$, respectively. The sampling interval δ was $1/12$ and $1/4$, and the order of the density approximation J was 1 and 2, respectively. For each δ and J , the sample size n was set at 500, 1,000 and 2,000, respectively. In addition to bias and standard deviation, we consider

$$\text{RMSD}(n, J, \delta) = \sqrt{\mathbb{E}\|\hat{\theta}_n^{(J)} - \hat{\theta}_n\|_2^2},$$

the square root of the expected square of modulated deviations between $\hat{\theta}_n^{(J)}$ and $\hat{\theta}_n$, as an overall performance measure.

Tables 2 and 3 summarize the simulation for the fixed δ case. They report the average bias and standard deviation (SD) for the full MLE and AMLEs with $J = 1$ and $J = 2$, as well as the RMSD between the AMLEs and the full MLE, for both the Vasicek and the CIR models. To give the simulation results more perspective and to confirm the derived approximate bias and variance formulas in Section 5, we also computed the asymptotic bias and standard deviation based on formulas (5.6) and (5.7). We observe from Tables 2 and 3 that at each δ ($1/12$ and $1/4$) experimented, the bias and the standard deviation of all the estimators for the three parameters became smaller as n increased. These confirmed the consistency of the estimators. The tables also showed that there was a good agreement among the three estimators in terms of the performance measures. It appeared that the bias and the variance of the AMLE with $J = 1$ and $J = 2$ were quite comparable to each other. However, by comparing RMSD, it was clear that in most of the cases (except for $n = 500$ of CIR model), the RMSD for $J = 2$ was smaller than $J = 1$, signaling the AMLE with $J = 2$ was closer to the full MLE than that of the AMLE with $J = 1$. This indicates that the AMLEs with $J = 2$ were indeed closer to those with $J = 1$, as confirmed by our early analysis. The asymptotic bias and standard deviation predicted for the AMLE with $J = 1$ and 2 offer more insights, and show good agreement between the simulated results and the predicted values by the theory, which is very assuring. We also observe that for $\delta = 1/4$, the AMLE with $J = 2$ performs better than AMLE with $J = 1$, which somehow reflects Table 1 which shows that $J = 2$ is preferred to $J = 1$ at this frequency. When δ was fixed at $1/12$, we see the performance between $J = 1$ and $J = 2$ was largely similar.

The second part of the simulation was devoted to the diminishing δ case. Here we wanted to confirm the differential behavior of the AMLEs in the limiting distribution between $J = 1$ and $J \geq 2$, as revealed in Section 5. The Vasicek model with $\theta = (\kappa, \alpha, \sigma)' = (0.892, 0.09, 0.1817)'$ was considered. We tried to create two scenarios: (i) $n\delta^3 \rightarrow \infty$ and (ii) $n\delta^3 \rightarrow 0$, while $\delta \rightarrow 0$. They were created by choosing $\delta = n^{-1/6}$ and $\delta = n^{-1/2}$, respectively, while selecting $n = 500, 1,000, 2,000, 4,000$ and $8,000$, respectively, to create two streams of asymptotic sequences. For each n and δ , we generated repeatedly the Vasicek sample paths 1,000 times. For each simulated sample path, we obtained the AMLEs $\hat{\theta}_n^{(J)}$ for $J = 1$ and 2, respectively, and computed the Wald statistics

$$W_n(J) = n(\hat{\theta}_n^{(J)} - \theta_0)' I(\delta)(\hat{\theta}_n^{(J)} - \theta_0).$$

TABLE 2
Simulated average bias (Bias) and standard deviations (SD) of the full MLE and two AMLEs with $J = 1$ and 2 for Vasicek model ($\kappa = 0.858, \alpha = 0.0891, \sigma = 0.0468$); A.Bias and A.SD are asymptotic bias and SD based on formulas (5.6) and (5.7); RMSD is the root of mean square deviation between $\hat{\theta}_n$ and $\hat{\theta}_n^{(J)}$

n	Statistics	$\delta = 1/12$			$\delta = 1/4$			
		MLE	$J = 1$	$J = 2$	MLE	$J = 1$	$J = 2$	
500	Bias	κ	0.0992	0.0896	0.0992	0.0380	0.0127	0.0396
		α	0.0002	0.0002	0.0002	4.09e-5	5.63e-5	4.17e-5
		σ	4.39e-5	4.14e-5	4.39e-5	9.12e-5	7.13e-5	9.43e-5
	A.Bias	κ		0.0908	0.1016		0.0174	0.0376
		α		0.0003	0.0002		0.0002	0.0001
		σ		4.55e-5	4.55e-5		0.0001	0.0001
	SD	κ	0.2307	0.2255	0.2309	0.1366	0.1290	0.1386
		α	0.0085	0.0085	0.0085	0.0050	0.0050	0.0050
		σ	0.0016	0.0016	0.0016	0.0016	0.0016	0.0016
	A.SD	κ		0.2251	0.2366		0.1215	0.1403
		α		0.0084	0.0085		0.0047	0.0050
		σ		0.0016	0.0016		0.0016	0.0016
	RMSD	κ		0.0173	0.0062		0.0332	0.0316
		α		0.0002	1.28e-5		0.0005	0.0002
		σ		1.36e-5	1.05e-5		0.0001	0.0001
1,000	Bias	κ	0.0518	0.0419	0.0520	0.0170	-0.0095	0.0186
		α	-0.0002	-0.0002	-0.0002	1.83e-5	2.81e-5	1.58e-5
		σ	7.05e-5	6.68e-5	7.06e-5	3.66e-5	6.83e-6	3.96e-5
	A.Bias	κ		0.0446	0.0529		-0.0097	0.0161
		α		-0.0001	-0.0002		1.69e-5	1.45e-5
		σ		0.0001	0.0001		3.29e-5	4.55e-5
	SD	κ	0.1624	0.1586	0.1625	0.0957	0.0905	0.0966
		α	0.0058	0.0058	0.0058	0.0034	0.0034	0.0034
		σ	0.0011	0.0011	0.0011	0.0012	0.0012	0.0012
	A.SD	κ		0.1585	0.1666		0.0849	0.0982
		α		0.0057	0.0058		0.0032	0.0034
		σ		0.0011	0.0011		0.0012	0.0012
	RMSD	κ		0.0100	0.0008		0.0316	0.0063
		α		0.0001	9.14e-6		0.0004	0.0001
		σ		7.39e-6	7.80e-7		0.0001	1.59e-5

If $\sqrt{n}I^{1/2}(\delta)(\hat{\theta}_n^{(J)} - \theta_0)$ is asymptotically standard normally distributed in \mathbb{R}^d , then the Wald statistic $W_n(J) \xrightarrow{d} \chi_3^2$. Based on the 1,000 Wald statistics from the simulations, we then performed the Kolmogorov–Smirnov (K–S) test to test $H_0: W_n(J) \sim \chi_3^2$, or not, for each of the designed sequences of (n, δ) generated

TABLE 2
(Continued)

<i>n</i>	Statistics	$\delta = 1/12$			$\delta = 1/4$			
		MLE	<i>J</i> = 1	<i>J</i> = 2	MLE	<i>J</i> = 1	<i>J</i> = 2	
2,000	Bias	κ	0.0245	0.0149	0.0246	0.0084	-0.0191	0.0100
		α	-3.97e-5	-3.34e-5	-4.01e-5	-5.72e-5	-4.90e-5	-5.80e-5
		σ	2.69e-5	2.30e-5	2.70e-5	4.00e-5	9.21e-6	4.34e-5
	A.Bias	κ		0.0179	0.0249		-0.0085	0.0071
		α		-2.63e-5	-2.98e-5		0.0001	-0.0001
		σ		4.55e-5	4.55e-5		4.55e-5	4.55e-5
	SD	κ	0.1114	0.1091	0.1115	0.0647	0.0611	0.0652
		α	0.0042	0.0041	0.0042	0.0024	0.0024	0.0024
		σ	0.0008	0.0008	0.0008	0.0008	0.0008	0.0008
	A.SD	κ		0.1088	0.1143		0.0576	0.0665
		α		0.0041	0.0042		0.0023	0.0024
		σ		0.0008	0.0008		0.0008	0.0008
	RMSD	κ		0.0100	0.0006		0.0300	0.0042
		α		0.0001	7.37e-6		0.0003	4.68e-5
		σ		6.27e-6	7.80e-7		0.0001	1.02e-5

under the two scenarios. Table 4 reports the *p*-values of the test, which show that for *J* = 1, under both scenarios, the *p*-values of the K-S test became smaller, and hence the above null hypothesis was rejected as *n* increased. For *J* = 2, the *p*-values of the K-S test were sharply different between the two scenarios. In particular, the *p*-values were mostly quite large under the scenario of $n\delta^3 \rightarrow \infty$, and they were largely significant (small) when δ was diminishing at the faster rate of $n^{-1/2}$ such that $n\delta^3 \rightarrow 0$. These were consistent with our theoretical findings in Section 5.

APPENDIX

We need the following technical assumptions in our analysis.

(A.1) (i) Θ is a compact set in \mathbb{R}^d , and the true parameter θ_0 is an interior point of Θ ; (ii) for all values of the parameters θ , Assumption 1–3 in Ait-Sahalia (2002) hold; (iii) the drift function $\mu(x; \theta)$ is a bona fide function of θ for each *x*.

(A.2) (i) For every $\delta > 0$, $\mathbb{E}\nabla_{\theta} \log f(X_t|X_{t-1}, \delta; \theta_0) = 0$, and θ_0 is the only root of $\mathbb{E}\nabla_{\theta} \log f(X_t|X_{t-1}, \delta; \theta) = 0$. (ii) the MLE $\hat{\theta}_n$ and the *J*-term approximate MLE $\hat{\theta}_n^{(J)}$ satisfy, respectively,

$$\sum_{t=1}^n \nabla_{\theta} \log f(X_t|X_{t-1}, \delta; \hat{\theta}_n) = 0 \quad \text{and} \quad \sum_{t=1}^n \nabla_{\theta} \log f^{(J)}(X_t|X_{t-1}, \delta; \hat{\theta}_n^{(J)}) = 0.$$

TABLE 3

Simulated average bias (Bias) and standard deviations (SD) of the full MLE and two AMLEs with $J = 1$ and 2 for CIR model ($\kappa = 0.892, \alpha = 0.09, \sigma = 0.1817$); A.Bias and A.SD are asymptotic bias and SD based on formulas (5.6) and (5.7); RMSD is the root of mean square deviation between $\hat{\theta}_n$ and $\hat{\theta}_n^{(J)}$

n	Statistics		$\delta = 1/12$			$\delta = 1/4$		
			MLE	$J = 1$	$J = 2$	MLE	$J = 1$	$J = 2$
500	Bias	κ	0.0980	0.0910	0.0978	0.0371	0.0234	0.0388
		α	0.0001	0.0004	0.0001	-6.38e-5	0.0008	-0.0001
		σ	0.0003	0.0003	0.0003	0.0004	0.0005	0.0003
	A.Bias	κ		0.0818	0.0984		0.0207	0.0513
		α		0.0005	0.0001		0.0008	-0.0001
		σ		0.0003	0.0003		0.0004	0.0002
	SD	κ	0.2389	0.2340	0.2405	0.1437	0.1338	0.2256
		α	0.0093	0.0093	0.0093	0.0055	0.0054	0.0055
		σ	0.0060	0.0060	0.0060	0.0065	0.0065	0.0069
	A.SD	κ		0.2169	0.2389		0.1159	0.1938
		α		0.0091	0.0093		0.0064	0.0055
		σ		0.0060	0.0060		0.0067	0.0065
	RMSD	κ		0.0200	0.0224		0.0447	0.1622
		α		0.0009	0.0004		0.0018	0.0004
		σ		0.0004	0.0004		0.0017	0.0021
1,000	Bias	κ	0.0521	0.0435	0.0521	0.0218	0.0070	0.0186
		α	-1.54e-5	0.0002	-2.22e-5	-0.0002	0.0007	-0.0003
		σ	3.86e-5	4.35e-5	3.81e-5	0.0003	0.0006	0.0003
	A.Bias	κ		0.0411	0.0525		0.0095	0.0262
		α		0.0004	-3.43e-5		0.0007	-0.0003
		σ		3.17e-5	2.69e-5		0.0003	0.0001
	SD	κ	0.1596	0.1558	0.1603	0.0968	0.0861	0.0980
		α	0.0067	0.0067	0.0067	0.0039	0.0037	0.0039
		σ	0.0043	0.0043	0.0043	0.0045	0.0045	0.0045
	A.SD	κ		0.1452	0.1596		0.0823	0.0969
		α		0.0066	0.0067		0.0044	0.0039
		σ		0.0040	0.0043		0.0047	0.0045
	RMSD	κ		0.0173	0.0141		0.0447	0.0200
		α		0.0003	2.66e-5		0.0020	0.0001
		σ		0.0002	3.91e-5		0.0021	0.0002

(A.3) There exist finite positive constants Δ and K_1 such that, for $l = 1, 2, 3$, any $\delta \in (0, \Delta]$, $i_1, i_2, i_3 \in \{1, \dots, d\}$ and $j = 1$ and 2,

$$\mathbb{E} \sup_{\theta \in \Theta} \left\{ \left| \frac{\partial^l A_j(X_t | X_{t-1}, \delta; \theta)}{\partial \theta_{i_1} \cdots \partial \theta_{i_l}} \right|^3 \right\} \leq K_1.$$

TABLE 3
(Continued)

<i>n</i>	Statistics	$\delta = 1/12$			$\delta = 1/4$			
		MLE	<i>J</i> = 1	<i>J</i> = 2	MLE	<i>J</i> = 1	<i>J</i> = 2	
2,000	Bias	κ	0.0295	0.0199	0.0294	0.0103	-0.0057	0.0069
		α	-0.0002	0.0001	-0.0002	-3.06e-5	0.0010	-9.87e-5
		σ	0.0002	0.0002	0.0002	3.05e-5	0.0006	1.33e-5
	A.Bias	κ		0.0213	0.0299		-0.0011	0.0147
		α		0.0002	-0.0002		0.0006	-0.0001
		σ		0.0002	0.0002		0.0005	1.06e-5
	SD	κ	0.1082	0.1053	0.1088	0.0696	0.0607	0.0698
		α	0.0048	0.0048	0.0048	0.0028	0.0027	0.0028
		σ	0.0030	0.0031	0.0030	0.0033	0.0037	0.0033
	A.SD	κ		0.1181	0.1105		0.0592	0.0697
		α		0.0047	0.0048		0.0027	0.0028
		σ		0.0030	0.0030		0.0034	0.0033
	RMSD	κ		0.0173	0.0068		0.0424	0.0100
		α		0.0004	0.0001		0.0020	0.0001
		σ		0.0005	0.0003		0.0027	0.0001

(A.4) There exist finite positive constants ν_l for $l = 0, 1, 2$ and 3 , $\Delta > 0$ and K_2 such that $\nu_0 > 3$, $\nu_2 > \nu_1 > 3$, $\nu_3 > 1$ and for any $i_1, \dots, i_3 \in \{1, \dots, d\}$ and $\delta \in (0, \Delta]$,

$$\mathbb{E} \left\{ \sup_{\theta \in \Theta} \left[\sum_{l=0}^{\infty} \left| \frac{\partial^q c_l(\gamma(X_t; \theta) | \gamma(X_{t-1}; \theta); \theta)}{\partial \theta_{i_1} \dots \partial \theta_{i_q}} \right| \frac{\Delta^l}{l!} \right]^{\nu_l} \right\} \leq K_2.$$

TABLE 4
p-values of Kolmogorov–Smirnov test for $W_n(J) \sim \chi_3^2$

Situation	<i>n</i>	δ	<i>J</i> = 1	<i>J</i> = 2
$\delta = n^{-1/6}$	500	0.3550	0.3524	0.0587
	1,000	0.3162	0.4595	0.5830
	2,000	0.2817	0.1149	0.2710
	4,000	0.2510	0.0019	0.8309
	8,000	0.2236	5.74e-8	0.6002
$\delta = n^{-1/2}$	500	0.0447	5.04e-7	2.45e-8
	1,000	0.0316	0.0003	9.72e-5
	2,000	0.0224	0.0006	0.0003
	4,000	0.0158	0.1109	0.0851
	8,000	0.0112	0.0470	0.0367

(A.5) For any $\delta > 0$, the Fisher information matrix

$$I(\delta) := -\mathbb{E}\nabla_{\theta\theta}^2 \log f(X_t|X_{t-1}, \delta; \theta_0)$$

is invertible and as $\delta \rightarrow 0$ the largest eigenvalues of $\delta I^{-1}(\delta)$ is bounded away from infinity.

(A.6) For each positive integer K , which may be infinite, and any $\delta \in (0, \Delta]$,

$$\mathbb{P}\left\{\inf_{\theta \in \Theta} \left| \sum_{l=0}^K c_l(\gamma(X_t; \theta)|\gamma(X_{t-1}; \theta); \theta) \frac{\delta^l}{l!} \right| = 0\right\} = 0.$$

(A.7) For any $\beta > 1$ and $\eta > 0$, there exists $\Delta(\beta, \eta) > 0$, then for any $\delta \in (0, \Delta(\beta, \eta)]$ and K , where K may be infinite,

$$\mathbb{P}\left\{\inf_{\theta \in \Theta} \left| \sum_{l=0}^K c_l(\gamma(X_t; \theta)|\gamma(X_{t-1}; \theta); \theta) \frac{\delta^l}{l!} \right| < \eta^{1/\beta}\right\} < \eta.$$

Assumptions (A.1) and (A.2) are standard requirements for maximum likelihood estimators. In particular, (A.1) (ii) contains conditions on the smoothness of the drift and the diffusion which ensures the existence of a unique solution to (2.1) as well as the infinite differentiability of the transition density $f(x|x_0, \delta; \theta)$ with respect to x , x_0 and δ , and three times differentiable with respect to θ [Friedman (1964)]. The second part of (A.2) is the simplified approach of Cramér (1946) assuming the MLEs are the solutions of the likelihood score equations. Assumption (A.3) is needed to guarantee the third derivative of $\log f(X_t|X_{t-1}, \delta; \theta)$ with respect to θ can be controlled by an integrable function, while condition (A.4) ensures the absolute convergence of the infinite series $\sum_{l=0}^{\infty} |c_l(\gamma(X_t; \theta)|\gamma(X_{t-1}; \theta))\delta^l/l!| = \exp\{\tilde{A}_3(x|x_0, \delta; \theta)\}$ as Ait-Sahalia (2002) has provided conditions on the nondegeneracy of the diffusion function and the boundary condition, which together with the third part of condition (A.1) leads to the convergence of the above infinite series $\exp\{\tilde{A}_3(x|x_0, \delta; \theta)\}$. Condition (A.4) is also needed to allow exchange of differentiation and summation for the infinite series. The first part of the (A.5) is of standard in likelihood inference. Its second part reflects the fact that for some processes $\lim_{\delta \rightarrow 0} I(\delta)$ may be singular, as conveyed in our discussion in Section 6 for the Vasicek process. Condition (A.6) is needed to guarantee the derivatives of log transition density and log approximate transition density exist with probability one. Condition (A.7) is needed to manage the denominators in the derivatives of the log approximate transition density, ensuring that the probability of their taking small values can be controlled uniformly.

We shall give the proofs for the propositions and theorems mentioned in Sections 3–6. We first present some lemmas about the true transition density and its approximations, which we will use in later proofs. The proofs for the lemmas can be found in Chang and Chen (2011).

LEMMA 1. Under (A.1) and (A.4), for any $\delta \in (0, \Delta)$, the infinite series

$$\sum_{l=0}^{\infty} c_l(\gamma(X_t; \theta) | \gamma(X_{t-1}; \theta)) \frac{\delta^l}{l!}$$

absolutely converges with probability 1, and for $k = 1, 2$ and 3 , and $i_1, i_2, i_3 \in \{1, \dots, d\}$,

$$\begin{aligned} & \frac{\partial^k}{\partial \theta_{i_1} \dots \partial \theta_{i_k}} \sum_{l=0}^{\infty} c_l(\gamma(X_t; \theta) | \gamma(X_{t-1}; \theta)) \frac{\delta^l}{l!} \\ &= \sum_{l=0}^{\infty} \frac{\partial^k}{\partial \theta_{i_1} \dots \partial \theta_{i_k}} c_l(\gamma(X_t; \theta) | \gamma(X_{t-1}; \theta)) \frac{\delta^l}{l!}. \end{aligned}$$

LEMMA 2. Under (A.6) and (A.7), for any positive $\beta > 1$, there exist two constants $m(\beta) < \infty$ and $\Delta_1(\beta) > 0$ such that for any $\delta \in (0, \Delta_1(\beta)]$ and J , where J can be infinity, then

$$\mathbb{E} \left\{ \sup_{\theta \in \Theta} \left| \sum_{l=0}^J c_l(\gamma(X_t; \theta) | \gamma(X_{t-1}; \theta)) \frac{\delta^l}{l!} \right|^{-\beta} \right\} < m(\beta).$$

LEMMA 3. Under (A.1), (A.3), (A.4), (A.6), (A.7), there exist two constants $M_1 < \infty$ and $\Delta_2 > 0$ such that, for any J , where J can be infinity, $\delta \in (0, \Delta_2)$ and $i, j, k \in \{1, \dots, d\}$,

$$\mathbb{E} \left\{ \sup_{\theta \in \Theta} \left| \frac{\partial^3 \log f^{(J)}(X_t | X_{t-1}, \delta; \theta)}{\partial \theta_i \partial \theta_j \partial \theta_k} \right| \right\} < M_1.$$

PROOF OF PROPOSITION 1. Using the same method in the proof of Lemma 3, we know (a) holds. On the other hand, Lemma 3 implies (b). \square

PROOF OF PROPOSITION 2. See the proof of Proposition 2 in [Chang and Chen \(2011\)](#). \square

PROOF OF PROPOSITION 3. Recall Proposition 2, then

$$\|I^{-1}(\delta)N(\theta_0, J, \delta) + E_d\|_2 \leq \|I^{-1}(\delta)\|_2 \cdot \|N(\theta_0, J, \delta) + I(\delta)\|_2 \leq C\delta^J.$$

If $C\delta^J < 1$, then

$$\|N^{-1}(\theta_0, J, \delta)I(\delta) + E_d\|_2 \leq \frac{\|I^{-1}(\delta)N(\theta_0, J, \delta) + E_d\|_2}{1 - \|I^{-1}(\delta)N(\theta_0, J, \delta) + E_d\|_2}.$$

From Proposition 2, if $C\delta^{J+1} < 1$, then

$$\|N^{-1}(\theta_0, J, \delta) + I^{-1}(\delta)\|_2 \leq \frac{\|I^{-1}(\delta)\|_2^2 \|N(\theta_0, J, \delta) + I(\delta)\|_2}{1 - \|I^{-1}(\delta)\|_2 \|N(\theta_0, J, \delta) + I(\delta)\|_2}.$$

On the other hand, using the same method in the proof of Proposition 2, we have

$$\|U(\theta_0, J, \delta)\|_2 \leq C\delta^{J+1}$$

for any positive J and $\delta \in (0, \bar{\Delta})$. Hence, we can find the constants C_1, C_2 and $\underline{\Delta} > 0$ such that

$$\|N^{-1}(\theta_0, J, \delta)I(\delta) + E_d\|_2 \leq C_1\delta^J \quad \text{and} \quad \|N^{-1}(\theta_0, J, \delta)U(\theta_0, J, \delta)\|_2 \leq C_2\delta^J$$

for any positive integer J and $\delta \in (0, \underline{\Delta})$. \square

PROOF OF PROPOSITION 4. Use the same method in the proof of Proposition 2. \square

PROOF OF PROPOSITION 5. We'll use Corollary 2.1 in Newey (1991) to prove this proposition. We only need to verify three conditions under two situations mentioned in Proposition 5:

(i) for any $i \in \{1, \dots, d\}$,

$$\mathbb{E}\left\{\frac{\partial}{\partial\theta_i} \log f(X_t|X_{t-1}, \delta; \theta)\right\} \quad \text{is equicontinuous;}$$

(ii) for any $i \in \{1, \dots, d\}$,

$$\sup_{\theta \in \Theta} \left\| \frac{1}{n} \sum_{t=1}^n \frac{\partial^2}{\partial\theta_i \partial\theta'} \log f(X_t|X_{t-1}, \delta; \theta) \right\|_2 = O_p(1);$$

(iii) for any $i \in \{1, \dots, d\}$ and $\theta \in \Theta$,

$$\frac{1}{n} \sum_{t=1}^n \frac{\partial}{\partial\theta_i} \log f(X_t|X_{t-1}, \delta; \theta) - \mathbb{E}\left\{\frac{\partial}{\partial\theta_i} \log f(X_t|X_{t-1}, \delta; \theta)\right\} \xrightarrow{p} 0.$$

For any $\theta^*, \theta^{**} \in \Theta$, note that

$$\begin{aligned} & \mathbb{E}\left\{\frac{\partial}{\partial\theta_i} \log f(X_t|X_{t-1}, \delta; \theta^*)\right\} - \mathbb{E}\left\{\frac{\partial}{\partial\theta_i} \log f(X_t|X_{t-1}, \delta; \theta^{**})\right\} \\ &= \mathbb{E}\left\{\frac{\partial^2}{\partial\theta_i \partial\theta'} \log f(X_t|X_{t-1}, \delta; \bar{\theta})\right\} \cdot (\theta^* - \theta^{**}), \end{aligned}$$

where $\bar{\theta}$ is on the joint line between θ^* and θ^{**} . Then

$$\begin{aligned} & \left| \mathbb{E}\left\{\frac{\partial}{\partial\theta_i} \log f(X_t|X_{t-1}, \delta; \theta^*)\right\} - \mathbb{E}\left\{\frac{\partial}{\partial\theta_i} \log f(X_t|X_{t-1}, \delta; \theta^{**})\right\} \right| \\ & \leq \left\| \mathbb{E}\left\{\frac{\partial^2}{\partial\theta_i \partial\theta'} \log f(X_t|X_{t-1}, \delta; \bar{\theta})\right\} \right\|_2 \cdot \|\theta^* - \theta^{**}\|_2. \end{aligned}$$

For any $j \in \{1, \dots, d\}$, use the same method in the proof of Lemma 3, we know that there exists a constant C , which is not dependent on J and δ , and $\hat{\Delta} > 0$ such that, for any J and $\delta \in (0, \hat{\Delta}]$,

$$\mathbb{E} \left\{ \sup_{\theta \in \Theta} \left| \frac{\partial^2}{\partial \theta_i \partial \theta_j} \log f(X_t | X_{t-1}, \delta; \theta) \right| \right\} < C.$$

Hence, (i) and (ii) can be established.

To verify (iii), from (A.3) [Lemmas 3 and 4 in Ait-Sahalia and Mykland (2004)], we know that there exists a positive constant κ such that for any $t_1 < t_2$,

$$\begin{aligned} & \left| \mathbb{E} \left\{ \left[\frac{\partial}{\partial \theta_i} \log f(X_{t_1} | X_{t_1-1}, \delta; \theta) - \mathbb{E} \left\{ \frac{\partial}{\partial \theta_i} \log f(X_{t_1} | X_{t_1-1}, \delta; \theta) \right\} \right] \right. \right. \\ & \quad \times \left. \left[\frac{\partial}{\partial \theta_i} \log f(X_{t_2} | X_{t_2-1}, \delta; \theta) - \mathbb{E} \left\{ \frac{\partial}{\partial \theta_i} \log f(X_{t_2} | X_{t_2-1}, \delta; \theta) \right\} \right] \right\} \Big| \\ & \leq C \cdot \exp\{-\kappa(t_2 - t_1)\delta\}, \end{aligned}$$

where

$$C = \mathbb{E} \left\{ \left[\frac{\partial}{\partial \theta_i} \log f(X_t | X_{t-1}, \delta; \theta) - \mathbb{E} \left\{ \frac{\partial}{\partial \theta_i} \log f(X_t | X_{t-1}, \delta; \theta) \right\} \right]^2 \right\}.$$

Then

$$\begin{aligned} & \mathbb{E} \left\{ \frac{1}{n} \sum_{t=1}^n \left[\frac{\partial}{\partial \theta_i} \log f(X_t | X_{t-1}, \delta; \theta) - \mathbb{E} \left\{ \frac{\partial}{\partial \theta_i} \log f(X_t | X_{t-1}, \delta; \theta) \right\} \right]^2 \right\} \\ & \leq \frac{C}{n} + \frac{C}{n} \cdot \frac{\exp\{-\kappa\delta\}}{1 - \exp\{-\kappa\delta\}} \\ & \leq 3 \left[2K_1 + K_2 \cdot m \left(\frac{2\nu_1}{\nu_1 - 2} \right) \right] \cdot \left\{ \frac{1}{n} + \frac{1}{n[\exp(\kappa\delta) - 1]} \right\} \rightarrow 0, \end{aligned}$$

under the two situations mentioned in the statement of Proposition 5. Hence we complete the proof. \square

PROOF OF PROPOSITION 6. From (A.2), we can get $n^{-1} \nabla_{\theta} \ell_{n,\delta}(\hat{\theta}_n) = 0$. Expanding it at θ_0 ,

$$0 = \frac{1}{n} \sum_{t=1}^n \nabla_{\theta} \log f(X_t | X_{t-1}, \delta; \theta_0) + \frac{1}{n} \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f(X_t | X_{t-1}, \delta; \tilde{\theta}) \cdot (\hat{\theta}_n - \theta_0).$$

Then

$$\hat{\theta}_n - \theta_0 = \left\{ -\frac{1}{n} \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f(X_t | X_{t-1}, \delta; \tilde{\theta}) \right\}^{-1} \cdot \frac{1}{n} \sum_{t=1}^n \nabla_{\theta} \log f(X_t | X_{t-1}, \delta; \theta_0).$$

Define $I_n(\delta) = -n^{-1} \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f(X_t|X_{t-1}, \delta; \theta_0)$. From Lemma 3, an $-n^{-1} \times \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f(X_t|X_{t-1}, \delta; \tilde{\theta}) = I_n(\delta) \cdot \{1 + o_p(1)\}$. Using the same way as that in the verification of (iii) in the proof of Proposition 5, we can get $I_n(\delta) - I(\delta) = O_p\{(n\delta)^{-1/2}\}$. If $n\delta^3 \rightarrow \infty$, by (A.5),

$$\begin{aligned} \left\{ -\frac{1}{n} \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f(X_t|X_{t-1}, \delta; \tilde{\theta}) \right\}^{-1} &= \{I(\delta) \cdot \{1 + o_p(1)\} + O_p\{(n\delta)^{-1/2}\}\}^{-1} \\ &= I^{-1}(\delta) \cdot \{1 + o_p(1)\}. \end{aligned}$$

Then

$$\sqrt{n}I^{1/2}(\delta)(\hat{\theta}_n - \theta_0) = I^{-1/2}(\delta) \frac{1}{n^{1/2}} \sum_{t=1}^n \nabla_{\theta} \log f(X_t|X_{t-1}, \delta; \theta_0) \cdot \{1 + o_p(1)\}.$$

We will use the martingale central limit theorem [Billingsley (1995), page 476] to show that the first part on the right-hand side of the above equation converges to a standard normal distribution. For any $\alpha \in \mathbb{R}^d$ with unit L_2 norm, to simplify notations, let $U_{n,m} = \alpha' I^{-1/2}(\delta) n^{-1/2} \nabla_{\theta} \log f(X_m|X_{m-1}, \delta; \theta_0)$ and $\mathcal{F}_{n,m} = \sigma(X_1, \dots, X_m)$. It is easy to check $(U_{n,m}, \mathcal{F}_{n,m})$ is a martingale difference array. By the Markov property and Birkhoff's Ergodic theorem, $V_{n,n} = \sum_{m=1}^n \mathbb{E}(U_{n,m}^2 | \mathcal{F}_{n,m}) \xrightarrow{p} \mathbb{E}U_{n,n}^2 = 1$. On the other hand, $\sum_{m=1}^n |U_{n,m}|^3 \leq C(n \times \delta^3)^{-1/2} \rightarrow 0$. This implies the asymptotic normality of $\sqrt{n}\alpha' I^{1/2}(\delta)(\hat{\theta}_n - \theta_0)$. Then we complete the proof. \square

PROOF OF THEOREM 1. From Propositions 4 and 5, we can get

$$\|\mathbb{E}\nabla_{\theta} \log f(X_t|X_{t-1}, \delta; \hat{\theta}_n^{(J)})\|_2 \xrightarrow{p} 0$$

for either: (i) $\delta \in (0, \tilde{\Delta} \wedge \dot{\Delta})$ being fixed, $J \rightarrow \infty$ and $n \rightarrow \infty$, or (ii) J being fixed, $n \rightarrow \infty$, $\delta \rightarrow 0$ but $n\delta \rightarrow \infty$. Hence, noting condition (A.2)(i), we have the consistency of the AMLE $\hat{\theta}_n^{(J)}$. \square

PROOF OF THEOREM 2. For fixed δ , from Theorem 1 and (4.1), we know that the leading order term of $\hat{\theta}_n^{(J)} - \theta_0$ contains two parts: one is $N^{-1}U_n$, and the other is $N^{-1}(N_n + F_n)(\hat{\theta}_n - \theta_0)$. Hence, $\hat{\theta}_n^{(J)} - \theta_0 = O_p\{\delta^{J+1} + (n\delta)^{-1/2}\}$.

For J fixed and $\delta \rightarrow 0$, Proposition 4 implies

$$\begin{aligned} \mathbb{E} \left\{ \left\| \frac{1}{n} \sum_{t=1}^n \nabla_{\theta} \log f(X_t|X_{t-1}, \delta; \hat{\theta}_n^{(J)}) - \frac{1}{n} \sum_{t=1}^n \nabla_{\theta} \log f(X_t|X_{t-1}, \delta; \hat{\theta}_n) \right\|_2 \right\} \\ \leq C\delta^{J+1}. \end{aligned}$$

This means that

$$\mathbb{E} \left\{ \left\| \frac{1}{n} \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f(X_t|X_{t-1}, \delta; \tilde{\theta}) \cdot (\hat{\theta}_n^{(J)} - \hat{\theta}_n) \right\|_2 \right\} \leq C\delta^{J+1},$$

where $\tilde{\theta}$ is on the joining line between $\hat{\theta}_n^{(J)}$ and $\hat{\theta}_n$. Hence

$$\frac{1}{n} \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f(X_t|X_{t-1}, \delta; \tilde{\theta}) \cdot (\hat{\theta}_n^{(J)} - \hat{\theta}_n) = O_p(\delta^{J+1}).$$

Since $\tilde{\theta} \xrightarrow{p} \theta_0$ and $\hat{\theta}_n^{(J)} - \hat{\theta}_n = o_p(1)$,

$$\frac{1}{n} \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f(X_t|X_{t-1}, \delta; \theta_0) \cdot (\hat{\theta}_n^{(J)} - \hat{\theta}_n) = O_p(\delta^{J+1}).$$

On the other hand, from Proposition 2, we know

$$\begin{aligned} & \frac{1}{n} \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f(X_t|X_{t-1}, \delta; \theta_0) - \frac{1}{n} \sum_{t=1}^n \nabla_{\theta\theta}^2 \log f^{(J)}(X_t|X_{t-1}, \delta; \theta_0) \\ &= O_p(\delta^{J+1}). \end{aligned}$$

Then $N_n(\hat{\theta}_n^{(J)} - \hat{\theta}_n) = O_p(\delta^{J+1})$. Using the same way of verifying (iii) in the proof of Proposition 5, we know $N_n - N = O_p\{(n\delta)^{-1/2}\}$. As $n\delta^3 \rightarrow \infty$, then $N(\hat{\theta}_n^{(J)} - \hat{\theta}_n) = O_p(\delta^{J+1})$. Hence, $\hat{\theta}_n^{(J)} - \hat{\theta}_n = O_p(\delta^J)$. At the same time, we know $\hat{\theta}_n - \theta_0 = O_p\{(n\delta)^{-1/2}\}$. Then

$$\hat{\theta}_n^{(J)} - \theta_0 = O_p\{\delta^J + (n\delta)^{-1/2}\}.$$

This completes the proof of Theorem 2. \square

PROOF OF THEOREM 4. We only need to prove following result:

$$\sqrt{n}\tilde{I}_n^{1/2}(\hat{\theta}_n^{(J)}, J, \delta)(\hat{\theta}_n^{(J)} - \theta_0) = \sqrt{n}I^{1/2}(\delta)(\hat{\theta}_n^{(J)} - \theta_0) + o_p(1)$$

under the two situations mentioned in Theorem 4. Using the approach in the proof of Lemma 3, we have $\tilde{I}_n(\hat{\theta}_n^{(J)}, J, \delta) - \tilde{I}_n(\theta_0, J, \delta) = O_p\{\|\hat{\theta}_n^{(J)} - \theta_0\|_2\}$. Also, using the same way of verifying (iii) in the proof of Proposition 5, $\tilde{I}_n(\theta_0, J, \delta) - \mathbb{E}\tilde{I}_n(\theta_0, J, \delta) = O_p\{(n\delta)^{-1/2}\}$. By the same argument in the proof of Proposition 2, $\mathbb{E}\tilde{I}_n(\theta_0, J, \delta) - I(\delta) = O(\delta^{J+1})$. Hence, if $n\delta^3 \rightarrow \infty$, under either asymptotic regime in Theorem 4,

$$\tilde{I}_n^{1/2}(\hat{\theta}_n^{(J)}, J, \delta) = I^{1/2}(\delta) \cdot \{1 + o_p(1)\}.$$

Then we complete the proof. \square

Acknowledgments. We thank the Associate Editor for very constructive comments and suggestions which have improved the presentation of the paper.

The first author thanks Department of Statistics at Iowa State University for hospitality during his visits.

REFERENCES

- AÏT-SAHALIA, Y. (1999). Transition densities for interest rate and other nonlinear diffusions. *J. Finance* **54** 1361–1395.
- AÏT-SAHALIA, Y. (2002). Maximum likelihood estimation of discretely sampled diffusions: A closed-form approximation approach. *Econometrica* **70** 223–262. [MR1926260](#)
- AÏT-SAHALIA, Y. (2008). Closed-form likelihood expansions for multivariate diffusions. *Ann. Statist.* **36** 906–937. [MR2396819](#)
- AÏT-SAHALIA, Y. and KIMMEL, R. (2007). Maximum likelihood estimation of stochastic volatility models. *Journal of Financial Economics* **83** 413–452.
- AÏT-SAHALIA, Y. and KIMMEL, R. (2010). Estimating affine multifactor term structure models using closed-form likelihood expansions. *Journal of Financial Economics* **98** 113–144.
- AÏT-SAHALIA, Y. and MYKLAND, P. A. (2004). Estimators of diffusions with randomly spaced discrete observations: A general theory. *Ann. Statist.* **32** 2186–2222. [MR2102508](#)
- AÏT-SAHALIA, Y., MYKLAND, P. A. and ZHANG, L. (2011). Ultra high frequency volatility estimation with dependent microstructure noise. *J. Econometrics* **160** 160–175. [MR2745875](#)
- BAKSHI, G., CAO, C. and CHEN, Z. (1997). Empirical performance of alternative option pricing models. *J. Finance* **52** 2003–2049.
- BESKOS, A., PAPASPILIOPOULOS, O., ROBERTS, G. O. and FEARNHEAD, P. (2006). Exact and computationally efficient likelihood-based estimation for discretely observed diffusion processes (with discussion). *J. R. Stat. Soc. Ser. B Stat. Methodol.* **68** 333–382. [MR2278331](#)
- BIBBY, B. M. and SØRENSEN, M. (1995). Martingale estimation functions for discretely observed diffusion processes. *Bernoulli* **1** 17–39. [MR1354454](#)
- BILLINGSLEY, P. (1995). *Probability and Measure*, 3rd ed. Wiley, New York. [MR1324786](#)
- BLACK, F. and SCHOLES, M. (1973). The pricing of options and corporate liabilities. *Journal of Political Economy* **81** 637–654.
- CHANG, J. and CHEN, S. X. (2011). On the approximate maximum likelihood estimation for diffusion processes. Technical Report 2011-08, Center for Statistical Science, Peking Univ.
- COX, J. C., INGERSOLL, J. E. and ROSS, S. A. (1985). A theory of term structure of interest rates. *Econometrica* **53** 385–407.
- CRAMÉR, H. (1946). *Mathematical Methods of Statistics*. Princeton Mathematical Series **9**. Princeton Univ. Press, Princeton, NJ. [MR0016588](#)
- DUMAS, B., FLEMING, J. and WHALEY, R. E. (1998). Implied volatility functions: Empirical tests. *J. Finance* **53** 2059–2106.
- FAN, J. (2005). A selective overview of nonparametric methods in financial econometrics. *Statist. Sci.* **20** 317–357. [MR2210224](#)
- FAN, J. and WANG, Y. (2007). Multi-scale jump and volatility analysis for high-frequency financial data. *J. Amer. Statist. Assoc.* **102** 1349–1362. [MR2372538](#)
- FAN, J. and ZHANG, C. (2003). A reexamination of diffusion estimators with applications to financial model validation. *J. Amer. Statist. Assoc.* **98** 118–134. [MR1965679](#)
- FELLER, W. (1952). The parabolic differential equations and the associated semi-groups of transformations. *Ann. of Math. (2)* **55** 468–519. [MR0047886](#)
- FRIEDMAN, A. (1964). *Partial Differential Equations of Parabolic Type*. Prentice Hall, Englewood Cliffs, NJ. [MR0181836](#)

- GOBET, E. (2002). LAN property for ergodic diffusions with discrete observations. *Ann. Inst. Henri Poincaré Probab. Stat.* **38** 711–737. [MR1931584](#)
- JACOBSEN, M. (2001). Discretely observed diffusions: Classes of estimating functions and small Δ -optimality. *Scand. J. Statist.* **28** 123–149. [MR1844353](#)
- KARATZAS, I. and SHREVE, S. E. (1991). *Brownian Motion and Stochastic Calculus*. Springer, New York.
- MERTON, R. C. (1973). Theory of rational option pricing. *Bell J. Econom. and Management Sci.* **4** 141–183. [MR0496534](#)
- MYKLAND, P. A. and ZHANG, L. (2009). Inference for continuous semimartingales observed at high frequency. *Econometrica* **77** 1403–1445. [MR2561071](#)
- NEWBY, W. K. (1991). Uniform convergence in probability and stochastic equicontinuity. *Econometrica* **59** 1161–1167. [MR1113551](#)
- ØKSENDAL, B. (2000). *Stochastic Differential Equations: An Introduction with Applications*, 5th ed. Springer, Berlin.
- PRAKASA RAO, B. L. S. (1999). *Semimartingales and Statistical Inference*. Chapman and Hall/CRC, London.
- PROTTER, P. E. (2004). *Stochastic Integration and Differential Equations: Stochastic Modelling and Applied Probability*, 2nd ed. *Applications of Mathematics (New York)* **21**. Springer, Berlin. [MR2020294](#)
- SØRENSEN, M. (2007). Efficient estimation for ergodic diffusions sampled at high frequency. Dept. Mathematical Sciences, Univ. Copenhagen.
- STRAMER, O. and YAN, J. (2007a). On simulated likelihood of discretely observed diffusion processes and comparison to closed-form approximation. *J. Comput. Graph. Statist.* **16** 672–691. [MR2351085](#)
- STRAMER, O. and YAN, J. (2007b). Asymptotics of an efficient Monte Carlo estimation for the transition density of diffusion processes. *Methodol. Comput. Appl. Probab.* **9** 483–496. [MR2404739](#)
- SUNDARESAN, S. M. (2000). Continuous time finance: A review and assessment. *J. Finance* **55** 1569–1622.
- TANG, C. Y. and CHEN, S. X. (2009). Parameter estimation and bias correction for diffusion processes. *J. Econometrics* **149** 65–81. [MR2515045](#)
- VASICEK, O. (1977). An equilibrium characterization of the term structure. *Journal of Financial Economics* **5** 177–188.
- WANG, Y. (2002). Asymptotic nonequivalence of Garch models and diffusions. *Ann. Statist.* **30** 754–783. [MR1922541](#)

GUANGHUA SCHOOL OF MANAGEMENT
PEKING UNIVERSITY
BEIJING, 100871
CHINA
E-MAIL: changjinyuan1986@pku.edu.cn

GUANGHUA SCHOOL OF MANAGEMENT
AND CENTER FOR STATISTICAL SCIENCE
PEKING UNIVERSITY
BEIJING, 100871
CHINA
AND
DEPARTMENT OF STATISTICS
IOWA STATE UNIVERSITY
AMES, IOWA 50011-1210
USA
E-MAIL: songchen@iastate.edu