

A MAXIMAL INEQUALITY FOR STOCHASTIC CONVOLUTIONS IN 2-SMOOTH BANACH SPACES

JAN VAN NEERVEN¹

Delft Institute of Applied Mathematics, Delft University of Technology, P.O. Box 5031, 2600 GA Delft, The Netherlands

email: J.M.A.M.vanNeerven@TUDelft.nl

JIAHUI ZHU²

Delft Institute of Applied Mathematics, Delft University of Technology, P.O. Box 5031, 2600 GA Delft, The Netherlands

email: Jiahui.Zhu@TUDelft.nl

Submitted May 24, 2011, accepted in final form October 24, 2011

AMS 2000 Subject classification: Primary 60H05; Secondary 60H15

Keywords: Stochastic convolutions, maximal inequality, 2-smooth Banach spaces, Itô formula.

Abstract

Let $(e^{tA})_{t \geq 0}$ be a C_0 -contraction semigroup on a 2-smooth Banach space E , let $(W_t)_{t \geq 0}$ be a cylindrical Brownian motion in a Hilbert space H , and let $(g_t)_{t \geq 0}$ be a progressively measurable process with values in the space $\gamma(H, E)$ of all γ -radonifying operators from H to E . We prove that for all $0 < p < \infty$ there exists a constant C , depending only on p and E , such that for all $T \geq 0$ we have

$$\mathbb{E} \sup_{0 \leq t \leq T} \left\| \int_0^t e^{(t-s)A} g_s dW_s \right\|^p \leq C \mathbb{E} \left(\int_0^T \|g_t\|_{\gamma(H, E)}^2 dt \right)^{\frac{p}{2}}.$$

For $p \geq 2$ the proof is based on the observation that $\psi(x) = \|x\|^p$ is Fréchet differentiable and its derivative satisfies the Lipschitz estimate $\|\psi'(x) - \psi'(y)\| \leq C(\|x\| + \|y\|)^{p-2} \|x - y\|$; the extension to $0 < p < 2$ proceeds via Lenglart's inequality.

1 Introduction

Let $(e^{tA})_{t \geq 0}$ be a C_0 -contraction semigroup on a 2-smooth Banach space E and let $(W_t)_{t \geq 0}$ be a cylindrical Brownian motion in a Hilbert space H . Let $(g_t)_{t \geq 0}$ be a progressively measurable process with values in the space $\gamma(H, E)$ of all γ -radonifying operators from H to E satisfying

$$\int_0^T \|g_t\|_{\gamma(H, E)}^2 dt < \infty \quad \mathbb{P}\text{-almost surely}$$

¹RESEARCH SUPPORTED BY VICI SUBSIDY 639.033.604 OF THE NETHERLANDS ORGANISATION FOR SCIENTIFIC RESEARCH (NWO)

²RESEARCH SUPPORTED BY VICI SUBSIDY 639.033.604 OF THE NETHERLANDS ORGANISATION FOR SCIENTIFIC RESEARCH (NWO)

for all $T \geq 0$. As is well known (see [6, 15, 16]), under these assumptions the stochastic convolution process

$$X_t = \int_0^t e^{(t-s)A} g_s dW_s, \quad t \geq 0,$$

is well-defined in E and provides the unique mild solution of the stochastic initial value problem

$$dX_t = AX_t dt + g_t dW_t, \quad X_0 = 0.$$

In order to obtain the existence of a continuous version of this process, one usually proves a maximal estimate of the form

$$\mathbb{E} \sup_{0 \leq t \leq T} \|X_t\|^p \leq C^p \mathbb{E} \left(\int_0^T \|g_t\|_{\gamma(H,E)}^2 dt \right)^{\frac{p}{2}}. \quad (1.1)$$

The first such estimate was obtained by Kotelenez [11, 12] for C_0 -contraction semigroups on Hilbert spaces E and exponent $p = 2$. Tubaro [19] extended this result to exponents $p \geq 2$ by a different method of proof which applies Itô's formula to the C^2 -mapping $x \mapsto \|x\|^p$. The case $p \in (0, 2)$ was covered subsequently by Ichikawa [10]. A very simple proof, still for C_0 -contraction semigroups on Hilbert spaces, which works for all $p \in (0, \infty)$, was obtained recently by Hausenblas and Seidler [9]. It is based on the Sz.-Nagy dilation theorem, which is used to reduce the problem to the corresponding problem for C_0 -contraction groups. Then, by using the group property, the maximal estimate follows from Burkholder's inequality. This proof shows, moreover, that the constant C in (1.1) may be taken equal to the constant appearing in Burkholder's inequality. In particular, this constant depends only on p .

The maximal inequality (1.1) has been extended by Brzeźniak and Peszat [4] to C_0 -contraction semigroups on Banach spaces E with the property that, for some $p \in [2, \infty)$, $x \mapsto \|x\|^p$ is twice continuously Fréchet differentiable and the first and second Fréchet derivatives are bounded by constant multiples of $\|x\|^{p-1}$ and $\|x\|^{p-2}$, respectively. Examples of spaces with this property, which we shall call (C_p^2) , are the spaces $L^q(\mu)$ for $q \in [p, \infty)$. Any (C_p^2) space is 2-smooth (the definition is recalled in Section 2), but the converse doesn't hold:

Example 1.1. Let F be a Banach space. The space $\ell^2(F)$ is 2-smooth whenever F is 2-smooth [8, Proposition 17]. On the other hand, the norm of $\ell^2(F)$ is twice continuously Fréchet differentiable away from the origin if and only if F is a Hilbert space [14, Theorem 3.9]. Thus, for $q \in (2, \infty)$, $\ell^2(\ell^q)$ and $\ell^2(L^q(0, 1))$ are examples of 2-smooth Banach spaces which fail property (C_p^2) for all $p \in [2, \infty)$.

To the best of our knowledge, the general problem of proving the maximal estimate (1.1) for C_0 -contraction semigroups on 2-smooth Banach space remains open. The present paper aims to fill this gap:

Theorem 1.2. *Let $(e^{tA})_{t \geq 0}$ be a C_0 -contraction semigroup on a 2-smooth Banach space E , let $(W_t)_{t \geq 0}$ be a cylindrical Brownian motion in a Hilbert space H , and let $(g_t)_{t \geq 0}$ be a progressively measurable process in $\gamma(H, E)$. If*

$$\int_0^T \|g_t\|_{\gamma(H,E)}^2 dt < \infty \quad \mathbb{P}\text{-almost surely,}$$

then the stochastic convolution process $X_t = \int_0^t e^{(t-s)A} g_s dW_s$ is well-defined and has a continuous version. Moreover, for all $0 < p < \infty$ there exists a constant C , depending only on p and E , such that

$$\mathbb{E} \sup_{0 \leq t \leq T} \|X_t\|^p \leq C^p \mathbb{E} \left(\int_0^T \|g_t\|_{\gamma(H,E)}^2 dt \right)^{\frac{p}{2}}.$$

For $p \geq 2$, the proof of Theorem 1.2 is based on a version of Itô’s formula (Theorem 3.1) which exploits the fact (proved in Lemma 2.1) that in 2-smooth Banach spaces the function $\psi(x) = \|x\|^p$ is Fréchet differentiable and satisfies the Lipschitz estimate

$$\|\psi'(x) - \psi'(y)\| \leq C(\|x\| + \|y\|)^{p-2}\|x - y\|.$$

The extension to exponents $0 < p < 2$ is obtained by applying Lenglar’s inequality (see (4.1)). We conclude this introduction with a brief discussion of some developments of the inequality (1.1) into different directions in the literature. Seidler [18] has proved the inequality (1.1) with optimal constant $C = O(\sqrt{p})$ as $p \rightarrow \infty$ for positive C_0 -contraction semigroups on the (2-smooth) space $E = L^q(\mu)$, $q \geq 2$. He also proved that the same result holds if the assumption ‘ e^{tA} is a positive contraction semigroup’ is replaced by ‘ $-A$ has a bounded H^∞ -calculus of angle strictly less than $\frac{1}{2}\pi$ ’. The latter result was subsequently extended by Veraar and Weis [20] to arbitrary UMD spaces E with type 2. In the same paper, still under the assumption that $-A$ has a bounded H^∞ -calculus of angle strictly less than $\frac{1}{2}\pi$, the following stronger estimate is obtained for UMD spaces E with Pisier’s property (α):

$$\mathbb{E} \sup_{0 \leq t \leq T} \|X_t\|^p \leq C^p \mathbb{E} \|g\|_{\gamma(L^2(0,T;H),E)}^p \tag{1.2}$$

with a constant C depending only on p and E . If, in addition, E has type 2, then the mapping $f \otimes (h \otimes x) \mapsto (f \otimes h) \otimes x$ extends to a continuous embedding $L^2(0, T; \gamma(H, E)) \hookrightarrow \gamma(L^2(0, T; H), E)$ and (1.2) implies (1.1).

Let us finally mention that, for $p > 2$, a weaker version of (1.1) for arbitrary C_0 -semigroups on Hilbert spaces has been obtained by Da Prato and Zabczyk [5]. Using the factorisation method they proved that

$$\mathbb{E} \sup_{0 \leq t \leq T} \|X_t\|^p \leq C^p \mathbb{E} \int_0^T \|g_t\|_{\gamma(H,E)}^p dt$$

with a constant C depending on p , E , and T . The proof extends *verbatim* to C_0 -semigroups on martingale type 2 spaces. This relates to the above results for 2-smooth spaces through a theorem of Pisier [17, Theorem 3.1], which states that a Banach space has martingale type p if and only if it is p -smooth.

2 The Fréchet derivative of $\|\cdot\|^p$

Let $1 < q \leq 2$. A Banach space E is q -smooth if the modulus of smoothness

$$\rho_{\|\cdot\|}(t) = \sup \left\{ \frac{1}{2}(\|x + ty\| + \|x - ty\|) - 1 : \|x\| = \|y\| = 1 \right\}$$

satisfies $\rho_{\|\cdot\|}(t) \leq Ct^q$ for all $t > 0$.

It is known (see [17, Theorem 3.1]) that E is q -smooth if and only if there exists a constant $K \geq 1$ such that for all $x, y \in E$,

$$\|x + y\|^q + \|x - y\|^q \leq 2\|x\|^q + K\|y\|^q. \tag{2.1}$$

Lemma 2.1. *Let E be a Banach space and let $1 < q \leq 2$ be given. For $p \geq q$ set $\psi_p(x) := \|x\|^p$.*

1. *E is q -smooth if and only if the Fréchet derivative of ψ_q is globally $(q - 1)$ -Hölder continuous on E .*

2. If E is q -smooth, then for $p > q$ the Fréchet derivative of ψ_p is locally $(q-1)$ -Hölder continuous on E .

Moreover, for all $p \geq q$ and $x, y \in E$ we have

$$\|\psi'_p(x) - \psi'_p(y)\| \leq C(\|x\| + \|y\|)^{p-q} \|x - y\|^{q-1}, \quad (2.2)$$

where C depends only on p, q and E .

Proof. If the Fréchet derivative of ψ_q is $(q-1)$ -Hölder continuous on E , then by the mean value theorem we can find $0 \leq \theta, \rho \leq 1$ such that for all $x, y \in E$,

$$\begin{aligned} \|x + y\|^q + \|x - y\|^q - 2\|x\|^q &= (\|x + y\|^q - \|x\|^q) + (\|x - y\|^q - \|x\|^q) \\ &\leq \|\psi'_q(x + \theta y) - \psi'_q(x - \rho y)\| \|y\| \\ &\leq L\|(x + \theta y) - (x - \rho y)\|^{q-1} \|y\| \leq 2^{q-1} L \|y\|^q. \end{aligned}$$

Hence the Banach space E is q -smooth.

Suppose now that the norm of E is q -smooth. Then for all $x, y \in E$ with $\|x\|, \|y\| = 1$ and all $t > 0$ we have

$$\|x + ty\| + \|x - ty\| - 2\|x\| \leq K \|ty\|^q. \quad (2.3)$$

Thus

$$\lim_{t \rightarrow 0} \frac{\|x + ty\| + \|x - ty\| - 2\|x\|}{\|ty\|} = 0,$$

which by [7, Lemma I.1.3] means that $\|\cdot\|$ is Fréchet differentiable on the unit sphere. Hence, by homogeneity, $\|\cdot\|$ is Fréchet differentiable on $E \setminus \{0\}$. Let us denote by f_x its Fréchet derivative at the point $x \neq 0$.

We begin by showing the $(q-1)$ -Hölder continuity of $x \mapsto f_x$ on the unit sphere of E , following the argument of [7, Lemma V.3.5]. We fix $x \neq y \in E$ such that $\|x\|, \|y\| = 1$ and $h \in E$ with $\|h\| = \|x - y\|$ and $x - y + h \neq 0$. Since the norm $\|\cdot\|$ is a convex function,

$$f_y(x - y) \leq \|x\| - \|y\|.$$

Similarly, we have

$$f_x(h) \leq \|x + h\| - \|x\|, \quad f_y(y - x - h) \leq \|2y - x - h\| - \|y\|.$$

By using above inequalities and the linearity of the function f_x , we have

$$\begin{aligned} f_x(h) - f_y(h) &\leq \|x + h\| - \|x\| - f_y(h) \\ &= \|x + h\| - \|y\| - f_y(x + h - y) + \|y\| - \|x\| + f_y(x - y) \\ &\leq \|x + h\| - \|y\| - f_y(x + h - y) \\ &= \|x + h\| - \|y\| + f_y(y - x - h) \\ &\leq \|x + h\| + \|2y - x - h\| - 2\|y\| \\ &= \left\| y + \|x + h - y\| \cdot \frac{x + h - y}{\|x + h - y\|} \right\| + \left\| y - \|x + h - y\| \cdot \frac{x + h - y}{\|x + h - y\|} \right\| - 2\|y\| \\ &\leq K \|x + h - y\|^q \leq K(\|x - y\| + \|h\|)^q = 2^q K \|x - y\|^q, \end{aligned}$$

where we also used (2.3). Since the roles of x and y may be reversed in this inequality, this implies

$$\|f_x - f_y\| = \sup_{\|h\|=\|x-y\|} \frac{|f_x(h) - f_y(h)|}{\|x - y\|} \leq 2^q K \|x - y\|^{q-1}$$

This proves the $(q-1)$ -Hölder continuity of the norm $\|\cdot\|$ on the unit sphere.

We proceed with the proof of (2.2); the $(q-1)$ -Hölder continuity of ψ_q as well as the local $(p-1)$ -Hölder continuity of ψ_p follow from it. For all $x, y \in E$ with $x \neq 0$ and $y \neq 0$ we have $\psi'_p(x) = p\|x\|^{p-1}f_x$.

It is easy to check that $f_x = f_{\frac{x}{\|x\|}}$ and $\|f_x\| = 1$. Following once more the argument of [7, Lemma V.3.5], this gives

$$\begin{aligned} \|\psi'_p(x) - \psi'_p(y)\| &= p \left\| \|x\|^{p-1}f_x - \|y\|^{p-1}f_y \right\| \\ &\leq p \left\| \|x\|^{p-1} \left(f_{\frac{x}{\|x\|}} - f_{\frac{y}{\|y\|}} \right) \right\| + p \left\| (\|x\|^{p-1} - \|y\|^{p-1}) f_{\frac{y}{\|y\|}} \right\| \\ &\leq p 2^q K \|x\|^{p-1} \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\|^{q-1} + p \left| \|x\|^{p-1} - \|y\|^{p-1} \right| \\ &\leq p 2^q K \|x\|^{p-q} \|y\|^{1-q} \left\| x\|y\| - y\|x\| \right\|^{q-1} + p \left| \|x\|^{p-1} - \|y\|^{p-1} \right| \\ &= p 2^q K \|x\|^{p-q} \|y\|^{1-q} \left\| \|y\|(x-y) + y(\|y\| - \|x\|) \right\|^{q-1} + p \left| \|x\|^{p-1} - \|y\|^{p-1} \right| \\ &\leq p 2^q K \|x\|^{p-q} \|y\|^{1-q} (2\|y\| \|x-y\|)^{q-1} + p \left| \|x\|^{p-1} - \|y\|^{p-1} \right| \\ &= p 2^{2q-1} K \|x\|^{p-q} \|x-y\|^{q-1} + p \left| \|x\|^{p-1} - \|y\|^{p-1} \right|. \end{aligned} \tag{2.4}$$

If $q \leq p \leq 2$, then by the inequality $|t^r - s^r| \leq |t - s|^r$, valid for $0 < r \leq 1$ and $s, t \in [0, \infty)$, we have

$$\left| \|x\|^{p-1} - \|y\|^{p-1} \right| \leq \left| \|x\| - \|y\| \right|^{p-1} \leq \|x - y\|^{p-1} \leq (\|x\| + \|y\|)^{p-q} \|x - y\|^{q-1}.$$

If $p > 2$, by applying the mean value theorem, for some $\theta \in [0, 1]$ we have

$$\begin{aligned} \left| \|x\|^{p-1} - \|y\|^{p-1} \right| &= (p-1) \left\| \theta x + (1-\theta)y \right\|^{p-2} f_{\theta x + (1-\theta)y}(x-y) \\ &\leq (p-1) (\|x\| + \|y\|)^{p-2} \|x-y\| \\ &\leq (p-1) (\|x\| + \|y\|)^{p-2} (\|x\| + \|y\|)^{2-q} \|x-y\|^{q-1} \\ &= (p-1) (\|x\| + \|y\|)^{p-q} \|x-y\|^{q-1}. \end{aligned}$$

Also, since $\psi'_p(0) = 0$, for $y \neq 0$ we have

$$\|\psi'_p(0) - \psi'_p(y)\| = p\|y\|^{p-1} = p\|y\|^{p-1} \left\| \frac{y}{\|y\|} \right\|^{p-1} \leq p\|y\|^{p-1} \left\| \frac{y}{\|y\|} \right\|^{q-1} = p\|y\|^{p-q} \|y\|^{q-1}.$$

□

The above lemma will be combined with the next one, which gives a first order Taylor formula with a remainder term involving the first derivative only.

Lemma 2.2. *Let E and F be Banach spaces, let $0 < \alpha \leq 1$, and let $\psi : E \rightarrow F$ be a Fréchet differentiable function whose Fréchet derivative $\psi' : E \rightarrow \mathcal{L}(E, F)$ is locally α -Hölder continuous. Then for all $x, y \in E$ we have*

$$\psi(y) = \psi(x) + \psi'(x)(y - x) + R(x, y),$$

where

$$R(x, y) = \int_0^1 (\psi'(x + r(y - x))(y - x) - \psi'(x)(y - x)) dr. \quad (2.5)$$

Proof. Pick $w \in E$ such that $\|w\| \leq 1$ and consider the function $f : \mathbb{R} \rightarrow F$ by

$$f(\theta) := \psi(x + \theta w).$$

For all $x^* \in F^*$, $\langle f', x^* \rangle$ is locally α -Hölder continuous. To see this, note that for $|\theta_1|, |\theta_2| \leq R$ and $\|x\| \leq R$ we have $\|x + \theta_1 w\|, \|x + \theta_2 w\| \leq 2R$, so by assumption there exists a constant C_{2R} such that

$$\begin{aligned} |\langle f'(\theta_1) - f'(\theta_2), x^* \rangle| &= |\langle \psi'(x + \theta_1 w)w, x^* \rangle - \langle \psi'(x + \theta_2 w)w, x^* \rangle| \\ &\leq \|\psi'(x + \theta_1 w) - \psi'(x + \theta_2 w)\| \|x^*\| \leq C_{2R} |\theta_1 - \theta_2|^\alpha \|x^*\|. \end{aligned}$$

Applying Taylor's formula and [1, Lemma 1, Theorem 3] to the function $\langle f, x^* \rangle$ we obtain

$$\langle f(t) - f(0), x^* \rangle = t \langle f'(0), x^* \rangle + \langle R_f(0, t), x^* \rangle,$$

where $R_f(0, t) = \int_0^1 t(f'(rt) - f'(0)) dr$. Now let $x, y \in E$ be given and set $t = \|y - x\|$ and $w = \frac{y-x}{\|y-x\|}$. With these choices we obtain

$$\begin{aligned} \langle \psi(y), x^* \rangle - \langle \psi(x), x^* \rangle - \langle \psi'(x)(y - x), x^* \rangle &= \langle \psi(x + tw), x^* \rangle - \langle \psi(x), x^* \rangle - t \langle \psi'(x)w, x^* \rangle \\ &= \langle f(t) - f(0) - tf'(0), x^* \rangle \\ &= \int_0^1 t \langle f'(rt) - f'(0), x^* \rangle dr \\ &= \int_0^1 \langle \psi'(x + r(y - x))(y - x) - \psi'(x)(y - x), x^* \rangle dr. \end{aligned}$$

Since $x^* \in F^*$ was arbitrary, this proves the lemma. \square

3 An Itô formula for $\|\cdot\|^p$

From now on we shall always assume that E is a 2-smooth Banach space. We fix $T \geq 0$ and let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space with a filtration $(\mathcal{F}_t)_{t \in [0, T]}$. Let H be a real Hilbert space, and denote by $\gamma(H, E)$ the Banach space of all γ -radonifying operators from H to E . We denote by $M([0, T]; \gamma(H, E))$ the space of all progressively measurable processes $\xi : [0, T] \times \Omega \rightarrow \gamma(H, E)$ such that

$$\int_0^T \|\xi_t\|_{\gamma(H, E)}^2 dt < \infty \quad \mathbb{P}\text{-almost surely.}$$

The space of all such ξ which satisfy

$$\mathbb{E} \left(\int_0^T \|\xi_t\|_{\gamma(H,E)}^2 dt \right)^{\frac{p}{2}} < \infty$$

is denoted by $M^p([0, T]; \gamma(H, E))$, $0 < p < \infty$.

On $(\Omega, \mathcal{F}, \mathbb{P})$, let $(W_t)_{t \in [0, T]}$ be an $(\mathcal{F}_t)_{t \in [0, T]}$ -cylindrical Brownian motion in H . For adapted simple processes $\xi \in M([0, T]; \gamma(H, E))$ of the form

$$\xi_t = \sum_{i=0}^{n-1} 1_{(t_i, t_{i+1}]}(t) \otimes A_i,$$

where $\Pi = \{0 = t_0 < t_1 < \dots < t_n = T\}$ is a partition of the interval $[0, T]$ and the random variables A_i are \mathcal{F}_{t_i} -measurable and take values in the space of all finite rank operators from H to E , we define the random variable $I(\xi) \in L^0(\Omega, \mathcal{F}_T; E)$ by

$$I(\xi) := \sum_{i=0}^{n-1} A_i(W_{t_{i+1}} - W_{t_i})$$

where $(h \otimes x)W_t := (W_t h) \otimes x$. It is well known that

$$\mathbb{E} \|I(\xi)\|^2 \leq C^2 \mathbb{E} \int_0^T \|\xi_t\|_{\gamma(H,E)}^2 dt,$$

where C depends on p and E only. It follows that I has a unique extension to a bounded linear operator $M^2([0, T]; \gamma(H, E))$ to $L^2(\Omega, \mathcal{F}_T; E)$. By a standard localisation argument, I extends continuous linear operator from $M([0, T]; \gamma(H, E))$ to $L^0(\Omega, \mathcal{F}_T; E)$. In what follows we write

$$\int_0^t \xi_s dW_s := I(1_{(0,t]} \xi), \quad t \in [0, T].$$

This stochastic integral has the following properties:

1. For all $\xi \in M([0, T]; \gamma(H, E))$ the process $t \rightarrow \int_0^t \xi_s dW_s$ is an E -valued continuous local martingale, which is a martingale if $\xi \in M^2([0, T]; \gamma(H, E))$.
2. For all $\xi \in M([0, T]; \gamma(H, E))$ and stopping times τ with values in $[0, T]$,

$$\int_0^\tau \xi_t dW_t = \int_0^T 1_{[0,\tau]}(t) \xi_t dW_t \quad \mathbb{P}\text{-almost surely.} \tag{3.1}$$

3. For all $\xi \in M^2([0, T]; \gamma(H, E))$ and $0 \leq u < t \leq T$,

$$\mathbb{E} \left(\left\| \int_u^t \xi_s dW_s \right\|^2 \middle| \mathcal{F}_u \right) \leq C \mathbb{E} \left(\int_u^t \|\xi_s\|_{\gamma(H,E)}^2 ds \middle| \mathcal{F}_u \right). \tag{3.2}$$

4. (Burkholder’s inequality [2, 6]) For all $0 < p < \infty$ there exists a constant C , depending only on p and E , such that for all $\xi \in M^p([0, T]; \gamma(H, E))$ and $t \in [0, T]$,

$$\mathbb{E} \sup_{s \in [0,t]} \left\| \int_0^s \xi_u dW_u \right\|^p \leq C \mathbb{E} \left(\int_0^t \|\xi_s\|_{\gamma(H,E)}^2 ds \right)^{\frac{p}{2}}. \tag{3.3}$$

An excellent survey of the theory of stochastic integration in 2-smooth Banach spaces with complete proofs is given in Ondreját's thesis [16], where also further references to the literature can be found.

In what follows we fix $p \geq 2$ and set $\psi(x) := \psi_p(x) = \|x\|^p$. Since we assume that E is 2-smooth, this function is Fréchet differentiable. Following the notation of Lemma 2.2 we set

$$R_\psi(x, y) := \int_0^1 (\psi'(x + r(y - x))(y - x) - \psi'(x)(y - x)) dr.$$

We have the following version of Itô's formula.

Theorem 3.1 (Itô formula). *Let E be a 2-smooth Banach space and let $2 \leq p < \infty$. Let $(a_t)_{t \in [0, T]}$ be an E -valued progressively measurable process such that*

$$\mathbb{E} \left(\int_0^T \|a_t\| dt \right)^p < \infty$$

and let $(g_t)_{t \in [0, T]}$ be a process in $M^p([0, T]; \gamma(H, E))$. Fix $x \in E$ and let $(X_t)_{t \in [0, T]}$ be given by

$$X_t = x + \int_0^t a_s ds + \int_0^t g_s dW_s.$$

The process $s \mapsto \psi'(X_s)g_s$ is progressively measurable and belongs to $M^1([0, T]; H)$, and for all $t \in [0, T]$ we have

$$\psi(X_t) = \psi(x) + \int_0^t \psi'(X_s)(a_s) ds + \int_0^t \psi'(X_s)(g_s) dW_s + \lim_{n \rightarrow \infty} \sum_{i=0}^{m(n)-1} R_\psi(X_{t_i^n \wedge t}, X_{t_{i+1}^n \wedge t}) \quad (3.4)$$

with convergence in probability, for any sequence of partitions $\Pi_n = \{0 = t_0^n < t_1^n < \dots < t_{m(n)}^n = T\}$ whose meshes $\|\Pi_n\| := \max_{0 \leq i \leq m(n)-1} |t_{i+1}^n - t_i^n|$ tend to 0 as $n \rightarrow \infty$. Moreover, there exists a constant C and, for each $\varepsilon > 0$, a constant C_ε , both independent of a and g , such that

$$\mathbb{E} \liminf_{n \rightarrow \infty} \sum_{i=0}^{m(n)-1} |R_\psi(X_{t_i^n \wedge t}, X_{t_{i+1}^n \wedge t})| \leq \varepsilon C \mathbb{E} \sup_{s \in [0, t]} \|X_s\|^p + C_\varepsilon \mathbb{E} \left(\int_0^t \|g_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{p}{2}}. \quad (3.5)$$

The proof shows that we may take $C_\varepsilon = C'(\varepsilon^{1-\frac{2}{p}} + 1)$ for some constant C' independent of a , g , and ε .

Before we start the proof of the theorem we state some lemmas. The first is an immediate consequence of Burkholder's inequality (3.3).

Lemma 3.2. *Under the assumptions of Theorem 3.1 we have*

$$\mathbb{E} \sup_{0 \leq t \leq T} \|X_t\|^p \leq C \mathbb{E} \left(\int_0^T \|a_s\| ds \right)^p + C \mathbb{E} \left(\int_0^T \|g_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{p}{2}}.$$

Lemma 3.3. *Under the assumptions of Theorem 3.1, the process $t \mapsto \psi'(X_t)(g_t)$ is progressively measurable and belongs to $M^1([0, T]; H)$.*

Proof. By the identity $\|\psi'(x)\| = p\|x\|^{p-1}$ and Hölder's inequality,

$$\begin{aligned} \mathbb{E}\left(\int_0^T \|\psi'(X_t)(g_t)\|_H^2 dt\right)^{\frac{1}{2}} &\leq \mathbb{E}\left(\int_0^T \|\psi'(X_t)\|^2 \|g_t\|_{\gamma(H,E)}^2 dt\right)^{\frac{1}{2}} \\ &\leq \mathbb{E} \sup_{t \in [0,T]} \|X_t\|^{p-1} \left(\int_0^T \|g_t\|_{\gamma(H,E)}^2 ds\right)^{\frac{1}{2}} \\ &\leq C \left(\mathbb{E} \sup_{t \in [0,T]} \|X_t\|^p\right)^{\frac{p-1}{p}} \left(\mathbb{E}\left(\int_0^T \|g_t\|_{\gamma(H,E)}^2 ds\right)^{\frac{p}{2}}\right)^{\frac{1}{p}}, \end{aligned}$$

and the right-hand side is finite by the previous lemma. The progressively measurability is clear. □

This lemma implies that the stochastic integral in (3.4) is well-defined.

Lemma 3.4. *Let $0 \leq u \leq t \leq T$ be arbitrary and fixed. Under the assumptions of Theorem 3.1, the process $s \mapsto \psi'(X_u)(g_s)$ is progressively measurable and belongs to $M^1([0, T]; H)$. Moreover, \mathbb{P} -almost surely,*

$$\psi'(X_u) \int_u^t g_s dW_s = \int_u^t \psi'(X_u)(g_s) dW_s.$$

Proof. By similar estimates as in the previous lemma,

$$\mathbb{E}\left(\int_u^t \|\psi'(X_u)(g_s)\|_H^2 ds\right)^{\frac{1}{2}} \leq C(\mathbb{E}\|X_u\|^p)^{\frac{p-1}{p}} \left(\mathbb{E}\left(\int_u^t \|g_s\|_{\gamma(H,E)}^2 ds\right)^{\frac{p}{2}}\right)^{\frac{1}{p}}.$$

The progressively measurability is again clear. To prove the identity we first assume that g is a simple adapted process of the form

$$g_s = \sum_{i=0}^{n-1} 1_{(t_i, t_{i+1}]}(s) A_i,$$

where $\Pi = \{u = t_0 < t_1 < \dots < t_n = t\}$ is a partition of the interval $[0, T]$ and the random variables are \mathcal{F}_{t_i} -measurable and take values in the space of all finite rank operators from H to E . Then,

$$\begin{aligned} \psi'(X_u) \int_u^t g_s dW_s &= \psi'(X_u) \left(\sum_{i=0}^{n-1} A_i (W_{t_{i+1}} - W_{t_i})\right) \\ &= \sum_{i=0}^{n-1} \psi'(X_u) (A_i (W_{t_{i+1}} - W_{t_i})) = \int_u^t \psi'(X_u)(g_s) dW_s. \end{aligned}$$

For general progressively measurable $g \in L^p(\Omega; L^2([0, T]; \gamma(H, E)))$, the identity follows by a routine approximation argument. □

Proof of Theorem 3.1. The proof of the theorem proceeds in two steps. All constants occurring in the proof may depend on E and p , even where this is not indicated explicitly, but not on T . The numerical value of the constants may change from line to line.

Step 1 – Applying Lemma 2.2 to the function $\psi(x) = \|x\|^p$ and the process X , we have, for every $t \in [0, T]$,

$$\begin{aligned} \psi(X_t) - \psi(x) &= \sum_{i=0}^{m(n)-1} \left(\psi(X_{t_{i+1}^n \wedge t}) - \psi(X_{t_i^n \wedge t}) \right) \\ &= \sum_{i=0}^{m(n)-1} \psi'(X_{t_i^n \wedge t})(X_{t_{i+1}^n \wedge t} - X_{t_i^n \wedge t}) + \sum_{i=0}^{m(n)-1} R_\psi(X_{t_i^n \wedge t}, X_{t_{i+1}^n \wedge t}). \end{aligned}$$

We shall prove the identity (3.4) by showing that

$$\lim_{n \rightarrow \infty} \sum_{i=0}^{m(n)-1} \psi'(X_{t_i^n \wedge t})(X_{t_{i+1}^n \wedge t} - X_{t_i^n \wedge t}) = \int_0^t \psi'(X_s)(a_s) ds + \int_0^t \psi'(X_s)(g_s) dW_s$$

with convergence in probability. In view of the definition of X_t , it is enough to show that

$$\lim_{n \rightarrow \infty} \left| \sum_{i=0}^{m(n)-1} \psi'(X_{t_i^n \wedge t}) \left(\int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} a_s ds \right) - \int_0^t \psi'(X_s)(a_s) ds \right| = 0 \quad \mathbb{P}\text{-almost surely}$$

and

$$\lim_{n \rightarrow \infty} \sum_{i=0}^{m(n)-1} \psi'(X_{t_i^n \wedge t}) \left(\int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} g_s dW_s \right) - \int_0^t \psi'(X_s)(g_s) dW_s = 0 \quad \text{in probability.} \quad (3.6)$$

By (2.2), \mathbb{P} -almost surely we have

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \left| \sum_{i=0}^{m(n)-1} \psi'(X_{t_i^n \wedge t}) \left(\int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} a_s ds \right) - \int_0^t \psi'(X_s)(a_s) ds \right| \\ & \leq \limsup_{n \rightarrow \infty} \sum_{i=0}^{m(n)-1} \left| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} (\psi'(X_{t_i^n \wedge t}) - \psi'(X_s))(a_s) ds \right| \\ & \leq C \sup_{s \in [0, T]} \|X_s\|^{p-2} \times \limsup_{n \rightarrow \infty} \sum_{i=0}^{m(n)-1} \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} \|X_{t_i^n \wedge t} - X_s\| \|a_s\| ds \\ & \leq C \sup_{s \in [0, T]} \|X_s\|^{p-2} \times \limsup_{n \rightarrow \infty} \left(\sup_{0 \leq i \leq m(n)-1} \sup_{s \in [t_i^n \wedge t, t_{i+1}^n \wedge t]} \|X_{t_i^n \wedge t} - X_s\| \right) \\ & \quad \times \left(\sum_{i=0}^{m(n)-1} \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} \|a_s\| ds \right) \\ & = 0, \end{aligned}$$

where we used the continuity of the process X in the last line.

Next, by Lemma 3.4 and the inequalities (3.2) and (2.2),

$$\sum_{i=0}^{m(n)-1} \psi'(X_{t_i^n \wedge t}) \left(\int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} g_s dW_s \right) - \int_0^t \psi'(X_s)(g_s) dW_s$$

$$\begin{aligned} &= \sum_{i=0}^{m(n)-1} \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} \psi'(X_{t_i^n \wedge t})(g_s) dW_s - \int_0^t \psi'(X_s)(g_s) dW_s \\ &= \int_0^t \sum_{i=0}^{m(n)-1} 1_{(t_i^n, t_{i+1}^n]}(s) (\psi'(X_{t_i^n \wedge t}) - \psi'(X_s))(g_s) dW_s. \end{aligned}$$

The localized stochastic integral being continuous from $M([0, t]; \gamma(H, E))$ into $L^0(\Omega, \mathcal{F}_t; E)$, in order to prove that the right-hand side converges to 0 in probability it suffices to prove that

$$\lim_{n \rightarrow \infty} \left\| s \mapsto \sum_{i=0}^{m(n)-1} 1_{(t_i^n, t_{i+1}^n]}(s) (\psi'(X_{t_i^n \wedge t}) - \psi'(X_s))(g_s) \right\|_{L^2([0, t]; H)} = 0 \text{ in probability.}$$

For this, in turn, it suffices to observe that \mathbb{P} -almost surely

$$\begin{aligned} &\lim_{n \rightarrow \infty} \left\| \sum_{i=0}^{m(n)-1} 1_{(t_i^n, t_{i+1}^n]}(s) (\psi'(X_{t_i^n \wedge t}) - \psi'(X_s)) \right\|_{L^\infty([0, t]; E^*)} \\ &= \lim_{n \rightarrow \infty} \sup_{0 \leq i \leq n-1} \sup_{s \in [t_i^n \wedge t, t_{i+1}^n \wedge t]} \|\psi'(X_{t_i^n \wedge t}) - \psi'(X_s)\| = 0 \end{aligned}$$

by the path continuity of X .

Step 2 – In this step we prove the estimate (3.5). By (2.2), for all $x, y \in E$ and $r \in [0, 1]$ we have

$$|\psi'(x + r(y - x)) - \psi'(x)| \leq (\|x\|^{p-2} \|x - y\| + \|x - y\|^{p-1}).$$

Combining this with (2.5) we obtain

$$|R_\psi(X_{t_i^n \wedge t}, X_{t_{i+1}^n \wedge t})| \leq C \|X_{t_i^n \wedge t}\|^{p-2} \|X_{t_{i+1}^n \wedge t} - X_{t_i^n \wedge t}\|^2 + C \|X_{t_{i+1}^n \wedge t} - X_{t_i^n \wedge t}\|^p. \tag{3.7}$$

We shall estimate the two terms on the right hand of (3.7) side separately.

For the first term, using the inequality $|a + b|^2 \leq 2|a|^2 + 2|b|^2$ we obtain

$$\begin{aligned} &\sum_{i=0}^{m(n)-1} \|X_{t_i^n \wedge t}\|^{p-2} \|X_{t_{i+1}^n \wedge t} - X_{t_i^n \wedge t}\|^2 \\ &\leq 2 \sum_{i=0}^{m(n)-1} \|X_{t_i^n \wedge t}\|^{p-2} \left\| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} a_s ds \right\|^2 + 2 \sum_{i=0}^{m(n)-1} \|X_{t_i^n \wedge t}\|^{p-2} \left\| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} g_s dW_s \right\|^2 \\ &=: I_1^n + I_2^n. \end{aligned}$$

For the first term we have

$$\begin{aligned} I_1^n &\leq 2C \sup_{s \in [0, t]} \|X_s\|^{p-2} \times \sup_i \left\| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} a_s ds \right\| \times \sum_{i=0}^{m(n)-1} \left\| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} a_s ds \right\| \\ &\leq 2C \sup_{s \in [0, t]} \|X_s\|^{p-2} \times \sup_i \left\| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} a_s ds \right\| \times \int_0^t \|a_s\| ds. \end{aligned}$$

By letting $n \rightarrow \infty$ we have $\max_{0 \leq i \leq m(n)-1} (t_{i+1}^n - t_i^n) \rightarrow 0$, so

$$\sup_{0 \leq i \leq m(n)-1} \left\| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} a_s ds \right\| \rightarrow 0$$

as $n \rightarrow \infty$. Therefore,

$$\lim_{n \rightarrow \infty} I_1^n = 0, \mathbb{P}\text{-almost surely.}$$

To estimate I_2 we use (3.2) and Young's inequality with $\varepsilon > 0$ to infer

$$\begin{aligned} \mathbb{E} \liminf_n I_2^n &\leq \liminf_n \mathbb{E} I_2^n = \liminf_n \mathbb{E} \sum_{i=0}^{m(n)-1} \|X_{t_i^n \wedge t}\|^{p-2} \left\| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} g_s dW_s \right\|^2 \\ &= \liminf_n \sum_{i=0}^{m(n)-1} \mathbb{E} \left(\|X_{t_i^n \wedge t}\|^{p-2} \mathbb{E} \left(\left\| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} g_s dW_s \right\|^2 \middle| \mathcal{F}_{t_i^n \wedge t} \right) \right) \\ &\leq C \liminf_n \sum_{i=0}^{m(n)-1} \mathbb{E} \left(\|X_{t_i^n \wedge t}\|^{p-2} \mathbb{E} \left(\int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} \|g_s\|_{\gamma(H,E)}^2 ds \middle| \mathcal{F}_{t_i^n \wedge t} \right) \right) \\ &\leq C \liminf_n \sum_{i=0}^{m(n)-1} \mathbb{E} \left(\|X_{t_i^n \wedge t}\|^{p-2} \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} \|g_s\|_{\gamma(H,E)}^2 ds \right) \\ &\leq C \liminf_n \mathbb{E} \left(\sup_{s \in [0,t]} \|X_s\|^{p-2} \sum_{i=0}^{m(n)-1} \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} \|g_s\|_{\gamma(H,E)}^2 ds \right) \\ &= C \mathbb{E} \left(\sup_{s \in [0,t]} \|X_s\|^{p-2} \int_0^t \|g_s\|_{\gamma(H,E)}^2 ds \right) \\ &\leq C \varepsilon \mathbb{E} \left(\sup_{s \in [0,t]} \|X_s\|^p \right) + C \varepsilon^{1-\frac{p}{2}} \mathbb{E} \left(\int_0^t \|g_s\|_{\gamma(H,E)}^2 ds \right)^{\frac{p}{2}}. \end{aligned}$$

Next we estimate the second term in (3.7). We have

$$\begin{aligned} \sum_{i=0}^{m(n)-1} \|X_{t_{i+1}^n \wedge t} - X_{t_i^n \wedge t}\|^p &\leq C \sum_{i=0}^{m(n)-1} \left\| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} a_s ds \right\|^p + C \sum_{i=0}^{m(n)-1} \left\| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} g_s dW_s \right\|^p \\ &=: I_3^n + I_4^n. \end{aligned}$$

A similar consideration as before yields

$$\lim_{n \rightarrow \infty} I_3^n \leq C \lim_{n \rightarrow \infty} \sup_{0 \leq i \leq m(n)-1} \left\| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} a_s ds \right\|^{p-1} \times \int_0^t \|a_s\| ds = 0.$$

Moreover, by Burkholder's inequality (3.3),

$$\mathbb{E} \liminf_n I_4^n \leq \liminf_n \mathbb{E} I_4^n = C \liminf_n \sum_{i=0}^{m(n)-1} \mathbb{E} \left\| \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} g_s dW_s \right\|^p$$

$$\begin{aligned}
&\leq C \liminf_n \sum_{i=0}^{m(n)-1} \mathbb{E} \left(\int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} \|g_s\|_{\gamma(H,E)}^2 ds \right)^{\frac{p}{2}} \\
&\leq C \liminf_n \mathbb{E} \left(\sum_{i=0}^{m(n)-1} \int_{t_i^n \wedge t}^{t_{i+1}^n \wedge t} \|g_s\|_{\gamma(H,E)}^2 ds \right)^{\frac{p}{2}} \\
&= C \mathbb{E} \left(\int_0^t \|g_s\|_{\gamma(H,E)}^2 ds \right)^{\frac{p}{2}}.
\end{aligned}$$

Collecting terms, for any $\varepsilon > 0$ we obtain the estimate

$$\begin{aligned}
&\mathbb{E} \liminf_{n \rightarrow \infty} \sum_{i=0}^{m(n)-1} |R_\psi(X_{t_i^n \wedge t}, X_{t_{i+1}^n \wedge t})| \\
&\leq C \varepsilon \mathbb{E} \left(\sup_{s \in [0, t]} \|X_s\|^p \right) + C(\varepsilon^{1-\frac{p}{2}} + 1) \mathbb{E} \left(\int_0^t \|g_s\|_{\gamma(H,E)}^2 ds \right)^{\frac{p}{2}}.
\end{aligned}$$

□

In the proof of Theorem 1.2 we will also need the following simple observation.

Lemma 3.5. \mathbb{P} -Almost surely we have

$$\liminf_{n \rightarrow \infty} \sup_{t \in [0, T]} \sum_{i=0}^{m(n)-1} |R_\psi(X_{t_i^n \wedge t}, X_{t_{i+1}^n \wedge t})| \leq \liminf_{n \rightarrow \infty} \sum_{i=0}^{m(n)-1} |R_\psi(X_{t_i^n}, X_{t_{i+1}^n})|. \quad (3.8)$$

Proof. Fix $t \in (0, T]$ and let $k(n)$ be the unique index such that $t \in (t_{k(n)}^n, t_{k(n)+1}^n]$. Then

$$\begin{aligned}
&\sum_{i=0}^{m(n)-1} |R_\psi(X_{t_i^n \wedge t}, X_{t_{i+1}^n \wedge t})| \\
&= \sum_{i=0}^{k(n)-1} |R_\psi(X_{t_i^n}, X_{t_{i+1}^n})| + |R_\psi(X_{t_{k(n)}^n}, X_t)| \\
&\leq \sum_{i=0}^{m(n)-1} |R_\psi(X_{t_i^n}, X_{t_{i+1}^n})| + |R_\psi(X_{t_{k(n)}^n}, X_t)| \\
&\leq \sum_{i=0}^{m(n)-1} |R_\psi(X_{t_i^n}, X_{t_{i+1}^n})| + C \|X_{t_{k(n)}^n}\|^{p-2} \|X_t - X_{t_{k(n)}^n}\|^2 + C \|X_t - X_{t_{k(n)}^n}\|^p \\
&\leq \sum_{i=0}^{m(n)-1} |R_\psi(X_{t_i^n}, X_{t_{i+1}^n})| + C \sup_{s \in [0, T]} \|X_s\|^{p-2} \|X_t - X_{t_{k(n)}^n}\|^2 + C \|X_t - X_{t_{k(n)}^n}\|^p.
\end{aligned}$$

Now (3.8) follows by taking the limes inferior for $n \rightarrow \infty$ and using path continuity. □

4 Proof of Theorem 1.2

We proceed in four steps. In Steps 1 and 2 we establish the estimate in the theorem for $g \in M^p([0, T]; \gamma(H, E))$ with $2 \leq p < \infty$. In order to be able to cover exponents $0 < p < 2$ in Step 3,

we need a stopped version of the inequalities proved in Steps 1 and 2. For reasons of economy of presentations, we therefore build in a stopping time τ from the start. In Step 4 we finally consider the case where $g \in M([0, T]; \gamma(H, E))$.

We shall apply (a special case of) Lenglart's inequality [13, Corollaire II] which states that if $(\xi_t)_{t \in [0, T]}$ and $(a_t)_{t \in [0, T]}$ are continuous non-negative adapted processes, the latter non-decreasing, such that $\mathbb{E}\xi_\tau \leq \mathbb{E}a_\tau$ for all stopping times τ with values in $[0, T]$, then for all $0 < r < 1$ one has

$$\mathbb{E} \sup_{0 \leq t \leq T} \xi_t^r \leq \frac{2-r}{1-r} \mathbb{E}a_T^r. \quad (4.1)$$

Step 1 – Fix $p \geq 2$ and suppose first that $g \in M^p([0, T]; \gamma(H, D(A)))$. As is well known (see [16]), under this condition the process $X_t = \int_0^t e^{(t-s)A} g_s dW_s$ is a strong solution to the equation

$$dX_t = AX_t dt + g_t dW_t, \quad t \geq 0; \quad X_0 = 0.$$

In other words, X satisfies

$$X_t = \int_0^t AX_s ds + \int_0^t g_s dW_s \quad \forall t \in [0, T] \quad \mathbb{P}\text{-almost surely.}$$

Hence if τ is a stopping time with values in $[0, T]$, then by (3.1),

$$X_{t \wedge \tau} = \int_0^t 1_{[0, \tau]}(s) AX_s ds + \int_0^t 1_{[0, \tau]}(s) g_s dW_s \quad \forall t \in [0, T], \quad \mathbb{P}\text{-almost surely.}$$

Let us check next that $a_t := 1_{[0, \tau]}(t) AX_t$ satisfies the assumptions of Theorem 3.1. Indeed, with $h_t := 1_{[0, \tau]}(t) Ag_t$ we have, using the contractivity of the semigroup S and Burkholder's inequality (3.3),

$$\begin{aligned} \mathbb{E} \left(\int_0^T \|a_t\| dt \right)^p &\leq \mathbb{E} \left(\int_0^T \left\| \int_0^t e^{(t-s)A} h_s dW_s \right\| dt \right)^p \\ &\leq CT^{p-1} \mathbb{E} \int_0^T \left\| \int_0^t e^{(t-s)A} h_s dW_s \right\|^p dt \\ &\leq CT^{p-1} \mathbb{E} \int_0^T \left(\int_0^t \|e^{(t-s)A} h_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{p}{2}} dt \\ &\leq CT^p \mathbb{E} \left(\int_0^T \|h_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{p}{2}} < \infty. \end{aligned}$$

Hence we may apply Theorem 3.1 and infer that

$$\begin{aligned} \|X_{t \wedge \tau}\|^p &= \int_0^t 1_{[0, \tau]}(s) \psi'(X_s)(AX_s) ds \\ &\quad + \int_0^t 1_{[0, \tau]}(s) \psi'(X_s)(g_s) dW_s + \lim_{n \rightarrow \infty} \sum_{i=0}^{m(n)-1} R_\psi(X_{t_i^n \wedge t \wedge \tau}, X_{t_{i+1}^n \wedge t \wedge \tau}) \\ &\leq \int_0^t 1_{[0, \tau]}(s) \psi'(X_s)(g_s) dW_s + \lim_{n \rightarrow \infty} \sum_{i=0}^{m(n)-1} R_\psi(X_{t_i^n \wedge t \wedge \tau}, X_{t_{i+1}^n \wedge t \wedge \tau}), \end{aligned}$$

since $\psi'(x)(Ax) \leq 0$ for all $x \in D(A)$ by the contractivity of e^{tA} (see [3, Lemma 4.2]). Hence, by Lemma 3.5,

$$\begin{aligned}
& \mathbb{E} \sup_{t \in [0, T]} \|X_{t \wedge \tau}\|^p \\
& \leq \mathbb{E} \sup_{t \in [0, T]} \int_0^t \mathbf{1}_{[0, \tau]}(s) \psi'(X_s)(g_s) dW_s + \mathbb{E} \sup_{t \in [0, T]} \liminf_{n \rightarrow \infty} \sum_{i=0}^{m(n)-1} |R_\psi(X_{t_i^n \wedge t \wedge \tau}, X_{t_{i+1}^n \wedge t \wedge \tau})| \\
& \leq \mathbb{E} \sup_{t \in [0, T]} \int_0^t \mathbf{1}_{[0, \tau]}(s) \psi'(X_s)(g_s) dW_s + \mathbb{E} \liminf_{n \rightarrow \infty} \sum_{i=0}^{m(n)-1} |R_\psi(X_{t_i^n \wedge \tau}, X_{t_{i+1}^n \wedge \tau})| \\
& \leq C \mathbb{E} \sup_{t \in [0, T]} \int_0^t \mathbf{1}_{[0, \tau]}(s) \psi'(X_s)(g_s) dW_s \\
& \quad + \varepsilon C \mathbb{E} \sup_{s \in [0, T]} \|X_{s \wedge \tau}\|^p + C_\varepsilon \mathbb{E} \left(\int_0^T \mathbf{1}_{[0, \tau]}(s) \|g_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{p}{2}}.
\end{aligned}$$

By Burkholder's inequality (3.3) and the identity $\|\psi'(y)\| = p\|y\|^{p-1}$,

$$\begin{aligned}
& \mathbb{E} \sup_{t \in [0, T]} \left| \int_0^t \mathbf{1}_{[0, \tau]}(s) \psi'(X_s)(g_s) dW_s \right| \\
& \leq C \mathbb{E} \left(\int_0^T \mathbf{1}_{[0, \tau]}(s) \|\psi'(X_s)\|^2 \|g_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{1}{2}} \\
& = C \mathbb{E} \left(\int_0^T \mathbf{1}_{[0, \tau]}(s) \|X_s\|^{2(p-1)} \|g_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{1}{2}} \\
& \leq C \mathbb{E} \left(\sup_{t \in [0, T]} \|X_{t \wedge \tau}\|^{p-1} \left(\int_0^T \mathbf{1}_{[0, \tau]}(s) \|g_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{1}{2}} \right) \\
& \leq C p^p \left(\mathbb{E} \sup_{t \in [0, T]} \|X_{t \wedge \tau}\|^p \right)^{\frac{p-1}{p}} \left(\mathbb{E} \left(\int_0^T \mathbf{1}_{[0, \tau]}(s) \|g_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{p}{2}} \right)^{\frac{1}{p}} \\
& \leq C \varepsilon \mathbb{E} \sup_{t \in [0, T]} \|X_{t \wedge \tau}\|^p + C_\varepsilon \mathbb{E} \left(\int_0^T \mathbf{1}_{[0, \tau]}(s) \|g_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{p}{2}},
\end{aligned}$$

where we also used the Hölder's inequality and Young's inequality.

Combining these estimates and taking $\varepsilon > 0$ small enough, we infer that

$$\mathbb{E} \sup_{t \in [0, T]} \|X_{t \wedge \tau}\|^p \leq C \mathbb{E} \left(\int_0^T \mathbf{1}_{[0, \tau]}(s) \|g_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{p}{2}}.$$

Step 2 – Now let $g \in M^p([0, T]; \gamma(H, E))$ be arbitrary. Set $g^n = n(nI - A)^{-1}g$, $n \geq 1$. These processes satisfy the assumptions of Step 1 and we have $\|g^n\|_{\gamma(H, E)} \leq \|g\|_{\gamma(H, E)}$ pointwise. Define $X_t^n = \int_0^t e^{(t-s)A} g_s^n ds$. From Step 1 we know that for any stopping time τ in $[0, T]$ we have

$$\mathbb{E} \sup_{t \in [0, T]} \|X_{t \wedge \tau}^n\|^p \leq C \mathbb{E} \left(\int_0^T \mathbf{1}_{[0, \tau]}(s) \|g_s^n\|_{\gamma(H, E)}^2 ds \right)^{\frac{p}{2}}.$$

In particular, as $n, m \rightarrow \infty$,

$$\mathbb{E} \sup_{t \in [0, T]} \|X_t^n - X_t^m\|^p \rightarrow 0.$$

In these circumstances there is a process \bar{X} such that $\lim_{n \rightarrow \infty} \mathbb{E} \sup_{t \in [0, T]} \|\bar{X}_t^n - X_t\|^p = 0$ and

$$\mathbb{E} \sup_{t \in [0, T]} \|\bar{X}_{t \wedge \tau}\|^p \leq C \mathbb{E} \left(\int_0^T \mathbf{1}_{[0, \tau]}(s) \|g_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{p}{2}}. \quad (4.2)$$

Also, notice that for every $t \in [0, T]$, we have

$$\mathbb{E} \|X_t^n - X_t\|^p = \mathbb{E} \left\| \int_0^t e^{(t-s)A} g_s^n ds - \int_0^t e^{(t-s)A} g_s ds \right\|^p \leq C \left(\mathbb{E} \int_0^t \|g_s^n - g_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{p}{2}}.$$

Hence $X_t^n \rightarrow X_t$ in $L^p(\Omega; E)$. Therefore, \bar{X} is a modification of X . This concludes the proof for $p \geq 2$.

Step 3 – In this step we extend the result to exponents $0 < p < 2$. First consider the case where $g \in M^2([0, T]; \gamma(H, E))$. By (4.2), for all stopping times τ in $[0, T]$ we have

$$\mathbb{E} \|X_\tau\|^2 \leq C \mathbb{E} \int_0^\tau \|g_s\|_{\gamma(H, E)}^2 ds.$$

It then follows from Lenglart's inequality (4.1) that for all $0 < p < 2$,

$$\mathbb{E} \sup_{t \in [0, T]} \|X_t\|^p \leq C \mathbb{E} \left(\int_0^T \|g_s\|_{\gamma(H, E)}^2 ds \right)^{\frac{p}{2}}.$$

For $g \in M^p([0, T]; \gamma(H, E))$ the result follows by approximation.

Step 4 – Finally, the existence of a continuous version for the process X under the assumption $g \in M([0, T]; \gamma(H, E))$ follows by a standard localisation argument.

References

- [1] G. A. Anastassiou and S. S. Dragomir, On some estimates of the remainder in Taylor's formula, *J. Math. Anal. Appl.* **263** (2001), no. 1, 246–263. [MR1865279](#) (2002g:26019)
- [2] P. Assouad, Espaces p -lisses et p -convexes, inégalités de Burkholder, *Sém. Maurey-Schwartz 1974-1975*, "Espaces L^p , applications radonifiantes, géométrie des espaces de Banach", Exposé XV, École polytechnique, Centre de Mathématiques, Paris, 1975.
- [3] Z. Brzeźniak, E. Hausenblas, J. Zhu, Maximal inequality of stochastic convolution driven by compensated Poisson random measures in Banach spaces. [arXiv:1005.1600](#).
- [4] Z. Brzeźniak and S. Peszat, Maximal inequalities and exponential estimates for stochastic convolutions in Banach spaces, in *Stochastic processes, physics and geometry: new interplays, I (Leipzig, 1999)*, 55–64, CMS Conf. Proc., 28 Amer. Math. Soc., Providence, RI. [MR1803378](#) (2001k:60084)

- [5] G. Da Prato and J. Zabczyk, A note on stochastic convolution, *Stochastic Anal. Appl.* **10** (1992), no. 2, 143–153. [MR1154532](#) (93e:60116)
- [6] E. Dettweiler, Stochastic integration relative to Brownian motion on a general Banach space, *Doğa Mat.* **15** (1991), no. 2, 58–97. [MR1115509](#) (93b:60112)
- [7] R. Deville, G. Godefroy and V. Zizler, *Smoothness and renormings in Banach spaces*, Pitman Monographs and Surveys in Pure and Applied Mathematics, 64, Longman Sci. Tech., Harlow, 1993. [MR1211634](#) (94d:46012)
- [8] T. Figiel, On the moduli of convexity and smoothness, *Studia Math.* **56** (1976), no. 2, 121–155. [MR0425581](#) (54 #13535)
- [9] E. Hausenblas and J. Seidler, A note on maximal inequality for stochastic convolutions, *Czechoslovak Math. J.* **51(126)** (2001), no. 4, 785–790. [MR1864042](#) (2002j:60092)
- [10] A. Ichikawa, Some inequalities for martingales and stochastic convolutions, *Stochastic Anal. Appl.* **4** (1986), no. 3, 329–339. [MR0857085](#) (87m:60105)
- [11] P. Kotelenez, A submartingale type inequality with applications to stochastic evolution equations, *Stochastics* **8** (1982/83), no. 2, 139–151. [MR0686575](#) (84h:60115)
- [12] P. Kotelenez, A stopped Doob inequality for stochastic convolution integrals and stochastic evolution equations, *Stochastic Anal. Appl.* **2** (1984), no. 3, 245–265. [MR0757338](#) (86k:60096)
- [13] E. Lenglart, Relation de domination entre deux processus, *Ann. Inst. H. Poincaré Sect. B (N.S.)* **13** (1977), no. 2, 171–179. [MR0471069](#) (57 #10810)
- [14] I. E. Leonard and K. Sundaresan, Geometry of Lebesgue-Bochner function spaces—smoothness, *Trans. Amer. Math. Soc.* **198** (1974), 229–251. [MR0367652](#) (51 #3894)
- [15] A.L. Neidhardt, “Stochastic integrals in 2-uniformly smooth Banach spaces”, Ph.D Thesis, University of Wisconsin, 1978.
- [16] M. Ondreját, Uniqueness for stochastic evolution equations in Banach spaces, *Dissertationes Math. (Rozprawy Mat.)* **426** (2004), 63 pp. [MR2067962](#) (2005e:60133)
- [17] G. Pisier, Martingales with values in uniformly convex spaces, *Israel J. Math.* **20** (1975), no. 3-4, 326–350. [MR0394135](#) (52 #14940)
- [18] J. Seidler, Exponential estimates for stochastic convolutions in 2-smooth Banach spaces, *Electron. J. Probab.* **15** (2010), no. 50, 1556–1573. [MR2735374](#) (2011k:60223)
- [19] L. Tubaro, An estimate of Burkholder type for stochastic processes defined by the stochastic integral, *Stochastic Anal. Appl.* **2** (1984), no. 2, 187–192. [MR0746435](#) (85h:60072)
- [20] M.C. Veraar, L.W. Weis, A note on maximal estimates for stochastic convolutions, *Czechoslovak Math. J.* **61** (2011), no. 3: 743-758.