

# More on the anti-automorphism of the Steenrod algebra

VINCE GIAMBALVO  
 HAYNES R MILLER

The relations of Barratt and Miller are shown to include all relations among the elements  $P^i \chi P^{n-i}$  in the mod  $p$  Steenrod algebra, and a minimal set of relations is given.

55S10

## 1 Introduction

Milnor [4] observed that the mod 2 Steenrod algebra  $\mathcal{A}$  forms a Hopf algebra with commutative diagonal determined by

$$(1) \quad \Delta \text{Sq}^n = \sum_i \text{Sq}^i \otimes \text{Sq}^{n-i} .$$

This allowed him to interpret the Cartan formula as the assertion that the cohomology of a space forms a module-algebra over  $\mathcal{A}$ . The anti-automorphism  $\chi$  in the Hopf algebra structure, defined inductively by

$$(2) \quad \chi \text{Sq}^0 = \text{Sq}^0, \quad \sum_i \text{Sq}^i \chi \text{Sq}^{n-i} = 0 \quad \text{for } n > 0,$$

has a topological interpretation too: If  $K$  is a finite complex then the homology of the Spanier–Whitehead dual  $DK_+$  of  $K_+$  is canonically isomorphic to the cohomology of  $K$ . Under this isomorphism the left action by  $\theta \in \mathcal{A}$  on  $H^*(K)$  corresponds to the right action of  $\chi\theta \in \mathcal{A}$  on  $H_*(DK_+)$ .

In 1974 Davis [3] proved that sometimes much more efficient ways exist to compute  $\chi \text{Sq}^n$ ; for example

$$(3) \quad \chi \text{Sq}^{2^r-1} = \text{Sq}^{2^r-1} \chi \text{Sq}^{2^r-1-1},$$

$$(4) \quad \chi \text{Sq}^{2^r-r-1} = \text{Sq}^{2^r-1-1} \chi \text{Sq}^{2^r-1-r} + \text{Sq}^{2^r-1} \chi \text{Sq}^{2^r-1-r-1} .$$

Similarly, Straffin [6] proved that if  $r \geq 0$  and  $b \geq 2$  then

$$(5) \quad \sum_i \text{Sq}^{2^r i} \chi \text{Sq}^{2^r(b-i)} = 0 .$$

Both authors give analogous identities among reduced powers and their images under  $\chi$  at an odd prime as well. Further relations among the Steenrod squares and their conjugates appear in these articles and elsewhere (eg Silverman [5]).

Barratt and Miller [1] found a general family of identities which includes (3), (4) and (5), and their odd-prime analogues, as special cases. We state it for the general prime. When  $p = 2$ ,  $P^n$  denotes  $Sq^n$ . Let  $\alpha(n)$  denote the sum of the  $p$ -adic digits of  $n$ .

**Theorem 1.1** [1; 2] *For any integer  $k$  and any integer  $l \geq 0$  such that  $pl - \alpha(l) < (p - 1)n$ ,*

$$(6) \quad \sum_i \binom{k-i}{l} P^i \chi P^{n-i} = 0.$$

The relations defining  $\chi$  occur with  $l = 0$ . Davis' formulas (for  $p = 2$ ) are the cases in which  $(n, l, k) = (2^r - 1, 2^{r-1} - 1, 2^r - 1)$  or  $(n, l, k) = (2^r - r - 1, 2^{r-1} - 2, 2^r - 2)$ . Straffin's identities (for  $p = 2$ ) occur as  $(n, l, k) = (2^r b, 2^r - 1, -1)$ .

Since  $\binom{(k+1)-i}{l} - \binom{k-i}{l} = \binom{k-i}{l-1}$ , the cases  $(l, k + 1)$  and  $(l, k)$  of (6) imply it for  $(l - 1, k)$ . Thus the relations for  $l = \phi(n) - 1$ , where

$$(7) \quad \phi(n) = 1 + \max\{j : pj - \alpha(j) < (p - 1)n\},$$

imply all the rest. Here we have adopted the notation  $\phi(n)$  used in [2]; we note that it is not the Euler function  $\varphi(n)$ .

When  $p = 2$ ,  $\phi(2^r - 1) = 2^{r-1}$  and  $\phi(2^r - r - 1) = 2^{r-1} - 1$ , so Davis's relations are among these basic relations.

Two questions now arise. To express them uniformly in the prime, let  $\mathcal{P}$  denote the algebra of Steenrod reduced powers (which is the full Steenrod algebra when  $p = 2$ ), but assign  $P^n$  degree  $n$ . Write

$$V_n = \text{Span}\{P^i \chi P^{n-i} : 0 \leq i \leq n\} \subseteq \mathcal{P}^n.$$

It is natural to ask:

- Are there yet other linear relations among the  $n + 1$  elements  $P^i \chi P^{n-i}$  in  $\mathcal{P}^n$ ?
- What is a basis for  $V_n$ ?

We answer these questions in Theorem 1.4 below.

Write  $e_i, 0 \leq i \leq n$ , for the  $i$ -th standard basis vector in  $\mathbb{F}_p^{n+1}$ .

**Proposition 1.2** For any integers  $l, m, n$ , with  $0 \leq l \leq n$ ,

$$(8) \quad \left\{ \sum_i \binom{k-i}{l} e_i : m \leq k \leq m+l \right\}$$

is linear independent in  $\mathbb{F}_p^{n+1}$ .

**Proposition 1.3** The set

$$(9) \quad \{P^i \chi P^{n-i} : \phi(n) \leq i \leq n\}$$

is linearly independent in  $\mathcal{P}^n$ .

Define a linear map

$$(10) \quad \mu: \mathbb{F}_p^{n+1} \rightarrow \mathcal{P}^n, \quad \mu e_i = P^i \chi P^{n-i}.$$

**Theorem 1.1** implies that if  $l = \phi(n) - 1$  the elements in (8) lie in  $\ker \mu$ , so Propositions 1.2 and 1.3 imply that (8) with  $l = \phi(n) - 1$  is a basis for  $\ker \mu$  and that (9) is a basis for  $V_n \subseteq \mathcal{P}^n$ . Thus:

**Theorem 1.4** Any  $\phi(n)$  consecutive relations from the set (6) with  $l = \phi(n) - 1$  form a basis of relations among the elements of  $\{P^i \chi P^{n-i} : 0 \leq i \leq n\}$ . The set  $\{P^i \chi P^{n-i} : \phi(n) \leq i \leq n\}$  is a basis for  $V_n$ .

**Acknowledgements** We thank Richard Stanley for the slick proof of Proposition 1.2. This material is based upon work supported by the National Science Foundation under grant number 0905950.

## 2 Independence of the relations

We wish to show that (8) is a linearly independent set. Regard elements of  $\mathbb{F}_p^{n+1}$  as column vectors, and arrange the  $l + 1$  vectors in (8) as columns in a matrix, which we claim is of rank  $l + 1$ . The top square portion is the mod  $p$  reduction of the  $(l + 1) \times (l + 1)$  integral Toeplitz matrix  $A_l(m)$  with  $(i, j)$ -th entry

$$\binom{m+j-i}{l}, \quad 0 \leq i, j \leq l.$$

**Lemma 2.1**  $\det A_l(m) = 1$ .

**Proof** By induction on  $m$ . Since  $\binom{-1}{l} = (-1)^l$  and  $\binom{-1+j}{l} = 0$  for  $0 < j \leq l$ ,  $A_l(-1)$  is lower triangular with determinant  $((-1)^l)^{l+1} = 1$ . Now we note the identity

$$BA_l(m) = A_l(m + 1)$$

where

$$B = \begin{bmatrix} \binom{l+1}{1} & -\binom{l+1}{2} & \dots & (-1)^{l-1} \binom{l+1}{l} & (-1)^l \binom{l+1}{l+1} \\ 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \dots & 1 & 0 \end{bmatrix}.$$

The matrix identity is an expression of the binomial identity

$$(11) \quad \sum_k (-1)^k \binom{l+1}{k} \binom{n-k}{l} = 0$$

(taking  $n = m + 1 - j$  and  $k = j + 1$ ). Since  $\det B = 1$ , the result follows for all  $m \in \mathbb{Z}$ . □

For completeness, we note that (11) is the case  $m = l + 1$  of the equation

$$(12) \quad \sum_k (-1)^k \binom{m}{k} \binom{n-k}{l} = \binom{n-m}{l-m}.$$

To prove this formula, note that the defining identity for binomial coefficients implies the case  $m = 1$ , and also that both sides satisfy the recursion  $C(l, m, n) - C(l, m, n - 1) = C(l, m + 1, n)$ .

### 3 Independence of the operations

We will prove Proposition 1.3 by studying how  $P^i \chi P^{n-i}$  pairs against elements in  $\mathcal{P}_*$ , the dual of the Hopf algebra of Steenrod reduced powers. According to Milnor [4], with our grading conventions

$$(13) \quad \begin{aligned} \mathcal{P}_* &= \mathbb{F}_p[\xi_1, \xi_2, \dots], & |\xi_j| &= \frac{p^j - 1}{p - 1}, \\ \Delta \xi_k &= \sum_{i+j=k} \xi_i^{p^j} \otimes \xi_j. \end{aligned}$$

For a finitely nonzero sequence of nonnegative integers  $R = (r_1, r_2, \dots)$  write  $\xi^R = \xi_1^{r_1} \xi_2^{r_2} \dots$  and let  $\|R\| = r_1 + pr_2 + p^2r_3 + \dots$  and

$$|R| = |\xi^R| = r_1 + \left(\frac{p^2 - 1}{p - 1}\right)r_2 + \left(\frac{p^3 - 1}{p - 1}\right)r_3 + \dots$$

The following clearly implies [Proposition 1.3](#).

**Proposition 3.1** *For any integer  $n > 0$  there exist sequences  $R_{n,j}$ ,  $0 \leq j \leq n - \phi(n)$ , such that  $|R_{n,j}| = n$  and*

$$\langle P^i \chi P^{n-i}, \xi^{R_{n,j}} \rangle = \begin{cases} \pm 1 & \text{for } i = n - j, \\ 0 & \text{for } i > n - j. \end{cases}$$

The starting point in proving this is the following result of Milnor.

**Lemma 3.2** [[4](#), Corollary 6]  $\langle \chi P^n, \xi^R \rangle = \pm 1$  for all sequences  $R$  with  $|R| = n$ .

In the basis of  $\mathcal{P}$  dual to the monomial basis of  $\mathcal{P}_*$ , the element corresponding to  $\xi_1^i$  is  $P^i$ . Since the diagonal in  $\mathcal{P}_*$  is dual to the product in  $\mathcal{P}$ , it follows from [\(13\)](#) and [Lemma 3.2](#) that

$$\langle P^i \chi P^{n-i}, \xi^R \rangle = \begin{cases} \pm 1 & \text{for } i = \|R\|, \\ 0 & \text{for } i > \|R\|. \end{cases}$$

So we wish to construct sequences  $R_{n,j}$ , for  $\phi(n) \leq j \leq n$ , such that  $|R_{n,j}| = n$  and  $\|R_{n,j}\| = j$ . We deal first with the case  $j = \phi(n)$ .

**Proposition 3.3** *For any  $n \geq 0$  there is a sequence  $M = (m_1, m_2, \dots)$  such that*

- (1)  $|M| = n$ ,
- (2)  $0 \leq m_i \leq p$  for all  $i$ ,
- (3) if  $m_j = p$  then  $m_i = 0$  for all  $i < j$ .

For any such sequence,  $\|M\| = \phi(n)$ .

**Proof** Give the set of sequences of dimension  $n$  the right-lexicographic order. We claim that the maximal sequence satisfies the hypotheses.

Suppose that  $R = (r_1, r_2, \dots)$  does not satisfy the hypotheses. If  $r_1 > p$  then the sequence  $(r_1 - (p + 1), r_2 + 1, r_3, \dots)$  is larger. If  $r_j > p$ , with  $j > 1$ , then the sequence  $(r_1, \dots, r_{j-2}, r_{j-1} + p, r_j - (p + 1), r_{j+1} + 1, r_{k+2}, \dots)$  is larger. This proves (2). To prove (3), suppose that  $r_j = p$  with  $j > 1$ , and suppose that some earlier entry is nonzero. Let  $i = \min\{k : r_k > 0\}$ . If  $i = 1$ , then the sequence

$(r_1 - 1, r_2, \dots, r_{j-1}, 0, r_{j+1} + 1, r_{j+2}, \dots)$  is larger. If  $i > 1$ , then  $S$  with  $s_k = 0$  for  $k < i - 1$  and  $i \leq k \leq j$ ,  $s_{i-1} = p$ ,  $s_{j+1} = r_{j+1} + 1$ , and  $s_k = r_k$  for  $k > j + 1$ , is larger. Let  $M$  be a sequence satisfying (1)–(3), and write  $l = \|M\| - 1$ . To see that  $l = \phi(n) - 1$  we must show that

$$(14) \quad p(l + 1) - \alpha(l + 1) \geq (p - 1)n,$$

$$(15) \quad pl - \alpha(l) < (p - 1)n.$$

The excess  $e(R)$  is the sum of the entries in  $R$ , so that  $p\|R\| - e(R) = (p - 1)|R|$ . The  $p$ -adic representation of a number minimizes excess, so for any sequence  $R$  we have  $e(R) \geq \alpha(\|R\|)$  and hence  $p\|R\| - \alpha(\|R\|) \geq (p - 1)|R|$ : so (14) holds for any sequence.

To see that (15) holds for  $M$ , let  $j = \min\{i : m_i > 0\}$ , so that  $(p - 1)n = (p^j - 1)m_j + (p^{j+1} - 1)m_{j+1} + \dots$  and  $l + 1 = p^{j-1}m_j + p^j m_{j+1} + \dots$ . The hypotheses imply that  $l$  has  $p$ -adic expansion

$$(1 + \dots + p^{j-2})(p - 1) + p^{j-1}(m_j - 1) + p^j m_{j+1} + \dots,$$

so 
$$\alpha(l) = (j - 1)(p - 1) + (m_j - 1) + m_{j+1} + \dots$$

from which we deduce

$$pl - \alpha(l) = (p - 1)(n - j) < (p - 1)n.$$

This completes the proof of Proposition 3.3. □

**Corollary 3.4** *The function  $\phi(n)$  is weakly increasing.*

**Proof** Let  $M$  be a sequence satisfying the conditions of Proposition 3.3, and note that the sequence  $R = (1, 0, 0, \dots) + M$  has  $|R| = n + 1$  and  $\|R\| = \|M\| + 1 = \phi(n) + 1$ . If  $p$  does not occur in  $M$ , then  $R$  satisfies the hypotheses of the proposition (in degree  $n + 1$ ) and hence  $\phi(n) \leq \phi(n + 1)$ . If  $p$  does occur in  $M$ , then the moves described above will lead to a sequence  $M'$  satisfying the hypotheses. None of the moves decrease  $\|-\|$ , so  $\phi(n) \leq \phi(n + 1)$ . □

**Remark 3.5** Properties (1)–(3) of Proposition 3.3 in fact determine  $M$  uniquely.

**Proof of Proposition 3.1** Define  $R_{n,\phi(n)}$  to be a sequence  $M$  as in Proposition 3.3. Then inductively define

$$R_{n,j} = (1, 0, 0, \dots) + R_{n-1,j-1} \quad \text{for } \phi(n) < j \leq n.$$

This makes sense by monotonicity of  $\phi(n)$ , and the elements clearly satisfy  $|R_{n,j}| = n$  and  $\|R_{n,j}\| = j$ . This completes the proof. □

## References

- [1] **M G Barratt, H R Miller**, *On the anti-automorphism of the Steenrod algebra*, *Contemp. Math.* 12 (1982) 47–52
- [2] **M C Crabb, M D Crossley, J R Hubbuck**,  *$K$ -theory and the anti-automorphism of the Steenrod algebra*, *Proc. Amer. Math. Soc.* 124 (1996) 2275–2281 [MR1328343](#)
- [3] **D M Davis**, *The antiautomorphism of the Steenrod algebra*, *Proc. Amer. Math. Soc.* 44 (1974) 235–236 [MR0328934](#)
- [4] **J Milnor**, *The Steenrod algebra and its dual*, *Ann. of Math. (2)* 67 (1958) 150–171 [MR0099653](#)
- [5] **J H Silverman**, *Conjugation and excess in the Steenrod algebra*, *Proc. Amer. Math. Soc.* 119 (1993) 657–661 [MR1152292](#)
- [6] **P D Straffin, Jr**, *Identities for conjugation in the Steenrod algebra*, *Proc. Amer. Math. Soc.* 49 (1975) 253–255 [MR0380796](#)

Department of Mathematics, University of Connecticut  
Storrs CT 06269, USA

Department of Mathematics, Massachusetts Institute of Technology  
77 Massachusetts Ave, Cambridge MA 02139-4307, USA

[vince@math.uconn.edu](mailto:vince@math.uconn.edu), [hrm@math.mit.edu](mailto:hrm@math.mit.edu)

Received: 7 June 2011      Revised: 18 July 2011