

ANALYSIS & PDE

Volume 12

No. 5

2019

BETSY STOVALL

**SCALE-INVARIANT FOURIER RESTRICTION TO A HYPERBOLIC
SURFACE**

SCALE-INVARIANT FOURIER RESTRICTION TO A HYPERBOLIC SURFACE

BETSY STOVALL

This result sharpens the bilinear-to-linear deduction of Lee and Vargas for extension estimates on the hyperbolic paraboloid in \mathbb{R}^3 to the sharp line, leading to the first scale-invariant restriction estimates, beyond the Stein–Tomas range, for a hypersurface on which the principal curvatures have different signs.

1. Introduction

We consider the Fourier restriction/extension problem for the hyperbolic paraboloid

$$S := \{(\tau, \xi) \in \mathbb{R}^{1+2} : \tau = \xi_1 \xi_2\}.$$

We denote by \mathcal{E} the extension operator,

$$\mathcal{E}f(t, x) := \int_{\mathbb{R}^2} e^{i(t,x)(\xi_1 \xi_2, \xi)} f(\xi) d\xi. \tag{1-1}$$

For consistency of exponents, we will consider the problem of establishing $L^r \rightarrow L^{2s}$ extension estimates for \mathcal{E} , and we are primarily interested in the case when $r = s'$.

Lee [2006] and Vargas [2005] independently established an essentially optimal L^2 -based bilinear adjoint restriction estimate for S . This result states that if f and g are supported in 1×1 axis-parallel rectangles that are separated from one another by a distance 1 in the horizontal direction and 1 in the vertical direction, then

$$\|\mathcal{E}f\mathcal{E}g\|_s \lesssim \|f\|_2 \|g\|_2, \quad s > \frac{5}{3}. \tag{1-2}$$

This two-parameter separation of the tiles is both necessary and troublesome. On the one hand, necessity can be seen by considering the case when each of f_{\pm} is supported on a $\frac{1}{2}$ -neighborhood of $(\pm 1, 0)$. On the other hand, the separation leads to difficulty in deducing linear restriction estimates from the bilinear ones. Indeed, the natural analogue of the Whitney decomposition approach of [Tao, Vargas, and Vega 1998] produces a sum in two scales, length and width, rather than a single distance scale, leading to a loss of the scaling line in the distinct approaches of [Lee 2006] and [Vargas 2005].

The purpose of this note is to overcome this obstacle and recover the sharp line.

Theorem 1.1. *With \mathcal{E} as in (1-1), assume that the estimate*

$$\|\mathcal{E}f\mathcal{E}g\|_s \lesssim \|f\|_r \|g\|_r \tag{1-3}$$

MSC2010: 42B20.

Keywords: Fourier restriction, bilinear restriction, hyperbolic restriction.

holds for some $\frac{3}{2} < s < 2$ and $\frac{r}{2} < s < r'$, whenever f and g are supported on 1×1 , axis-parallel rectangles that are separated from one another by a distance 1 in both the horizontal and vertical directions. Then \mathcal{E} is of restricted strong type $(s', 2s)$, and consequently of strong type $(\tilde{s}', 2\tilde{s})$ for all $\tilde{s} > s$.

To put the hypothesis on s in context, we recall that for $s \leq \frac{3}{2}$, linear extension estimates are known to be impossible; that for $s > \frac{3}{2}$, $2s > s'$; and that for $s \geq 2$, linear extension estimates are already known, [Tomas 1975].

As is well known, a (local, linear) $L^{r_0} \rightarrow L^{2s_0}$ extension estimate for some $r_0 > s'_0$ allows us, by interpolation with the L^2 -based bilinear extension estimate (1-2), to establish the L^r -based bilinear extension estimate (1-3) for some $s > s_0$ and $\frac{r}{2} < s < r'$. Replacing s_0 with s is a loss (whose magnitude depends on the distance from (r_0^{-1}, s_0^{-1}) to the scaling line), but $r < s'$ is a gain in the sense that the corresponding linear extension estimate $\mathcal{E} : L^r \rightarrow L^{2s}$ is false.

Lee [2006] and Vargas [2005] independently used the bilinear extension estimate (1-2) to prove that

$$\|\mathcal{E}f\|_{2s} \lesssim \|f\|_{L^r} \tag{1-4}$$

for all $s > \frac{5}{3}$, $r > s'$, and f supported in the unit ball. Cho and Lee [2017] used the polynomial partitioning argument from [Guth 2016] to prove (1-4) for f supported in the unit ball and $2s = r > 3.25$; this was subsequently improved by Kim [2017] to the range $2s > 3.25$ and $r > s'$. Using these results and the discussion in the preceding paragraph, Theorem 1.1 immediately yields the following slight improvement on Kim's result.

Corollary 1.2. *For $2s > 3.25$, the extension operator \mathcal{E} is bounded from $L^{s'}$ to L^{2s} .*

To the author's knowledge, this is the first scalable restriction estimate for a negatively curved hypersurface, beyond the Stein–Tomas range ($s = 2$).

Terminology. A constant will be said to be admissible if it depends only on s, r . The inequality $A \lesssim B$ means that $A \leq CB$ for some implicit, admissible constant C , and implicit constants will be allowed to change from line to line. A dyadic interval is an interval of the form $[m2^{-n}, (m+1)2^{-n}]$ for some $m, n \in \mathbb{Z}$, and \mathcal{I}_n denotes the set of all dyadic intervals of length 2^{-n} . A tile is a product of two dyadic intervals, and $\mathcal{D}_{J,K}$ denotes the set of all $2^{-J} \times 2^{-K}$ tiles. We denote by π_1, π_2 the projections $\pi_j : \mathbb{R}^2 \rightarrow \mathbb{R}$, $\pi_j(x) = x_j$. We use \mathcal{H}^1 for the one-dimensional Hausdorff measure. Finally, we use \log to denote the base-2 logarithm.

Outline of proof. To prove our restricted strong-type estimate, it suffices to bound the extension of a characteristic function. Our starting point is the bilinear-to-linear deduction of [Vargas 2005], which shows that, under the hypotheses of Theorem 1.1, the extension of the characteristic function of a set Ω with roughly constant (vertical) fiber length obeys the scalable restriction estimate $\|\mathcal{E}\chi_\Omega\|_{2s} \lesssim |\Omega|^{\frac{1}{s'}}$. In [Vargas 2005], off-scaling estimates are obtained by subdividing a set Ω in the unit cube into subsets having constant fiber length. Off-scaling contributions from those subsets with very short fibers are small (because the sets themselves are small), and adding these amounts to summing a convergent geometric series.

We wish to remain on the sharp line, so we must be more careful. Our first step, taken in [Section 2](#), is to understand when Vargas’s constant fiber length estimate can be improved. To this end, we prove a dichotomy result: If Ω has constant fiber length, then either Ω is highly structured (more precisely, Ω is nearly a tile), or we have a better bound on the extension of χ_Ω . Roughly speaking, this reduces matters to controlling the extension of a union of tiles τ_k each having height 2^{-k} , which is the task of [Section 3](#). We can estimate

$$\|\mathcal{E}\chi_{\cup \tau_k}\|_{2s} \lesssim \left(\sum \|\mathcal{E}\chi_{\tau_k}\|_{2s}^{2s}\right)^{\frac{1}{2s}} + \text{off-diagonal terms},$$

where the off-diagonal terms involve products $\mathcal{E}\chi_{\tau_k}\mathcal{E}\chi_{\tau_{k'}}$, with $|k - k'|$ large. Boundedness of the main term follows from Vargas’s estimate and convexity ($2s > s'$). It remains to bound the off-diagonal terms, for which it suffices to prove a bilinear estimate with decay:

$$\|\mathcal{E}\chi_{\tau_k}\mathcal{E}\chi_{\tau_{k'}}\|_s \lesssim 2^{-c_0|k-k'|} \max\{|\tau_k|, |\tau_{k'}|\}^{\frac{1}{s'}},$$

and we prove this by combining the bilinear extension estimate for separated tiles with a further decomposition.

Of course, we have lied. In [Section 2](#), our dichotomy is not that a constant fiber length set Ω is either a tile or has zero extension, and so we still have remainder terms that must be summed. To address this, we argue more quantitatively than has been suggested above: Any constant fiber length set can be approximated by a union of tiles, where the number of tiles and tightness of the approximation depends on the sharpness of our estimate $\|\mathcal{E}\chi_\Omega\|_{2s} \lesssim |\Omega|^{\frac{1}{s'}}$; then we must bound extensions of sets $\bigcup_k \bigcup_{\tau \in \mathcal{T}_k} \tau$, where $\mathcal{T}_k \subseteq \mathcal{D}_{j(k),k}$ may be large (but fortunately, not too large).

2. An inverse problem related to Vargas’s linear estimate

To prove [Theorem 1.1](#), it suffices to prove that $\|\mathcal{E}\chi_\Omega\|_{2s} \lesssim |\Omega|^{\frac{1}{s'}}$ for all measurable sets Ω . By scaling, it suffices to consider Ω contained in the unit cube $[-1, 1]^2$. Vargas [\[2005\]](#) proved the following.

Theorem 2.1 [\[Vargas 2005\]](#). *For each $K \geq 0$, let*

$$\Omega(K) := \{\xi \in \Omega : \mathcal{H}^1(\pi_1^{-1}(\xi_1) \cap \Omega) \sim 2^{-K}\}. \tag{2-1}$$

Then under the hypotheses of [Theorem 1.1](#), for any measurable set $\Omega' \subseteq \Omega(K)$,

$$\|\mathcal{E}\chi_{\Omega'}\|_{2s} \lesssim |\Omega(K)|^{\frac{1}{s'}}. \tag{2-2}$$

This version differs slightly from the one stated in [\[Vargas 2005\]](#), but it follows from the same proof. In proving the next proposition, we will review Vargas’s argument, so the reader may verify the above-stated version below.

Our first step is to solve an inverse problem: Characterize those sets $\Omega = \Omega(K)$ for which the inequality in [\(2-2\)](#) can be reversed.

Proposition 2.2. *Assume that the hypotheses of [Theorem 1.1](#) hold. Let $\Omega \subseteq [-1, 1]^2$ be a measurable set, and assume that $\Omega = \Omega(K)$ for some integer $K \geq 0$. Choose a nonnegative integer J such that*

$|\pi_1(\Omega)| \sim 2^{-J}$, and let $\varepsilon \lesssim 1$ denote the smallest dyadic number such that

$$\|\mathcal{E}\chi_{\Omega'}\|_{2s} \leq \varepsilon|\Omega|^{\frac{1}{s'}}$$

for every measurable $\Omega' \subseteq \Omega$. Then $\Omega = \bigcup_{0 < \delta \leq \varepsilon} \Omega_\delta$, with the union taken over dyadic δ . For each δ , $\Omega_\delta \subseteq \bigcup_{\tau \in \mathcal{T}_\delta} \tau$, where $\mathcal{T}_\delta \subseteq \mathcal{D}_{J,K}$ has cardinality at most $O(\delta^{-C})$, with C an admissible constant. For each subset $\Omega' \subseteq \Omega_\delta$, $\|\mathcal{E}\chi_{\Omega'}\|_{2s} \lesssim \delta|\Omega|^{\frac{1}{s'}}$.

Proof of Proposition 2.2. It suffices to produce a union that contains almost every point of Ω , as a set of measure zero makes no contribution to the extension. Our decomposition will be done in three stages. Our first decomposition will be of Ω into sets Ω_η^1 , with $\pi_1(\Omega_\eta^1)$ nearly an interval, $I \in \mathcal{I}_J$. Our second decomposition will be of Ω_η^1 into sets $\Omega_{\eta,\rho}^2$, $\rho \leq \eta$, each of which is nearly a product of I with a set of measure 2^{-K} . Our third decomposition will be of $\Omega_{\eta,\rho}^2$ into sets $\Omega_{\eta,\rho,\delta}^3$, $\delta \leq \rho$, each of which is nearly a product of I with an interval in \mathcal{I}_K . The product of two dyadic intervals is a tile, so we take $\Omega_\delta := \bigcup_{\rho \geq \delta} \bigcup_{\eta \geq \rho} \Omega_{\eta,\rho,\delta}^3$; the $(\log \delta^{-1})^2$ factor that arises from taking this union is harmless.

Let $S := \pi_1(\Omega)$. We know that $|S| \sim 2^{-J}$ and that $S \subseteq [-1, 1]$. Let $\xi_1 \in S$, and for each $0 < \eta < \varepsilon$, let $I_\eta(\xi_1)$ be the maximal dyadic interval $I \ni \xi_1$ satisfying $|I \cap S| \geq \eta^C |I|$, if such an interval exists. We record that $|I_\eta(\xi_1)| \leq \eta^{-C} 2^{-J}$, and if ξ_1 is a Lebesgue point of S , then $|I_\eta(\xi_1)| > 0$. Let

$$T_\eta := \{\xi_1 \in S : |I_\eta(\xi_1)| \geq \eta^C 2^{-J}\},$$

and let $S_\varepsilon := T_\varepsilon$, $S_\eta := T_\eta \setminus T_{2\eta}$ for dyadic $0 < \eta < \varepsilon$. Then a.e. (indeed, every Lebesgue) point of S is contained in a unique S_η . We set $\Omega_\eta^1 := \Omega \cap \pi_1^{-1}(S_\eta)$.

Lemma 2.3. *For each $0 < \eta \leq \varepsilon$, S_η is contained in a union of $O(\eta^{-3C})$ dyadic intervals $I \in \mathcal{I}_J$, and for each $\eta < \varepsilon$ and each subset $\Omega' \subseteq \Omega_\eta^1$,*

$$\|\mathcal{E}\chi_{\Omega'}\|_{2s} \lesssim \eta^2 |\Omega|^{\frac{1}{s'}}. \tag{2-3}$$

Proof of Lemma 2.3. By construction, S_η is covered by dyadic intervals I of length $|I| \geq \eta^C |S|$, in which S has density $|I \cap S| \geq \eta^C |I|$. The density of each such interval in S is $|I \cap S| \geq \eta^{2C} |S|$, and so the collection of maximal (hence pairwise disjoint) dyadic intervals with these properties has cardinality at most η^{-2C} . Moreover, from the density estimate, we see that $|I| \leq \eta^{-C} 2^{-J}$, so these intervals are covered by a total of η^{-3C} intervals in \mathcal{I}_J .

To establish (2-3), we will optimize Vargas’s proof of Theorem 2.1. Performing a Whitney decomposition in each variable ξ_1, ξ_2 separately and applying the almost orthogonality lemma from [Tao, Vargas, and Vega 1998] (for which it is important that $s \leq 2$),

$$\|\mathcal{E}\chi_{\Omega'}\|_{2s}^2 \lesssim \sum_{k,j} \left(\sum_{\tau \sim \tau' \in \mathcal{D}_{j,k}} \|\mathcal{E}\chi_{\Omega' \cap \tau} \mathcal{E}\chi_{\Omega' \cap \tau'}\|_s^s \right)^{\frac{1}{s}},$$

where we say that $\tau \sim \tau'$ if τ and τ' are 2^{-j} separated in the horizontal direction and 2^{-k} separated in the vertical direction.

By rescaling our hypothesis, (1-3), for f, g supported on tiles in $\mathcal{D}_{j,k}$ that are separated by a distance 2^{-k} in the vertical direction and 2^{-j} in the horizontal direction,

$$\|\mathcal{E}f\mathcal{E}g\|_s \lesssim 2^{(j+k)(\frac{2}{s}+\frac{2}{r}-2)} \|f\|_r \|g\|_r. \tag{2-4}$$

Thus

$$\begin{aligned} \|\mathcal{E}\chi_{\Omega'}\|_{2s}^2 &\lesssim \sum_{k,j} 2^{(j+k)(\frac{2}{s}+\frac{2}{r}-2)} \left(\sum_{\tau \in \mathcal{D}_{j,k}} |\Omega' \cap \tau| \frac{2^s}{r} \right)^{\frac{1}{s}} \\ &\lesssim \sum_{k,j} 2^{(j+k)(\frac{2}{s}+\frac{2}{r}-2)} \max_{\tau \in \mathcal{D}_{j,k}} |\Omega' \cap \tau|^{\frac{2}{r}-\frac{1}{s}} |\Omega'|^{\frac{1}{s}}. \end{aligned} \tag{2-5}$$

Our hypotheses on r, s imply that all exponents in the above sum are positive. To bound this double sum, Vargas used the inequality

$$|\Omega' \cap \tau| \lesssim \min\{2^{-j}, 2^{-J}\} \min\{2^{-k}, 2^{-K}\}. \tag{2-6}$$

The definition of Ω_η^1 will allow us to improve on this bound.

For $I_j \in \mathcal{I}_j$, we trivially have $|I_j \cap S_\eta| \leq \min\{|I_j|, |S_\eta|\} \leq \min\{2^{-j}, 2^{-J}\}$, but when $|j - J| < \frac{C}{4} \log \eta^{-1}$, we can do rather better. Suppose that $|j - J| \leq \frac{C}{4} \log \eta^{-1}$. Since

$$|I_j| = 2^{-j} \geq \eta^{\frac{C}{4}} 2^{-J} \geq (2\eta)^C 2^{-J}$$

(provided η is sufficiently small), $I_j \cap S_\eta \neq \emptyset$ implies that $I_j \cap S_\eta \not\subseteq T_{2\eta}$, whence

$$|I_j \cap S_\eta| \leq |I_j \cap S| \leq (2\eta)^C |I_j| = (2\eta)^C 2^{-j} \lesssim \eta^{\frac{3C}{4}} \min\{2^{-j}, 2^{-J}\},$$

where for the last inequality, we used $2^{-j} \leq \eta^{-\frac{C}{4}} 2^{-J}$.

Inserting this gain and $|\Omega' \cap (I_j \times I_k)| \leq |S_\eta \cap I_j| \min\{2^{-k}, 2^{-K}\}$ into (2-5), and summing the resulting geometric series gives

$$\|\mathcal{E}\chi_{\Omega'}\|_{2s} \lesssim \eta^{C'} 2^{-(J+K)(1-\frac{3}{2s})} |\Omega'|^{\frac{1}{2s}} \lesssim \eta^{C'} |\Omega'|^{\frac{1}{s'}}$$

for $C' > 0$ some admissible constant dictated by C, r, s ; we can reverse engineer C so that $C' = 2$. \square

We now turn to our second decomposition. Although $\pi_1(\Omega_\eta^1)$ may be (roughly) thought of as a union of a small number of intervals, an individual horizontal slice $\pi_2^{-1}(\xi_2) \cap \Omega_\eta^1$ might be much smaller. Our next step is to decompose into sets where the size of a nonempty slice is roughly comparable to the size of the projection of the whole. (Sets with this property are nearly products.)

Fix $0 < \eta \leq \varepsilon$. For dyadic $0 < \rho \leq \eta$, we define

$$V_\rho = \{\xi_2 \in \pi_2(\Omega_\eta^1) : \mathcal{H}^1(\pi_2^{-1}(\xi_2) \cap \Omega_\eta^1) \geq \rho^C 2^{-J}\},$$

and set $U_\eta := V_\eta$, $U_\rho := V_\rho \setminus V_{2\rho}$ for $\rho < \eta$. We define $\Omega_{\eta,\rho}^2 := \pi_2^{-1}(U_\rho) \cap \Omega_\eta^1$.

Lemma 2.4. *For each $0 < \rho < \eta \leq \varepsilon$, and each subset $\Omega' \subseteq \Omega_{\eta,\rho}^2$, we have $\|\mathcal{E}\chi_{\Omega'}\|_{2s} \lesssim \rho^2 |\Omega'|^{\frac{1}{s'}}$.*

Proof of Lemma 2.4. Let $\tau \in \mathcal{D}_{j,k}$ and $\Omega' \subseteq \Omega_{\eta,\rho}^2$. Then $\tau \cap \Omega'$ has vertical and horizontal fiber lengths at most

$$\int \chi_{\tau \cap \Omega'}(\xi_1, \xi_2) d\xi_2 \lesssim \min\{2^{-K}, 2^{-k}\}, \quad \int \chi_{\tau \cap \Omega'}(\xi_1, \xi_2) d\xi_1 \lesssim \min\{\rho^C 2^{-J}, 2^{-j}\},$$

respectively, and projections of size at most

$$|\pi_1(\tau \cap \Omega')| \lesssim \min\{2^{-J}, 2^{-j}\}, \quad |\pi_2(\tau \cap \Omega')| \lesssim 2^{-k}.$$

By Fubini, we can bound $|\tau \cap \Omega'|$ by the measure of the projection times the maximum fiber length, so

$$|\tau \cap \Omega'| \lesssim \min\{2^{-(J+K)}, 2^{-(j+K)}, 2^{-(j+k)}, \rho^C 2^{-(J+k)}\}. \tag{2-7}$$

To utilize (2-7), we let $C' = \frac{C}{2}$ and subdivide $\mathbb{Z}^2 = R_1 \cup R_2 \cup R_3 \cup R_4$, where

$$\begin{aligned} R_1 &:= \{(j, k) : J - C' \log \rho^{-1} \geq j, K \geq k\} \cup \{(j, k) : J \geq j, K - C' \log \rho^{-1} \geq k\}, \\ R_2 &:= \{(j, k) : j \geq J + C' \log \rho^{-1}, K \geq k\} \cup \{(j, k) : j \geq J, K - C' \log \rho^{-1} \geq k\}, \\ R_3 &:= \{(j, k) : j \geq J + C' \log \rho^{-1}, k \geq K\} \cup \{(j, k) : j \geq J, k \geq K + C' \log \rho^{-1}\}, \\ R_4 &:= \{(j, k) : J + C' \log \rho^{-1} \geq j, k + C' \log \rho^{-1} \geq K\}. \end{aligned}$$

Now we insert (2-7) into (2-5) to obtain

$$\begin{aligned} \|\mathcal{E} \chi_{\Omega'}\|_{2_s}^2 &\lesssim \sum_{R_1} 2^{(j+k)(\frac{2}{s} + \frac{2}{r} - 2)} 2^{-\frac{2(J+K)}{r}} + \sum_{R_2} 2^{-j(2 - \frac{3}{s})} 2^{k(\frac{2}{s} + \frac{2}{r} - 2)} 2^{-\frac{J}{s}} 2^{-\frac{2K}{r}} \\ &\quad + \sum_{R_3} 2^{-(j+k)(2 - \frac{3}{s})} 2^{-\frac{J+K}{s}} + \rho^{C(\frac{2}{r} - \frac{1}{s})} \sum_{R_4} 2^{-k(2 - \frac{3}{s})} 2^{j(\frac{2}{s} + \frac{2}{r} - 2)} 2^{-\frac{2J}{r}} 2^{-\frac{K}{s}}. \end{aligned}$$

As $\frac{2}{s} + \frac{2}{r} - 2$ and $2 - \frac{3}{s}$ are both positive, it is a simple matter to sum each of these terms, obtaining

$$\|\mathcal{E} \chi_{\Omega'}\|_{2_s}^2 \lesssim (\rho^{(\frac{2}{s} + \frac{2}{r} - 2)C'} + \rho^{(2 - \frac{3}{s})C'} + \rho^{C(\frac{2}{r} - \frac{1}{s}) - C'(2 - \frac{3}{s}) - C'(\frac{2}{s} + \frac{2}{r} - 2)}) 2^{-\frac{2(J+K)}{s'}}.$$

Since $\frac{2}{r} - \frac{1}{s} > 0$ and $|\Omega| \sim 2^{-(J+K)}$, we obtain the bound claimed in the lemma by choosing C sufficiently large. □

Now we turn to our third decomposition. A single $\Omega_{\eta,\rho}^2$ is “nearly” a product, but $\pi_2(\Omega_{\eta,\rho}^2)$ might be far from an interval. However, we may simply perform the first decomposition, with the roles of the coordinate indices reversed. Indeed, our sets satisfy analogous hypotheses to those in Lemma 2.3 (i.e., the hypotheses of Proposition 2.2 with the indices reversed) when $\rho < \eta$, because $|\mathcal{H}^1(\pi_2^{-1}(\xi_2) \cap \Omega_{\eta,\rho}^2)| \sim \rho^C 2^{-J}$ for all $\xi_2 \in \pi_2(\Omega_{\eta,\rho}^2)$; when $\rho = \eta$, we may abuse notation slightly by decomposing $\Omega_{\eta,\eta}^2$ into $\log \eta^{-1}$ sets $\Omega_{\eta,\rho}^2$ with the same property.

We complete the proof of Proposition 2.2 by taking unions as described at the outset. The factors of η^2 and ρ^2 in Lemmas 2.3 and 2.4 (and the factor of δ^2 in the analogue for $\Omega_{\eta,\rho,\delta}^3$) mean that the resulting factor of $(\log \delta^{-1})^2$ is indeed harmless. □

3. Extensions of characteristic functions of near tiles

We recall the definition (2-1) of $\Omega(K)$. For each K , we define $J(K)$ to be an integer such that $|\Omega(K)| \sim 2^{-J(K)-K}$. Let $\mathcal{K}(\varepsilon)$ denote the collection of all $K \in \mathbb{Z}_{\geq 0}$ for which ε is the smallest dyadic number such that $\|\mathcal{E}\chi_{\Omega'}\|_{2s} \leq \varepsilon |\Omega(K)|^{\frac{1}{s'}}$ holds for all $\Omega' \subseteq \Omega(K)$. Then Proposition 2.2 gives us a decomposition $\Omega(K) = \bigcup_{0 < \delta \leq \varepsilon} \Omega_\delta(K)$, where for each δ , we have $\Omega_\delta(K) \subseteq \bigcup_{\tau \in \mathcal{T}_\delta(K)} \tau$ for some $\mathcal{T}_\delta(K) \subseteq \mathcal{D}_{J(K), K}$ of cardinality $\#\mathcal{T}_\delta(K) \lesssim \delta^{-C}$.

Lemma 3.1. *For $0 < \delta \leq \varepsilon$, under the hypotheses of Theorem 1.1,*

$$\left\| \sum_{K \in \mathcal{K}(\varepsilon)} \mathcal{E}\chi_{\Omega_\delta(K)} \right\|_{2s}^{2s} \lesssim (\log \delta^{-1})^{4s} \sum_{K \in \mathcal{K}(\varepsilon)} \|\mathcal{E}\chi_{\Omega_\delta(K)}\|_{2s}^{2s} + \delta |\Omega|^{\frac{2s}{s'}}.$$

Proof of Lemma 3.1. To prove the lemma, it suffices to prove

$$\left\| \sum_{K \in \mathcal{K}} \mathcal{E}\chi_{\Omega_\delta(K)} \right\|_{2s}^{2s} \lesssim \sum_{K \in \mathcal{K}} \|\mathcal{E}\chi_{\Omega_\delta(K)}\|_{2s}^{2s} + \delta^2 |\Omega|^{\frac{2s}{s'}},$$

with $\mathcal{K} \subseteq \mathcal{K}(\varepsilon)$ chosen so that \mathcal{K} and $J(\mathcal{K})$ are both $A \log \delta^{-1}$ -separated, with A a sufficiently large admissible constant. (A will be much larger than the constant C in Proposition 2.2.) Since $s < 2$, the triangle inequality gives

$$\left\| \sum_{K \in \mathcal{K}} \mathcal{E}\chi_{\Omega_\delta(K)} \right\|_{2s}^{2s} = \int \left| \sum_{K \in \mathcal{K}^4} \prod_{i=1}^4 \mathcal{E}\chi_{\Omega_\delta(K_i)} \right|^{\frac{s}{2}} \lesssim \sum_{K \in \mathcal{K}} \|\mathcal{E}\chi_{\Omega_\delta(K)}\|_{2s}^{2s} + \sum' \left\| \prod_{i=1}^4 \mathcal{E}\chi_{\Omega_\delta(K_i)} \right\|_{\frac{s}{2}}^{\frac{s}{2}},$$

where \sum' indicates a sum taken on quadruples $\mathbf{K} = (K_1, K_2, K_3, K_4) \in \mathcal{K}^4$, with at least two entries distinct. The following lemma will be useful in controlling this sum.

Lemma 3.2. *If $K, K' \in \mathcal{K}$, and $J := J(K), J' := J(K')$, then*

$$\|\mathcal{E}\chi_{\Omega_\delta(K)} \mathcal{E}\chi_{\Omega_\delta(K')}\|_s \lesssim 2^{-c_0|K-K'|} \max\{|\Omega(K)|, |\Omega(K')|\}^{\frac{2}{s'}} \tag{3-1}$$

for some admissible constant $c_0 > 0$.

Proof of Lemma 3.2. Inequality (3-1) is an immediate consequence of Cauchy–Schwarz and (2-2) whenever

$$|\Omega(K)|^{\frac{1}{s'}} |\Omega(K')|^{\frac{1}{s'}} \lesssim 2^{-\frac{|K-K'|}{s'}} \max\{|\Omega(K)|, |\Omega(K')|\}^{\frac{2}{s'}}.$$

This inequality holds whenever $K = K', J = J', J < J'$ and $K < K'$, or $J > J'$ and $K > K'$.

By symmetry, this leaves us to prove (3-1) when $K < K'$ and $J > J'$. By Proposition 2.2 and the separation condition on \mathcal{K} , it suffices to prove that

$$\|\mathcal{E}\chi_{\tau \cap \Omega_\delta(K)} \mathcal{E}\chi_{\tau' \cap \Omega_\delta(K')}\|_s \lesssim \delta^{-C} 2^{-c_0|K-K'|} |\Omega(K)|^{\frac{1}{s'}} |\Omega(K')|^{\frac{1}{s'}} \tag{3-2}$$

for tiles $\tau \in \mathcal{T}_\delta(K), \tau' \in \mathcal{T}_\delta(K')$.

Note that our conditions on J, J', K, K' mean that τ is taller than τ' , and τ' is wider than τ . By translating, we may assume that the y -axis forms the center line of τ and that the x -axis forms the center line of τ' . Now our tiles are contained in $[-2, 2]^2$, and we decompose:

$$\tau = \bigcup_{k=0}^{K'} \tau_k, \quad \tau' = \bigcup_{j=0}^J \tau'_j,$$

where

$$\tau_k = \begin{cases} \tau \cap \{\xi : |\xi_2| \sim 2^{-k}\}, & k < K', \\ \tau \cap \{\xi : |\xi_2| \lesssim 2^{-K'}\}, & k = K' \end{cases} \quad \text{and} \quad \tau'_j = \begin{cases} \tau' \cap \{\xi : |\xi_1| \sim 2^{-j}\}, & j < J, \\ \tau' \cap \{\xi : |\xi_1| \lesssim 2^{-J}\}, & j = J. \end{cases}$$

By the (2-parameter) Littlewood–Paley square function estimate (the two-parameter version can be proved using Khintchine’s inequality), the fact that $s < 2$, and the triangle inequality,

$$\begin{aligned} \|\mathcal{E}\chi_{\tau \cap \Omega_\delta(K)} \mathcal{E}\chi_{\tau' \cap \Omega_\delta(K')}\|_s^s &\lesssim \int \left(\sum_{k=0}^{K'} \sum_{j=1}^J |\mathcal{E}\chi_{\tau_k \cap \Omega(K)} \mathcal{E}\chi_{\tau'_j \cap \Omega(K')}|^2 \right)^{\frac{s}{2}} \\ &\lesssim \sum_{k=0}^{K'} \sum_{j=0}^J \|\mathcal{E}\chi_{\tau_k \cap \Omega(K)} \mathcal{E}\chi_{\tau'_j \cap \Omega(K')}\|_s^s. \end{aligned} \tag{3-3}$$

We begin with the sum over those terms with $k = K'$. By Cauchy–Schwarz and (2-2),

$$\sum_{j=0}^J \|\mathcal{E}\chi_{\tau_{K'} \cap \Omega(K)} \mathcal{E}\chi_{\tau'_j \cap \Omega(K')}\|_s^s \lesssim \sum_{j=0}^J \|\mathcal{E}\chi_{\tau_{K'} \cap \Omega(K)}\|_{2s}^s \|\mathcal{E}\chi_{\tau'_j \cap \Omega(K')}\|_{2s}^s \lesssim \sum_{j=0}^J |\tau_{K'}|_{\frac{s}{s'}}^{\frac{s}{s'}} |\tau'_j|_{\frac{s}{s'}}^{\frac{s}{s'}}.$$

Because of the way the τ'_j were defined, we have at most two nonempty τ'_j with $j \leq J'$. This, combined with the bound $|\tau'_j| \leq \min\{2^{-(j-J')}, 1\}|\tau'|$ gives $\sum_j |\tau'_j|_{\frac{s}{s'}}^{\frac{s}{s'}} \lesssim |\tau'|_{\frac{s}{s'}}^{\frac{s}{s'}}$ (despite the fact that $s < s'$). Since $|\tau_{K'}| \sim 2^{-(K'-K)}|\tau|$, $|\tau| \sim |\Omega(K)|$, and $|\tau'| \sim |\Omega(K')|$,

$$\sum_{j=0}^J \|\mathcal{E}\chi_{\tau_{K'} \cap \Omega(K)} \mathcal{E}\chi_{\tau'_j \cap \Omega(K')}\|_s^s \lesssim 2^{-(K'-K)\frac{s}{s'}} |\Omega(K)|_{\frac{s}{s'}}^{\frac{s}{s'}} |\Omega(K')|_{\frac{s}{s'}}^{\frac{s}{s'}}.$$

In the case $j = J$, a similar argument implies that

$$\sum_{k=0}^{K'} \|\mathcal{E}\chi_{\tau_k \cap \Omega(K)} \mathcal{E}\chi_{\tau'_J \cap \Omega(K')}\|_s^s \lesssim 2^{-(J-J')\frac{s}{s'}} |\Omega(K)|_{\frac{s}{s'}}^{\frac{s}{s'}} |\Omega(K')|_{\frac{s}{s'}}^{\frac{s}{s'}} \sim 2^{-(K'-K)\frac{s}{s'}} |\Omega(K)|_{\frac{s}{s'}}^{\frac{2s}{s'}}.$$

In the cases $k < K'$ and $j < J$, we have a gain, due to our bilinear extension estimate. If $k < K'$ and $j < J$, then τ_k is a (subset of four) tile(s) in $\mathcal{D}_{J, \max\{k, K\}}$, τ_j is a (subset of four) tile(s) in $\mathcal{D}_{\max\{j, J'\}, K'}$, and these tiles are separated by a distance 2^{-k} in the vertical direction 2^{-j} in the horizontal direction. These tiles are thus contained in separated tiles in $\mathcal{D}_{j,k}$, so by (2-4),

$$\|\mathcal{E}\chi_{\tau_k \cap \Omega(K)} \mathcal{E}\chi_{\tau'_j \cap \Omega(K')}\|_s \lesssim 2^{(j+k)(\frac{2}{s} + \frac{2}{r} - 2)} |\tau_k \cap \Omega(K)|_{\frac{1}{r}} |\tau'_j \cap \Omega(K')|_{\frac{1}{s'}}.$$

From our observation above that we have at most two values of j (resp. k) in our sum with $j \leq J'$ (resp. $k \leq K$), our assumption that $r < s'$ gives

$$\begin{aligned} \sum_{j=0}^J \sum_{k=0}^{K'} 2^{(j+k)(2+\frac{2s}{r}-2s)} |\tau_k \cap \Omega(K)|^{\frac{s}{r}} |\tau'_j \cap \Omega(K')|^{\frac{s}{r}} \\ \leq \sum_{j=0}^J \sum_{k=0}^{K'} 2^{(j+k)(2+\frac{2s}{r}-2s)} |\tau_k|^{\frac{s}{r}} |\tau'_j|^{\frac{s}{r}} \\ \lesssim 2^{(J'+K)(2+\frac{2s}{r}-2s)} |\tau|^{\frac{s}{r}} |\tau'|^{\frac{s}{r}} \sim \delta^{-C} 2^{(J'+K)(2+\frac{2s}{r}-2s)} |\Omega(K)|^{\frac{s}{r}} |\Omega(K')|^{\frac{s}{r}} \\ \lesssim \delta^{-C} 2^{(J-J'+K'-K)(1+\frac{s}{r}-s)} |\Omega(K)|^{\frac{s}{s'}} |\Omega(K')|^{\frac{s}{s'}}, \end{aligned}$$

which, by (3-3) and $\frac{1}{s} + \frac{1}{r} - 1 > 0$, is stronger than (3-2). □

We return to the proof of Lemma 3.1.

Let $K_1, K_2, K_3, K_4 \in \mathcal{K}$, not all equal. Rearranging indices if needed, we may assume that $N_1 := K_1 + J(K_1)$ is minimal among all $N_i := K_i + J(K_i)$ and that $|K_1 - K_4| \geq \frac{1}{2}|K_i - K_j|$ for all i, j . Thus $|\Omega(K_1)|$ is maximal. By Hölder's inequality and Lemma 3.2,

$$\left\| \prod_{i=1}^4 \mathcal{E} \chi_{\Omega_\delta(K_i)} \right\|_{\frac{s}{2}} \lesssim 2^{-c_0|K_1-K_4|} |\Omega(K_1)|^{\frac{4}{s'}}.$$

Therefore

$$\sum' \left\| \prod_{i=1}^4 \mathcal{E} \chi_{\Omega_\delta(K_i)} \right\|_{\frac{s}{2}}^{\frac{s}{2}} \lesssim \sum_{K_1 \in \mathcal{K}} \sum_{K_1 \neq K_4 \in \mathcal{K}} |K_4 - K_1|^2 2^{-c_0|K_4-K_1|} |\Omega(K_1)|^{\frac{2s}{s'}}.$$

Because $2s > s'$ and \mathcal{K} is $A \log \delta^{-1}$ -separated for some very large A , this error term is bounded by $\delta^C |\Omega|^{\frac{2s}{s'}}$. □

Proof of Theorem 1.1. We decompose Ω by fiber length as in (2-1), $\Omega = \bigcup \Omega(K)$, then decompose the fiber lengths according to the exactness of Vargas's estimate as at the beginning of Section 3, $\mathbb{Z}_{\geq 0} = \bigcup_{0 \leq \varepsilon \leq 1} \mathcal{K}(\varepsilon)$, and finally apply the decomposition in Proposition 2.2, $\Omega(K) = \bigcup_{0 < \delta \leq \varepsilon} \Omega_\delta(K)$. By the triangle inequality,

$$\|\mathcal{E} \chi_\Omega\|_{2s} \leq \sum_{0 < \varepsilon \leq 1} \sum_{0 < \delta \leq \varepsilon} \left\| \sum_{K \in \mathcal{K}(\varepsilon)} \mathcal{E} \chi_{\Omega_\delta(K)} \right\|_{2s}.$$

Thus by Lemma 3.1 and Proposition 2.2,

$$\begin{aligned} \|\mathcal{E} \chi_\Omega\|_{2s} &\lesssim \sum_{0 < \varepsilon \leq 1} \sum_{0 < \delta \leq \varepsilon} \left\{ (\log \delta^{-1})^{4s} \sum_{K \in \mathcal{K}(\varepsilon)} \|\mathcal{E} \chi_{\Omega_\delta(K)}\|_{2s}^{2s} + \delta |\Omega|^{\frac{2s}{s'}} \right\}^{\frac{1}{2s}} \\ &\lesssim \left\{ \sum_{0 < \varepsilon \leq 1} \sum_{0 < \delta \leq \varepsilon} (\log \delta^{-1})^2 \left(\sum_{K \in \mathcal{K}(\varepsilon)} \delta^{2s} |\Omega(K)|^{\frac{2s}{s'}} \right)^{\frac{1}{2s}} \right\} + |\Omega|^{\frac{1}{s'}}. \end{aligned}$$

Since $2s > s'$ and the $\Omega(K)$ are disjoint, we may use the triangle inequality for $\ell^{\frac{2s}{s'}}$ to sum the volumes of the $\Omega(K)$ in the preceding, and, finally, we sum a geometric series to obtain

$$\|\mathcal{E}\chi_{\Omega}\|_{2s} \lesssim \sum_{0 < \varepsilon \lesssim 1} \sum_{0 < \delta \leq \varepsilon} (\log \delta^{-1})^2 \delta |\Omega|^{\frac{1}{s'}} \lesssim |\Omega|^{\frac{1}{s'}}. \quad \square$$

Acknowledgements

This work has been supported in part by a grant from the National Science Foundation (DMS-1600458), and was carried out in part while the author was in residence at the Mathematical Sciences Research Institute (MSRI) in Berkeley, California, during the Spring of 2017, a visit that was supported in part by a National Science Foundation grant to MSRI (DMS-1440140). The author would like to thank Sanghyuk Lee, Andreas Seeger, and Ana Vargas, from whom she learned of this problem and some of its history. She would also like to thank the anonymous referees and Benjamin Bruce, as the manuscript benefited greatly from their detailed comments and suggestions.

References

- [Cho and Lee 2017] C.-H. Cho and J. Lee, “Improved restriction estimate for hyperbolic surfaces in \mathbb{R}^3 ”, *J. Funct. Anal.* **273**:3 (2017), 917–945. [MR](#) [Zbl](#)
- [Guth 2016] L. Guth, “A restriction estimate using polynomial partitioning”, *J. Amer. Math. Soc.* **29**:2 (2016), 371–413. [MR](#) [Zbl](#)
- [Kim 2017] J. Kim, “Some remarks on Fourier restriction estimates”, preprint, 2017. [arXiv](#)
- [Lee 2006] S. Lee, “Bilinear restriction estimates for surfaces with curvatures of different signs”, *Trans. Amer. Math. Soc.* **358**:8 (2006), 3511–3533. [MR](#) [Zbl](#)
- [Tao, Vargas, and Vega 1998] T. Tao, A. Vargas, and L. Vega, “A bilinear approach to the restriction and Kakeya conjectures”, *J. Amer. Math. Soc.* **11**:4 (1998), 967–1000. [MR](#) [Zbl](#)
- [Tomas 1975] P. A. Tomas, “A restriction theorem for the Fourier transform”, *Bull. Amer. Math. Soc.* **81** (1975), 477–478. [MR](#) [Zbl](#)
- [Vargas 2005] A. Vargas, “Restriction theorems for a surface with negative curvature”, *Math. Z.* **249**:1 (2005), 97–111. [MR](#) [Zbl](#)

Received 8 Sep 2017. Revised 6 Aug 2018. Accepted 17 Sep 2018.

BETSY STOVALL: stovall@math.wisc.edu

Department of Mathematics, University of Wisconsin, Madison, WI, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Alessio Figalli	ETH Zurich, Switzerland alessio.figalli@math.ethz.ch	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbb@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zvorski@math.berkeley.edu

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2019 is US \$310/year for the electronic version, and \$520/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 12 No. 5 2019

On the Luzin N -property and the uncertainty principle for Sobolev mappings ADELE FERONE, MIKHAIL V. KOROBKOV and ALBA ROVIELLO	1149
Unstable normalized standing waves for the space periodic NLS NILS ACKERMANN and TOBIAS WETH	1177
Scale-invariant Fourier restriction to a hyperbolic surface BETSY STOVALL	1215
Steady three-dimensional rotational flows: an approach via two stream functions and Nash–Moser iteration BORIS BUFFONI and ERIK WAHLÉN	1225
Sparse bounds for the discrete cubic Hilbert transform AMALIA CULIUC, ROBERT KESLER and MICHAEL T. LACEY	1259
On the dimension and smoothness of radial projections TUOMAS ORPONEN	1273
Cartan subalgebras of tensor products of free quantum group factors with arbitrary factors YUSUKE ISONO	1295
Commutators of multiparameter flag singular integrals and applications XUAN THINH DUONG, JI LI, YUMENG OU, JILL PIPHER and BRETT D. WICK	1325
Rokhlin dimension: absorption of model actions GÁBOR SZABÓ	1357