

Desingularization of G_2 manifolds with isolated conical singularities

SPIRO KARIGIANNIS

We present a method to desingularize a compact G_2 manifold M with isolated conical singularities by cutting out a neighbourhood of each singular point x_i and gluing in an asymptotically conical G_2 manifold N_i . Controlling the error on the overlap gluing region enables us to use a result of Joyce to conclude that the resulting compact smooth 7-manifold \tilde{M} admits a torsion-free G_2 structure, with full G_2 holonomy.

There are topological obstructions for this procedure to work, which arise from the degree 3 and degree 4 cohomology of the asymptotically conical G_2 manifolds N_i which are glued in at each conical singularity. When a certain necessary topological condition on the manifold M with isolated conical singularities is satisfied, we can introduce correction terms to the gluing procedure to ensure that it still works. In the case of degree 4 obstructions, these correction terms are trivial to construct, but in the case of degree 3 obstructions we need to solve an elliptic equation on a noncompact manifold. For this we use the Lockhart–McOwen theory of weighted Sobolev spaces on manifolds with ends. This theory is also used to obtain a good asymptotic expansion of the G_2 structure on an asymptotically conical G_2 manifold N under an appropriate gauge-fixing condition, which is required to make the gluing procedure work.

[53C29](#); [58J05](#)

1 Introduction

This is the first of two papers in which we study conical singularities for G_2 manifolds. The present paper deals with the *desingularization* of G_2 manifolds with conical singularities by gluing. The next paper [24] is about the *deformation theory* (moduli spaces) of such manifolds and of the closely related asymptotically conical G_2 manifolds. We use the same notation and occasionally mention some results from [24]. However, this paper is completely self-contained. All the results we use from other sources are carefully stated.

The main theorem we prove in this paper is the following. The notation and terminology used in this theorem will all be defined in Sections 2 and 3.

Main Theorem (Theorem 3.28) *Let M be a compact G_2 manifold with isolated conical singularities, with singularities x_1, \dots, x_n , cones C_1, \dots, C_n and rates μ_1, \dots, μ_n , respectively. Suppose that we have asymptotically conical G_2 manifolds N_1, \dots, N_n , with the same cones C_1, \dots, C_n and rates ν_1, \dots, ν_n , respectively, with each $\nu_i \leq -3$. If the topological conditions (74) and (75) of Theorem 3.10 are satisfied, then there exists a one-parameter family \tilde{M}_s of smooth, compact G_2 manifolds, for $0 < s < \kappa$, (with holonomy exactly equal to G_2), which desingularize M .*

The study of manifolds with isolated conical singularities (ICS) or of asymptotically conical (AC) manifolds in the context of special holonomy and calibrated geometry was initiated by Joyce in his series of papers [20; 21; 22; 23; 19] about special Lagrangian submanifolds with ICS. Marshall [33] and Pacini [36] studied AC special Lagrangian submanifolds of \mathbb{C}^m , and Lotay [31; 32; 30] studied coassociative AC and ICS submanifolds of G_2 manifolds. The results in the present paper most closely relate to the work of Chan [11; 10; 12] on AC and ICS Calabi–Yau 3–folds.

For those readers who are familiar with the papers [11; 12], the main differences between the G_2 case of the present paper and the Calabi–Yau 3–fold case studied by Chan are the following. There are two sources of topological obstructions, coming from the 3–form φ and the 4–form ψ of a G_2 structure, compared to just the one source coming from Ω in the Calabi–Yau case. The solution to 3–form obstructions is very similar to the approach of Chan [12], except that we can use the full $d + d^*$ operator rather than having to restrict to a subcomplex, but we need to also obtain excluded ranges for the orders of homogeneity for harmonic 2–forms, in addition to the 0–forms and 1–forms which are needed in the Calabi–Yau 3–fold case. The solution to 4–form obstructions cannot be obtained in the same way, for technical reasons, but there turns out to be a much easier way to deal with those. Also, Chan assumes that the asymptotically conical manifolds used in the gluing already admit a good asymptotic expansion near infinity (and he shows that this does hold in the examples he discusses.) We prove that under a natural gauge–fixing condition, a good asymptotic expansion which allows the gluing construction to succeed always exists.

We now discuss the organization of this paper. Section 1.1 is a review of the main facts about G_2 manifolds which we require. More thorough discussions of G_2 structures can be found in Bryant [7] and Joyce [18].

Section 2 is about G_2 manifolds modelled on cones. In Section 2.1 we treat G_2 cones, and in Section 2.2 we present several results about differential forms on G_2 cones which we will need later. Sections 2.3 and 2.4 discuss, respectively, G_2 manifolds with isolated conical singularities (ICS) and asymptotically conical G_2 manifolds (AC), including the known explicit examples in the AC case.

In [Section 3](#) we present the details of the desingularization procedure as three steps. Step one in [Section 3.1](#) is concerned with constructing a compact smooth manifold \widetilde{M}_s for all s sufficiently small. Then in step two in [Section 3.2](#) we construct a family φ_s of closed G_2 structures on \widetilde{M}_s with small torsion. It is at this stage where we require two analytic results, which are described here, but whose proof is postponed until later in the paper. The first result involves the existence of a good asymptotic expansion near infinity for the G_2 structure of an asymptotically conical G_2 manifold. The second result is concerned with topological obstructions to the gluing procedure that can arise and their resolution. Finally in step three of the desingularization procedure in [Section 3.3](#) we obtain estimates to show that the torsion of φ_s is small enough to invoke a theorem of Joyce that gives the existence of a torsion-free G_2 structure $\widetilde{\varphi}_s$ on \widetilde{M}_s for s sufficiently small.

In [Section 4](#), we briefly review and summarize the relevant results which we will need from the Fredholm theory on weighted Sobolev spaces for noncompact manifolds with ends. [Section 4.1](#) is about manifolds with isolated conical singularities, while [Section 4.2](#) is about asymptotically conical manifolds.

In [Section 5](#) we discuss how to overcome the topological obstructions to our gluing procedure of [Section 3.2](#) when a certain necessary condition is satisfied, by explaining how to explicitly construct “correction forms” to be able to carry out the construction in the obstructed case. In [Section 5.1](#) we deal with the case of 3-form obstructions, and in [Section 5.2](#) we handle the 4-form obstructions.

Finally, in [Section 6](#) we prove that a good asymptotic expansion exists for an asymptotically conical G_2 manifold which satisfies a natural gauge-fixing condition.

Conventions There are two sign conventions in G_2 geometry. The convention we choose to use is the one used in Bryant–Salamon [\[8\]](#) and in Harvey–Lawson [\[16\]](#), but differs from the convention used in Bryant [\[7\]](#) or Joyce [\[18\]](#). A detailed discussion of sign conventions and orientations in G_2 geometry can be found in the author’s note [\[25\]](#).

The letter C is used in two ways: to denote a *cone* or to denote a *positive constant*. The use will always be clear from the context. When we are estimating various quantities, the value of $C > 0$ will change from step to step in the calculations, but we will always use the same letter C to denote it, to avoid a proliferation of notation. We use $f = O(r)$ to mean that $f \leq Cr$ for some $C > 0$ as $r \rightarrow 0$ (when we are working near a singularity), or as $r \rightarrow \infty$ (when we are working on an asymptotic end at infinity.)

One fact we will use repeatedly is that for any form ω , the exterior derivative $d\omega$ is obtained by the skew-symmetrization of the covariant derivative $\nabla\omega$, so there is some

$C > 0$ such that

$$(1) \quad |d\omega| \leq C|\nabla\omega|, \quad |\nabla d\omega| \leq C|\nabla^2\omega|.$$

Also, we have

$$|d^*\omega| = |*d*\omega| = |d*\omega| \leq C|\nabla*\omega| = C|*\nabla\omega| = C|\nabla\omega|$$

using the fact that $d^* = \pm *d*$ and that $*$ is an isometry and commutes with ∇ . Thus we also have

$$(2) \quad |d^*\omega| \leq C|\nabla\omega|.$$

Acknowledgements The author is extremely grateful to Dominic Joyce for his tireless patience in explaining the technical details about weighted Banach spaces and gluing constructions, for his many useful suggestions and for providing the initial overall direction for this project. Useful discussions were also had with Mark Haskins, Tommaso Pacini and Yinan Song.

The author's research has been supported by a Marie Curie Fellowship of the European Commission under contract number MIF1-CT-2006-039113. The contents of this work reflect only the author's views and not the views of the European Commission.

1.1 Review of G_2 manifolds

Recall that a G_2 structure on a connected smooth 7-manifold M is described by a smooth 3-form φ satisfying a certain “nondegeneracy” condition. When such a structure exists, there is an open subbundle Ω_+^3 of the bundle Ω^3 of 3-forms consisting of nondegenerate 3-forms, also called *positive* or *stable* 3-forms. A G_2 structure φ determines a Riemannian metric g_φ and an orientation vol_φ in a nonlinear way. Thus φ determines a Hodge star operator $*_\varphi$, and $\psi = *_\varphi\varphi$ is the dual 4-form which is an element of the bundle Ω_+^4 of positive 4-forms.

Definition 1.1 A G_2 manifold is a connected manifold with a G_2 structure (M, φ) such that φ is *parallel* with respect to the Levi-Civita connection ∇ determined by g_φ . That is, $\nabla_{g_\varphi}\varphi = 0$. Such a G_2 structure is also called *torsion-free*. In this case the Riemannian holonomy $\text{Hol}_g(M)$ of (M, g_φ) is contained in the group $G_2 \subseteq \text{SO}(7)$. If the fundamental group $\pi_1(M)$ is finite, then the holonomy of g_φ is exactly G_2 . (See Joyce [18] for more details.)

Remark 1.2 A G_2 manifold is always *Ricci-flat*, and by Fernández–Gray [13], a G_2 structure φ is torsion-free if and only if it is both closed and coclosed: $d\varphi = 0$ and $d\psi = 0$.

The space Ω^3 of 3-forms on a manifold with G_2 structure φ decomposes as

$$(3) \quad \Omega^3 = \Omega_1^3 \oplus \Omega_7^3 \oplus \Omega_{27}^3$$

into irreducible G_2 representations. Similarly we have a decomposition of the space Ω^2 as

$$(4) \quad \Omega^2 = \Omega_7^2 \oplus \Omega_{14}^2,$$

as well as isomorphic splittings of Λ^4 and Λ^5 given by the Hodge star of the above decompositions: $\Lambda_l^k = *_\varphi(\Lambda_l^{7-k})$. When the G_2 structure is torsion-free, these decompositions are preserved by the Hodge Laplacian $\Delta = dd^* + d^*d$. The essential aspect of this fact that we will need is the following. Suppose f is any function and ω is any 1-form on a G_2 manifold M . Then

$$(5) \quad \Delta(f\varphi) = (\Delta f)\varphi, \quad \Delta(f\psi) = (\Delta f)\psi,$$

$$(6) \quad \Delta(\omega \wedge \varphi) = (\Delta\omega) \wedge \varphi, \quad \Delta(\omega \wedge \psi) = (\Delta\omega) \wedge \psi.$$

The identities in (5) can be proved using just the fact that φ and ψ are parallel, while the identities in (6) also require the fact that G_2 manifolds have vanishing Ricci curvature.

Let $\Theta: \Lambda_+^3 \rightarrow \Lambda_+^4$ be the nonlinear map which associates to any G_2 structure φ , the dual 4-form $\psi = \Theta(\varphi) = *_\varphi\varphi$ with respect to the metric g_φ associated to φ . One result which will be crucial is the following. This is Proposition 10.3.5 in Joyce [18], adapted to suit our present purposes.

Lemma 1.3 *Suppose that φ is a G_2 structure with induced metric g_φ and dual 4-form $\psi = *_\varphi\varphi$. Let ξ be a 3-form which has sufficiently small pointwise norm with respect to g_φ , so that $\varphi + \xi$ is still nondegenerate. Then we have*

$$(7) \quad \Theta(\varphi + \xi) = \psi + *_\varphi \left(\frac{4}{3}\pi_1(\xi) + \pi_7(\xi) - \pi_{27}(\xi) \right) + F_\varphi(\xi),$$

where π_k is the projection onto the subspace Ω_k^3 with respect to the G_2 structure φ . The nonlinear function $F_\varphi: \Omega^3 \rightarrow \Omega^4$ satisfies

$$(8) \quad F_\varphi(0) = 0, \quad |F_\varphi(\xi)| \leq C|\xi|^2, \quad |\nabla F_\varphi(\xi)| \leq C(|\xi|^2|\nabla\varphi| + |\xi||\nabla\xi|),$$

for some $C > 0$, where the norms and the covariant derivatives are taken with respect to g_φ .

Remark 1.4 It is clear that the same sort of result holds for the map $\Theta^{-1}: \Omega_+^4 \rightarrow \Omega_+^3$. That is, if η is a 4–form with sufficiently small pointwise norm, then

$$(9) \quad \Theta^{-1}(\psi + \eta) = \varphi + *_{\varphi} \left(\frac{3}{4} \pi_1(\eta) + \pi_7(\eta) - \pi_{27}(\eta) \right) + G_{\varphi}(\eta),$$

where π_k is the projection onto Ω_k^4 with respect to φ and $G_{\varphi}: \Omega^4 \rightarrow \Omega^3$ satisfies

$$(10) \quad G_{\varphi}(0) = 0, \quad |G_{\varphi}(\eta)| \leq C|\eta|^2, \quad |\nabla G_{\varphi}(\eta)| \leq C(|\eta|^2|\nabla\varphi| + |\eta||\nabla\eta|),$$

for some $C > 0$.

It will be convenient to define the linear operator $J: \Omega^4 \rightarrow \Omega^3$ given by

$$(11) \quad J_{\varphi}(\eta) = *_{\varphi} \left(\frac{3}{4} \pi_1(\eta) + \pi_7(\eta) - \pi_{27}(\eta) \right).$$

Then Equation (9) becomes $\Theta^{-1}(\psi + \eta) = \varphi + J_{\varphi}(\eta) + G_{\varphi}(\eta)$. Given explicit formulas for π_1 and π_7 , as can be found for example in [26], it is not difficult to verify that

$$(12) \quad |J_{\varphi}(\eta)| \leq C|\eta|, \quad |\nabla J_{\varphi}(\eta)| \leq C(|\eta||\nabla\varphi| + |\nabla\eta|),$$

which we will need in Section 3.3 to estimate the torsion of the G_2 structure we construct by gluing in Section 3.2.

2 G_2 manifolds modeled on cones

In this section we discuss G_2 manifolds which are modeled on cones. Specifically these are: G_2 cones, G_2 manifolds with isolated conical singularities (ICS) and asymptotically conical G_2 manifolds (AC).

2.1 G_2 cones

Let Σ^6 be a compact, connected, smooth 6–manifold. An $SU(3)$ structure on Σ is described by a Riemannian metric g_{Σ} , an almost complex structure J which is orthogonal with respect to g_{Σ} , the associated Kähler form $\omega(u, v) = g_{\Sigma}(Ju, v)$ which is real and of type $(1, 1)$ with respect to J and a nonvanishing complex $(3, 0)$ –form Ω . The two forms are related by the normalization condition

$$(13) \quad \text{vol}_{\Sigma} = \frac{1}{6} \omega^3 = \frac{i}{8} \Omega \wedge \bar{\Omega} = \frac{1}{4} \text{Re}(\Omega) \wedge \text{Im}(\Omega).$$

Such a structure exists whenever Σ admits an almost complex structure and $c_1(\Sigma) = 0$, so that the canonical bundle $\Lambda^{3,0}$ of Σ is topologically trivial and a nonvanishing

section Ω exists. Then we can always scale Ω by a nonzero complex valued function to ensure that (13) holds.

Definition 2.1 The manifold Σ^6 with $SU(3)$ structure (ω, Ω) is called *strictly nearly Kähler*, or a *weak $SU(3)$ manifold*, if the following equations are satisfied:

$$(14) \quad d_\Sigma \omega = -3 \operatorname{Re}(\Omega), \quad d_\Sigma \operatorname{Im}(\Omega) = 4 \frac{\omega^2}{2}.$$

Remark 2.2 All strictly nearly Kähler 6-manifolds can be shown to be *Einstein*, with *positive* scalar curvature. It then follows from the Weitzenböck formula that in the compact case, the first Betti number vanishes: $b_1(\Sigma) = 0$. Some references for strictly nearly Kähler manifolds are Bär [3], Reyes-Carrión-Salamon [37] and Gray [14; 15].

The following result is proved in Bär [3].

Proposition 2.3 *Let Σ^6 be strictly nearly Kähler. Then there exists a torsion-free G_2 structure (φ_C, ψ_C, g_C) on $C' = (0, \infty) \times \Sigma$. This structure is defined by*

$$(15) \quad \varphi_C = r^3 \operatorname{Re}(\Omega) - r^2 dr \wedge \omega,$$

$$(16) \quad \psi_C = -r^3 dr \wedge \operatorname{Im}(\Omega) - r^4 \frac{\omega^2}{2},$$

$$(17) \quad g_C = dr^2 + r^2 g_\Sigma,$$

where r is the coordinate on $(0, \infty)$. The metric g_C in (17) is called a *cone metric* and we say $C = C' \cup \{\mathbf{0}\}$ is a G_2 cone. In fact for any Riemannian cone $C = (0, \infty) \times \Sigma \cup \{\mathbf{0}\}$ with holonomy contained in G_2 , the holonomy is either trivial, in which case Σ is the standard round sphere S^6 and C is the Euclidean \mathbb{R}^7 , or else the holonomy is exactly equal to G_2 , in which case the link Σ is strictly nearly Kähler, but not equal to S^6 .

In this paper, a G_2 cone will always have holonomy exactly G_2 , so we exclude the case where the link is S^6 . The space $C = C' \cup \{\mathbf{0}\}$ is a G_2 cone, C' is the *smooth part* of C , the point $\mathbf{0}$ is the *singular point* of the cone, and Σ is called the *link* of the cone. Notice that $\mathbf{0} = \lim_{r \rightarrow 0^+} (r, \sigma)$ for any point σ in Σ . Now $\det(g_C) = r^{12} \det(g_\Sigma)$, and we choose the orientation on C' so that

$$(18) \quad \operatorname{vol}_C = r^6 dr \wedge \operatorname{vol}_\Sigma$$

is the volume form on C' .

For any $t > 0$, we have a *dilation* map $\mathbf{t}: C \rightarrow C$ defined by

$$(19) \quad \mathbf{t}(\mathbf{0}) = \mathbf{0}, \quad \mathbf{t}(r, \sigma) = (tr, \sigma).$$

If we pull back φ_C , ψ_C , g_C and vol_C with respect to \mathbf{t} , we see that

$$(20) \quad \begin{aligned} \mathbf{t}^*(\varphi_C) &= t^3 \varphi_C, & \mathbf{t}^*(\psi_C) &= t^4 \psi_C, \\ \mathbf{t}^*(g_C) &= t^2 g_C, & \mathbf{t}^*(\text{vol}_C) &= t^7 \text{vol}_C, \end{aligned}$$

and we say that the conical G_2 structure is *dilation-equivariant*. Similarly any contravariant tensor ω of degree k such that $\mathbf{t}^*(\omega) = t^k \omega$ is called dilation-equivariant. A useful property of dilations which we will use frequently is the following. Since $g_{C_i}(r, \sigma) = t^{-2} g_{C_i}(tr, \sigma)$ for vector fields, we see that

$$(21) \quad |\mathbf{t}^*(\omega)(r, \sigma)|_{g_{C_i}(r, \sigma)} = t^k |\omega(tr, \sigma)|_{g_{C_i}(tr, \sigma)}$$

whenever ω is a contravariant tensor of degree k .

Proposition 2.4 *The forms φ_C and ψ_C on the smooth part C' of a G_2 cone C are exact, and hence the cohomology classes $[\varphi_C]$ and $[\psi_C]$ are trivial in $H^3(C', \mathbb{R}) \cong H^3(\Sigma, \mathbb{R})$ and $H^4(C', \mathbb{R}) \cong H^4(\Sigma, \mathbb{R})$, respectively.*

Proof By using the relations (14) defining a strictly nearly Kähler structure on the link Σ , it is easy to check that (15) and (16) can be written as

$$\varphi_C = d\left(-\frac{r^3}{3}\omega\right), \quad \psi_C = d\left(-\frac{r^4}{4}\text{Im}(\Omega)\right),$$

which is exactly what we needed to show. □

There are three known compact strictly nearly Kähler manifolds (other than the round S^6), and hence three known G_2 cones, which we now discuss. They are all obtained by taking the biinvariant metric on a compact Lie group G and descending this to the normal metric on G/H for an appropriate Lie subgroup H . In particular, all these examples are homogeneous spaces. See Bär [3] for more details.

Example 2.5 (The complex projective space $\mathbb{C}P^3$) We can view $\mathbb{C}P^3$ (diffeomorphically) as the homogeneous space $\text{Sp}(2)/(\text{Sp}(1) \times \text{U}(1))$. There is a natural nonintegrable complex structure on this space, different from the standard one, and the normal metric is not the Fubini–Study metric, so this $\mathbb{C}P^3$ is *not* Kähler. This space is also the *twistor space* of the standard round S^4 , which is the unit sphere subbundle of the bundle $\Lambda_-^2(S^4)$ of anti-self-dual 2-forms on S^4 .

Example 2.6 (The complex flag manifold $F_{1,2} = \text{SU}(3)/T^2$) This is the homogeneous space $\text{SU}(3)/T^2$ where $T^2 = \text{S}(\text{U}(1) \times \text{U}(1) \times \text{U}(1)) \subset \text{SU}(3)$ is the maximal torus. This is the space of pairs (V_1, V_2) where V_1 and V_2 are complex linear subspaces

of \mathbb{C}^3 with $V_1 \subset V_2$. This space is also the *twistor space* of the standard Fubini–Study $\mathbb{C}P^2$, which is the unit sphere subbundle of the bundle $\Lambda_-^2(\mathbb{C}P^2)$ of anti-self-dual 2-forms on $\mathbb{C}P^2$.

Example 2.7 (The product of two 3-spheres $S^3 \times S^3$) View S^3 as the Lie group of unit quaternions, which is isomorphic to $SU(2)$. Then let $S^3 \times S^3 = (S^3 \times S^3 \times S^3)/S^3$ where we embed S^3 into $S^3 \times S^3 \times S^3$ as the diagonal subgroup. This space is also the unit sphere subbundle of the *spinor bundle* $\mathcal{S}(S^3)$ of the standard round S^3 .

Remark 2.8 We note for later use that Examples 2.5 and 2.6 both have $H^3(\Sigma, \mathbb{R}) = 0$ and $H^4(\Sigma, \mathbb{R}) \neq 0$ and that Example 2.7 has $H^3(\Sigma, \mathbb{R}) \neq 0$ and $H^4(\Sigma, \mathbb{R}) = 0$.

Remark 2.9 In Butruille [9] it is proved that the above examples are the only *homogeneous* strictly nearly Kähler compact manifolds. It is expected that there should exist many nonhomogeneous examples, but as far as the author is aware, none have been found yet.

2.2 Differential forms on G_2 cones

We will denote by $*_\Sigma, \nabla_\Sigma, d_\Sigma, d_\Sigma^*$ and Δ_Σ the Hodge star, Levi-Civita connection, exterior derivative, coderivative and Hodge Laplacian of Σ . Similarly $*_C, \nabla_C, d_C, d_C^*$ and Δ_C will denote the analogous operators for the smooth part C' of the cone C . We will often abuse notation and say “form on the cone C ” when we really mean “form on the smooth part C' of C .” The following proposition is a simple exercise.

Proposition 2.10 *Let ω be a smooth k -form on C . Then we can write*

$$(22) \quad \omega = dr \wedge \alpha + \beta$$

where α is a $(k-1)$ -form and β is a k -form on Σ , both depending on r as a parameter. We use $'$ to denote differentiation with respect to the parameter r . The following formulas hold:

$$(23) \quad d_C(dr \wedge \alpha + \beta) = dr \wedge (\beta' - d_\Sigma \alpha) + (d_\Sigma \beta),$$

$$(24) \quad *_C(dr \wedge \alpha + \beta) = dr \wedge ((-1)^k r^{6-2k} *_\Sigma \beta) + (r^{8-2k} *_\Sigma \alpha),$$

$$(25) \quad d_C^*(dr \wedge \alpha + \beta) = dr \wedge \left(-\frac{1}{r^2} d_\Sigma^* \alpha \right) + \left(-\frac{(8-2k)}{r} \alpha - \alpha' + \frac{1}{r^2} d_\Sigma^* \beta \right),$$

and finally

$$(26) \quad \Delta_C(dr \wedge \alpha + \beta) = dr \wedge \left(\frac{1}{r^2} \Delta_\Sigma \alpha + \frac{(8-2k)}{r^2} \alpha - \frac{(8-2k)}{r} \alpha' - \alpha'' - \frac{2}{r^3} d_\Sigma^* \beta \right) + \left(\frac{1}{r^2} \Delta_\Sigma \beta - \frac{(6-2k)}{r} \beta' - \beta'' - \frac{2}{r} d_\Sigma \alpha \right).$$

Suppose now that α is a $(k-1)$ -form on Σ and β is a k -form on Σ . Then from (17) it is easy to see that

$$(27) \quad |dr \wedge \alpha|_{g_C} = r^{-(k-1)} |\alpha|_{g_\Sigma} \quad \text{and} \quad |\beta|_{g_C} = r^{-k} |\beta|_{g_\Sigma},$$

from which it follows that

$$(28) \quad |r^{k-1} dr \wedge \alpha + r^k \beta|_{g_C}^2 = |\alpha|_{g_\Sigma}^2 + |\beta|_{g_\Sigma}^2.$$

For this reason, we will always write a k -form on the cone C' in the form $\omega = r^{k-1} dr \wedge \alpha + r^k \beta$ for some α and β , which are forms on Σ possibly depending on the parameter r . Note that if α and β were independent of r , then ω would be dilation-equivariant, as defined above.

The next lemma is an important result about closed differential forms on a cone C with certain growth rates near 0 or ∞ .

Lemma 2.11 *Let ω be a smooth closed k -form on $C' = (0, \infty) \times \Sigma$. Suppose that either*

- i) $|\omega|_{g_C} = O(r^\lambda)$ on $(0, \varepsilon) \times \Sigma$, for $\lambda > -k$, or
- ii) $|\omega|_{g_C} = O(r^\lambda)$ on $(R, \infty) \times \Sigma$, for $\lambda < -k$.

for some small ε or some large R . Then for each case respectively we have that

- i) $\omega = d\Omega$ for some $(k-1)$ -form Ω on $(0, \varepsilon) \times \Sigma$, or
- ii) $\omega = d\Omega$ for some $(k-1)$ -form Ω on $(R, \infty) \times \Sigma$,

where in each case, on its domain of definition, Ω satisfies $|\Omega|_{g_C} = O(r^{\lambda+1})$. If we suppose further that $0 < \lambda < 1$ in case i), then for either case if we also have that $|\nabla_C^j \omega|_{g_C} = O(r^{\lambda-j})$, then for this Ω we have $|\nabla_C^j \Omega|_{g_C} = O(r^{\lambda+1-j})$ for all $j \geq 0$.

Proof Write $\omega(r, \sigma) = dr \wedge \alpha(r, \sigma) + \beta(r, \sigma)$, where α and β are a $(k-1)$ -form and a k -form on Σ , respectively. Since $d\omega = 0$, by Equation (23) we have

$$(29) \quad d_\Sigma \beta = 0, \quad d_\Sigma \alpha = \beta'.$$

For case i) let us define

$$\Omega(r, \sigma) = \int_0^r \alpha(t, \sigma) dt.$$

This integral converges since $|\Omega(r, \sigma)|_{g_\Sigma} \leq \int_0^r |\alpha(t, \sigma)|_{g_\Sigma} dt \leq C \int_0^r t^{k-1+\lambda} dt < \infty$, where we have used (27) and the facts that $|\alpha(r, \sigma)|_{g_C(r, \sigma)} \leq Cr^\lambda$ and $\lambda + k > 0$ by hypothesis. Note that these hypotheses also show that

$$(30) \quad \lim_{r \rightarrow 0} \beta(r, \sigma) = 0.$$

It is clear (since $\lambda \neq -k$) that $|\Omega|_{g_C} = O(r^{\lambda+1})$, and so in particular also

$$(31) \quad |d\Omega|_{g_C} = O(r^\lambda).$$

Now we compute that

$$\begin{aligned} d\Omega(r, \sigma) &= dr \wedge \alpha(r, \sigma) + \int_{r_0}^r d_\Sigma \alpha(t, \sigma) dt \\ &= dr \wedge \alpha(r, \sigma) + \int_{r_0}^r \beta'(t, \sigma) dt = dr \wedge \alpha(r, \sigma) + \beta(r, \sigma) - \lim_{r \rightarrow 0} \beta(r, \sigma) \\ &= \omega(r, \sigma) \end{aligned}$$

using (29) and (30). The fact that $|\nabla_C^j \Omega|_{g_C} = O(r^{\lambda+1-j})$ if we know that $|\nabla_C^j \omega|_{g_C} = O(r^{\lambda-j})$ is a simple exercise. The constraints on λ are sufficient to ensure no logarithmic terms arise from having to integrate t^{-1} . The proof of case ii) is analogous, defining in this case $\Omega(r, \sigma) = \int_\infty^r \alpha(t, \sigma) dt$ and using the fact that $\lambda + k < 0$ to ensure that the integral converges and that $\lim_{r \rightarrow \infty} \beta(r, \sigma) = 0$. \square

Definition 2.12 We say that a smooth k form ω on C' is *homogeneous of order λ* if

$$(32) \quad \omega = r^\lambda (r^{k-1} dr \wedge \alpha + r^k \beta)$$

where α and β are forms on Σ , independent of r . Then we see that

$$|\mathbf{t}^*(\omega)(r, \sigma)|_{g_C(tr, \sigma)} = |t^{\lambda+k} \omega(r, \sigma)|_{g_C(tr, \sigma)} = t^{\lambda+k} t^{-k} |\omega(r, \sigma)|_{g_C(r, \sigma)},$$

which we can write more concisely as

$$(33) \quad \mathbf{t}^*|\omega|_{g_C} = t^\lambda |\omega|_{g_C},$$

so the function $|\omega|_{g_C}$ on C' is homogeneous of order λ in the variable r in the usual sense.

Remark 2.13 It is easy to see that a homogeneous k -form ω of order λ is dilation-equivariant ($\mathbf{t}^*(\omega) = t^k \omega$) if $\lambda = 0$. Similarly it is *dilation-invariant* ($\mathbf{t}^*(\omega) = \omega$) if $\lambda = -k$.

Remark 2.14 It follows directly from (24) that if ω is homogeneous of order λ , then $*_C \omega$ is also homogeneous of the same order λ .

Using (23), (25) and (26), for a homogenous k -form $\omega = r^\lambda(r^{k-1} dr \wedge \alpha + r^k \beta)$ of order λ , we find that

$$(34) \quad d_C \omega = r^{\lambda+k-1} dr \wedge ((\lambda + k)\beta - d_\Sigma \alpha) + r^{\lambda+k} d_\Sigma \beta,$$

$$(35) \quad d_C^* \omega = r^{\lambda+k-3} dr \wedge (-d_\Sigma^* \alpha) + r^{\lambda+k-2} (-(\lambda - k + 7)\alpha + d_\Sigma^* \beta),$$

$$(36) \quad \Delta_C \omega = r^{\lambda+k-3} dr \wedge (\Delta_\Sigma \alpha - (\lambda + k - 2)(\lambda - k + 7)\alpha - 2d_\Sigma^* \beta) + r^{\lambda+k-2} (\Delta_\Sigma \beta - (\lambda + k)(\lambda - k + 5)\beta - 2d_\Sigma \alpha).$$

In Section 5.1, we will need to consider the possible order λ of a homogeneous k -form ω_k on a cone C which is in the kernel of Δ_C , or of a mixed degree form $\omega = \sum_{k=0}^7 \omega_k$ which is in the kernel of $d_C + d_C^*$. The fact that we will need to use repeatedly in the rest of this section is that the Laplacian Δ_Σ on the link Σ has nonnegative eigenvalues, so whenever we have an expression of the form $\Delta_\Sigma \gamma = \mu \gamma$ for some $\mu < 0$ and γ a form on Σ , we must have $\gamma = 0$.

Proposition 2.15 *Let ω be a homogeneous k -form of order λ which is harmonic on the cone: $\Delta_C \omega = 0$. Then we have:*

$$(37) \quad \text{For } k = 0, 7, \quad \omega = 0 \text{ if } \lambda \in (-5, 0),$$

$$(38) \quad \text{For } k = 1, 6, \quad \omega = 0 \text{ if } \lambda \in (-4, -1),$$

$$(39) \quad \text{For } k = 2, 5, \quad \omega = 0 \text{ if } \lambda \in (-3, -2).$$

This gives an excluded range of orders of homogeneity for harmonic forms on the cone C .

Proof Since the star operator $*_C$ commutes with the Laplacian Δ_C and preserves the order of homogeneity by Remark 2.14, it suffices to check the cases $k = 0, 1, 2$. Let $k = 0$ in Equation (36), and noting that $\alpha_0 = 0$ and $d_\Sigma^* \beta_0 = 0$, we see that $\Delta_C \omega = 0$ implies

$$\Delta_\Sigma \beta_0 = \lambda(\lambda + 5)\beta_0,$$

and the result now follows for $\lambda \in (-5, 0)$. We need to work a little harder for $k = 1$ and $k = 2$.

For $k = 1$, we get the following system of equations:

$$(40) \quad \Delta_\Sigma \alpha_1 = (\lambda - 1)(\lambda + 6)\alpha_1 + 2d_\Sigma^* \beta_1, \quad \Delta_\Sigma \beta_1 = (\lambda + 1)(\lambda + 4)\beta_1 + 2d_\Sigma \alpha_1.$$

Now suppose that $\lambda \in (-4, -1)$. If we take d_Σ of the second equation, since Δ_Σ commutes with d_Σ , we get $\Delta_\Sigma(d_\Sigma \beta_1) = (\lambda + 1)(\lambda + 4)d_\Sigma \beta_1$ and hence $d_\Sigma \beta_1 = 0$. Similarly taking d_Σ^* of the first equation gives $d_\Sigma^* \alpha_1 = 0$. Also, taking d_Σ of the first equation gives

$$\Delta_\Sigma(d_\Sigma \alpha_1) = (\lambda - 1)(\lambda + 6)d_\Sigma \alpha_1 + 2\Delta_\Sigma \beta_1 = (\lambda^2 + 5\lambda - 2)d_\Sigma \alpha_1 + (2\lambda^2 + 10\lambda + 8)\beta_1.$$

Now an easy calculation reveals that

$$\Delta_\Sigma(d_\Sigma \alpha_1 + (\lambda + 4)\beta_1) = (\lambda + 1)(\lambda + 6)(d_\Sigma \alpha_1 + (\lambda + 4)\beta_1),$$

and hence $d_\Sigma \alpha_1 + (\lambda + 4)\beta_1 = 0$ for $\lambda \in (-4, -1)$. This relation can now be substituted into each equation in (40) to yield

$$\Delta_\Sigma \alpha_1 = (\lambda - 1)(\lambda + 4)\alpha_1, \quad \Delta_\Sigma \beta_1 = (\lambda - 1)(\lambda + 4)\beta_1,$$

from which it follows that $\alpha_1 = 0$ and $\beta_1 = 0$.

The proof for $k = 2$ is analogous, but we present it for completeness. We get the following system of equations:

$$(41) \quad \Delta_\Sigma \alpha_2 = \lambda(\lambda + 5)\alpha_2 + 2d_\Sigma^* \beta_2, \quad \Delta_\Sigma \beta_2 = (\lambda + 2)(\lambda + 3)\beta_2 + 2d_\Sigma \alpha_2.$$

Suppose that $\lambda \in (-3, -2)$. We get $\Delta_\Sigma(d_\Sigma \beta_2) = (\lambda + 2)(\lambda + 3)d_\Sigma \beta_2$, so $d_\Sigma \beta_2 = 0$. Similarly taking d_Σ^* of the first equation gives $d_\Sigma^* \alpha_2 = 0$. We also have

$$\Delta_\Sigma(d_\Sigma \alpha_2) = \lambda(\lambda + 5)d_\Sigma \alpha_2 + 2\Delta_\Sigma \beta_2 = (\lambda^2 + 5\lambda + 4)d_\Sigma \alpha_2 + (2\lambda^2 + 10\lambda + 12)\beta_2.$$

This time one can check that

$$\Delta_\Sigma(d_\Sigma \alpha_2 + (\lambda + 3)\beta_2) = (\lambda + 2)(\lambda + 5)(d_\Sigma \alpha_2 + (\lambda + 3)\beta_2),$$

and hence $d_\Sigma \alpha_2 + (\lambda + 3)\beta_2 = 0$ for $\lambda \in (-3, -2)$. This relation can now be substituted into each equation in (41) to yield

$$\Delta_\Sigma \alpha_2 = \lambda(\lambda + 3)\alpha_2, \quad \Delta_\Sigma \beta_2 = \lambda(\lambda + 3)\beta_2,$$

from which it follows that $\alpha_2 = 0$ and $\beta_2 = 0$.

The reason we have gone through all three cases carefully is the following. The reader may now be tempted to try the same trick to get an excluded range of orders for

homogeneous harmonic 3-forms. It does not take long to verify that this procedure breaks down for $k = 3$. See [Corollary 2.17](#), however, for what we can say about homogeneous 3-forms. □

To study the operator $d_C + d_C^*$ on a cone, we begin with the following lemma.

Lemma 2.16 *Let $\omega = \sum_{k=0}^7 \omega_k$, where $\omega_k = r^{\lambda+k-1} dr \wedge \alpha_k + r^{\lambda+k} \beta_k$ is a homogeneous k -form of order λ . Then if ω is in the kernel of $d_C + d_C^*$, the following equations hold:*

$$(42) \quad (\lambda + k - 1)\beta_{k-1} = d_\Sigma \alpha_{k-1} + d_\Sigma^* \alpha_{k+1},$$

$$(43) \quad (\lambda - k + 6)\alpha_{k+1} = d_\Sigma \beta_{k-1} + d_\Sigma^* \beta_{k+1},$$

for all $k = 0, \dots, 7$.

Proof By (34) and (35), we can compute that

$$\begin{aligned} & (d_C + d_C^*)\omega \\ &= \sum_{k=0}^7 (r^{\lambda+k-1} dr \wedge ((\lambda + k)\beta_k - d_\Sigma \alpha_k) + r^{\lambda+k} d_\Sigma \beta_k) \\ & \quad + \sum_{k=0}^7 (r^{\lambda+k-3} dr \wedge (-d_\Sigma^* \alpha_k) + r^{\lambda+k-2}(-(\lambda - k + 7)\alpha_k + d_\Sigma^* \beta_k)) \\ &= \sum_{l=1}^8 (r^{\lambda+l-2} dr \wedge ((\lambda + l - 1)\beta_{l-1} - d_\Sigma \alpha_{l-1}) + r^{\lambda+l-1} d_\Sigma \beta_{l-1}) \\ & \quad + \sum_{l=-1}^6 (r^{\lambda+l-2} dr \wedge (-d_\Sigma^* \alpha_{l+1}) + r^{\lambda+l-1}(-(\lambda - l + 6)\alpha_{l+1} + d_\Sigma^* \beta_{l+1})). \end{aligned}$$

Using the fact that $\alpha_0 = 0$ and $\beta_7 = 0$, and $d_\Sigma \alpha_7 = 0$ and $d_\Sigma^* \beta_0 = 0$, both these sums can be taken from $l = 0$ to 7. Relabelling the l to a k again and combining terms, we find that

$$\begin{aligned} (d_C + d_C^*)\omega &= \sum_{k=0}^7 (r^{\lambda+k-2} dr \wedge ((\lambda + k - 1)\beta_{k-1} - d_\Sigma \alpha_{k-1} - d_\Sigma^* \alpha_{k+1})) \\ & \quad + \sum_{k=0}^7 (r^{\lambda+k-1}(-(\lambda - k + 6)\alpha_{k+1} + d_\Sigma^* \beta_{k+1} + d_\Sigma \beta_{k-1})) \end{aligned}$$

where it is understood that α_k and β_k vanish when $k < 0$ or $k > 7$. Setting this expression equal to zero yields equations (42) and (43). □

Corollary 2.17 Suppose that ω is a homogeneous k -form of order λ which is closed and coclosed: $d_C \omega = 0$, $d_C^* \omega = 0$. Then we have:

$$(44) \quad \text{For } k = 0, 7, \quad \omega = 0 \text{ if } \lambda \in (-7, 0),$$

$$(45) \quad \text{For } k = 1, 6, \quad \omega = 0 \text{ if } \lambda \in (-6, -1),$$

$$(46) \quad \text{For } k = 2, 5, \quad \omega = 0 \text{ if } \lambda \in (-5, -2),$$

$$(47) \quad \text{For } k = 3, 4, \quad \omega = 0 \text{ if } \lambda \in (-4, -3).$$

This gives an excluded range of orders of homogeneity for closed and coclosed forms on the cone C .

Proof Let $\alpha_l = 0$ and $\beta_l = 0$ for all $l \neq k$ in equations (42) and (43). Thus we have that $d_C \omega = 0$ and $d_C^* \omega = 0$ together imply the following equations:

$$(48) \quad d_\Sigma \alpha_k = (\lambda + k) \beta_k, \quad d_\Sigma^* \beta_k = (\lambda - k + 7) \alpha_k, \quad d_\Sigma^* \alpha_k = 0, \quad d_\Sigma \beta_k = 0.$$

Since ω is closed and coclosed, it is also harmonic: $\Delta_C \omega = 0$. Therefore, Equation (36) shows that

$$\begin{aligned} \Delta_\Sigma \alpha_k &= (\lambda + k - 2)(\lambda - k + 7) \alpha_k + 2d_\Sigma^* \beta_k, \\ \Delta_\Sigma \beta_k &= (\lambda + k)(\lambda - k + 5) \beta_k + 2d_\Sigma \alpha_k. \end{aligned}$$

Now substituting in the relations from (48) and simplifying, we obtain

$$\Delta_\Sigma \alpha_k = (\lambda + k)(\lambda - k + 7) \alpha_k, \quad \Delta_\Sigma \beta_k = (\lambda + k)(\lambda - k + 7) \beta_k.$$

Since the Laplacian has nonnegative eigenvalues, we see that both α_k and β_k must vanish (and hence $\omega = 0$) if $(\lambda + k)(\lambda - k + 7) < 0$, which occurs exactly when λ is between $-k$ and $k - 7$. □

Remark 2.18 Corollary 2.17 should be compared to Proposition 2.15. Closed and coclosed forms are harmonic, but not always conversely. The above result says that when the form is known to be closed and coclosed, we can get a bigger range of excluded orders of homogeneity. Also, Proposition 2.15 tells us nothing about 3-forms.

Corollary 2.19 Suppose that ω is a homogeneous k -form of order $-k$ which is closed and coclosed: $d_C \omega = 0$, $d_C^* \omega = 0$. Then $\omega = \beta$, where β is a harmonic k -form on the link Σ : $\Delta_\Sigma \beta = 0$.

Proof We substitute $\lambda = -k$ in the relations (48) to get

$$d_\Sigma \alpha_k = 0, \quad d_\Sigma^* \beta_k = (7 - 2k) \alpha_k, \quad d_\Sigma^* \alpha_k = 0, \quad d_\Sigma \beta_k = 0.$$

Since Σ is compact and oriented, we can use Hodge theory. The first and third equations above say that α_k is harmonic, but since $(7 - 2k)$ is never zero, the second equations says that α_k is also coexact. Thus $\alpha_k = 0$, and then the second and fourth equations say that β_k is harmonic on Σ . \square

The next three propositions are needed in [Section 5.1](#) for the solution of the obstruction problem and also in [Section 6](#) for the asymptotic expansion of the G_2 structure on an asymptotically conical G_2 manifold.

Proposition 2.20 *Let $\omega = \sum_{k=0}^3 \omega_{2k}$ be a mixed even-degree form on the cone $C' = (0, \infty) \times \Sigma$, which is homogeneous of order $\lambda = -3$. That is, each $\omega_{2k} = r^{2k-4} dr \wedge \alpha_{2k} + r^{2k-3} \beta_{2k}$, where α_{2k} and β_{2k} are $(2k-1)$ -forms and $2k$ -forms on Σ , respectively, independent of r . If $(d_C + d_C^*)(\omega) = 0$, then $\omega = dr \wedge \alpha_4$, where α_4 is a harmonic 3-form on Σ .*

Proof Substituting $\lambda = -3$ into [Lemma 2.16](#), we obtain

$$(49) \quad (k - 4)\beta_{k-1} = d_\Sigma \alpha_{k-1} + d_\Sigma^* \alpha_{k+1}, \quad (3 - k)\alpha_{k+1} = d_\Sigma \beta_{k-1} + d_\Sigma^* \beta_{k+1}.$$

If we take d_Σ and d_Σ^* of these equations and relabel indices, we get

$$\begin{aligned} d_\Sigma d_\Sigma^* \alpha_k &= (k - 5)d_\Sigma \beta_{k-2}, & d_\Sigma^* d_\Sigma \alpha_k &= (k - 3)d_\Sigma^* \beta_k, \\ d_\Sigma d_\Sigma^* \beta_k &= (4 - k)d_\Sigma \alpha_k, & d_\Sigma^* d_\Sigma \beta_k &= (2 - k)d_\Sigma^* \alpha_{k+2}. \end{aligned}$$

These can now be combined to yield

$$\Delta_\Sigma \alpha_k = (k - 3)d_\Sigma^* \beta_k + (k - 5)d_\Sigma \beta_{k-2}, \quad \Delta_\Sigma \beta_k = (2 - k)d_\Sigma^* \alpha_{k+2} + (4 - k)d_\Sigma \alpha_k.$$

We can now use (49) again to eliminate $d_\Sigma^* \beta_k$ and $d_\Sigma^* \alpha_{k+2}$ in the above expressions to finally obtain

$$(50) \quad \Delta_\Sigma \alpha_k = -(k - 3)(k - 4)\alpha_k - 2d_\Sigma \beta_{k-2}, \quad \Delta_\Sigma \beta_k = -(k - 3)(k - 2)\beta_k + 2d_\Sigma \alpha_k.$$

Now $\alpha_0 = 0$, so the second equation in (50) gives $\Delta_\Sigma \beta_0 = -6\beta_0$, so $\beta_0 = 0$. Then the first equation in (50) gives $\Delta_\Sigma \alpha_2 = -2\alpha_2$, so $\alpha_2 = 0$. We continue in this way to alternate between the equations. Next we get $\Delta_\Sigma \beta_2 = 0$, but from (49) we see that β_2 is also coexact, so $\beta_2 = 0$. Then $\Delta_\Sigma \alpha_4 = 0$, but this time (49) gives no information about α_4 so all we can say is that it is a harmonic 3-form on Σ , hence closed and coclosed. Then $\Delta_\Sigma \beta_4 = -2\beta_4$, so $\beta_4 = 0$. Continuing we get $\Delta_\Sigma \alpha_6 = -6\alpha_6$, so $\alpha_6 = 0$, and finally $\Delta_\Sigma \beta_6 = -12\beta_6$, so $\beta_6 = 0$. \square

Proposition 2.21 *Let $\omega = \sum_{l=0}^m (\log r)^l \sum_{k=0}^3 \omega_{2k,l}$ be an even-degree mixed form on the cone $C' = (0, \infty) \times \Sigma$, satisfying $(d_C + d_C^*)(\omega) = 0$, where each $\omega_{2k,l}$ is homogeneous of order -3 , and $m \geq 0$. Then in fact necessarily $m = 0$ and ω is as given in Proposition 2.20.*

Proof We prove this by contradiction. Suppose that $m > 0$. Hence $\sum_{k=0}^3 \omega_{2k,m} \neq 0$. Each $\omega_{2k,l}$ is homogenous of order -3 , so it can be written as

$$(51) \quad \omega_{2k,l} = r^{2k-4} dr \wedge \alpha_{2k,l} + r^{2k-3} \beta_{2k,l}$$

where for each l , $\alpha_{2k,l}$ and $\beta_{2k,l}$ are $(2k-1)$ -forms and $2k$ -forms on Σ , respectively, independent of r . It is easy to check that if ω_k is any k -form on the cone C , then

$$\begin{aligned} & (d_C + d_C^*)((\log r)^l \omega_k) \\ &= (\log r)^l (d_C + d_C^*)(\omega_k) + \frac{l}{r} (\log r)^{l-1} (dr \wedge \omega_k) - \frac{l}{r} (\log r)^{l-1} \left(\frac{\partial}{\partial r} \lrcorner \omega_k \right). \end{aligned}$$

Using this identity, one can now compute that

$$\begin{aligned} & (d_C + d_C^*) \left(\sum_{l=0}^m (\log r)^l \sum_{k=0}^3 \omega_{2k,l} \right) \\ &= (\log r)^m (d_C + d_C^*) \left(\sum_{k=0}^3 \omega_{2k,m} \right) \\ & \quad + \sum_{l=0}^{m-1} (\log r)^l \sum_{k=0}^3 \left(d_C(\omega_{2k,l}) + d_C^*(\omega_{2k,l}) \right. \\ & \quad \quad \left. + \frac{(l+1)}{r} dr \wedge \omega_{2k,l+1} - \frac{(l+1)}{r} \frac{\partial}{\partial r} \lrcorner \omega_{2k,l+1} \right). \end{aligned}$$

The above expression must vanish as a polynomial in $\log r$. Setting the coefficient of $(\log r)^m$ equal to zero, we see that $\sum_{k=0}^3 \omega_{2k,m}$ is in the kernel of $d_C + d_C^*$ and homogeneous of order -3 , so by Proposition 2.20 we have

$$(52) \quad \sum_{k=0}^3 \omega_{2k,m} = dr \wedge \alpha_{4,m}$$

where $\alpha_{4,m}$ is a nonzero harmonic 3-form on the link Σ . Now consider the coefficient of the next highest term, $(\log r)^{m-1}$. Setting it equal to zero and using (52) gives

$$\sum_{k=0}^3 (d_C(\omega_{2k,m-1}) + d_C^*(\omega_{2k,m-1})) - \frac{m}{r} \alpha_{4,m} = 0.$$

Using (34) and (35) with $\lambda = -3$, this becomes

$$\sum_{k=0}^3 (r^{2k-4} dr \wedge ((2k-3)\beta_{2k,m-1} - d_\Sigma \alpha_{2k,m-1}) + r^{2k-3} d_\Sigma \beta_{2k,m-1}) + \sum_{k=0}^3 (r^{2k-6} dr \wedge (-d_\Sigma^* \alpha_{2k,m-1}) + r^{2k-5} ((2k-4)\alpha_{2k,m-1} + d_\Sigma^* \beta_{2k,m-1})) - \frac{m}{r} \alpha_{4,m} = 0.$$

Taking the 3-form component of the above equation and simplifying gives

$$r^{-2} dr \wedge (-\beta_{2,m-1} - d_\Sigma \alpha_{2,m-1} - d_\Sigma^* \alpha_{4,m-1}) + r^{-1} (d_\Sigma \beta_{1,m-1} + d_\Sigma^* \beta_{4,m-1} - m\alpha_{4,m}) = 0,$$

which in turn says that

$$\alpha_{4,m} = d_\Sigma \left(\frac{1}{m} \beta_{1,m-1} \right) + d_\Sigma^* \left(\frac{1}{m} \beta_{4,m-1} \right).$$

Since Σ is compact, Hodge theory says that harmonic forms are orthogonal to the image of d_Σ and to the image of d_Σ^* , so we must have $\alpha_{4,m} = 0$, giving us our contradiction. Therefore $m = 0$. □

Proposition 2.22 *Let $\omega = \sum_{l=0}^m (\log r)^l \omega_{4,l}$ be a pure 4-form on the cone $C' = (0, \infty) \times \Sigma$, satisfying $(d + d_C^*)(\omega) = 0$, where each $\omega_{4,l}$ is homogeneous of order -4 , and $m \geq 0$. Then in fact necessarily $m = 0$ and ω is as given in Corollary 2.19.*

Proof The proof of this proposition is exactly analogous to the proof of Proposition 2.21 and is omitted. However, we note here that it is essential that ω be a pure 4-form for this result to be true. In a general even-degree mixed form in the kernel of $d + d_C^*$ which is a polynomial in $\log(r)$ with coefficients being homogeneous forms of order -4 , there can occur $\log(r)$ terms. □

The next result is about the type decomposition of 3-forms on a G_2 cone, which are homogeneous of order -3 .

Proposition 2.23 *Let (C, φ_C) be a G_2 cone. Let ξ be a homogeneous 3-form on C of order -3 which is harmonic. Then ξ is in Λ_{27}^3 with respect to φ_C .*

Proof Let $\xi = \xi_1 + \xi_7 + \xi_{27}$ be the decomposition of ξ into components. Now $\xi_1 = f\varphi_C$ for some function f on C , and since, up to a constant, $f = *_C(\xi \wedge \psi_C)$, we see that f is homogenous of order -3 . Similarly, we have $\xi_7 = *_C(\omega \wedge \varphi_C)$ for some 1-form ω on C , and, up to a constant, $\omega = *_C(\xi \wedge \varphi_C)$, so ω is also homogenous of order -3 . But ξ is harmonic, and the Laplacian commutes with the projections. Therefore $\Delta_C(\xi_1) = 0$ and $\Delta_C(\xi_7) = 0$. By (5) and (6) we then see that $\Delta_C(f) = 0$ and $\Delta_C(\omega) = 0$. Finally, Proposition 2.15 says that both f and ω are zero, since the order $\lambda = -3$ lies in the excluded range for both functions and 1-forms. Thus $\xi = \xi_{27}$ as claimed. \square

We close this section with an observation about representing cohomology classes of $(a, b) \times \Sigma$, where (a, b) is any open subinterval of $(0, \infty)$.

Proposition 2.24 *Suppose B is a cohomology class in $H^k((a, b) \times \Sigma, \mathbb{R})$. Then there exists a k -form β on $(a, b) \times \Sigma$, harmonic with respect to the cone metric and homogeneous of order $-k$ such that $[\beta] = B$.*

Proof The projection $\pi: (a, b) \times \Sigma \rightarrow \Sigma$ induces an isomorphism $\pi^*: H^k(\Sigma, \mathbb{R}) \rightarrow H^k((a, b) \times \Sigma, \mathbb{R})$ on cohomology by pullback. Then $(\pi^*)^{-1}(B)$ is a class in $H^k(\Sigma, \mathbb{R})$. Since Σ is compact and oriented, by Hodge theory there exists a unique harmonic k -form β on Σ such that $[\beta] = (\pi^*)^{-1}(B)$. Then $\pi^*(\beta) = B$, as a form on $(a, b) \times \Sigma$ and represents the class B . Now from Equation (26), with $\alpha = 0$ and $\beta' = 0$, we see that $\Delta_C \beta = (1/r^2)\Delta_\Sigma \beta = 0$, so β is a harmonic k -form on $(a, b) \times \Sigma$, homogeneous of order $-k$. \square

2.3 Compact G_2 manifolds with isolated conical singularities

Let M be a compact connected Hausdorff topological space, and let x_1, \dots, x_n be a finite set of points in M . Assume that $M' = M \setminus \{x_1, \dots, x_n\}$ is a smooth noncompact 7-dimensional manifold which we call the *smooth part* of M and $\{x_1, \dots, x_n\}$ will be called the *singular points* of M .

Definition 2.25 The space M is called a G_2 manifold with isolated conical singularities, with cones C_1, \dots, C_n at x_1, \dots, x_n and rates μ_1, \dots, μ_n , where each $\mu_i > 0$, if all of the following holds:

- The smooth part M' is a G_2 manifold with torsion-free G_2 structure φ_M and metric g_M .
- There are G_2 cones $(C_i, \varphi_{C_i}, g_{C_i})$ with links Σ_i for all $i = 1, \dots, n$.

- There is a compact subset $K \subset M'$ such that $M' \setminus K$ is a union of open sets S_1, \dots, S_n whose closures $\bar{S}_1, \dots, \bar{S}_n$ in M are all *disjoint* in M , and there exists an $\varepsilon \in (0, 1)$ such that for each $i = 1, \dots, n$, there is a smooth function $f_i: (0, \varepsilon) \times \Sigma_i \rightarrow M'$ that is a *diffeomorphism* of $(0, \varepsilon) \times \Sigma_i$ onto S_i .
- The pullback $f_i^*(\varphi_M)$ is a torsion-free G_2 structure on the subset $(0, \varepsilon) \times \Sigma_i$ of C_i . We require that this approach the torsion-free G_2 structure φ_{C_i} in a C^∞ sense, with rate $\mu_i > 0$. This means that

$$(53) \quad |\nabla_{C_i}^j (f_i^*(\varphi_M) - \varphi_{C_i})|_{g_{C_i}} = O(r^{\mu_i - j}) \quad \forall j \geq 0$$

in $(0, \varepsilon) \times \Sigma_i$. Note that all norms and derivatives are computed using the cone metric g_{C_i} .

It is easy to see that the holonomy necessarily has to be exactly G_2 , because the holonomy of the asymptotic cones is exactly G_2 , and the holonomy of M must be at least as big as the holonomy of its asymptotic cones, but it is contained in G_2 since φ_M is a torsion-free G_2 structure.

Since the metric g_M and the 4-form ψ_M are pointwise smooth functions of φ_M , by Taylor's theorem we also have

$$(54) \quad \begin{aligned} |\nabla_{C_i}^j (f_i^*(g_M) - g_{C_i})|_{g_{C_i}} &= O(r^{\mu_i - j}) \quad \forall j \geq 0, \\ |\nabla_{C_i}^j (f_i^*(\psi_M) - \psi_{C_i})|_{g_{C_i}} &= O(r^{\mu_i - j}) \quad \forall j \geq 0. \end{aligned}$$

A G_2 manifold with isolated conical singularities will sometimes be called a G_2 manifold with ICS for brevity. [Figure 1](#) shows a compact manifold with isolated conical singularities.




Figure 1: A compact manifold M with isolated conical singularities

Remark 2.26 It can be shown that if we assume that (53) holds only for $j = 0$ and $j = 1$, then there exists *particular* diffeomorphisms f_i that satisfy (53) for all $j \geq 0$. These special diffeomorphisms satisfy a *gauge-fixing condition* which forces them to solve (in some sense) an elliptic equation, and this results in their improved regularity. This will be discussed in the sequel [24] to this paper. For our present purposes, it suffices to assume from the outset that (53) holds for all $j \geq 0$. See also Remark 3.4.

Next we discuss an important fact about G_2 manifolds with ICS.

Proposition 2.27 *The closed 3-forms $f_i^*(\varphi_M)$ and φ_{C_i} on $(0, \varepsilon) \times \Sigma_i$ represent the same cohomology class in $H^3((0, \varepsilon) \times \Sigma_i, \mathbb{R}) \cong H^3(\Sigma_i, \mathbb{R})$. Similarly the closed 4-forms $f_i^*(\psi_M)$ and ψ_{C_i} on $(0, \varepsilon) \times \Sigma_i$ represent the same cohomology class in $H^4((0, \varepsilon) \times \Sigma_i, \mathbb{R}) \cong H^4(\Sigma_i, \mathbb{R})$. Therefore by Proposition 2.4, the cohomology classes $[f_i^*(\varphi_M)]$ in $H^3(\Sigma_i, \mathbb{R})$ and $[f_i^*(\psi_M)]$ in $H^4(\Sigma_i, \mathbb{R})$ are always trivial.*

Proof Using $|(f_i^*(\varphi_M) - \varphi_{C_i})|_{g_{C_i}} = O(r^{\mu_i})$ and $|(f_i^*(\psi_M) - \psi_{C_i})|_{g_{C_i}} = O(r^{\mu_i})$ and the fact that $\mu_i > 0$, this follows immediately from Lemma 2.11. \square

Remark 2.28 As discussed in Proposition 2.37 and before Definition 2.38, the analogous statement will be *false* for asymptotically conical G_2 manifolds.

On several occasions we will need to compare $f_i^*(*_M\omega)$ with $*_{C_i}f_i^*(\omega)$ near x_i .

Lemma 2.29 *Let ω be a smooth k -form on M' , with*

$$|f_i^*(\omega)|_{g_{C_i}} = O(r^\lambda)$$

near the i -th singular point x_i . Then the $(7-k)$ -forms $f_i^(*_M\omega)$ and $*_{C_i}f_i^*(\omega)$ satisfy*

$$(55) \quad |f_i^*(*_M\omega) - *_{C_i}f_i^*(\omega)|_{g_{C_i}} = O(r^{\lambda+\mu_i}).$$

Proof We begin by computing that

$$\begin{aligned} |f_i^*(*_M\omega) - *_{C_i}f_i^*(\omega)|_{g_{C_i}} &= |*_i f_i^*(g_M)(f_i^*(\omega)) - *_{C_i}(f_i^*(\omega))|_{g_{C_i}} \\ &= |(*_i f_i^*(g_M) - *_{C_i})f_i^*(\omega)|_{g_{C_i}}. \end{aligned}$$

But by (54), we have $|*_i f_i^*(g_M) - g_{C_i}|_{g_{C_i}} = O(r^{\mu_i})$, and thus

$$|(*_i f_i^*(g_M) - *_{C_i})f_i^*(\omega)|_{g_{C_i}} \leq Cr^{\mu_i} |f_i^*(\omega)|_{g_{C_i}} \leq Cr^{\lambda+\mu_i},$$

and the proof is complete. \square

Remark 2.30 There are at present no known examples of compact G_2 manifolds with isolated conical singularities, although they are expected to exist in abundance. Our main theorem in this paper can be interpreted as further evidence for the likelihood of their existence and that they should arise as “boundary points” in the moduli space of smooth compact G_2 manifolds. The author is currently working in collaboration with Dominic Joyce on a new construction of compact G_2 manifolds which should also be able to produce the first examples of compact G_2 manifolds with ICS.

2.4 Asymptotically conical G_2 manifolds

In this section we define an *asymptotically conical G_2 manifold* and discuss three explicit examples. Let N be a noncompact, connected smooth 7–dimensional manifold.

Definition 2.31 The manifold N is called an *asymptotically conical G_2 manifold* with cone C and rate $\nu < 0$ if all of the following holds:

- The manifold N is a G_2 manifold with torsion-free G_2 structure φ_N and metric g_N .
- There is a G_2 cone (C, φ_C, g_C) with link Σ .
- There is a compact subset $L \subset N$, and there exists an $R > 1$ and a smooth function $h: (R, \infty) \times \Sigma \rightarrow N$ that is a *diffeomorphism* of $(R, \infty) \times \Sigma$ onto $N \setminus L$.
- The pullback $h^*(\varphi_N)$ is a torsion-free G_2 structure on the subset $(R, \infty) \times \Sigma$ of C . We require that this approach the torsion-free G_2 structure φ_C in a C^∞ sense, with rate $\nu < 0$. This means that

$$(56) \quad |\nabla_C^j (h^*(\varphi_N) - \varphi_C)|_{g_C} = O(r^{\nu-j}) \quad \forall j \geq 0$$

in $(R, \infty) \times \Sigma$. Note that all norms and derivatives are computed using the cone metric g_C .

This should be compared to [Definition 2.25](#). An asymptotically conical G_2 manifold also has holonomy exactly equal to G_2 , by the same argument as in the ICS case. Also, an asymptotically conical G_2 manifold always has only one asymptotic end. This follows from the *Cheeger–Gromoll splitting theorem*, which says that a complete noncompact Ricci-flat manifold with more than one end isometrically splits into a Riemannian product, and hence if we had more than one end, the holonomy would be reducible. Hence the link Σ of the asymptotic cone of N must be connected. This is why we defined the link of a G_2 cone to be connected in [Section 2.1](#). See Besse [\[5\]](#) for more details.

Since the metric g_N and the 4-form ψ_N are pointwise smooth functions of φ_N , by Taylor's theorem we also have

$$(57) \quad \begin{aligned} |\nabla_C^j (h^*(g_N) - g_C)|_{g_C} &= O(r^{\nu-j}) & \forall j \geq 0, \\ |\nabla_C^j (h^*(\psi_N) - \psi_C)|_{g_C} &= O(r^{\nu-j}) & \forall j \geq 0. \end{aligned}$$

An asymptotically conical G_2 manifold will sometimes be called an AC G_2 manifold for brevity. Figure 2 shows an asymptotically conical manifold.




Figure 2: An asymptotically conical manifold N

Remark 2.32 As in Remark 2.26, it can be shown that if we assume that (56) holds only for $j = 0$ and $j = 1$, then there exists a *particular* diffeomorphism h that satisfies (56) for all $j \geq 0$.

There are three known examples of asymptotically conical G_2 manifolds, whose asymptotic cones have links given by the strictly nearly Kähler manifolds of Examples 2.5, 2.6 and 2.7, respectively. They are all total spaces of vector bundles over a compact base. They were discovered by Bryant–Salamon [8] and were the first examples of complete G_2 manifolds.

Example 2.33 (The bundle $\Lambda_-^2(S^4)$ of anti-self-dual 2-forms over the 4-sphere) This is a nontrivial rank 3 vector bundle over the standard round S^4 . This AC G_2 manifold is asymptotic to the cone over the non-Kähler $\mathbb{C}\mathbb{P}^3$ of Example 2.5, with rate $\nu = -4$.

Example 2.34 (The bundle $\Lambda_-^2(\mathbb{C}\mathbb{P}^2)$ of anti-self-dual 2-forms over the complex projective plane) This is a nontrivial rank 3 vector bundle over the standard Fubini–Study $\mathbb{C}\mathbb{P}^2$. This AC G_2 manifold is asymptotic to the cone over the complex flag manifold $F_{1,2}$ of Example 2.6, also with rate $\nu = -4$.

Example 2.35 (The spinor bundle $\mathcal{S}(S^3)$ of the 3–sphere) This is a trivial rank 4 vector bundle over the standard round S^3 , hence is topologically $S^3 \times \mathbb{R}^4$. This AC G_2 manifold is asymptotic to the cone over the nearly Kähler $S^3 \times S^3$ of [Example 2.7](#), with rate $\nu = -3$.

Remark 2.36 Explicit formulas for the asymptotically conical G_2 structures of [Examples 2.33](#), [2.34](#) and [2.35](#), as well as the fact that their rates are -4 , -4 and -3 , respectively, can be found in Bryant–Salamon [\[8\]](#) and also in Atiyah–Witten [\[1\]](#). We will not have need for these explicit formulas.

Next we discuss the AC analogue of [Proposition 2.27](#), which is *different* in a very important way which will lead to topological obstructions to our desingularization procedure.

Proposition 2.37 *The two closed 3–forms $h^*(\varphi_N)$ and φ_C on $(R, \infty) \times \Sigma$ represent the same cohomology class in $H^3((R, \infty) \times \Sigma, \mathbb{R}) \cong H^3(\Sigma, \mathbb{R})$ if the rate ν satisfies $\nu < -3$. Similarly the two closed 4–forms $h^*(\psi_N)$ and ψ_C on $(R, \infty) \times \Sigma$ represent the same cohomology class in $H^4((R, \infty) \times \Sigma, \mathbb{R}) \cong H^4(\Sigma, \mathbb{R})$ if the rate ν satisfies $\nu < -4$.*

Proof This follows immediately from $|(h^*(\varphi_M) - \varphi_C)|_{g_C} = O(r^\nu)$ and $|(h^*(\psi_M) - \psi_C)|_{g_C} = O(r^\nu)$, using [Lemma 2.11](#). \square

Therefore, in contrast to the case of G_2 manifolds with ICS as discussed in [Proposition 2.27](#), we *cannot* conclude that the cohomology classes $[h^*(\varphi_N)]$ and $[h^*(\psi_N)]$ are trivial in $H^3(\Sigma, \mathbb{R})$ and $H^4(\Sigma, \mathbb{R})$, respectively, and indeed in general when the rate ν is not sufficiently negative, they will not be. This will introduce some *obstructions* to our gluing procedure, which are discussed in [Section 3.2](#). These observations lead us to make the following definition.

Definition 2.38 For an AC G_2 manifold $(N, \varphi_N, \psi_N, g_N)$, we define the cohomological invariants $\Phi(N) \in H^3(\Sigma, \mathbb{R})$ and $\Psi(N) \in H^4(\Sigma, \mathbb{R})$ to be the cohomology classes $[h^*(\varphi_N)]$ and $[h^*(\psi_N)]$, respectively. By [Proposition 2.37](#), $\Phi(N) = 0$ if the rate $\nu < -3$ and $\Psi(N) = 0$ if $\nu < -4$.

Remark 2.39 For the Bryant–Salamon examples of asymptotically conical G_2 manifolds described in [Examples 2.33](#), [2.34](#) and [2.35](#), *only one* of these invariants can be (and is) nonzero in each case, by [Remark 2.8](#) and [Definition 2.38](#). For the spaces $\Lambda_-^2(S^4)$ and $\Lambda_-^2(\mathbb{C}P^2)$, we have $\Phi(N) = 0$ and $\Psi(N) \neq 0$, whereas for the space $\mathcal{S}(S^3)$ we have $\Phi(N) \neq 0$ and $\Psi(N) = 0$.

3 Desingularization of compact G_2 manifolds with ICS

Suppose M is a compact G_2 manifold with isolated conical singularities x_1, \dots, x_n , cones C_1, \dots, C_n and rates $\mu_1, \dots, \mu_n > 0$. Assume also that we have asymptotically conical G_2 manifolds N_1, \dots, N_n , with the same cones C_1, \dots, C_n and rates $\nu_1, \dots, \nu_n \leq -3$. (We will see later why we need to assume that each $\nu_i \leq -3$ rather than just $\nu_i < 0$.) We want to *desingularize* M to obtain a *smooth* compact G_2 manifold. The idea is to cut out a neighbourhood of each singularity x_i and *glue* in N_i . In this way we obtain a smooth compact manifold \tilde{M} , and then we need to show using analysis that \tilde{M} admits a torsion-free G_2 structure.

The main tool that we will require is the following theorem of Joyce, which says that if one can find a closed G_2 structure φ on a compact manifold M whose torsion is sufficiently small, then there exists a *torsion-free* G_2 structure $\tilde{\varphi}$ on M which is *close* to φ in some sense. It was used by Joyce in [17] to construct the first compact examples of manifolds with G_2 holonomy. (Another gluing construction of smooth compact G_2 manifolds is due to Kovalev [27].) The precise statement of Joyce’s theorem is as follows.

Theorem 3.1 (Joyce [18, Theorem 11.6.1]) *Let κ, D_1, D_2 and D_3 be any positive constants. Then there exists $s_0 \in (0, 1]$ and $D_4 > 0$, such that whenever $0 < s \leq s_0$, the following holds:*

Let M be a smooth compact 7–manifold, with G_2 structure φ and associated metric g satisfying $d\varphi = 0$. Suppose there is a smooth 3–form χ on M satisfying $d_g^ \chi = d_g^* \varphi$ such that:*

- i) $\|\chi\|_{C^0} \leq D_1 s^\kappa, \quad \|\chi\|_{L^2} \leq D_1 s^{7/2+\kappa}, \quad \|d_g^* \chi\|_{L^{14}} \leq D_1 s^{-1/2+\kappa}.$
- ii) *The injectivity radius $\mathcal{I}(g)$ satisfies $\mathcal{I}(g) \geq D_2 s$.*
- iii) *The Riemann curvature $\mathcal{R}(g)$ satisfies $\|\mathcal{R}(g)\|_{C^0} \leq D_3 s^{-2}$.*

Then there exists a smooth, torsion-free G_2 structure $\tilde{\varphi}$ with metric \tilde{g} on M and such that:

- $\|\varphi - \tilde{\varphi}\|_{C^0} \leq D_4 s^\kappa.$
- $[\varphi] = [\tilde{\varphi}]$ in $H^3(M, \mathbb{R})$.

Here all norms are computed using the original metric g .

Remark 3.2 The exponents in i) of [Theorem 3.1](#) are the best possible for the theorem to be true and are improvements to those presented in [Theorem 11.6.1](#) of [Joyce \[18\]](#). See the discussion at the bottom of page 296 of [Joyce \[18\]](#) for more details.

In [Section 3.1](#) we construct a one parameter family \widetilde{M}_s of compact, smooth 7-manifolds for small $s > 0$. These manifolds will be actually all diffeomorphic to each other, so we really have just one compact smooth manifold \widetilde{M} up to diffeomorphism. Nevertheless it will be convenient to consider each \widetilde{M}_s separately. In [Section 3.2](#) we construct a closed G_2 structure φ_s on \widetilde{M}_s , and in [Section 3.3](#) we show that for s sufficiently small, there exists a torsion-free G_2 structure $\widetilde{\varphi}_s$ on \widetilde{M}_s close to φ_s .

3.1 Construction of the smooth compact manifolds M_s

By letting $R = \max(R_1, \dots, R_n)$, we can assume that the parameter R in [Definition 2.31](#) is the same for all the AC G_2 manifolds N_1, \dots, N_n . For now we will consider only those $s > 0$ which are small enough so that $2sR < \varepsilon$. Later, in [Section 3.2](#), we will need to further restrict the values of s .

We first apply a *homothety* (scaling) to each asymptotically conical G_2 manifold $(N_i, \varphi_{N_i}, \psi_{N_i}, g_{N_i})$ as follows. We have a diffeomorphism $h_i: (R, \infty) \rightarrow N_i \setminus L_i$, such that $h_i^*(\varphi_{N_i}) = \varphi_{C_i} + O(r^{\nu_i})$ as $r \rightarrow \infty$. For a fixed constant s , we define

$$(58) \quad N_{i,s} = N_i, \quad \varphi_{N_{i,s}} = s^3 \varphi_{N_i}, \quad \psi_{N_{i,s}} = s^4 \psi_{N_i}, \quad g_{N_{i,s}} = s^2 g_{N_i}.$$

It is clear that $(N_{i,s}, \varphi_{N_{i,s}}, \psi_{N_{i,s}}, g_{N_{i,s}})$ is again a G_2 manifold, as we have simply scaled the G_2 structure by a constant. We claim that $(N_{i,s}, \varphi_{N_{i,s}}, \psi_{N_{i,s}}, g_{N_{i,s}})$ is still asymptotically conical with the same asymptotic G_2 cone $(C_i, \varphi_{C_i}, \psi_{C_i}, g_{C_i})$ and the same rate ν_i . To see this, define $h_{i,s}: (sR, \infty) \times \Sigma_i \rightarrow N_i \setminus L_i$ by $h_{i,s}(r, \sigma) = h_i(s^{-1}r, \sigma)$. That is, $h_{i,s} = h_i \circ (s^{-1})$, where s^{-1} is a dilation as defined in [\(19\)](#). Then we have

$$(59) \quad \begin{aligned} h_{i,s}^*(\varphi_{N_{i,s}}) - \varphi_{C_i} &= (s^{-1})^* \circ h_i^*(s^3 \varphi_{N_i}) - \varphi_{C_i} \\ &= s^3 (s^{-1})^*(h_i^*(\varphi_{N_i})) - s^3 (s^{-1})^*(\varphi_{C_i}) = s^3 (s^{-1})^*(h_i^*(\varphi_{N_i}) - \varphi_{C_i}) \end{aligned}$$

where we have used the fact that φ_{C_i} is a dilation-equivariant 3-form. Now using [\(21\)](#), we see that

$$|(h_{i,s}^*(\varphi_{N_{i,s}}) - \varphi_{C_i})|_{g_{C_i}} = O(r^{\nu_i}),$$

and similarly

$$|\nabla_C^j (h_{i,s}^*(\varphi_{N_{i,s}}) - \varphi_{C_i})|_{g_{C_i}} = O(r^{\nu_i - j}) \quad \forall j \geq 0,$$

as claimed. Essentially, all we have done here is to rescale each asymptotically conical G_2 manifold N_i so that the value R where $N_i \setminus L_i$ resembles $(R, \infty) \times \Sigma_i$ is changed to sR . By taking s small enough so that $sR < \varepsilon$, we will be able to identify the subset $(2sR, \varepsilon) \times \Sigma_i$ of $N_{i,s}$ with the subset $(2sR, \varepsilon) \times \Sigma_i$ of a neighbourhood of the singularity x_i of M in order to be able to glue the two manifolds together. This will all be made more precise now.

For each asymptotically conical G_2 manifold $N_{i,s}$, we want to keep the compact part L_i together with a little bit of the part which looks like a cone. Specifically, for each $i = 1, \dots, n$, define

$$(60) \quad P_{i,s} = L_i \cup h_{i,s} \left((sR', \varepsilon) \times \Sigma_i \right),$$

where $R' = 2R$. The reason for using $2R$ rather than R will be apparent in Section 3.2. See Figure 3 for a picture of $P_{i,s}$. Similarly for the G_2 manifold M with isolated




Figure 3: The region $P_{i,s}$ of the scaled AC manifold $N_{i,s}$

conical singularities, we want to keep the compact part $K = M' \setminus \bigsqcup_{i=1}^n S_i$ together with a little bit of each S_i which looks like a cone. Specifically, we define

$$(61) \quad Q_s = K \cup \bigsqcup_{i=1}^n f_i \left((sR', \varepsilon) \times \Sigma_i \right).$$

See Figure 4 for a picture of the part of Q_s near x_i . The exponent γ in both Figures 3 and 4 will be explained in Section 3.2. Taking the composition $h_{i,s} \circ f_i^{-1}$ of the two diffeomorphisms gives an identification which allows us to define the i -th overlapping region $U_{i,s}$ by

$$(62) \quad U_{i,s} = (sR', \varepsilon) \times \Sigma_i \cong f_i \left((sR', 2\varepsilon) \times \Sigma_i \right) \cong h_{i,s} \left((sR', 2\varepsilon) \times \Sigma_i \right).$$




Figure 4: Part of the region Q_s of the manifold M with ICS

We use this identification to perform our gluing, by defining the *smooth, compact* 7-manifold \tilde{M}_s to be

$$(63) \quad \tilde{M}_s = \left(\bigsqcup_{i=1}^n (P_{i,s} \setminus U_{i,s}) \right) \sqcup \left(Q_s \setminus \bigsqcup_{i=1}^n U_{i,s} \right) \sqcup \left(\bigsqcup_{i=1}^n U_{i,s} \right).$$

This defines \tilde{M}_s as a smooth manifold. See Figure 5 for a picture of \tilde{M}_s . It is clear that for all small s the \tilde{M}_s 's are diffeomorphic. Essentially, we have done the topology of the desingularization, and now we need to do the geometry. The next step is to construct a closed G_2 structure φ_s on \tilde{M}_s with small torsion. This is where the rates μ_i and ν_i will be important.

3.2 Construction of the G_2 structure φ_s on M_s

We will now construct a closed G_2 structure φ_s on \tilde{M}_s , by patching together the torsion-free G_2 structures on M' and the N_i 's. Strictly speaking, we will actually construct a pair (φ_s, ψ_s) , which are nondegenerate closed 3-forms and 4-forms on \tilde{M}_s , respectively, but ψ_s will not equal $*_{g_{\varphi_s}} \varphi_s$, although it will be “close” in a sense to be made precise, when s is sufficiently small. It is from this pair that we will be able to measure the torsion of φ_s . We will see that sometimes the construction of the pair (φ_s, ψ_s) will only be possible if certain topological conditions are satisfied. Finally in Section 3.3 we will show that for s sufficiently small, φ_s has small enough torsion to apply Theorem 3.1.




Figure 5: The smooth compact manifold \tilde{M}_s , for a fixed small s

In this section we will require two analytic results, Theorems 3.6 and 3.10, whose proofs will be postponed until Sections 6 and 5, respectively, as the proofs require Fredholm theory on weighted Sobolev spaces and are best treated separately.

The basic idea behind the construction of the pair (φ_s, ψ_s) on \tilde{M}_s is that we need to smoothly interpolate between the pair $(\varphi_{N_{i,s}}, \psi_{N_{i,s}})$ on $N_{i,s}$ and the pair (φ_M, ψ_M) for each $i = 1, \dots, n$. To be able to do this, we first need a good asymptotic expansion of the G_2 structure near infinity for an asymptotically conical G_2 manifold.

Definition 3.3 Let N_i be an AC G_2 manifold, with diffeomorphism $h_i: (R, \infty) \times \Sigma_i \rightarrow N_i \setminus L_i$. We say that h_i satisfies the *gauge-fixing condition* if $h_i^*(\varphi_{N_i}) - \varphi_{C_i}$ lies in Ω_{27}^3 with respect to φ_{C_i} .

Remark 3.4 The reason for making this definition is the following. One can show that if the gauge-fixing condition is satisfied, the forms $h_i^*(\varphi_N) - \varphi_{C_i}$ and $h_i^*(\psi_N) - \psi_{C_i}$ satisfy *elliptic* equations, and as such they possess improved regularity. This will

become apparent in Section 6. In the sequel [24] we show that one can always find a diffeomorphism h_i for which the gauge-fixing condition is satisfied.

Remark 3.5 For the Bryant–Salamon examples of asymptotically conical G_2 manifolds, given by Examples 2.33, 2.34 and 2.35, it is trivial to verify that one can always find a diffeomorphism for which the gauge-fixing condition is satisfied. This is because these examples are *cohomogeneity one*, which means the G_2 structure is described by functions of one variable. The gauge-fixing condition is then equivalent to a change of variables (on the asymptotic end) satisfying a linear ordinary differential equation, which can be integrated exactly.

For our purposes, the importance of the gauge-fixing condition is the following theorem.

Theorem 3.6 *Suppose that N_i is an asymptotically conical G_2 manifold with rate $\nu_i \leq -3$ and that h_i satisfies the gauge-fixing condition given in Definition 3.3. Then on the subset $(2R, \infty) \times \Sigma_i$ of the cone C_i we can write*

$$(64) \quad h_i^*(\varphi_{N_i}) = \varphi_{C_i} + \xi_i + d\zeta_i,$$

$$(65) \quad h_i^*(\psi_{N_i}) = \psi_{C_i} + \eta_i - *_{C_i}\xi_i + d\theta_i,$$

where ξ_i is a harmonic 3-form, homogeneous of order -3 , and in Ω_{27}^3 with respect to φ_{C_i} , η_i is a harmonic 4-form, homogeneous of order -4 , and ζ_i and θ_i are 2-forms and 3-forms on $(2R, \infty) \times \Sigma_i$, respectively, satisfying

$$(66) \quad |\nabla_{C_i}^j \xi_i|_{g_{C_i}} = O(r^{\nu_i+1-j}), \quad |\nabla_{C_i}^j \theta_i|_{g_{C_i}} = O(r^{\nu_i+1-j}), \quad \forall j \geq 0,$$

where $\nu_i' = -4$. Furthermore, $[\xi_i] = \Phi(N_i)$ and $[\eta_i] = \Psi(N_i)$, where $\Phi(N_i)$ and $\Psi(N_i)$ are the cohomological invariants of the AC G_2 manifold N_i from Definition 2.38.

Theorem 3.6 is proved in Section 6. From now on we will assume that all of our AC G_2 manifolds N_1, \dots, N_n have been gauge-fixed, so the asymptotic expansions in (64) and (65) apply. As noted in Remark 3.5, this can always be done for all the known examples of asymptotically conical G_2 manifolds.

Remark 3.7 It may seem perverse to write ν_i' instead of its exact value -4 , but we do so for the following reason. We want to emphasize that ν_i' is a decay rate at infinity for the left over parts to the asymptotically conical G_2 structure $(\varphi_{N_i}, \psi_{N_i})$, after subtracting topological components. In fact our gluing construction will actually succeed for any $\nu_i' < -\frac{7}{2}$, of which $\nu_i' = -4$ is just a special case which we can achieve using Theorem 3.6 when the gauge-fixing condition is satisfied. One may be able to

avoid the need to satisfy the hypotheses for [Theorem 3.6](#) in certain explicit examples, if one can show directly that $v'_i < -\frac{7}{2}$. See also [Remark 3.26](#).

Remark 3.8 It is because we must use [Theorem 3.6](#) which requires each $v_i \leq -3$, that we need to make this restriction on the v_i 's. If some $v_{i_0} \in (-3, 0)$, then the question of whether or not the gluing construction could be possible would depend on the spectrum of the Laplacian on the particular link Σ_{i_0} . Therefore, only if all $v_i \leq -3$ can we hope to be able to prove a general result like our [Theorem 3.28](#). See also [Remark 6.6](#).

Using the scaling in [\(58\)](#) and the fact that ξ_i and η_i are dilation-invariant (see [Remark 2.13](#)), we can now argue just as in [Equation \(59\)](#) to obtain

$$(67) \quad h_{i,s}^*(\varphi_{N_{i,s}}) = \varphi_{C_i} + s^3 \xi_i + d\zeta_{i,s},$$

$$(68) \quad h_{i,s}^*(\psi_{N_{i,s}}) = \psi_{C_i} + s^4 \eta_i - s^3 *_C \xi_i + d\theta_{i,s},$$

where $\zeta_{i,s}(r, \sigma) = s^3 \zeta_i(s^{-1}r, \sigma)$ and $\theta_{i,s}(r, \sigma) = s^4 \theta_i(s^{-1}r, \sigma)$. Now we can compute

$$\begin{aligned} |\nabla_{C_i}^j \zeta_{i,s}(r, \sigma)|_{g_{C_i}(r, \sigma)} &= s^3 |\nabla_{C_i}^j \zeta_i(s^{-1}r, \sigma)|_{g_{C_i}(r, \sigma)} \\ &= s^3 s^{-2-j} |\nabla_{C_i}^j \zeta_i(s^{-1}r, \sigma)|_{g_{C_i}(s^{-1}r, \sigma)} \\ &= s^{1-j} O((s^{-1}r)^{v'_i+1-j}) = s^{-v'_i} O(r^{v'_i+1-j}), \end{aligned}$$

where we have used [\(21\)](#) and the fact that $\nabla_{C_i}^j \zeta_i \in (T^*)^j \otimes \Lambda^2(T^*)$. We can compute similarly for $\theta_{i,s}$ and find that

$$(69) \quad |\nabla_{C_i}^j \zeta_{i,s}|_{g_{C_i}} = s^{-v'_i} O(r^{v'_i+1-j}), \quad |\nabla_{C_i}^j \theta_{i,s}|_{g_{C_i}} = s^{-v'_i} O(r^{v'_i+1-j}), \quad \forall j \geq 0,$$

on $(R', \infty) \times \Sigma_i$, where $R' = 2R$.

Now consider the i -th conical singularity x_i of M . By [Proposition 2.27](#), on $(0, \varepsilon) \times \Sigma_i$ we can write

$$(70) \quad f_i^*(\varphi_M) = \varphi_{C_i} + d\alpha_i,$$

$$(71) \quad f_i^*(\psi_M) = \psi_{C_i} + d\beta_i,$$

where α_i is a 2-form and β_i is a 3-form on $(0, \varepsilon) \times \Sigma_i$, and by [Lemma 2.11](#) we know that

$$(72) \quad |\nabla_{C_i}^j \alpha_i|_{g_{C_i}} = O(r^{\mu_i+1-j}), \quad |\nabla_{C_i}^j \beta_i|_{g_{C_i}} = O(r^{\mu_i+1-j}), \quad \forall j \geq 0.$$

In order to be able to patch together the G_2 structures on M' and the $N_{i,s}$'s smoothly to obtain a G_2 structure φ_s on \widetilde{M}_s , we will arrange that in the i -th overlap region $U_{i,s}$, the forms will be in the same cohomology class, and hence differ by an exact piece.

Comparing equations (67) and (68) with equations (70) and (71), we see that we need to replace the forms φ_M and ψ_M on M' with $\varphi_M + s^3\xi$ and $\psi_M + s^4\eta - s^3 *_M \xi$ that are asymptotic, at the appropriate rate, to $\varphi_{C_i} + s^3\xi_i$ and $\psi_{C_i} + s^4\eta_i - s^3 *_M \xi_i$ near x_i , respectively. In fact, for the construction to work, we need η to be closed but ξ to be both closed *and* coclosed. There are topological obstructions that can prevent this from being possible. To understand this, we need to consider the algebraic topology of the noncompact smooth part M' of a compact G_2 manifold M with isolated conical singularities.

Fix some $r_0 \in (0, \varepsilon)$. The map $\iota_i: \Sigma_i \rightarrow (0, \varepsilon) \times \Sigma_i$ given by $\iota_i(\sigma) = (r_0, \sigma)$ embeds the link Σ_i as a submanifold of $(0, \varepsilon) \times \Sigma_i$, and the pullback map $\iota_i^*: H^k((0, \varepsilon) \times \Sigma_i, \mathbb{R}) \rightarrow H^k(\Sigma_i, \mathbb{R})$ is an isomorphism, which is just restriction on the level of forms. We also have the smooth maps $f_i: (0, \varepsilon) \times \Sigma_i \rightarrow M'$, and thus for each i we get a map $(f_i \circ \iota_i)^*: H^k(M', \mathbb{R}) \rightarrow H^k(\Sigma_i, \mathbb{R})$.

Definition 3.9 For $0 \leq k \leq 7$, define a map $\Upsilon^k: H^k(M', \mathbb{R}) \rightarrow \bigoplus_{i=1}^n H^k(\Sigma_i, \mathbb{R})$ by

$$(73) \quad \Upsilon^k([\omega]) = \bigoplus_{i=1}^n (f_i \circ \iota_i)^*([\omega])$$

for any cohomology class $[\omega] \in H^k(M', \mathbb{R})$. Essentially, at each end we restrict the k -form ω to the link Σ_i and take its cohomology in $H^k(\Sigma_i, \mathbb{R})$.

Theorem 3.10 Let M be a compact G_2 manifold with isolated conical singularities. For each i , let ξ_i and η_i be 3-forms and 4-forms, homogeneous of orders -3 and -4 , respectively, and closed and coclosed on each cone C_i , with respect to g_{C_i} . Suppose that

$$(74) \quad \bigoplus_{i=1}^n [\xi_i] \in \bigoplus_{i=1}^n H^3(\Sigma_i, \mathbb{R}) \text{ lies in the image of } \Upsilon^3: H^3(M', \mathbb{R}) \rightarrow \bigoplus_{i=1}^n H^3(\Sigma_i, \mathbb{R}),$$

$$(75) \quad \bigoplus_{i=1}^n [\eta_i] \in \bigoplus_{i=1}^n H^4(\Sigma_i, \mathbb{R}) \text{ lies in the image of } \Upsilon^4: H^4(M', \mathbb{R}) \rightarrow \bigoplus_{i=1}^n H^4(\Sigma_i, \mathbb{R}),$$

where the maps Υ^k are given in [Definition 3.9](#). Then for $\delta > 0$ sufficiently small, there exists a smooth 3-form ξ and a smooth 4-form η on M' such that

$$(76) \quad \begin{aligned} d\xi &= 0, & d_{g_M}^* \xi &= 0, & d\eta &= 0, \\ |\nabla_{C_i}^j (f_i^*(\xi) - \xi_i)|_{g_{C_i}} &= O(r^{-3+\delta-j}) & \forall j \geq 0, \end{aligned}$$

$$(77) \quad |\nabla_{C_i}^j (f_i^*(\eta) - \eta_i)|_{g_{C_i}} = O(r^{-4+\delta-j}) \quad \forall j \geq 0.$$

Notice that we do not say that η is coclosed. Furthermore, the 3-form ξ is in Λ_{27}^3 with respect to the G_2 structure φ_M .

[Theorem 3.10](#) is proved in [Section 5](#). Assuming this result for the moment, we now continue our construction of the pair (φ_s, ψ_s) .

Corollary 3.11 *Near the i -th singular point x_i , the 3-form $f_i^*(\xi)$ and the 4-form $f_i^*(\eta)$ satisfy*

$$(78) \quad |f_i^*(\xi)|_{g_{C_i}} = O(r^{-3}), \quad |f_i^*(\eta)|_{g_{C_i}} = O(r^{-4}).$$

Proof By [Equation \(76\)](#) we know that $|f_i^*(\xi)|_{g_{C_i}} \leq |\xi_i|_{g_{C_i}} + O(r^{-3+\delta})$, and also $|\xi_i|_{g_{C_i}} = O(r^{-3})$, so $|f_i^*(\xi)|_{g_{C_i}} = O(r^{-3})$. The proof for η is identical using [\(77\)](#). \square

Corollary 3.12 *If $\delta > 0$ is sufficiently small, then near the i -th singular point x_i , the 4-form $f_i^*(*_M \xi)$ and the 4-form $*_{C_i} \xi_i$ satisfy*

$$(79) \quad |f_i^*(*_M \xi) - *_{C_i} \xi_i|_{g_{C_i}} = O(r^{-3+\delta}),$$

and as a result we can say that

$$(80) \quad f_i^*(*_M \xi) = *_{C_i} \xi_i - dE_i,$$

for some smooth 3-form E_i on $(0, \varepsilon) \times \Sigma_i$, where

$$(81) \quad |\nabla_{C_i}^j E_i|_{g_{C_i}} = O(r^{-2+\delta-j}).$$

Proof We compute that

$$\begin{aligned} |f_i^*(*_M \xi) - *_{C_i} \xi_i|_{g_{C_i}} &\leq |f_i^*(*_M \xi) - *_{C_i} (f_i^*(\xi))|_{g_{C_i}} + |*_{C_i} (f_i^*(\xi)) - *_{C_i} \xi_i|_{g_{C_i}} \\ &= |f_i^*(*_M \xi) - *_{C_i} (f_i^*(\xi))|_{g_{C_i}} + |f_i^*(\xi) - \xi_i|_{g_{C_i}} \\ &= O(r^{-3+\mu_i}) + O(r^{-3+\delta}) \end{aligned}$$

where we have used [\(78\)](#) and [Lemma 2.29](#) on the first term and [\(76\)](#) and the fact that $*_{C_i}$ is an isometry with respect to g_{C_i} on the second term. This gives [\(79\)](#) as long

as we choose δ less than each μ_i . Now the 3-form $f_i^*(\ast_M \xi) - \ast_{C_i} \xi_i$ is closed by Theorem 3.10, and so equations (80) and (81) now follow from Lemma 2.11. \square

Remark 3.13 The sign on E_i in (80) is chosen for later convenience.

Recall that if each $\nu_i < -4$, then all the ξ_i 's and η_i 's vanish, so we can just take ξ and η to both be zero. This is the *unobstructed case*. From now on, we will assume that the situation is either unobstructed or that the conditions (74) and (75) are satisfied. By Lemma 2.11 and Equation (76), since $\delta > 0$, we see that for each i there exists a smooth 2-form A_i on $(0, \varepsilon) \times \Sigma_i$ such that

$$(82) \quad f_i^*(\varphi_M + s^3 \xi) = \varphi_{C_i} + s^3 \xi_i + d\alpha_i + s^3 dA_i,$$

where we have also used Equation (70). Similarly using equations (77), (80) and (71), we have that for each i there exists a smooth 3-form B_i on $(0, \varepsilon) \times \Sigma_i$ such that

$$(83) \quad f_i^*(\psi_M + s^4 \eta - s^3 \ast_M \xi) = \psi_{C_i} + s^4 \eta_i - s^3 \ast_{C_i} \xi_i + dB_i + s^4 dB_i + s^3 dE_i.$$

Also, by Lemma 2.11 and equations (72), we have

$$(84) \quad |\nabla_{C_i}^j \alpha_i|_{g_{C_i}} = O(r^{\mu_i+1-j}), \quad |\nabla_{C_i}^j A_i|_{g_{C_i}} = O(r^{-2+\delta-j}), \quad \forall j \geq 0,$$

$$(85) \quad |\nabla_{C_i}^j \beta_i|_{g_{C_i}} = O(r^{\mu_i+1-j}), \quad |\nabla_{C_i}^j B_i|_{g_{C_i}} = O(r^{-3+\delta-j}), \quad \forall j \geq 0.$$

We are now ready to construct a smooth 3-form φ_s and a smooth 4-form ψ_s on \widetilde{M}_s . Let us take $\gamma \in (0, 1)$, which will be chosen later. It will become clear later why we need to do this. From now on we will consider only those values of $s > 0$ which are small enough so that

$$(86) \quad sR' < s^\gamma < 2s^\gamma < \varepsilon.$$

Recall that $R' = 2R$. These inequalities hold for all $s < \max((\varepsilon/2)^{1/\gamma}, (1/R')^{1/(1-\gamma)})$. This enables us to split up each overlapping region $U_{i,s}$ into three disjoint pieces:

$$U_{i,s} = (sR', s^\gamma) \times \Sigma_i \sqcup [s^\gamma, 2s^\gamma] \times \Sigma_i \sqcup (2s^\gamma, \varepsilon) \times \Sigma_i.$$

The reader should refer again to Figures 3 and 4.

We can write the smooth compact manifold \widetilde{M}_s as the (nondisjoint) union of the *inner piece* $K \cup \bigsqcup_{i=1}^n f_i((2s^\gamma, \varepsilon) \times \Sigma_i)$ with an *outer piece* $L_i \cup h_{i,s}((sR', s^\gamma) \times \Sigma_i)$ and an overlapping region $U_{i,s} = (sR', \varepsilon) \times \Sigma_i$ for each $i = 1, \dots, n$. Considering it in this way, the inner piece intersects each $U_{i,s}$ in the set $f_i((2s^\gamma, \varepsilon) \times \Sigma_i) \cong (2s^\gamma, \varepsilon) \times \Sigma_i$, and each inner piece intersects its associated $U_{i,s}$ in the set $h_{i,s}((sR', s^\gamma) \times \Sigma_i) \cong (sR', s^\gamma) \times \Sigma_i$.

Let $u: (0, \infty) \rightarrow \mathbb{R}$ be any smooth increasing function such that

$$u(r) = \begin{cases} 0 & \text{for } 0 < r \leq 1, \\ 1 & \text{for } 2 \leq r < \infty. \end{cases}$$

We will use $u(r)$ to *interpolate* between the pair $(\varphi_M + s^3\xi, \psi_M + s^4\eta - s^3 *_M \xi)$ on M' and the pairs $(\varphi_{N_{i,s}}, \psi_{N_{i,s}})$ on the $N_{i,s}$'s. Define $u_s(r) = u(s^{-\gamma}r)$. Then clearly

$$u_s(r) = \begin{cases} 0 & \text{for } 0 < r \leq s^\gamma, \\ 1 & \text{for } 2s^\gamma \leq r < \infty. \end{cases}$$

Using u_s , we define a smooth 2-form ρ_i and a smooth 3-form τ_i on each $(sR', \varepsilon) \times \Sigma_i$ as follows:

$$(87) \quad \rho_i = (\alpha_i + s^3 A_i)u_s + \zeta_{i,s}(1 - u_s),$$

$$(88) \quad \tau_i = (\beta_i + s^4 B_i + s^3 E_i)u_s + \theta_{i,s}(1 - u_s).$$

Definition 3.14 We define a smooth 3-form φ_s on \widetilde{M}_s as follows:

$$(89) \quad \varphi_s = \begin{cases} \varphi_M + s^3\xi & \text{on } K \cup \bigsqcup_{i=1}^n f_i ((2s^\gamma, \varepsilon) \times \Sigma_i), \\ \varphi_{C_i} + s^3\xi_i + d\rho_i & \text{on } (sR', \varepsilon) \times \Sigma_i, \\ \varphi_{N_{i,s}} & \text{on } L_i \cup h_{i,s} ((sR', s^\gamma) \times \Sigma_i), \end{cases}$$

and we define a smooth 4-form ψ_s on \widetilde{M}_s as follows:

$$(90) \quad \psi_s = \begin{cases} \psi_M + s^4\eta - s^3 *_M \xi & \text{on } K \cup \bigsqcup_{i=1}^n f_i ((2s^\gamma, \varepsilon) \times \Sigma_i), \\ \psi_{C_i} + s^4\eta_i - s^3 *_C \xi_i + d\tau_i & \text{on } (sR', \varepsilon) \times \Sigma_i, \\ \psi_{N_{i,s}} & \text{on } L_i \cup h_{i,s} ((sR', s^\gamma) \times \Sigma_i). \end{cases}$$

We need to check that φ_s is well defined, in the regions where we have given it two definitions. When $r \leq s^\gamma$, then $\rho_i = \zeta_{i,s}$ and therefore by (67) we have $\varphi_s = h_{i,s}^*(\varphi_{N_{i,s}})$ on $(sR', s^\gamma) \times \Sigma_i$ and $\varphi_s = \varphi_{N_{i,s}}$ on $h_{i,s}((sR', s^\gamma) \times \Sigma_i)$, and these are indeed identified with each other under the diffeomorphism $h_{i,s}$. Whereas when $r \geq 2s^\gamma$, then $\rho_i = \alpha_i + s^3 A_i$ and therefore by (82) we have $\varphi_s = f_i^*(\varphi_M + s^3\xi)$ on $(2s^\gamma, \varepsilon) \times \Sigma_i$ and $\varphi_s = \varphi_M + s^3\xi$ on $f_i((2s^\gamma, \varepsilon) \times \Sigma_i)$, and these are identified with each other under the diffeomorphism f_i . Similarly one checks using (68) and (83) that ψ_s is also well defined. It is clear that φ_s and ψ_s are both *closed* forms, for all $s > 0$.

It is intuitively clear that as $s \rightarrow 0$, the manifold with G_2 structure $(\widetilde{M}_s, \varphi_s)$ ‘‘approaches’’ the original compact G_2 manifold (M, φ_M) with isolated conical singulari-

ties. This can be made precise using the notion of Gromov–Hausdorff convergence, but we will not have occasion to use this.

3.3 Existence of the torsion-free G_2 structure $\tilde{\varphi}_s$ on M_s

In this section, we will show that from the pair (φ_s, ψ_s) constructed in Section 3.2, we can satisfy the hypotheses of Joyce’s Theorem 3.1 to conclude that \tilde{M}_s admits a torsion-free G_2 structure $\tilde{\varphi}_s$.

Since nondegeneracy is an open condition, it will follow from the estimates in Proposition 3.20 that for s sufficiently small, φ_s and ψ_s are nondegenerate 3–forms and 4–forms, respectively. However, ψ_s is *not* equal to $*_{g_s}\varphi_s$, where g_s is the metric determined by φ_s , so φ_s is not coclosed with respect to its induced metric, and hence not torsion-free. However, for s sufficiently small, we will see that ψ_s is close to $*_{g_s}\varphi_s$ in a suitable sense, and this difference, together with its derivative, is a measure of the torsion of φ_s .

Definition 3.15 For each s , we define a smooth 3–form χ_s on \tilde{M}_s by

$$(91) \quad \chi_s = \varphi_s - *_{g_s}\psi_s.$$

Remark 3.16 In the notation of Lemma 1.3 and Remark 1.4, we have $\chi_s = \varphi_s - \Theta^{-1}(\psi_s)$.

Lemma 3.17 The form χ_s satisfies $d_{g_s}^*\chi_s = d_{g_s}^*\varphi_s$ for all s .

Proof We compute

$$d_{g_s}^*\chi_s = d_{g_s}^*(\varphi_s - *_{g_s}\psi_s) = d_{g_s}^*\varphi_s - *_{g_s}d*_{g_s}*_{g_s}\psi_s = d_{g_s}^*\varphi_s - *_{g_s}d\psi_s = d_{g_s}^*\varphi_s$$

using the fact that $d_{g_s}^* = *_{g_s}d*_{g_s}$ on 4–forms, that $*_{g_s}^2 = 1$ and that $d\psi_s = 0$. \square

This 3–form χ_s will play the role of χ in Theorem 3.1. To apply this result, we need to estimate various norms of χ_s in the different regions of \tilde{M}_s .

Proposition 3.18 The 3–form χ_s on \tilde{M}_s is given by

$$(92) \quad \chi_s = \begin{cases} -J_{\varphi_M}(s^4\eta) - G_{\varphi_M}(s^4\eta - s^3 *_{M}\xi) & \text{on } K \cup \bigsqcup_{i=1}^n f_i((2s^\nu, \varepsilon) \times \Sigma_i), \\ \begin{bmatrix} d\rho_i - J_{\varphi_{C_i}}(s^4\eta_i + d\tau_i) \\ -G_{\varphi_{C_i}}(s^4\eta_i - s^3 *_{C_i}\xi_i + d\tau_i) \end{bmatrix} & \text{on } [s^\nu, 2s^\nu] \times \Sigma_i, \\ 0 & \text{on } L_i \cup h_{i,s}((sR', s^\nu) \times \Sigma_i), \end{cases}$$

in the different regions used in Definition 3.14. Here the maps J and G are as defined in Remark 1.4 and Equation (11).

Proof On the region $L_i \cup h_{i,s} ((sR', s^\nu) \times \Sigma_i)$, we have $\psi_s = \psi_{N_{i,s}}$ and $\varphi_s = \varphi_{N_{i,s}}$, so

$$\chi_s = \varphi_{N_{i,s}} - \Theta^{-1}(\psi_{N_{i,s}}) = 0.$$

In other words, the G_2 structure φ_s is already torsion-free on these regions. Now on the inner region $K \cup \bigsqcup_{i=1}^n f_i ((2s^\nu, \varepsilon) \times \Sigma_i)$, we have $\psi_s = \psi_M + s^4\eta - s^3 *_M \xi$, and hence by (9) we have

$$\Theta^{-1}(\psi_s) = \varphi_M + J_{\varphi_M}(s^4\eta - s^3 *_M \xi) + G_{\varphi_M}(s^4\eta - s^3 *_M \xi).$$

However, since by [Theorem 3.10](#) we know that ξ is in Ω_{27}^3 with respect to φ_M , [Equation \(11\)](#) tells us that $J_{\varphi_M}(*_M \xi) = -\xi$. Therefore

$$\Theta^{-1}(\psi_s) = \varphi_M + s^3\xi + J_{\varphi_M}(s^4\eta) + G_{\varphi_M}(s^4\eta - s^3 *_M \xi).$$

Now since $\varphi_s = \varphi_M + s^3\xi$, we obtain the expression for χ_s in the first line of (92). Finally, for the overlap regions $[s^\nu, 2s^\nu] \times \Sigma_i$, the calculation is similar, this time using the fact that ξ_i is in Ω_{27}^3 with respect to φ_{C_i} , which was shown in [Proposition 2.23](#). \square

Remark 3.19 Notice in (92) we are now writing the overlap regions as $[s^\nu, 2s^\nu] \times \Sigma_i$ as opposed to $(sR', \varepsilon) \times \Sigma_i$ as was used in [Definition 3.14](#), so that now all the regions are disjoint. This will make is easier to estimate various norms in what follows, so we will decompose \widetilde{M}_s this way from now on.

Note that in the unobstructed case, we have $\xi = 0$ and $\eta = 0$. Thus χ_s also vanishes on the inner region, so φ_s only fails to be torsion-free in the annuli $[s^\nu, 2s^\nu] \times \Sigma_i$ where we have interpolated between two torsion-free G_2 structures. However in the obstructed case, the G_2 structure φ_s also has torsion on the inner region coming from the corrections ξ and η to φ_M and ψ_M .

We are now ready to begin estimating various norms of χ_s in order to apply [Theorem 3.1](#). We begin with estimates of $d\rho_i$ and $d\tau_i$.

Proposition 3.20 *Consider the 2-forms ρ_i and the 3-forms τ_i given by equations (87) and (88). In the region $s^\nu \leq r \leq 2s^\nu$, and with respect to the cone metric g_{C_i} , we have the following estimates:*

$$(93) \quad |d\rho_i|_{g_{C_i}} \leq C(s^{\nu\mu_i} + s^{-\nu'_i(1-\nu)} + s^{3(1-\nu)+\delta\nu}),$$

$$(94) \quad |\nabla_{C_i} d\rho_i|_{g_{C_i}} \leq C(s^{\nu\mu_i-\nu} + s^{-\nu'_i(1-\nu)-\nu} + s^{3(1-\nu)+\delta\nu-\nu}),$$

$$(95) \quad |d\tau_i|_{g_{C_i}} \leq C(s^{\nu\mu_i} + s^{-\nu'_i(1-\nu)} + s^{3(1-\nu)+\delta\nu}),$$

$$(96) \quad |\nabla_{C_i} d\tau_i|_{g_{C_i}} \leq C(s^{\nu\mu_i-\nu} + s^{-\nu'_i(1-\nu)-\nu} + s^{3(1-\nu)+\delta\nu-\nu}),$$

where each C denotes some positive constant.

Proof Notice that $\frac{d}{dr}u_s(r) = s^{-\gamma}u'_s(r)$. Now from (87), we see that

$$(97) \quad d\rho_i = s^{-\gamma}u'_s dr \wedge (\alpha_i + s^3 A_i) + u_s(d\alpha_i + s^3 dA_i) - s^{-\gamma}u'_s dr \wedge \zeta_{i,s} + (1 - u_s)d\zeta_{i,s}.$$

Since u_s and u'_s are uniformly bounded functions of r (independent of s), we have

$$|d\rho_i|_{g_{C_i}} \leq C s^{-\gamma} |\alpha_i + s^3 A_i|_{g_{C_i}} + C |d\alpha_i + s^3 dA_i|_{g_{C_i}} + C s^{-\gamma} |\zeta_{i,s}|_{g_{C_i}} + C |d\zeta_{i,s}|_{g_{C_i}}.$$

Using equations (1), (84) and (69), we compute

$$|d\rho_i|_{g_{C_i}} \leq C(s^{-\gamma}r^{\mu_i+1} + s^{3-\gamma}r^{-2+\delta} + r^{\mu_i} + s^3r^{-3+\delta} + s^{-\gamma}s^{-\nu'_i}r^{\nu'_i+1} + s^{-\nu'_i}r^{\nu'_i}).$$

Since $s^\gamma \leq r \leq 2s^\gamma$, we have $r^a \leq C s^{\gamma a}$ for any $a > 0$. Therefore the above expression becomes

$$|d\rho_i|_{g_{C_i}} \leq C(s^{\gamma\mu_i} + s^{3(1-\gamma)+\delta\gamma} + s^{\gamma\mu_i} + s^{3(1-\gamma)+\delta\gamma} + s^{-\nu'_i(1-\gamma)} + s^{-\nu'_i(1-\gamma)}),$$

which is (93). To obtain (94), take the covariant derivative of (97) and proceed as before.

For the estimates on τ_i , Equation (88) shows that

$$(98) \quad d\tau_i = s^{-\gamma}u'_s dr \wedge (\beta_i + s^4 B_i + s^3 E_i) + u_s(d\beta_i + s^4 dB_i + s^3 dE_i) - s^{-\gamma}u'_s dr \wedge \theta_{i,s} + (1 - u_s)d\theta_{i,s}.$$

Now we use equations (1), (85), (81) and (69), and get

$$|d\tau|_{g_{C_i}} \leq C(s^{\gamma\mu_i} + s^{4(1-\gamma)+\delta\gamma} + s^{3(1-\gamma)+\delta\gamma} + s^{-\nu'_i(1-\gamma)}),$$

where the $O(s^{4(1-\gamma)+\delta\gamma})$ terms come from B_i and the $O(s^{3(1-\gamma)+\delta\gamma})$ terms come from E_i . But since $s < 1$, the former terms are smaller than, and can be absorbed by, the latter terms. This gives (95). The proof of (96) is similar. \square

Remark 3.21 The estimates in Proposition 3.20 explain why we needed to introduce the parameter $\gamma \in (0, 1)$ to construct φ_s and ψ_s . If, instead of (86), we had divided the interval (sR', ε) into $sR' < R_1 < R_2 < \varepsilon$, which corresponds to $\gamma = 0$, or into $sR' < sR_1 < sR_2 < s\varepsilon$, which corresponds to $\gamma = 1$, then the estimates of various norms of χ_s would include terms of size $O(1)$, and they would not be small enough for us to apply Theorem 3.1 later.

In Theorem 3.1, all the norms are measured with respect to the metric g_s induced by φ_s . However, it is easier for us to estimate various terms using the metrics g_M from φ_M or g_{C_i} from φ_{C_i} . To translate these to estimates using g_s , we need the following observation.

Lemma 3.22 *For s sufficiently small, the metric g_s is uniformly close to the metric g_M (in the g_M metric) on the inner region $K \cup \bigsqcup_{i=1}^n f_i ((2s^\gamma, \varepsilon) \times \Sigma_i)$. This means that on this region*

$$\sup |g_M - g_s|_{g_M} \leq e, \quad \sup |g_M^{-1} - g_s^{-1}|_{g_M} \leq e,$$

for some small $e < \frac{1}{2}$. As a result, it follows easily that for any tensor ω defined on the inner region,

$$(99) \quad |\omega|_{g_s} \leq C|\omega|_{g_M}$$

for some $C > 0$, independent of s and ω .

Similarly for s sufficiently small, the metric g_s is uniformly close to the metric g_{C_i} (in the g_{C_i} metric) on the overlap region $[s^\gamma, \varepsilon) \times \Sigma_i$, and hence

$$(100) \quad |\omega|_{g_s} \leq C|\omega|_{g_{C_i}}.$$

Proof On the inner region $K \cup \bigsqcup_{i=1}^n f_i ((2s^\gamma, \varepsilon) \times \Sigma_i)$, from [Definition 3.14](#) we have $\varphi_s - \varphi_M = s^3\xi$. On the subset K , which is a compact manifold with boundary, $|\xi|_{g_M}$ is bounded, so $s^3|\xi|_{g_M} = O(s^3)$. Meanwhile on the subset $\bigsqcup_{i=1}^n f_i ((2s^\gamma, \varepsilon) \times \Sigma_i)$, we claim that $s^3|\xi|_{g_M} = O(s^{3(1-\gamma)})$. To see this, first we note that by [\(54\)](#), we have $|f_i^*(g_M) - g_{C_i}|_{g_{C_i}} = O(r^{\mu_i})$, which is uniformly bounded as $r \leq \varepsilon$. Therefore

$$|\xi|_{g_M} = |f_i^*(\xi)|_{f_i^*(g_M)} \leq C|f_i^*(\xi)|_{g_{C_i}}$$

for some $C > 0$. But by [\(76\)](#) we know that

$$|f_i^*(\xi)|_{g_{C_i}} \leq |\xi_i|_{g_{C_i}} + O(r^{-3+\delta}),$$

and also $|\xi_i|_{g_{C_i}} = O(r^{-3})$, so $s^3|\xi|_{g_M} \leq Cs^3r^{-3} \leq Cs^{3(1-\gamma)}$ as claimed, since $r^{-1} \leq s^{-\gamma}$. Therefore on the inner region we have

$$\sup |\varphi_s - \varphi_M|_{g_M} \leq C(s^3 + s^{3(1-\gamma)})$$

which can be made as small as we want by taking s sufficiently small. Therefore we can make $|g_s - g_M|_{g_M} < e$ for some $e < \frac{1}{2}$. The claim now follows.

On the region $[s^\gamma, \varepsilon) \times \Sigma_i$, we have from [Definition 3.14](#) that $\varphi_s - \varphi_{C_i} = s^3\xi_i + d\rho_i$. Now $|\xi_i|_{g_{C_i}} = O(r^{-3})$, so $s^3|\xi_i|_{g_{C_i}} \leq Cs^{3(1-\gamma)}$ since $r^{-1} \leq Cs^{-\gamma}$. Using [\(93\)](#), we have

$$\sup |\varphi_s - \varphi_{C_i}|_{g_{C_i}} \leq C(s^{3(1-\gamma)} + s^{\gamma\mu_i} + s^{-\nu_i'(1-\gamma)} + s^{3(1-\gamma)+\delta\gamma}),$$

and now the claim follows as in the first case. □

We will use (99) and (100) repeatedly in what follows, without explicit mention. We can now estimate the pointwise norms of χ_s and $\nabla_s \chi_s$, where ∇_s is the covariant derivative with respect to the metric g_s of φ_s .

Proposition 3.23 *Let χ_s be as given in Proposition 3.18. Then we have the following pointwise estimates for $|\chi_s|_{g_s}$ and $|\nabla_s \chi_s|_{g_s}$, on the different regions that were used in (92). On the subset K of the inner region, we have*

$$(101) \quad |\chi_s|_{g_s} \leq Cs^4, \quad |\nabla_s \chi_s|_{g_s} \leq Cs^4.$$

On each subset $f_i((2s^\nu, \varepsilon) \times \Sigma_i)$ of the inner region, we have

$$(102) \quad |\chi_s|_{g_s} \leq Cs^4 r^{-4}, \quad |\nabla_s \chi_s|_{g_s} \leq Cs^4 r^{-4}.$$

On the overlap regions $[s^\nu, 2s^\nu] \times \Sigma_i$ we have

$$(103) \quad \begin{aligned} |\chi_s|_{g_s} &\leq C(s^{\nu\mu_i} + s^{-\nu'_i(1-\nu)} + s^{3(1-\nu)+\delta\nu}), \\ |\nabla_s \chi_s|_{g_s} &\leq C(s^{\nu\mu_i-\nu} + s^{-\nu'_i(1-\nu)-\nu} + s^{3(1-\nu)+\delta\nu-\nu}), \end{aligned}$$

and finally on the outer regions $L_i \cup h_{i,s}((sR', s^\nu) \times \Sigma_i)$ we have

$$(104) \quad |\chi_s|_{g_s} = 0, \quad |\nabla_s \chi_s|_{g_s} = 0.$$

Proof We start with the inner region. Using (92) and (12) and (10), we see that

$$(105) \quad \begin{aligned} |\chi_s|_{g_s} &\leq C|\chi_s|_{g_M} \leq C|J_{\varphi_M}(s^4\eta)|_{g_M} + C|G_{\varphi_M}(s^4\eta - s^3 *_M \xi)|_{g_M} \\ &\leq Cs^4|\eta|_{g_M} + C|s^4\eta - s^3 *_M \xi|_{g_M}^2 \\ &\leq Cs^4|\eta|_{g_M} + C(s^4|\eta|_{g_M} + s^3|\xi|_{g_M})^2, \end{aligned}$$

where we have used the fact that $*_M$ is an isometry. A similar computation yields

$$(106) \quad \begin{aligned} |\nabla_s \chi_s|_{g_s} &\leq Cs^4|\nabla_M \eta|_{g_M} \\ &\quad + C(s^4|\nabla_M \eta|_{g_M} + s^3|\nabla_M \xi|_{g_M})(s^4|\eta|_{g_M} + s^3|\xi|_{g_M}). \end{aligned}$$

To obtain the above estimate we need to use the fact that ∇_M commutes with $*_M$ and that φ_M is torsion-free, so that the $|\nabla_M \varphi_M|_{g_M}$ terms in (12) and (10) are zero. Now the subset K of the inner region is a compact manifold with boundary, so $|\xi|_{g_M}$, $|\eta|_{g_M}$, $|\nabla_M \xi|_{g_M}$ and $|\nabla_M \eta|_{g_M}$ are all bounded there. Hence by (105) we have

$$|\chi_s|_{g_s} \leq C(s^4 + s^6 + s^7 + s^8) \leq Cs^4$$

by absorbing smaller terms into the dominant term. Similarly by (106) we get $|\nabla_s \chi_s|_{g_s} \leq Cs^4$. This proves (101). Now consider a subset $f_i((2s^\nu, \varepsilon) \times \Sigma_i)$ of the inner region. By (54), we have $|f_i^*(g_M) - g_{C_i}|_{g_{C_i}} = O(r^{\mu_i})$, which is uniformly

bounded as $r \leq \varepsilon$. Therefore

$$|\xi|_{g_M} = |f_i^*(\xi)|_{f_i^*(g_M)} \leq C|f_i^*(\xi)|_{g_{C_i}} \leq C|\xi_i|_{g_{C_i}} + O(r^{-3+\delta}) \leq Cr^{-3},$$

using (76) and the fact that ξ_i is homogenous of order -3 . In exactly the same way, we can show that on $f_i((2s^\gamma, \varepsilon) \times \Sigma_i)$, we have

$$(107) \quad |\nabla_M \xi|_{g_M} \leq Cr^{-4}, \quad |\eta|_{g_M} \leq Cr^{-4}, \quad |\nabla_M \eta|_{g_M} \leq Cr^{-5}.$$

Hence (105) and (106) then give

$$|\chi_s|_{g_s} \leq Cs^4r^{-4}, \quad |\nabla_s \chi_s|_{g_s} \leq Cs^4r^{-4},$$

which is (102). Here we have used the fact that the $s^k r^{-k}$ terms for $k > 4$ are dominated by $s^4 r^{-4}$ since $sr^{-1} < Cs^{1-\gamma} < C$, because $s < 1$ and $\gamma \in (0, 1)$.

We move on to the overlap regions. Again using (92) and (12) and (10), we get

$$\begin{aligned} |\chi_s|_{g_s} &\leq C|\chi_s|_{g_{C_i}} \\ &\leq C|d\rho_i|_{g_{C_i}} + C|J_{\varphi_{C_i}}(s^4\eta_i + d\tau_i)|_{g_{C_i}} + C|G_{\varphi_{C_i}}(s^4\eta_i - s^3 *_i \xi_i + d\tau_i)|_{g_{C_i}} \\ &\leq C|d\rho_i|_{g_{C_i}} + C|s^4\eta_i + d\tau_i|_{g_{C_i}} + C|s^4\eta_i - s^3 *_i \xi_i + d\tau_i|_{g_{C_i}}^2. \end{aligned}$$

Now we use (93) and (95) to obtain

$$(108) \quad |\chi_s|_{g_s} \leq C(s^{\gamma\mu_i} + s^{-\nu'_i(1-\gamma)} + s^{3(1-\gamma)+\delta\gamma}) + Cs^4|\eta_i|_{g_{C_i}} + C(s^4|\eta_i|_{g_{C_i}} + s^3|\xi_i|_{g_{C_i}} + |d\tau_i|_{g_{C_i}})^2.$$

We claim that the second and third terms above are smaller than, and can thus be absorbed by, the first term. To see this, recall that $|\xi_i|_{g_{C_i}} = O(r^{-3})$ and $|\eta_i|_{g_{C_i}} = O(r^{-4})$ and that $r^a \leq Cr^{a\gamma}$ for $r \in [s^\gamma, 2s^\gamma]$, and thus

$$s^4|\eta_i|_{g_{C_i}} \leq Cs^{4(1-\gamma)}, \quad s^3|\xi_i|_{g_{C_i}} \leq Cs^{3(1-\gamma)}.$$

Since δ is close to zero, the $O(s^{4(1-\gamma)})$ term can be absorbed in $O(s^{3(1-\gamma)+\delta\gamma})$. Now the final term in (108) is

$$(O(s^{3(1-\gamma)}) + O(s^{\gamma\mu_i}) + O(s^{-\nu'_i(1-\gamma)}))^2.$$

Since $s < 1$, and δ is close to zero, every product of two terms in the above expression can be absorbed by one of the summands in the first term of (108). This proves the first estimate in (103). The second estimate in (103) is obtained similarly to the estimate in (106), this time using the fact that ∇_{C_i} commutes with $*_{C_i}$ and that φ_{C_i} is torsion-free, so the $|\nabla_{C_i} \varphi_{C_i}|_{g_{C_i}}$ terms in (12) and (10) are zero.

Finally the equations in (104) are immediate from (92). □

We can now use [Proposition 3.23](#) to estimate the norms needed in part i) of [Theorem 3.1](#).

Theorem 3.24 *Let χ_s be as given in [Proposition 3.18](#). Then the following estimates hold for norms of χ_s on the smooth compact manifold \widetilde{M}_s .*

$$(109) \quad \|\chi_s\|_{C^0} \leq C(s^{\gamma\mu_i} + s^{-\nu'_i(1-\gamma)} + s^{3(1-\gamma)+\delta\gamma}),$$

$$(110) \quad \|\chi_s\|_{L^2} \leq C(s^{\gamma\mu_i+\frac{7}{2}\gamma} + s^{-\nu'_i(1-\gamma)+\frac{7}{2}\gamma} + s^{3(1-\gamma)+\delta\gamma+\frac{7}{2}\gamma}),$$

$$(111) \quad \|d_{g_s}^* \chi_s\|_{L^{14}} \leq C(s^{\gamma\mu_i-\frac{1}{2}\gamma} + s^{-\nu'_i(1-\gamma)-\frac{1}{2}\gamma} + s^{3(1-\gamma)+\delta\gamma-\frac{1}{2}\gamma}).$$

Proof We begin with the C^0 norms of χ_s and of $\nabla_{g_s} \chi_s$ with respect to g_s . Using the estimates in [\(101\)](#), [\(102\)](#), [\(103\)](#) and [\(104\)](#), we find

$$(112) \quad \|\chi_s\|_{C^0} = \sup_{p \in \widetilde{M}_s} |\chi_s(p)|_{g_s(p)} \leq C(s^{\gamma\mu_i} + s^{-\nu'_i(1-\gamma)} + s^{3(1-\gamma)+\delta\gamma}),$$

$$(113) \quad \begin{aligned} \|\nabla_{g_s} \chi_s\|_{C^0} &= \sup_{p \in \widetilde{M}_s} |(\nabla_{g_s} \chi_s)(p)|_{g_s(p)} \\ &\leq C(s^{\gamma\mu_i-\gamma} + s^{-\nu'_i(1-\gamma)-\gamma} + s^{3(1-\gamma)+\delta\gamma-\gamma}), \end{aligned}$$

where in both cases the terms from [\(101\)](#) and [\(102\)](#) are absorbed by the last terms of [\(103\)](#), using the fact that $r^{-4} \leq C s^{-4\gamma}$ in [\(102\)](#). The estimate [\(112\)](#) is [\(109\)](#).

We now move on to the L^2 norm of χ_s . This is

$$\|\chi_s\|_{L^2} = \left(\int_{\widetilde{M}_s} |\chi_s|_{g_s}^2 \text{vol}_{g_s} \right)^{\frac{1}{2}}.$$

We will consider the contributions to the integral coming from the different regions in [\(92\)](#). By [\(104\)](#), there is no contribution from the outer regions. By [Lemma 3.22](#) and [Equation \(18\)](#), in the overlap region $[s^\gamma, 2s^\gamma] \times \Sigma_i$ we have $\text{vol}_{g_s} \leq C \text{vol}_C = C r^6 dr \wedge \text{vol}_\Sigma$. Now the estimate [\(103\)](#) gives

$$\begin{aligned} \int_{[s^\gamma, 2s^\gamma] \times \Sigma_i} |\chi_s|_{g_s}^2 \text{vol}_{g_s} &\leq C \text{vol}(\Sigma_i) \int_{s^\gamma}^{2s^\gamma} |\chi_s|_{g_s}^2 r^6 dr \\ &\leq C(s^{\gamma\mu_i} + s^{-\nu'_i(1-\gamma)} + s^{3(1-\gamma)+\delta\gamma})^2 \int_{s^\gamma}^{2s^\gamma} r^6 dr \\ &\leq C(s^{\gamma\mu_i} + s^{-\nu'_i(1-\gamma)} + s^{3(1-\gamma)+\delta\gamma})^2 s^{7\gamma}, \end{aligned}$$

and hence the contribution to $\|\chi_s\|_{L^2}$ from the overlap region is bounded by

$$(114) \quad C(s^{\gamma\mu_i+\frac{7}{2}\gamma} + s^{-\nu'_i(1-\gamma)+\frac{7}{2}\gamma} + s^{3(1-\gamma)+\delta\gamma+\frac{7}{2}\gamma}).$$

On a subset $f_i((2s^\gamma, \varepsilon) \times \Sigma_i)$ of the inner region, by Lemma 3.22 we again have $\text{vol}_{g_s} \leq Cr^6 dr \wedge \text{vol}_\Sigma$, and so by (102) we see

$$\begin{aligned} \int_{(2s^\gamma, \varepsilon) \times \Sigma_i} |\chi_s|_{g_s}^2 \text{vol}_{g_s} &\leq C \text{vol}(\Sigma_i) \int_{2s^\gamma}^\varepsilon |\chi_s|_{g_s}^2 r^6 dr \\ &\leq Cs^8 \int_{2s^\gamma}^\varepsilon r^{-8} r^6 dr \\ &\leq C(s^8 + s^{8-\gamma}) \leq Cs^{8-\gamma}, \end{aligned}$$

and so the contribution to $\|\chi_s\|_{L^2}$ from this region is bounded by

$$(115) \quad Cs^{4-\frac{1}{2}\gamma} = Cs^{4(1-\gamma)+\frac{7}{2}\gamma}.$$

Finally the subset K of the inner region is compact, so by (101), the contribution to $\|\chi_s\|_{L^2}$ on this region is bounded by Cs^4 , which is smaller than, and hence absorbed by, (115). Putting (114) and (115) together, we find

$$\|\chi_s\|_{L^2} \leq C(s^{\gamma\mu_i+\frac{7}{2}\gamma} + s^{-\nu'_i(1-\gamma)+\frac{7}{2}\gamma} + s^{3(1-\gamma)+\delta\gamma+\frac{7}{2}\gamma} + s^{4(1-\gamma)+\frac{7}{2}\gamma}).$$

It is easy to check that the fourth term above is smaller than, and absorbed by, the third term, exactly when

$$(116) \quad \delta < \frac{1-\gamma}{\gamma},$$

so by taking δ smaller if necessary, we ensure that (110) holds.

Lastly we need to consider the L^{14} norm of $d_{g_s}^* \chi_s$. This is

$$\|d_{g_s}^* \chi_s\|_{L^{14}} = \left(\int_{\tilde{M}_s} |d_{g_s}^* \chi_s|_{g_s}^{14} \text{vol}_{g_s} \right)^{\frac{1}{14}}.$$

By Equation (2), we have $|d_{g_s}^* \chi_s|_{g_s} \leq C|\nabla_{g_s} \chi_s|_{g_s}$. Now we proceed exactly as in the derivation of the L^2 estimates, using the pointwise estimates on $|\nabla_{g_s} \chi_s|$ given in (101), (102) and (103). We sketch the details. On the overlap regions, one obtains that the contribution to $\|d_{g_s}^* \chi_s\|_{L^{14}}$ is bounded by

$$(117) \quad C(s^{\gamma\mu_i-\gamma+\frac{1}{2}\gamma} + s^{-\nu'_i(1-\gamma)-\gamma+\frac{1}{2}\gamma} + s^{3(1-\gamma)+\delta\gamma-\gamma+\frac{1}{2}\gamma}),$$

whereas on the inner region the contribution to $\|d_{g_s}^* \chi_s\|_{L^{14}}$ is bounded by

$$Cs^{4-\frac{7}{2}\gamma} = Cs^{4(1-\gamma)+\frac{1}{2}\gamma}.$$

This term is absorbed by the third term in (117) whenever $\delta < \frac{1}{\gamma}$, which is automatic from (116). Thus we obtain the estimate (111). \square

Corollary 3.25 *There exists $\gamma \in (0, 1)$ and $\kappa > 0$ such that, for s sufficiently small, the closed G_2 structure φ_s constructed in Definition 3.14 satisfies part i) of Joyce’s Theorem 3.1:*

$$\|\chi_s\|_{C^0} \leq D_1 s^\kappa, \quad \|\chi_s\|_{L^2} \leq D_1 s^{\frac{7}{2}+\kappa}, \quad \|d_{g_s}^* \chi_s\|_{L^{14}} \leq D_1 s^{-\frac{1}{2}+\kappa}.$$

Proof By Theorem 3.24, the three estimates in part i) of Theorem 3.1 will be satisfied if and only if

$$(118) \quad \mu_i \gamma \geq \kappa, \quad -v'_i(1-\gamma) \geq \kappa, \quad 3(1-\gamma) + \delta \gamma \geq \kappa,$$

$$(119) \quad \begin{aligned} \mu_i \gamma + \frac{7}{2} \gamma &\geq \frac{7}{2} + \kappa, \quad -v'_i(1-\gamma) + \frac{7}{2} \gamma \geq \frac{7}{2} + \kappa, \\ 3(1-\gamma) + \delta \gamma + \frac{7}{2} \gamma &\geq \frac{7}{2} + \kappa, \end{aligned}$$

$$(120) \quad \begin{aligned} \mu_i \gamma - \frac{1}{2} \gamma &\geq -\frac{1}{2} + \kappa, \quad -v'_i(1-\gamma) - \frac{1}{2} \gamma \geq -\frac{1}{2} + \kappa, \\ 3(1-\gamma) + \delta \gamma - \frac{1}{2} \gamma &\geq -\frac{1}{2} + \kappa. \end{aligned}$$

It is trivial to verify that for $\gamma < 1$, the inequalities in (119) automatically imply the inequalities in (118) and (120). That is, the L^2 estimate implies the C^0 and L^{14} estimates. The three inequalities in (119) can be rearranged to yield

$$(121) \quad \gamma \geq \frac{\frac{7}{2} + \kappa}{\frac{7}{2} + \mu_i}, \quad \gamma \geq 1 + \frac{\kappa}{\frac{7}{2} + v'_i}, \quad \gamma \geq \frac{1 + 2\kappa}{1 + 2\delta}.$$

We want to ensure that there exists a $\kappa > 0$ such that if we define γ by the three expressions above, then γ will be in $(0, 1)$. Recall that $\mu_i > 0$, $\delta > 0$ and $v'_i = -4$ (from Theorem 3.6.) For the first inequality in (121), we see that $\gamma > 0$ for any $\kappa > 0$, and $\gamma < 1$ if $\kappa < \mu_i$. For the third inequality in (121), we have $\gamma > 0$ for any $\kappa > 0$, and $\gamma < 1$ if $\kappa < \delta$. Finally, for the middle inequality in (121), we see that $\gamma < 1$ for any $\kappa > 0$, and $\gamma > 0$ if $\kappa < -(\frac{7}{2} + v'_i) = \frac{1}{2}$. Therefore it suffices to choose κ satisfying

$$\kappa < \delta, \quad \kappa < \frac{1}{2}, \quad \text{and} \quad \kappa < \mu_i, \quad \text{for all } i = 1, \dots, n.$$

The intersection of the three inequalities in (121) with the regions $0 < \gamma < 1$ and $\kappa > 0$ is represented by the shaded region in Figure 6. □

Remark 3.26 We can see from (121) that there is a *critical rate* $v_c = -\frac{7}{2}$ such that gluing in an AC G_2 manifold with decay rate $v' < v_c$ works, while for rate $v' > v_c$ it *does not* work. The determining factor is the size of the L^2 norm of the error caused by




Figure 6: Intersection of the inequalities in the γ - κ plane

smoothing off the AC decay, which is contained in χ_s . This shows the importance of [Theorem 3.6](#), which we were able to use to show that for any *initial* AC rate $\nu \leq -3$, we can extract cohomological parts, in equations (64) and (65) and the left over parts then have a *faster* rate of decay $\nu' = -4$, which is less than the critical rate $\nu_c = -\frac{7}{2}$, so that our gluing construction can actually be made to work for any initial rate $\nu \leq -3$. In effect, [Theorem 3.6](#) allows us to extend the AC rates which allow the desingularization to succeed from $\nu < -\frac{7}{2}$ to $\nu \leq -3$. The reader should also review again [Remark 3.7](#).

Next we need to estimate the Riemann curvature and the injectivity radius of g_s .

Proposition 3.27 *For s sufficiently small, the metric g_s induced by the closed G_2 structure φ_s constructed in [Definition 3.14](#) satisfies parts ii) and iii) of [Joyce’s Theorem 3.1](#):*

$$\mathcal{I}(g_s) \geq D_2 s, \quad \|\mathcal{R}(g_s)\|_{C^0} \leq D_3 s^{-2}.$$

Proof We begin by noting that under a conformal scaling of metric $\tilde{g} = s^2 g$, the Riemann curvature tensor \mathcal{R} and the injectivity radius \mathcal{I} change by

$$\|\mathcal{R}(\tilde{g})\|_{C^0(\tilde{g})} = s^{-2} \|\mathcal{R}(g)\|_{C^0(g)} \quad \text{and} \quad \mathcal{I}(\tilde{g}) = s \mathcal{I}(g).$$

We will consider the three regions of \tilde{M}_s as given in [Definition 3.14](#).

On the inner region $K \cup \bigsqcup_{i=1}^n f_i((2s^\nu, \varepsilon) \times \Sigma_i)$, we have $|g_s - g_M|_{g_M} \leq \varepsilon$, for some $\varepsilon < \frac{1}{2}$ by [Lemma 3.22](#). Hence by Taylor’s theorem we have $|\mathcal{R}(g_s) - \mathcal{R}(g_M)|_{g_M} \leq C\varepsilon$.

Therefore

$$|\mathcal{R}(g_s)|_{g_s} \leq C|\mathcal{R}(g_s)|_{g_M} \leq C|\mathcal{R}(g_M)|_{g_M} + C|\mathcal{R}(g_s) - \mathcal{R}(g_M)|_{g_M} \leq C,$$

using (99) and the fact that $\mathcal{R}(g_M)$ is smooth and hence bounded on any compact subset of M containing the inner region. On the overlap regions $[s^\gamma, 2s^\gamma] \times \Sigma_i$ we can again use the fact that g_s is uniformly close to g_{C_i} by Lemma 3.22, so as above we can say

$$|\mathcal{R}(g_s)|_{g_s} \leq C|\mathcal{R}(g_{C_i})|_{g_{C_i}} + C|\mathcal{R}(g_s) - \mathcal{R}(g_{C_i})|_{g_{C_i}} \leq C|\mathcal{R}(g_{C_i})|_{g_{C_i}} + C.$$

Since $g_{C_i}(tr, \sigma) = t^2 g_{C_i}(t, \sigma)$, we have

$$|\mathcal{R}(g_{C_i})(tr, \sigma)|_{g_{C_i}(tr, \sigma)} = t^{-2} |\mathcal{R}(g_{C_i})(r, \sigma)|_{g_{C_i}(r, \sigma)}$$

on the cone C_i . Thus since $r \in [s^\gamma, 2s^\gamma]$ on these regions, we have

$$|\mathcal{R}(g_{C_i})|_{g_{C_i}} \leq Cs^{-2\gamma}.$$

Finally we consider the outer regions $L_i \cup h_{i,s}((sR, s^\gamma) \times \Sigma_i)$. Here we have $g_s = s^2 g_{N_i}$ exactly, so

$$|\mathcal{R}(g_s)|_{g_s} = s^{-2} |\mathcal{R}(g_{N_i})|_{g_{N_i}} \leq Cs^{-2}$$

as $\mathcal{R}(g_{N_i})$ is smooth and thus bounded on any compact subset of N_i containing that outer region. Putting all three estimates together gives iii) of Theorem 3.1.

The estimate on the injectivity radius in ii) of Theorem 3.1 is proved in essentially the same way. On the inner region, we get $\mathcal{I}(g_s) \geq C$, on the overlap regions we get $\mathcal{I}(g_s) \geq Cs^\gamma$, and on the outer regions we get $\mathcal{I}(g_s) \geq Cs$. Thus when s is sufficiently small, putting these all together gives ii) of Theorem 3.1.

Note that in both cases, the dominant contribution comes from the conformally scaled metrics $g_s = s^2 g_{N_i}$ on the outer regions which were glued into M to obtain \widetilde{M}_s . \square

Finally we can prove our main theorem.

Theorem 3.28 *Let M be a compact G_2 manifold with isolated conical singularities, with singularities x_1, \dots, x_n , cones C_1, \dots, C_n and rates μ_1, \dots, μ_n , respectively. Suppose that we have asymptotically conical G_2 manifolds N_1, \dots, N_n , with the same cones C_1, \dots, C_n and rates ν_1, \dots, ν_n , respectively, with each $\nu_i \leq -3$. If the conditions (74) and (75) of Theorem 3.10 are satisfied, then there exists a one-parameter family \widetilde{M}_s of smooth, compact G_2 manifolds, for $0 < s < \kappa$, (with holonomy exactly equal to G_2), which desingularize M .*

Proof From [Corollary 3.25](#) and [Proposition 3.27](#), we see that the hypotheses of [Theorem 3.1](#) are satisfied with $\varphi = \varphi_s$ and $\chi = \chi_s$ on the smooth compact manifold \tilde{M}_s . Hence there exists a torsion-free G_2 structure $\tilde{\varphi}_s$ on \tilde{M}_s , for $0 < s < \kappa$. The Riemannian holonomy of the metric \tilde{g}_s induced by $\tilde{\varphi}_s$ must be exactly G_2 , since the holonomy of M' and of the N_i 's was exactly G_2 , and the holonomy of a manifold obtained by gluing must be at least as large as the holonomy of its constituent pieces. \square

In fact we can say informally that “ $\lim_{s \rightarrow 0} \tilde{M}_s = M$ ”, since as $s \rightarrow 0$, the asymptotically conical G_2 manifolds $N_{i,s}$ shrink to points, corresponding to the original conical singularities x_i , and hence the manifold \tilde{M}_s approaches the original G_2 manifold M with isolated conical singularities. Therefore compact G_2 manifolds with isolated conical singularities can be thought of as possible “boundary points” in the moduli spaces of compact smooth G_2 manifolds.

4 Analysis on manifolds with ends

In this section we summarize some aspects of analysis on noncompact manifolds with ends. We will only need parts of the theory for weighted Sobolev spaces. It applies equally well to weighted Hölder spaces. Analysis on noncompact manifolds with ends using weighted Banach spaces has been discussed by many authors, including Maz'ja–Plamenevskii [34], Bartnik [4], Biquard [6] and Aubin [2]. We choose to follow the approach developed in Lockhart–McOwen [29] and Lockhart [28]. A detailed exposition of this approach can also be found in Marshall [33], and a summary in the context of manifolds with ICS was first presented in Joyce [20]. Weighted Sobolev space techniques have also been applied by Nordström [35] to study asymptotically cylindrical G_2 manifolds.

4.1 Analysis on manifolds with isolated conical singularities

Let M be a compact G_2 manifold with isolated conical singularities, as in [Definition 2.25](#). In order to be able to define sensible “weighted” Banach spaces on the noncompact smooth manifold M' , we need to introduce the concept of a radius function.

Definition 4.1 A *radius function* ϱ on M' is a smooth function on M' that satisfies the following conditions. On the compact subset $K = M' \setminus \bigsqcup_{i=1}^n S_i$, we define $\varrho \equiv 1$. Let x be a point in the neighbourhood $f_i((0, \frac{1}{2}\varepsilon) \times \Sigma_i)$ of the i -th singularity x_i . Then $f_i^{-1}(x) = (r, \sigma)$ for some $r \in (0, \frac{1}{2}\varepsilon)$, and we set $\varrho(x) = r$ for such a point. Finally, in the regions $f_i((\frac{1}{2}\varepsilon, \varepsilon) \times \Sigma_i)$, the function ϱ is defined by interpolating smoothly in an increasing fashion between these two definitions.

Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be an n -tuple of real numbers. We can add such n -tuples and multiply them by real numbers using the vector space structure of \mathbb{R}^n . We also define $\lambda + j = (\lambda_1 + j, \dots, \lambda_n + j)$ for any $j \in \mathbb{R}$. Now define ϱ^λ to equal ϱ^{λ_i} on $f_i((0, \varepsilon) \times \Sigma_i)$ and to equal 1 on K . Then ϱ^λ is a smooth function on M' which equals r^{λ_i} on the neighbourhood $f_i((0, \frac{1}{2}\varepsilon) \times \Sigma_i)$ of x_i .

Definition 4.2 Let $p > 1$, $l \geq 0$ and $\lambda \in \mathbb{R}^n$. We define the *weighted Sobolev space* $L^p_{l,\lambda}(\Lambda^k(T^*M'))$ of k -forms on M' as follows. Consider the space $C^\infty_{cs}(\Lambda^k(T^*M'))$ of smooth compactly supported k -forms on M' . For such forms the quantity

$$(122) \quad \|\omega\|_{L^p_{l,\lambda}} = \left(\sum_{j=0}^l \int_{M'} |\varrho^{-\lambda+j} \nabla_M^j \omega|_{g_M}^p \varrho^{-7 \text{vol}_M} \right)^{1/p}$$

is clearly finite and is a norm. We define the Banach space $L^p_{l,\lambda}(\Lambda^k(T^*M'))$ to be the completion of $C^\infty_{cs}(\Lambda^k(T^*M'))$ with respect to this norm. As a topological vector space, $L^p_{l,\lambda}(\Lambda^k(T^*M'))$ is independent of the choice of radius function ϱ , and any two such choices lead to equivalent norms.

Remark 4.3 There are several observations to be made about this definition.

- i) It is clear that $L^p_{l,\lambda}(\Lambda^k(T^*M')) \subseteq L^p_{l,\lambda'}(\Lambda^k(T^*M'))$ if $\lambda_i > \lambda'_i$ for all $i = 1, \dots, n$.
- ii) It is easy to show that if $|\omega|_{g_M} = O(r^{\lambda_i})$ near x_i , then $\omega \in L^p_{0,\lambda-e}(\Lambda^k(T^*M'))$ for any $e > 0$.
- iii) Because of the factor of ϱ^{-7} in (122), for $l = 0$, the space $L^p_{0,-\frac{7}{p}}(\Lambda^k(T^*M'))$ is the usual $L^p(\Lambda^k(T^*M'))$ space, and in particular we have

$$(123) \quad L^2_{0,-\frac{7}{2}}(\Lambda^k(T^*M')) = L^2(\Lambda^k(T^*M')).$$

Here it is understood that $\frac{7}{p}$ denotes the ‘‘constant’’ n -tuple $(\frac{7}{p}, \dots, \frac{7}{p})$.

- iv) The differential operators d and d^*_M , defined on $C^\infty_{cs}(\Lambda^k(T^*M'))$, clearly extend to bounded linear operators

$$\begin{aligned} d: L^p_{l+1,\lambda}(\Lambda^k(T^*M')) &\rightarrow L^p_{l,\lambda-1}(\Lambda^{k+1}(T^*M')), \\ d^*_M: L^p_{l+1,\lambda}(\Lambda^k(T^*M')) &\rightarrow L^p_{l,\lambda-1}(\Lambda^{k-1}(T^*M')). \end{aligned}$$

Proposition 4.4 Let q satisfy $\frac{1}{p} + \frac{1}{q} = 1$. There is a Banach space isomorphism

$$(L^p_{0,\lambda}(\Lambda^k(T^*M')))^\ast \cong L^q_{0,-\lambda-7}(\Lambda^k(T^*M')),$$

given by the L^2 inner product pairing. That is, if $\alpha \in L^p_{0,\lambda}(\Lambda^k(T^*M'))$ and $\beta \in L^q_{0,-\lambda-7}(\Lambda^k(T^*M'))$, then

$$(124) \quad |\langle \alpha, \beta \rangle_{L^2}| \leq C \|\alpha\|_{L^p_{0,\lambda}} \|\beta\|_{L^q_{0,-\lambda-7}}.$$

Hence $\langle \cdot, \cdot \rangle_{L^2}$ is well-defined and continuous on

$$L^p_{0,\lambda}(\Lambda^k(T^*M')) \times L^q_{0,-\lambda-7}(\Lambda^k(T^*M')).$$

Corollary 4.5 Suppose $\alpha \in L^p_{1,\lambda+1}(\Lambda^k(T^*M'))$ and $\beta \in L^q_{1,-\lambda-7}(\Lambda^{k+1}(T^*M'))$, where $\frac{1}{p} + \frac{1}{q} = 1$. Then we can “integrate by parts” to say that

$$(125) \quad \langle d\alpha, \beta \rangle_{L^2} = \langle \alpha, d^*_M \beta \rangle_{L^2}.$$

This also holds if $\beta \in L^q_{1,\lambda'}(\Lambda^{k+1}(T^*M'))$ whenever $\lambda + \lambda' > -7$.

Proof Equation (125) clearly holds when α and β are both smooth, compactly supported forms. By item iv) of Remark 4.3, we know that $d\alpha \in L^p_{0,\lambda}(\Lambda^{k+1}(T^*M'))$ and $d^*_M \beta \in L^q_{0,-\lambda-8}(\Lambda^k(T^*M'))$. But then by the continuity of the pairing $\langle \cdot, \cdot \rangle_{L^2}$ from (124) and the fact that smooth compactly supported forms are *dense* in the weighted Sobolev spaces, Equation (125) follows by taking limits. The last part of the claim follows immediately from item i) of Remark 4.3. □

Let $\Lambda^*(T^*M') = \sum_{k=0}^7 \Lambda^k(T^*M')$. Define

$$L^p_{l,\lambda}(\Lambda^*(T^*M')) = \sum_{k=0}^7 L^p_{l,\lambda}(\Lambda^k(T^*M')).$$

We will be interested in the following two differential operators:

$$(126) \quad (d + d^*_M)^p_{l+1,\lambda}: L^p_{l+1,\lambda}(\Lambda^*(T^*M')) \rightarrow L^p_{l,\lambda-1}(\Lambda^*(T^*M')),$$

$$(127) \quad (\Delta_M)^p_{l+2,\lambda}: L^p_{l+2,\lambda}(\Lambda^k(T^*M')) \rightarrow L^p_{l,\lambda-2}(\Lambda^k(T^*M')).$$

They are defined by extending the operators $d + d^*_M$ and Δ_M from smooth compactly supported forms to the Sobolev spaces.

Definition 4.6 Let C be a cone. The set $\mathcal{D}_{d+d^*_C}$ of *critical rates* of the operator $d + d^*_C$ on $\Lambda^*(T^*C)$ is defined as follows:

$$(128) \quad \mathcal{D}_{d+d^*_C} = \left\{ \begin{array}{l} \lambda \in \mathbb{R}; \exists \text{ a nonzero } \omega = \sum_{k=0}^7 \omega_k \in \Lambda^*(T^*C), \\ \text{homogeneous of order } \lambda, \text{ such that } (d + d^*_C)(\omega) = 0 \end{array} \right\}.$$

Similarly the set \mathcal{D}_{Δ_C} of critical rates of Δ_C on $\Lambda^k(T^*C)$ is defined as

$$(129) \quad \mathcal{D}_{\Delta_C} = \left\{ \lambda \in \mathbb{R}; \exists \text{ a nonzero } \omega \in \Lambda^k(T^*C), \right. \\ \left. \text{homogeneous of order } \lambda, \text{ such that } \Delta_C \omega = 0 \right\}.$$

Both $\mathcal{D}_{d+d_C^*}$ and \mathcal{D}_{Δ_C} are countable, discrete subsets of \mathbb{R} .

Our manifold M with isolated conical singularities has n singular points, with cones C_1, \dots, C_n . The relevance of the critical rates is that they are related to the rates λ for which the operators $(d + d_M^*)_{l+1,\lambda}^p$ and $(\Delta_M)_{l+2,\lambda}^p$ of (126) and (127) are Fredholm, by the following theorem, which is taken from Lockhart–McOwen [29, Theorem 1.1].

Theorem 4.7 *The map*

$$(d + d_M^*)_{l+1,\lambda}^p : L_{l+1,\lambda}^p(\Lambda^*(T^*M')) \rightarrow L_{l,\lambda-1}^p(\Lambda^*(T^*M'))$$

is Fredholm if and only if $\lambda_i \notin \mathcal{D}_{d+d_{C_i}^*}$ for all $i = 1, \dots, n$. Similarly the map

$$(\Delta_M)_{l+2,\lambda}^p : L_{l+2,\lambda}^p(\Lambda^k(T^*M')) \rightarrow L_{l,\lambda-2}^p(\Lambda^k(T^*M'))$$

is Fredholm if and only if $\lambda_i \notin \mathcal{D}_{\Delta_{C_i}}$ for all $i = 1, \dots, n$.

We have the following *elliptic regularity* result for the operators $d + d_M^*$ and Δ_M , which can be found in Lockhart–McOwen [29, inequality (2.4)].

Theorem 4.8 *Suppose that ω and v are locally integrable sections of $\Lambda^*(T^*M')$ and that ω is a weak solution of the equation $(d + d_M^*)(\omega) = v$. If $\omega \in L_{0,\lambda}^p(\Lambda^*(T^*M'))$ and $v \in L_{l,\lambda-1}^p(\Lambda^*(T^*M'))$, then $\omega \in L_{l+1,\lambda}^p(\Lambda^*(T^*M'))$, and ω is a strong solution of $(d + d_M^*)(\omega) = v$. Furthermore, we have*

$$(130) \quad \|\omega\|_{L_{l+1,\lambda}^p} \leq C(\|(d + d_M^*)(\omega)\|_{L_{l,\lambda-1}^p} + \|\omega\|_{L_{0,\lambda}^p})$$

for some constant $C > 0$ independent of ω . That is, ω has at least one more derivative's worth of regularity than $v = (d + d_M^*)(\omega)$.

Similarly suppose ω and v are locally integrable sections of $\Lambda^k(T^*M')$, and ω is a weak solution of the equation $\Delta_M \omega = v$. If $\omega \in L_{0,\lambda}^p(\Lambda^k(T^*M'))$ and $v \in L_{l,\lambda-2}^p(\Lambda^k(T^*M'))$, then $\omega \in L_{l+2,\lambda}^p(\Lambda^k(T^*M'))$, and ω is a strong solution of $\Delta_M \omega = v$. Furthermore, we have

$$(131) \quad \|\omega\|_{L_{l+2,\lambda}^p} \leq C(\|\Delta_M \omega\|_{L_{l,\lambda-2}^p} + \|\omega\|_{L_{0,\lambda}^p})$$

for some $C > 0$ independent of ω .

We also need the following important result from Lockhart–McOwen [29, Lemma 7.3].

Theorem 4.9 *The kernel of $(d + d_M^*)_{l+1, \lambda}^p$ is independent of $p > 1$ and independent of l . Hence we can denote it unambiguously as $\ker(d + d_M^*)_{\lambda}$. This kernel is also invariant as we change the rates λ , as long as we do not hit any critical rates. That is, if $\lambda = (\lambda_1, \dots, \lambda_n)$ and $\lambda' = (\lambda'_1, \dots, \lambda'_n)$, with the interval $[\lambda_i, \lambda'_i]$ contained in $\mathbb{R} \setminus \mathcal{D}_{d+d_{C_i}^*}$ for each i , then*

$$\ker(d + d_M^*)_{\lambda'} = \ker(d + d_M^*)_{\lambda}.$$

Similarly the kernel of $(\Delta_M)_{l+2, \lambda}^p$ is independent of $p > 1$ and independent of l . Hence we can denote it unambiguously as $\ker(\Delta_M)_{\lambda}$. This kernel is also invariant as we change the rates λ , as long as we do not hit any critical rates. That is, if $\lambda = (\lambda_1, \dots, \lambda_n)$ and $\lambda' = (\lambda'_1, \dots, \lambda'_n)$, with the interval $[\lambda_i, \lambda'_i]$ contained in $\mathbb{R} \setminus \mathcal{D}_{\Delta_{C_i}}$ for each i , then

$$\ker(\Delta_M)_{\lambda'} = \ker(\Delta_M)_{\lambda}.$$

Remark 4.10 *There is a ‘‘Sobolev embedding theorem’’ in the context of manifolds with ICS, which (for large enough p and l) embeds the Sobolev space $L_{l, \lambda}^p$ into an appropriate Hölder space $C_{\lambda'}^{k, \alpha}$ having k continuous derivatives, where k , α and λ' depend on p , l and λ . It follows from this theorem and the elliptic regularity of [Theorem 4.8](#) that elements in the kernel of $d + d_M^*$ or Δ_M are smooth, and the independence of the kernels on l follows from this. In particular, it now follows that*

$$\begin{aligned} \omega \in \ker(d + d_M^*)_{\lambda} &\implies |f_i^*(\omega)|_{f_i^*(g_M)} \leq C r^{\lambda_i} && \text{on } \left(0, \frac{1}{2}\varepsilon\right) \times \Sigma_i, \\ \omega \in \ker(\Delta_M)_{\lambda} &\implies |f_i^*(\omega)|_{f_i^*(g_M)} \leq C r^{\lambda_i} && \text{on } \left(0, \frac{1}{2}\varepsilon\right) \times \Sigma_i. \end{aligned} \tag{132}$$

The independence of the kernels on $p > 1$ is more complicated and can be found in Lockhart–McOwen [\[29, Sections 7–8\]](#). We will not need the embedding theorem explicitly in this paper.

Lemma 4.11 *Let $p > 1$. Suppose that $\omega \in L_{l+2, \lambda}^p(\Lambda^k(T^*M'))$ and that $\Delta_M \omega = 0$. If $\lambda_i > -\frac{5}{2}$ for all $i = 1, \dots, n$, then $d_M^* \omega = 0$ and $d\omega = 0$.*

Proof *By [Theorem 4.9](#), if $\omega \in \ker(\Delta_M)_{l+2, \lambda}^p$, then $\omega \in \ker(\Delta_M)_{l+2, \lambda}^2$. Now the elliptic regularity of [Theorem 4.8](#) tells us that $\omega \in L_{l+2, \lambda}^2(\Lambda^k(T^*M'))$, and hence $d\omega$ and $d_M^* \omega$ are both elements of $L_{l+1, \lambda-1}^2(\Lambda^*(T^*M'))$. Because each $\lambda_i > -\frac{5}{2}$, [Corollary 4.5](#) says that the following integration by parts computation is valid:*

$$0 = \langle \Delta_M \omega, \omega \rangle_{L^2} = \langle dd_M^* \omega, \omega \rangle_{L^2} + \langle d_M^* d\omega, \omega \rangle_{L^2} = |d_M^* \omega|_{L^2}^2 + |d\omega|_{L^2}^2,$$

and so we have $d_M^* \omega = 0$ and $d\omega = 0$. □

Corollary 4.12 Suppose that $\omega_k \in L^2_{l+2,\lambda}(\Lambda^k(T^*M'))$ and that $\Delta_M \omega_k = 0$. Then we have:

- (133) For $k = 0, 7$, if each $\lambda_i > -5$, then $d^*_M \omega = 0$ and $d\omega = 0$.
- (134) For $k = 1, 6$, if each $\lambda_i > -4$, then $d^*_M \omega = 0$ and $d\omega = 0$.
- (135) For $k = 2, 5$, if each $\lambda_i > -3$, then $d^*_M \omega = 0$ and $d\omega = 0$.

Proof By Proposition 2.15 we see that in all three cases, there are no critical rates until at least $\lambda_i = -2$. So using Theorem 4.9, in all cases we can say that $\omega_k \in \ker(\Delta_M)_{-\frac{5}{2}+e}$ for some small $e > 0$. The claims now all follow from Lemma 4.11. □

The following proposition (in a general setting) originally appeared in Lockhart [28, Example 0.16] and a version in the setting of manifolds with ICS is stated in Lotay [31, Theorem 6.5.2].

Proposition 4.13 Let $\mathcal{H}^k_{L^2}$ denote the subspace of $L^2(\Lambda^k(T^*M'))$ consisting of closed and coclosed k -forms. Then for $0 \leq k \leq 3$, the map

$$[\cdot]: \mathcal{H}^k_{L^2} \rightarrow H^k(M', \mathbb{R})$$

$$\omega \mapsto [\omega]$$

is an isomorphism. Here $[\omega]$ denotes the cohomology class of the closed form ω .

The rest of the results in this section also all have analogues for Δ_M , but we will only require them for the operator $d + d^*_M$. First we will need to consider the formal adjoint of the map

$$(d + d^*_M)^p_{l+1,\lambda}: L^p_{l+1,\lambda}(\Lambda^*(T^*M')) \rightarrow L^p_{l,\lambda-1}(\Lambda^*(T^*M')).$$

By Proposition 4.4, the formal adjoint extends to a map

$$(136) \quad (d + d^*_M)^q_{m+1,-\lambda-6}: L^q_{m+1,-\lambda-6}(\Lambda^*(T^*M')) \rightarrow L^q_{m,-\lambda-7}(\Lambda^*(T^*M')),$$

where $\frac{1}{p} + \frac{1}{q} = 1$, and $l, m \geq 0$.

The next result is the version of the ‘‘Fredholm Alternative’’ for manifolds with ICS, adapted from Lockhart–McOwen [29, Theorem 1.1].

Theorem 4.14 Suppose that λ_i is not in $\mathcal{D}_{d+d^*_M}$ for all $i = 1, \dots, n$, so that by Theorem 4.7, the map

$$(d + d^*_M)^p_{l+1,\lambda}: L^p_{l+1,\lambda}(\Lambda^*(T^*M')) \rightarrow L^p_{l,\lambda-1}(\Lambda^*(T^*M'))$$

is Fredholm and also uniformly elliptic. Then the image $\text{im}(d + d_M^*)_{l+1,\lambda}^p$ of this map is given by

$$\begin{aligned} &\text{im}(d + d_M^*)_{l+1,\lambda}^p \\ &= \{v \in L_{l,\lambda-1}^p(\Lambda^*(T^*M')); \langle v, \omega \rangle_{L^2} = 0, \forall \omega \in \ker(d + d_M^*)_{-\lambda-6}\}. \end{aligned}$$

That is, the equation $(d + d_M^*)(\omega) = v$ is solvable for $\omega \in L_{l+1,\lambda}^p(\Lambda^*(T^*M'))$ if and only if the right hand side v is orthogonal (with respect to the L^2 inner product) to every element of $\ker(d + d_M^*)_{-\lambda-6}$, the kernel of the formal adjoint map.

We will need to understand how, near each singular point x_i , we can asymptotically expand elements of the kernel of $(d + d_M^*)_{l+1,\lambda}^p$ in terms of certain special solutions of $(d + d_{C_i}^*)(\omega) = 0$.

Definition 4.15 Let C be a cone. For $\lambda \in \mathbb{R}$, we define the space $\mathcal{K}(\lambda)_{d+d_C^*}$ to be (137)

$$\mathcal{K}(\lambda)_{d+d_C^*} := \left\{ \begin{array}{l} \omega = \sum_{k=0}^7 \sum_{j=0}^m (\log(r))^j \omega_{k,j}; \text{ such that } (d + d_C^*)(\omega) = 0, \\ \text{and each } \omega_{k,j} \text{ is a homogeneous } k\text{-form of order } \lambda \end{array} \right\}.$$

That is, $\mathcal{K}(\lambda)_{d+d_C^*}$ is the set mixed-degree forms on C in the kernel of $d + d_C^*$, which are polynomials in $\log(r)$ with coefficients being homogeneous forms of order λ .

We will use $\ker_i(d + d_M^*)_\lambda$ to denote the restriction of the kernel of $(d + d_M^*)_{l+1,\lambda}^p$ to the subset $f_i((0, \frac{1}{2}\varepsilon) \times \Sigma_i)$ of the i -th end of M' (recall by [Theorem 4.9](#) this kernel is independent of $p > 1$ and $l \geq 0$.) The following result appears (in a slightly disguised form) in Lockhart–McOwen [[29](#), Section 5].

Proposition 4.16 Let β, β' be two n -tuples of rates. Fix i and suppose that β_i and β'_i are in $\mathbb{R} \setminus \mathcal{D}_{d+d_{C_i}^*}$ and that $\beta_i < \beta'_i$. Let $\alpha_1 < \alpha_2 < \dots < \alpha_N$ be all the critical rates in $\mathcal{D}_{d+d_{C_i}^*}$ between β_i and β'_i . If $\omega \in \ker_i(d + d_M^*)_\beta$, then there exist $v_j \in \mathcal{K}(\alpha_j)_{d+d_{C_i}^*}$, for $j = 1, \dots, N$, and an ω' defined near x_i with $|\omega'|_{g_{C_i}} = O(r^{\beta_i + \mu_i})$ such that

$$\omega - \sum_{j=1}^N (f_i^{-1})^*(v_j) - \omega' \in \ker_i(d + d_M^*)_{\beta'}.$$

That is, when restricted to the subset $f_i((0, \frac{1}{2}\varepsilon) \times \Sigma_i)$ of the i -th end of M' , an element in the kernel of $d + d_M^*$ with noncritical rate β_i admits an expansion in terms of elements in $\mathcal{K}(\alpha_j)_{d+d_{C_i}^*}$ for each critical rate α_j between β_i and β'_i , plus a remainder term which is of order $O(r^{\beta_i + \mu_i})$ and another term which is in the kernel of $d + d_M^*$ with noncritical rate $\beta'_i > \beta_i$.

Remark 4.17 The $O(r^{\beta_i + \mu_i})$ term arises from comparing a solution to the equation $(d + d_M^*)(\omega) = 0$ to a solution of $(d + d_{C_i}^*)(\omega) = 0$, using the relation (54) between the metrics g_{C_i} and g_M near x_i .

We will use Proposition 4.16 to prove the following.

Corollary 4.18 Let $\lambda = (\lambda, \dots, \lambda)$ be a constant n -tuple. Choose $\delta > 0$ small enough so that in the closed interval $[\lambda - \delta, \lambda + \delta]$, at most only λ itself is a critical rate for each end. That is, $[\lambda - \delta, \lambda + \delta] \cap \mathcal{D}_{d+d_{C_i}^*} \subseteq \{\lambda\}$ for each i . This is possible since each \mathcal{D}_i is a discrete subset of \mathbb{R} . Further assume that $\delta > 0$ is small enough so that $\delta < \frac{1}{2}\mu_i$ for each i , where μ_i are the rates of the singularities of the manifold M with ICS. If $\omega \in \Lambda^*(T^*M')$ is in the kernel of $(d + d_M^*)_{i+1, \lambda - \delta}^P$, then

$$(138) \quad |f_i^*(\omega) - v_i|_{g_{C_i}} = O(r^{\lambda + \delta}) \quad \text{on} \quad \left(0, \frac{1}{2}\varepsilon\right) \times \Sigma_i,$$

for some v_i in $\mathcal{K}(\lambda)_{d+d_{C_i}^*}$, which may be zero (if $\lambda \notin \mathcal{D}_{d+d_{C_i}^*}$.)

Proof We apply Proposition 4.16, with $\beta_i = \lambda - \delta$ and $\beta'_i = \lambda + \delta$. Because of the hypotheses on δ , for some $v_i \in \mathcal{K}(\lambda)_{d+d_{C_i}^*}$, on $f_i((0, \frac{1}{2}\varepsilon) \times \Sigma_i)$ we have that $\omega - (f_i^{-1})^*(v_i) - \omega'$ lies in $\ker_i(d + d_M^*)_{\lambda + \delta}$, where $|\omega'| = O(r^{\lambda - \delta + \mu_i})$. Hence by (132), we have

$$|f_i^*(\omega) - v_i|_{f_i^*(g_M)} \leq Cr^{\lambda + \delta} + Cr^{\lambda - \delta + \mu_i} \quad \text{on} \quad \left(0, \frac{1}{2}\varepsilon\right) \times \Sigma_i.$$

Since by hypothesis we have $\delta < \frac{1}{2}\mu_i$ and $r < 1$, the second term on the right is smaller than and can be absorbed by, the first term. □

There are many more results that are known in this setting. For example, the index of $d + d_M^*$ will stay constant (like the kernel) unless we cross a critical rate, and when we do, the change in the index can be explicitly computed using the dimensions of the spaces $\mathcal{K}(\lambda)_{d+d_{C_i}^*}$. We will not need these results here. The interested reader is referred to Marshall [33] for a comprehensive review.

4.2 Analysis on asymptotically conical manifolds

Let N be an asymptotically conical G_2 manifold, as in Definition 2.31. All of the analytic results about manifolds with ICS also hold for AC manifolds. The only differences are that there is only one end this time, and since $r \rightarrow \infty$ instead of $r \rightarrow 0$ at the end, all the inequalities involving rates are reversed. In fact these results can

be shown to follow from those of [Section 4.1](#) because an AC metric (restricted to the asymptotic end) is actually conformal to an ICS metric. However, for completeness we prefer to state precisely those results in the AC setting which we will require.

Definition 4.19 A *radius function* ϱ on N is a smooth function on N that satisfies the following conditions. On the compact subset L of N , we define $\varrho \equiv 1$. Let x be a point in $h((2R, \infty) \times \Sigma)$. Then $h^{-1}(x) = (r, \sigma)$ for some $r \in (2R, \infty)$, and we set $\varrho(x) = r$ for such a point. Finally, in the region $h((R, 2R) \times \Sigma)$, the function ϱ is defined by interpolating smoothly between these two definitions.

Definition 4.20 Let $p > 1$, $l \geq 0$ and $\lambda \in \mathbb{R}$. We define the *weighted Sobolev space* $L^p_{l,\lambda}(\Lambda^k(T^*N))$ of k -forms on N as follows. Consider the space $C^\infty_{cs}(\Lambda^k(T^*N))$ of smooth compactly supported k -forms on N . For such forms the quantity

$$(139) \quad \|\omega\|_{L^p_{l,\lambda}} = \left(\sum_{j=0}^l \int_N |\varrho^{-\lambda+j} \nabla_N^j \omega|_{g_N}^p \varrho^{-7} \text{vol}_N \right)^{1/p}$$

is a norm, and we let $L^p_{l,\lambda}(\Lambda^k(T^*N))$ be the completion of $C^\infty_{cs}(\Lambda^k(T^*N))$ with respect to this norm.

Remark 4.21 Here are the observations analogous to those of [Remark 4.3](#).

- i) This time we have $L^p_{l,\lambda}(\Lambda^k(T^*N)) \subseteq L^p_{l,\lambda'}(\Lambda^k(T^*N))$ if $\lambda < \lambda'$. (The inequality is in the opposite direction to the ICS case.)
- ii) It is easy to show that if $|\omega|_{g_N} = O(r^\lambda)$ near infinity, $\omega \in L^p_{0,\lambda+e}(\Lambda^k(T^*N))$ for any $e > 0$.
- iii) The space $L^p_{0,-\frac{7}{p}}(\Lambda^k(T^*N))$ is the usual $L^p(\Lambda^k(T^*N))$ space, and in particular we have

$$(140) \quad L^2_{0,-\frac{7}{2}}(\Lambda^k(T^*N)) = L^2(\Lambda^k(T^*N)).$$

- iv) The differential operators d and d^*_N , defined on $C^\infty_{cs}(\Lambda^k(T^*N))$, clearly extend to bounded linear operators

$$\begin{aligned} d: L^p_{l+1,\lambda}(\Lambda^k(T^*N)) &\rightarrow L^p_{l,\lambda-1}(\Lambda^{k+1}(T^*N)), \\ d^*_N: L^p_{l+1,\lambda}(\Lambda^k(T^*N)) &\rightarrow L^p_{l,\lambda-1}(\Lambda^{k-1}(T^*N)). \end{aligned}$$

Proposition 4.22 Let q satisfy $\frac{1}{p} + \frac{1}{q} = 1$. There is a Banach space isomorphism

$$(L^p_{0,\lambda}(\Lambda^k(T^*N)))^* \cong L^q_{0,-\lambda-7}(\Lambda^k(T^*N)),$$

given by the L^2 inner product pairing.

Corollary 4.23 Suppose $\alpha \in L^p_{1,\lambda+1}(\Lambda^k(T^*N))$ and $\beta \in L^q_{1,-\lambda-7}(\Lambda^{k+1}(T^*N))$, where $\frac{1}{p} + \frac{1}{q} = 1$. Then we can “integrate by parts” to say that

$$(141) \quad \langle d\alpha, \beta \rangle_{L_2} = \langle \alpha, d^*_M \beta \rangle_{L_2}.$$

This also holds if $\beta \in L^q_{1,\lambda'}(\Lambda^{k+1}(T^*N))$ whenever $\lambda + \lambda' < -7$.

For AC manifolds we will only be interested in the operator:

$$(142) \quad (d + d^*_N)^p_{l+1,\lambda}: L^p_{l+1,\lambda}(\Lambda^*(T^*N)) \rightarrow L^p_{l,\lambda-1}(\Lambda^*(T^*N)).$$

The AC analogues of Theorems 4.7, 4.8 and 4.9 are the following.

Theorem 4.24 The map $(d + d^*_N)^p_{l+1,\lambda}: L^p_{l+1,\lambda}(\Lambda^*(T^*N)) \rightarrow L^p_{l,\lambda-1}(\Lambda^*(T^*N))$ is Fredholm if and only if $\lambda \notin \mathcal{D}_{d+d^*_C}$, where the set of critical rates $\mathcal{D}_{d+d^*_C}$ is as given in Definition 4.6.

Theorem 4.25 Suppose that ω and ν are both locally integrable sections of $\Lambda^*(T^*N)$ and that ω is a weak solution of the equation $(d + d^*_N)(\omega) = \nu$. If $\omega \in L^p_{0,\lambda}(\Lambda^*(T^*N))$ and $\nu \in L^p_{l,\lambda-1}(\Lambda^*(T^*N))$, then $\omega \in L^p_{l+1,\lambda}(\Lambda^*(T^*N))$, and ω is a strong solution of $(d + d^*_N)(\omega) = \nu$. Furthermore, we have

$$(143) \quad \|\omega\|_{L^p_{l+1,\lambda}} \leq C (\|(d + d^*_N)(\omega)\|_{L^p_{l,\lambda-1}} + \|\omega\|_{L^p_{0,\lambda}})$$

for some constant $C > 0$ independent of ω .

Theorem 4.26 The kernel of $(d + d^*_N)^p_{l+1,\lambda}$ is independent of $p > 1$ and independent of l . Hence we can denote it unambiguously as $\ker(d + d^*_N)_\lambda$. This kernel is also invariant as we change the rate λ , as long as we do not hit any critical rates. That is, if the interval $[\lambda, \lambda']$ is contained in $\mathbb{R} \setminus \mathcal{D}_{d+d^*_C}$, then

$$\ker(d + d^*_N)_{\lambda'} = \ker(d + d^*_N)_\lambda.$$

Remark 4.27 As in Remark 4.10, it follows that

$$(144) \quad \omega \in \ker(d + d^*_N)_\lambda \implies |h^*(\omega)|_{h^*(g_N)} \leq Cr^\lambda \quad \text{on } (2R, \infty) \times \Sigma.$$

We will need to consider the formal adjoint of the map

$$(d + d^*_N)^p_{l+1,\lambda}: L^p_{l+1,\lambda}(\Lambda^*(T^*N)) \rightarrow L^p_{l,\lambda-1}(\Lambda^*(T^*N)).$$

By Proposition 4.22, the formal adjoint extends to a map

$$(145) \quad (d + d^*_N)^q_{m+1,-\lambda-6}: L^q_{m+1,-\lambda-6}(\Lambda^*(T^*N)) \rightarrow L^q_{m,-\lambda-7}(\Lambda^*(T^*N)),$$

where $\frac{1}{p} + \frac{1}{q} = 1$ and $l, m \geq 0$.

Next we have the ‘‘Fredholm Alternative’’ for AC manifolds.

Theorem 4.28 Suppose that λ is not in $\mathcal{D}_{d+d_C^*}$, so that by [Theorem 4.24](#), the map

$$(d + d_N^*)_{l+1,\lambda}^p : L_{l+1,\lambda}^p(\Lambda^*(T^*N)) \rightarrow L_{l,\lambda-1}^p(\Lambda^*(T^*N))$$

is Fredholm and also uniformly elliptic. Then the image $\text{im}(d + d_N^*)_{l+1,\lambda}^p$ of this map is given by

$$\begin{aligned} &\text{im}(d + d_N^*)_{l+1,\lambda}^p \\ &= \{v \in L_{l,\lambda-1}^p(\Lambda^*(T^*N)); \langle v, \omega \rangle_{L^2} = 0, \forall \omega \in \ker(d + d_N^*)_{-\lambda-6}\}. \end{aligned}$$

Let $\mathcal{K}_{d+d_C^*}$ be as given in [Definition 4.15](#). We will use $\ker_\infty(d + d_N^*)_\lambda$ to denote the restriction of the kernel of $(d + d_N^*)_{l+1,\lambda}^p$ to the subset $h((2R, \infty) \times \Sigma)$ of the asymptotic end of N . The following is the AC analogue of [Proposition 4.16](#).

Proposition 4.29 Suppose that β and β' are in $\mathbb{R} \setminus \mathcal{D}_{d+d_C^*}$ and that $\beta > \beta'$. Let $\alpha_1 < \alpha_2 < \dots < \alpha_N$ be all the critical rates in $\mathcal{D}_{d+d_C^*}$ between β and β' . If $\omega \in \ker_\infty(d + d_N^*)_\beta$, then there exist $v_j \in \mathcal{K}(\alpha_j)_{d+d_C^*}$, for $j = 1, \dots, N$, and an ω' defined near infinity with $|\omega'|_{g_C} = O(r^{\beta+\nu})$ such that

$$\omega - \sum_{j=1}^N (h^{-1})^*(v_j) - \omega' \in \ker_\infty(d + d_N^*)_{\beta'}.$$

That is, when restricted to the subset $h((2R, \infty) \times \Sigma)$ of the asymptotic end of N , an element in the kernel of $d + d_N^*$ with noncritical rate β admits an expansion in terms of elements in $\mathcal{K}(\alpha_j)_{d+d_C^*}$ for each critical rate α_j between β and β' , plus a remainder term which is of order $O(r^{\beta+\nu})$ and another term which is in the kernel of $d + d_N^*$ with noncritical rate $\beta' < \beta$.

Remark 4.30 The $O(r^{\beta+\nu})$ term arises from comparing a solution to $(d + d_N^*)(\omega) = 0$ to a solution of $(d + d_C^*)(\omega) = 0$, using the relation [\(57\)](#) between the metrics g_C and g_N near infinity.

From [Proposition 4.29](#) we can prove the following, exactly as in the proof of [Corollary 4.18](#).

Corollary 4.31 Suppose that $\lambda_1 < \lambda_2$. Choose $\delta > 0$ small enough so that in the closed interval $[\lambda_1 - \delta, \lambda_2 + \delta]$, the only critical rates for the asymptotic end lie in the closed interval $[\lambda_1, \lambda_2]$. That is, $[\lambda_1 - \delta, \lambda_2 + \delta] \cap \mathcal{D}_{d+d_C^*} \subseteq [\lambda_1, \lambda_2]$. Further assume

that $\delta > 0$ is small enough so that $\delta < \frac{1}{2}(\lambda_1 - \lambda_2 - \nu)$, where ν is the rate of the AC manifold N . (This will only be possible when $\nu < \lambda_1 - \lambda_2$.) If $\omega \in \Lambda^*(T^*N)$ is in $\ker_\infty(d + d_N^*)_{\lambda_2 + \delta}$, then

$$(146) \quad |h^*(\omega) - \sum_{j=1}^N v_j|_{g_C} = O(r^{\lambda_1 - \delta}) \quad \text{on } (2R, \infty) \times \Sigma,$$

for some v_j in $\mathcal{K}(\alpha_j)_{d+d_C^*}$, where $\alpha_1, \dots, \alpha_N$ are all the critical rates for $d + d_C^*$ in $[\lambda_1, \lambda_2]$.

5 Solution of the obstruction problem

In this section we prove [Theorem 3.10](#), which gives the existence of η and ξ on M' with the required properties to perform the desingularization and which was used in [Section 3.2](#). We restate the theorem here for the convenience of the reader.

Theorem 3.10 *Let M be a compact G_2 manifold with isolated conical singularities. For each i , let ξ_i and η_i be 3-forms and 4-forms, homogeneous of orders -3 and -4 , respectively, and closed and coclosed on each cone C_i , with respect to g_{C_i} . Suppose that*

$$(147) \quad \bigoplus_{i=1}^n [\xi_i] \in \bigoplus_{i=1}^n H^3(\Sigma_i, \mathbb{R}) \text{ lies in the image of } \Upsilon^3: H^3(M', \mathbb{R}) \rightarrow \bigoplus_{i=1}^n H^3(\Sigma_i, \mathbb{R}),$$

$$(148) \quad \bigoplus_{i=1}^n [\eta_i] \in \bigoplus_{i=1}^n H^4(\Sigma_i, \mathbb{R}) \text{ lies in the image of } \Upsilon^4: H^4(M', \mathbb{R}) \rightarrow \bigoplus_{i=1}^n H^4(\Sigma_i, \mathbb{R}).$$

where the maps Υ^k are given in [Definition 3.9](#). Then for $\delta > 0$ sufficiently small, there exists a smooth 3-form ξ and a smooth 4-form η on M' such that

$$(149) \quad \begin{aligned} d\xi &= 0, & d_{g_M}^* \xi &= 0, & d\eta &= 0, \\ |\nabla_{C_i}^j (f_i^*(\xi) - \xi_i)|_{g_{C_i}} &= O(r^{-3+\delta-j}) & \forall j \geq 0, \end{aligned}$$

$$(150) \quad |\nabla_{C_i}^j (f_i^*(\eta) - \eta_i)|_{g_{C_i}} = O(r^{-4+\delta-j}) \quad \forall j \geq 0.$$

Notice that we do not say that η is coclosed. Furthermore, the 3-form ξ is in Λ^3_{27} with respect to the G_2 structure φ_M .

Notice that if (149) and (150) hold for $j = 0$, then since ξ , η , ξ_i , and η_i are all closed and $\delta > 0$, Lemma 2.11 shows that the cohomology classes $[f_i^*(\xi) - \xi_i]$ and $[f_i^*(\eta) - \eta_i]$ vanish in $H^3(\Sigma_i, \mathbb{R})$ and $H^4(\Sigma_i, \mathbb{R})$, respectively. But by the definition of the maps Υ^k in Definition 3.9, this precisely says that $\Upsilon^3([\xi]) = \bigoplus_{i=1}^n [\xi_i]$ and $\Upsilon^4([\eta]) = \bigoplus_{i=1}^n [\eta_i]$. Hence the conditions in (147) and (148) are necessary for closed forms ξ and η to exist satisfying (149) and (150). These are therefore global topological conditions for the desingularization to be possible, which relate the different singular points. We will see in Sections 5.1 and 5.2 that the conditions in (147) and (148) are also sufficient to construct a ξ and η satisfying all the conclusions of Theorem 3.10.

5.1 Construction of the 3-form correction ξ

Let ξ' be any smooth 3-form on M' satisfying

$$(151) \quad |\nabla_{C_i}^j (f_i^*(\xi') - \xi_i)|_{g_{C_i}} = O(r^{-3+\delta-j}) \quad \forall j \geq 0.$$

This can clearly be done by defining $f_i^*(\xi')$ to be equal to ξ_i in a neighbourhood $(0, \frac{1}{2}\varepsilon) \times \Sigma_i$ of x_i , defining ξ' to be zero in the compact core K of M' , and by smoothly interpolating between these definitions on the annuli $[\frac{1}{2}\varepsilon, \varepsilon] \times \Sigma_i$. The goal of this section is to modify ξ' to produce a 3-form ξ with the properties given in Theorem 3.10.

Let $\Lambda^{\text{odd}}(T^*M')$ and $\Lambda^{\text{even}}(T^*M')$ denote the space of odd and even degree forms on M' , respectively. Since the operator $d + d_M^*$ interchanges the odd and even forms, all the results about $d + d_M^*$ from Section 4.1 remain true when we consider $d + d_M^*$ as mapping just from odd to even forms or conversely. We need to consider the map

$$(152) \quad P := (d + d_M^*)_{l+1, -3+\delta}^{\text{odd}, p} : L_{l+1, -3+\delta}^p(\Lambda^{\text{odd}}(T^*M')) \rightarrow L_{l, -4+\delta}^p(\Lambda^{\text{even}}(T^*M')).$$

By (136), the formal adjoint P^* of P extends to a map

$$(153) \quad P^* := (d + d_M^*)_{m+1, -3-\delta}^{\text{even}, q} : L_{m+1, -3-\delta}^q(\Lambda^{\text{even}}(T^*M')) \rightarrow L_{m, -4-\delta}^q(\Lambda^{\text{odd}}(T^*M')).$$

First, we have to describe the kernel of P^* explicitly.

Proposition 5.1 *If $v = \sum_{k=0}^3 v_{2k} \in L_{m+1,-3-\delta}^q(\Lambda^{\text{even}}(T^*M'))$ satisfies $P^*(v) = 0$, then on the neighbourhood $(0, \frac{1}{2}\varepsilon) \times \Sigma_i$ of x_i , we have*

$$(154) \quad f_i^*(v) = dr \wedge \alpha_{4,i} + O(r^{-3+\delta}),$$

where each $\alpha_{4,i}$ is a harmonic 3-form on the link Σ_i , respectively. Furthermore, each component v_{2k} of v is closed and coclosed with respect to the metric g_M on M' .

Proof The fact that $f_i^*(v)$ takes the form shown in (154) follows from Corollary 4.18 (with $\lambda = -3$) and Proposition 2.21. It remains to show that each v_{2k} is closed and coclosed. The equation $(d + d_M^*)(v) = 0$ breaks up into

$$(155) \quad dv_0 + d_M^*v_2 = 0, \quad dv_2 + d_M^*v_4 = 0, \quad dv_4 + d_M^*v_6 = 0, \quad dv_6 = 0.$$

Also, $(d + d_M^*)(v) = 0$ implies that $\Delta_M v_{2k} = 0$ for all $k = 0, \dots, 3$. Since $-3 - \delta > -4$, we see that v_0 and v_6 are both closed and coclosed, from (133) and (134), respectively. Therefore (155) simplifies to

$$(156) \quad d_M^*v_2 = 0, \quad dv_2 + d_M^*v_4 = 0, \quad dv_4 = 0.$$

Thus we will be done if we can show that $dv_2 = 0$. But dv_2 is a closed 3-form, and by the middle equation in (156), it is also coclosed. Now by Proposition 4.13 for $k = 3$, the map $dv_2 \mapsto [dv_2] \in H^3(M', \mathbb{R})$ is injective. But $[dv_2] = 0$, so $dv_2 = 0$ and the proof is complete. \square

Theorem 5.2 *Let ξ' be a smooth 3-form on M' , satisfying (151), and further suppose that the condition (147) is satisfied. Then there exists $\omega = \sum_{k=0}^3 \omega_{2k+1} \in L_{l+1,-3+\delta}^p(\Lambda^{\text{odd}}(T^*M'))$ satisfying the equation*

$$(157) \quad (d + d_M^*)(\omega) = -d_M^*\xi' - d\xi'.$$

Moreover, using this solution, if we define $\tilde{\xi} = \xi' + \omega_3$, then $\tilde{\xi} \in L_{l+1,-3}^p(\Lambda^3(T^*M'))$ and satisfies $d\tilde{\xi} = 0$ and $d_M^*\tilde{\xi} = 0$.

Proof By Theorem 4.14, we need to check that the right hand side of the equation is orthogonal (with respect to the L^2 inner product) to the kernel of the formal adjoint map. Let $v = \sum_{k=0}^3 v_{2k}$ be in the kernel of P^* . Then

$$(158) \quad \langle -d_M^*\xi' - d\xi', v \rangle_{L^2} = \int_{M'} \langle -d_M^*\xi', v_2 \rangle_{g_M} \text{vol}_M + \int_{M'} \langle -d\xi', v_4 \rangle_{g_M} \text{vol}_M.$$

On 3-forms we have $d_M^* = -*_M d*_M$, and $*_M$ is an isometry with respect to g_M , so therefore we can write $\langle -d_M^*\xi', v_2 \rangle_{g_M} = \langle d*_M\xi', *_M v_2 \rangle_{g_M}$. Hence Equation (158)

simplifies to

$$\begin{aligned}
 \langle -d_M^* \xi' - d \xi', v \rangle_{L^2} &= \int_{M'} (d *_{M'} \xi' \wedge v_2) - \int_{M'} (d \xi' \wedge *_{M'} v_4) \\
 (159) \qquad \qquad \qquad &= \int_{M'} d(*_{M'} \xi' \wedge v_2) - \int_{M'} d(\xi' \wedge *_{M'} v_4),
 \end{aligned}$$

where we have used the fact that $d v_2 = 0$ and $d *_{M'} v_4 = 0$ from [Proposition 5.1](#). Let us define $M'_r = \{x \in M'; \varrho(x) \geq r\}$, where ϱ is the radius function on M' from [Definition 4.1](#). Then M'_r is a manifold with boundary, and $\partial M'_r = \bigsqcup_{i=1}^n f_i(\{r\} \times \Sigma_i)$. Then using Stokes' Theorem, [Equation \(159\)](#) becomes

$$\begin{aligned}
 (160) \quad \langle -d_M^* \xi' - d \xi', v \rangle_{L^2} &= \lim_{r \rightarrow 0} \sum_{i=1}^n \int_{\{r\} \times \Sigma_i} f_i^*(*_{M'} \xi') \wedge f_i^*(v_2) \\
 &\quad - \lim_{r \rightarrow 0} \sum_{i=1}^n \int_{\{r\} \times \Sigma_i} f_i^*(\xi') \wedge f_i^*(*_{M'} v_4).
 \end{aligned}$$

By [\(151\)](#) and the fact that $|\xi_i|_{g_{C_i}} = O(r^{-3})$, we have $|f_i^*(\xi')|_{g_{C_i}} = O(r^{-3})$. The proof of [\(79\)](#) also works for ξ' , since only the property [\(151\)](#) is used. Therefore we also have $|f_i^*(*_{M'} \xi')|_{g_{C_i}} = O(r^{-3})$. [Proposition 5.1](#) tells us that $|f_i^*(v_2)|_{g_{C_i}} = O(r^{-3+\delta})$, and $|f_i^*(v_4) - dr \wedge \alpha_{4,i}|_{g_{C_i}} = O(r^{-3+\delta})$ on $\{r\} \times \Sigma_i$, where $\alpha_{4,i}$ is a harmonic 3-form on Σ_i . Now we can mimic the proof of [\(79\)](#) using v_4 and $dr \wedge \alpha_{4,i}$ in place of ξ and ξ_i , respectively, to obtain that $|f_i^*(*_{M'} v_4) - *_{C_i}(dr \wedge \alpha_{4,i})|_{g_{C_i}} = O(r^{-3+\delta})$. Using [\(24\)](#) we see that $*_{C_i}(dr \wedge \alpha_{4,i}) = *_{\Sigma_i} \alpha_{4,i}$. In summary, we have:

$$\begin{aligned}
 (161) \quad &|f_i^*(*_{M'} \xi')|_{g_{C_i}} = O(r^{-3}), & |f_i^*(v_2)|_{g_{C_i}} &= O(r^{-3+\delta}), \\
 &|f_i^*(\xi') - \xi_i|_{g_{C_i}} = O(r^{-3+\delta}), & |f_i^*(*_{M'} v_4) - *_{\Sigma_i} \alpha_{4,i}|_{g_{C_i}} &= O(r^{-3+\delta}), \\
 &|f_i^*(\xi')|_{g_{C_i}} = O(r^{-3}), & |*_{\Sigma_i} \alpha_{4,i}|_{g_{C_i}} &= O(r^{-3}).
 \end{aligned}$$

Writing $f_i^*(\xi') = \xi_i + (f_i^*(\xi') - \xi_i)$ and $f_i^*(*_{M'} v_4) = *_{\Sigma_i} \alpha_{4,i} + (f_i^*(*_{M'} v_4) - *_{\Sigma_i} \alpha_{4,i})$, we can then apply the estimates in [\(161\)](#) to [Equation \(160\)](#), to obtain

$$\begin{aligned}
 \langle -d_M^* \xi' - d \xi', v \rangle_{L^2} &\leq \lim_{r \rightarrow 0} \sum_{i=1}^n \int_{\{r\} \times \Sigma_i} C r^{-6+\delta} \text{vol}_{\{r\} \times \Sigma_i} \\
 &\quad + \left| \lim_{r \rightarrow 0} \sum_{i=1}^n \int_{\{r\} \times \Sigma_i} \xi_i \wedge *_{\Sigma_i} \alpha_{4,i} \right| \\
 (162) \qquad \qquad \qquad &\leq \lim_{r \rightarrow 0} C r^\delta + \left| \lim_{r \rightarrow 0} \sum_{i=1}^n \int_{\{r\} \times \Sigma_i} \xi_i \wedge *_{\Sigma_i} \alpha_{4,i} \right|.
 \end{aligned}$$

using the fact that the volume of $\{r\} \times \Sigma_i$ is equal to Cr^6 , where C is the volume of $\{1\} \times \Sigma_i$. The first limit in (162) is zero, so to conclude the solvability of Equation (157) it remains to prove that

$$(163) \quad \lim_{r \rightarrow 0} \sum_{i=1}^n \int_{\{r\} \times \Sigma_i} \xi_i \wedge *_{\Sigma_i} \alpha_{4,i} = 0.$$

This is where we will need to use the hypothesis that the condition (147) is satisfied. This condition says that there exists a smooth closed 3-form ξ'' on M' such that $f_i^*(\xi'')$ differs from ξ_i on $(0, \varepsilon) \times \Sigma_i$ by an exact piece. That is,

$$(164) \quad \xi_i = f_i^*(\xi'') + dV_i \quad \text{on } (0, \varepsilon) \times \Sigma_i$$

for some smooth 2-forms V_i defined on $(0, \varepsilon) \times \Sigma_i$. We also know that $*_{\Sigma_i} \alpha_{4,i}$ is closed (since $\alpha_{4,i}$ is harmonic on Σ_i) and also that $f_i^*(\nu_4)$ is closed by Proposition 5.1. Therefore the fourth equation in (161) and Lemma 2.11 together imply that $f_i^>(*_M \nu_4)$ differs from $*_{\Sigma_i} \alpha_{4,i}$ on $(0, \varepsilon) \times \Sigma_i$ by an exact piece. That is,

$$(165) \quad *_{\Sigma_i} \alpha_{4,i} = f_i^>(*_M \nu_4) + dW_i \quad \text{on } (0, \varepsilon) \times \Sigma_i$$

for some smooth 2-forms W_i defined on $(0, \varepsilon) \times \Sigma_i$. Using (164) and (165), the left hand side of Equation (163) now becomes

$$\begin{aligned} \lim_{r \rightarrow 0} \left(\sum_{i=1}^n \int_{\{r\} \times \Sigma_i} f_i^*(\xi'') \wedge f_i^>(*_M \nu_4) \right) \\ + \lim_{r \rightarrow 0} \left(\sum_{i=1}^n \int_{\{r\} \times \Sigma_i} (dV_i \wedge *_{\Sigma_i} \alpha_{4,i} + \xi_i \wedge dW_i + dV_i \wedge dW_i) \right). \end{aligned}$$

Since ξ_i and $*_{\Sigma_i} \alpha_{4,i}$ are closed, the integrands in the second sum of integrals above are all exact, and hence these integrals all vanish by Stokes' Theorem, because $\{r\} \times \Sigma_i$ is compact and without boundary. Therefore the left hand side of Equation (163) has been simplified to

$$(166) \quad \lim_{r \rightarrow 0} \sum_{i=1}^n \int_{f_i(\{r\} \times \Sigma_i)} \xi'' \wedge *_M \nu_4 = \lim_{r \rightarrow 0} \int_{\partial M'_r} \xi'' \wedge *_M \nu_4 = \lim_{r \rightarrow 0} \int_{M'_r} d(\xi'' \wedge *_M \nu_4) \\ = \int_{M'} d(\xi'' \wedge *_M \nu_4),$$

using Stokes' Theorem again. But this integral vanishes since both ξ'' and $*_M \nu_4$ are closed, by Proposition 5.1 and the hypothesis (147) which ensures the existence of such a closed ξ'' . Hence the right hand side of (157) is L^2 -orthogonal to the

kernel of P^* , and therefore there exists a solution $\omega = \sum_{k=0}^3 \omega_{2k+1}$ to Equation (157). It still remains to show that $\tilde{\xi} = \xi' + \omega_3$ is closed and coclosed. The equation $(d + d_M^*)(\omega) = -d_M^* \xi' - d\xi'$ breaks up into

$$(167) \quad \begin{aligned} d_M^* \omega_1 &= 0, & d\omega_1 + d_M^* \omega_3 &= -d_M^* \xi', \\ d\omega_3 + d_M^* \omega_5 &= -d\xi', & d\omega_5 + d_M^* \omega_7 &= 0. \end{aligned}$$

Now (167) implies that $\Delta_M \omega_{2k+1} = 0$ for $k = 0, 2, 3$. Since $-3 + \delta > -3$, we see that ω_1, ω_5 , and ω_7 are all closed and coclosed, from (134), (135) and (133), respectively. Therefore (167) simplifies to

$$d_M^*(\omega_3 + \xi') = 0, \quad d(\omega_3 + \xi') = 0.$$

Also, $f_i^*(\xi') = \xi_i$ near x_i , and $|\xi_i|_{g_{C_i}} = O(r^{-3})$. Therefore we have

$$(168) \quad |\nabla_{C_i}^j (f_i^*(\tilde{\xi}) - \xi_i)|_{g_{C_i}} = O(r^{-3+\delta-j}) \quad \forall j \geq 0,$$

and hence $\tilde{\xi} = \omega_3 + \xi'$ is in $L_{l+1,-3}^p(\Lambda^3(T^*M'))$ and is closed and coclosed. \square

We have constructed a smooth 3-form $\tilde{\xi}$ on M' which is closed and coclosed and satisfies (149). The final step is to modify $\tilde{\xi}$ to a 3-form ξ which lies in Ω_{27}^3 with respect to φ_M .

Proposition 5.3 *Let $\tilde{\xi}$ be as given in Theorem 5.2. Denote by $\tilde{\xi} = \tilde{\xi}_1 + \tilde{\xi}_7 + \tilde{\xi}_{27}$ the decomposition of $\tilde{\xi}$ into components of the subspaces $\Omega^3 = \Omega_1^3 \oplus \Omega_7^3 \oplus \Omega_{27}^3$ determined by φ_M . Then the 3-form $\xi = \tilde{\xi}_{27}$ is closed and coclosed and satisfies (149).*

Proof The decomposition of Ω^3 is orthogonal with respect to g_M and preserved by the covariant derivative (since φ_M is torsion-free), so each $\tilde{\xi}_1, \tilde{\xi}_7$ and $\tilde{\xi}_{27}$ is in $L_{l+2,-3+\delta}^p(\Lambda^3(T^*M'))$. Since $d\tilde{\xi} = 0$ and $d_M^* \tilde{\xi} = 0$, we have $\Delta_M \tilde{\xi} = 0$. Again because φ_M is torsion-free, the Laplacian commutes with the projections onto the G_2 invariant subspaces, so $\Delta_M \tilde{\xi}_1 = 0$. But $\tilde{\xi}_1 = f\varphi_M$ for some function f in $L_{l+2,-3+\delta}^p(\Lambda^0(T^*M'))$, and the first equation in (5) says that $\Delta_M f = 0$. Hence by (133) we see that f is closed (hence constant), and therefore $\tilde{\xi}_1 = f\varphi_M$ is closed and coclosed. Similarly $\Delta_M \tilde{\xi}_7 = 0$, and since $\tilde{\xi}_7 = *\alpha$ for some 1-form α in $L_{l+2,-3+\delta}^p(\Lambda^1(T^*M'))$, the first equation in (6) says that $\Delta_M \alpha = 0$. Proposition 2.15 says there are no critical rates for the Laplacian on 1-forms in the interval $(-4, -1)$, so by Theorem 4.9, we can say α is in $\ker(\Delta_M)_{-1-e}$ for some small $e > 0$, and then Theorem 4.8 tells us $\alpha \in L_{l+2,-1-e}^p(\Lambda^1(T^*M'))$. Therefore $\nabla_M \alpha$ lies in $L_{l+1,-2-e}^p(T^*M' \otimes T^*M')$, which is in L^2 because $-2 - e > -\frac{7}{2}$. (Here we are using an $L_{l,\lambda}^p$ norm on sections of the tensor bundle $T^*M' \otimes T^*M'$, and the

analogous statement to iii) from Remark 4.3.) On a manifold with vanishing Ricci curvature, the usual Bochner–Weitzenböck formula reduces to $\Delta = \nabla^* \nabla$ on 1-forms. Therefore (using an obvious analogue of Corollary 4.5) we can compute that

$$0 = \langle \Delta_M \alpha, \alpha \rangle_{L^2} = \langle \nabla_M^* \nabla_M \alpha, \alpha \rangle_{L^2} = |\nabla_M \alpha|_{L^2}^2,$$

and thus we see that $\nabla_M \alpha = 0$. But a manifold with holonomy exactly equal to G_2 has no nonzero parallel 1-forms [8, Theorem 2], so $\alpha = 0$, and hence $\tilde{\xi}_7 = 0$.

Thus we have $\tilde{\xi}_{27} = \tilde{\xi} - \tilde{\xi}_1$, and since $\tilde{\xi}$ and $\tilde{\xi}_1$ are both closed and coclosed, we see that $\xi = \tilde{\xi}_{27}$ is also closed and coclosed. We still need to show that ξ satisfies (149). Let π_{27}^M denote the projection onto the Ω_{27}^3 subspace of $\Omega^3(T^*M')$ with respect to φ_M and $\pi_{27}^{C_i}$ denote the projection onto the Ω_{27}^3 subspace of $\Omega^3(T^*C_i)$ with respect to φ_{C_i} . Because of (53), the G_2 structures $f_i^*(\varphi_M)$ and φ_{C_i} on $(0, \varepsilon) \times \Sigma_i$ agree up to order $O(r^{\mu_i})$, and thus we have

$$|f_i^*(\pi_{27}^M \omega) - \pi_{27}^{C_i} f_i^*(\omega)|_{g_{C_i}} = |f_i^*(\omega)|_{g_{C_i}} O(r^{\mu_i})$$

for any 3-form ω on M' . Since $\xi = \pi_{27}^M \tilde{\xi}$, the above equation becomes

$$|f_i^*(\xi) - \pi_{27}^{C_i} f_i^*(\tilde{\xi})|_{g_{C_i}} = |f_i^*(\tilde{\xi})|_{g_{C_i}} O(r^{\mu_i}) = O(r^{-3+\mu_i}).$$

But then we have

$$\begin{aligned} |f_i^*(\xi) - \xi_i|_{g_{C_i}} &\leq |f_i^*(\xi) - \pi_{27}^{C_i} f_i^*(\tilde{\xi})|_{g_{C_i}} + |\pi_{27}^{C_i} f_i^*(\tilde{\xi}) - \xi_i|_{g_{C_i}} \\ &= O(r^{-3+\mu_i}) + |\pi_{27}^{C_i} (f_i^*(\tilde{\xi}) - \xi_i)|_{g_{C_i}} \\ &\leq O(r^{-3+\mu_i}) + O(r^{-3+\delta}) = O(r^{-3+\delta}), \end{aligned}$$

where we have used the fact that $\pi_{27}^{C_i} \xi_i = \xi_i$ from Proposition 2.23, Equation (168) and the fact that $\delta < \mu_i$. Similarly we can show that

$$|\nabla_{C_i}^j (f_i^*(\xi) - \xi_i)|_{g_{C_i}} = O(r^{-3+\delta-j}) \quad \forall j \geq 0,$$

and the proof is complete. □

We have succeeded in constructing the 3-form ξ of Theorem 3.10 needed to correct both φ_M and ψ_M for the gluing of Section 3.3 to work. We still need to construct the 4-form correction η .

5.2 Construction of the 4-form correction η

We can try to follow the same procedure as in Section 5.1 to construct the 4-form correction η . However, in this case one encounters some difficulties. The main problem

is that the analogue of Proposition 2.21 fails. The relevant rate in this case is -2 , and it turns out that there *can* be $\log(r)$ terms in the kernel of $d + d_C$ with order -2 on odd forms. In fact it turns out that the \log terms arise only for the 3-form components of the kernel. Because of this, the proof of an exact 4-form analogue to Theorem 5.2 would break down, but one can carry out this procedure partially to produce a 4-form η which is merely closed, but *not* coclosed. The arguments become much more involved, however. And then the analogous argument to Proposition 5.3 for projection onto the Ω_{27}^4 component would also break down.

However, we saw in Section 3.3 that it was unnecessary for η to be coclosed or to be in Ω_{27}^4 , just that it be closed. Since that and condition (150) are our only requirements on η , there is a much simpler argument that avoids all the technical analysis of Section 5.1.

Proposition 5.4 *Suppose that condition (148) is satisfied. Then there exists a smooth closed 4-form η on M' satisfying*

$$|\nabla_{C_i}^j (f_i^*(\eta) - \eta_i)|_{g_{C_i}} = O(r^{-4+\delta-j}) \quad \forall j \geq 0,$$

on $(0, \varepsilon) \times \Sigma_i$, for any $\delta > 0$.

Proof As in the discussion before Equation (164), the fact that condition (148) is satisfied says that there exists a smooth *closed* 4-form η'' on M' such that $f_i^*(\eta'')$ differs from η_i on $(0, \varepsilon) \times \Sigma_i$ by an exact piece. That is,

$$(169) \quad \eta_i = f_i^*(\eta'') + dU_i \quad \text{on } (0, \varepsilon) \times \Sigma_i$$

for some smooth 3-forms U_i defined on $(0, \varepsilon) \times \Sigma_i$. Let $w: (0, \infty) \rightarrow \mathbb{R}$ be any smooth decreasing function such that

$$w(r) = \begin{cases} 1 & \text{for } 0 < r \leq \frac{4}{6}\varepsilon, \\ 0 & \text{for } \frac{5}{6}\varepsilon \leq r < \infty. \end{cases}$$

Now we define the smooth 4-form η on M' as follows:

$$(170) \quad \eta = \begin{cases} \eta'' & \text{on the compact core } K \text{ of } M', \\ \eta'' + (f_i^{-1})^*(d(wU_i)) & \text{on } f_i([\frac{1}{2}\varepsilon, \varepsilon] \times \Sigma_i), \\ (f_i^{-1})^*(\eta_i) & \text{on } f_i((0, \frac{1}{2}\varepsilon] \times \Sigma_i). \end{cases}$$

It is clear that η is closed, and by Equation (169) and the definition of w it is also smooth and well-defined. Also, on $f_i((0, \frac{1}{2}\varepsilon) \times \Sigma_i)$, we have $f_i^*(\eta) = \eta_i$, so

$$|\nabla_{C_i}^j (f_i^*(\eta) - \eta_i)|_{g_{C_i}} = O(r^{-4+\delta-j}) \quad \forall j \geq 0,$$

for any $\delta > 0$, since it is identically equal to zero for $r < \frac{1}{2}\varepsilon$. □

We have succeeded in constructing the 4–form η of [Theorem 3.10](#) needed to correct ψ_M for the gluing of [Section 3.3](#) to work. Thus the proof of [Theorem 3.10](#) is now complete.

6 Asymptotic expansion of G_2 structures on AC G_2 manifolds

In this section we prove [Theorem 3.6](#), which says that under an appropriate *gauge-fixing condition* we can obtain a nice asymptotic expansion of the G_2 structure (φ_N, ψ_N) on an AC G_2 manifold N . This theorem was used in [Section 3.2](#) for the gluing construction. We restate the theorem here for the convenience of the reader and drop the i subscripts to simplify notation.

Theorem 3.6 *Suppose that N is an asymptotically conical G_2 manifold with rate $\nu \leq -3$ and that h satisfies the gauge-fixing condition given in [Definition 3.3](#). Then on the subset $(2R, \infty) \times \Sigma$ of the cone C we can write*

$$(171) \quad h^*(\varphi_N) = \varphi_C + \xi + d\zeta,$$

$$(172) \quad h^*(\psi_N) = \psi_C + \eta - *\zeta\xi + d\theta.$$

where ξ is a harmonic 3–form, homogeneous of order -3 , and in Ω_{27}^3 with respect to φ_C , η is a harmonic 4–form, homogeneous of order -4 , and ζ and θ are 2–forms and 3–forms on $(2R, \infty) \times \Sigma$, respectively, satisfying

$$(173) \quad |\nabla_C^j \zeta|_{g_C} = O(r^{\nu'+1-j}), \quad |\nabla_C^j \theta|_{g_C} = O(r^{\nu'+1-j}), \quad \forall j \geq 0,$$

where $\nu' = -4$. Furthermore, $[\xi] = \Phi(N)$ and $[\eta] = \Psi(N)$, where $\Phi(N)$ and $\Psi(N)$ are the cohomological invariants of the AC G_2 manifold N from [Definition 2.38](#).

We will begin by showing that the gauge-fixing condition leads to an elliptic equation. Let N be an asymptotically conical G_2 manifold with rate $\nu \leq -3$. Then we have that

$$|h^*(\varphi_N) - \varphi_C|_{g_C} = O(r^\nu),$$

by [Equation \(56\)](#). Since ν can equal -3 , the invariant $\Phi(N)$ in $H^3(\Sigma, \mathbb{R})$ of [Definition 2.38](#) can be nonzero. By [Proposition 2.24](#), we can represent the class $\Phi(N)$ by a 3–form ξ on $(R, \infty) \times \Sigma$ which is homogeneous of order -3 and harmonic with respect to the cone metric. Furthermore, [Proposition 2.23](#) says that ξ is in Λ_{27}^3 with respect to φ_C . Now $h^*(\varphi_N) - \varphi_C - \xi$ is exact, so we can write

$$h^*(\varphi_N) - \varphi_C = \xi + d\zeta$$

for some 2–form ζ on $(R, \infty) \times \Sigma$, with $|d\zeta|_{g_C} = O(r^{-3})$.

Lemma 6.1 *Suppose that N is an AC G_2 manifold with rate $\nu \leq -3$ and that h satisfies the gauge-fixing condition of Definition 3.3. Then we have*

$$(174) \quad |(d + d_C^*)(h^*(\psi_N) - \psi_C)|_{g_C} = O(r^{-7}) \quad \text{on } (R, \infty) \times \Sigma.$$

Proof We have $h^*(\varphi_N) = \varphi_C + \xi + d\zeta$, where by hypothesis $\xi + d\zeta \in \Omega_{2,7}^3$ with respect to φ_C . By Lemma 1.3, we see that

$$\begin{aligned} h^*(\psi_N) &= \Theta(h^*(\varphi_N)) = \psi_C - *_C(\xi + d\zeta) + F_{\varphi_C}(\xi + d\zeta) \\ &= \psi_C - *_C\xi - d_C^*(*_C\zeta) + F_{\varphi_C}(\xi + d\zeta), \end{aligned}$$

where we have used the fact that $*_C d = d_C^* *_C$ on 2-forms. From Equation (25), we know that ξ is coclosed, since ξ is a closed and coclosed form on the link Σ . Therefore if we take d_C^* of both sides of the above equation, we get $d_C^*(h^*(\psi_N)) = d_C^*(F_{\varphi_C}(\xi + d\zeta))$, and from (8) and the fact that $|\xi + d\zeta|_{g_C} = O(r^{-3})$, we see that

$$|d_C^*(h^*(\psi_N))|_{g_C} = |d_C^*(F_{\varphi_C}(\xi + d\zeta))|_{g_C} \leq Cr^{-7}.$$

Equation (174) now follows since $(d + d_C^*)(h^*(\psi_N) - \psi_C) = d_C^*(h^*(\psi_N))$. □

Corollary 6.2 *Suppose that N is an AC G_2 manifold with rate $\nu \leq -3$ and that h satisfies the gauge-fixing condition. Then $\psi_N - (h^{-1})^*(\psi_C)$ satisfies the equation*

$$(175) \quad (d + d_N^*)(\psi_N - (h^{-1})^*(\psi_C)) = d_N^* \nu \quad \text{on } h((R, \infty) \times \Sigma),$$

for some exact 4-form ν defined on the subset $h((R, \infty) \times \Sigma)$ of N , with $|d_N^* \nu|_{g_N} = O(r^{-7})$.

Proof We begin by observing that

$$|(d + d_N^*)(\psi_N - (h^{-1})^*(\psi_C))|_{g_N} = |(d + d_{h^*(N)}^*)(h^*(\psi_N) - \psi_C)|_{h^*(g_N)}$$

where $d_{h^*(N)}^*$ is the coderivative with respect to the metric $h^*(g_N)$ on $(R, \infty) \times \Sigma$. Now by (57), the tensor $h^*(g_N) - g_C$ is uniformly bounded with respect to the g_C metric, on $(R, \infty) \times \Sigma$. Thus $|\omega|_{h^*(g_N)} \leq C|\omega|_{g_C}$ for any tensor ω , and therefore

$$|(d + d_N^*)(\psi_N - (h^{-1})^*(\psi_C))|_{g_N} \leq C|(d + d_{h^*(N)}^*)(h^*(\psi_N) - \psi_C)|_{g_C}.$$

But we also have

$$\begin{aligned} |(d + d_{h^*(N)}^*)(h^*(\psi_N) - \psi_C)|_{g_C} \\ \leq |(d + d_C^*)(h^*(\psi_N) - \psi_C)|_{g_C} + |(d_{h^*(N)}^* - d_C^*)(h^*(\psi_N) - \psi_C)|_{g_C}. \end{aligned}$$

The first term on the right hand side is $O(r^{-7})$ by Lemma 6.1. For the second term, we use (57) again to see that

$$|(d_{h^*(N)}^* - d_C^*)(h^*(\psi_N) - \psi_C)|_{g_C} \leq Cr^{\nu-1} |h^*(\psi_N) - \psi_C|_{g_C} = O(r^{-7}),$$

since $|h^*(\psi_N) - \psi_C|_{g_C} = O(r^{-3})$ and $\nu \leq -3$. Putting these all together gives

$$|(d + d_N^*)(\psi_N - (h^{-1})^*(\psi_C))|_{g_N} \leq Cr^{-7}.$$

It is clear that $(d + d_N^*)(\psi_N - (h^{-1})^*(\psi_C)) = d_N^* \nu$ with $\nu = (h^{-1})^*(\psi_C)$, and we know that ν is exact since ψ_C is exact by Proposition 2.4. \square

In order to obtain a nice asymptotic expansion of $h^*(\varphi_N)$ and $h^*(\psi_N)$, we will use Corollary 4.31, but first we require one more preliminary result.

Lemma 6.3 *There exists a closed 4-form ω_4 in $L_{1,-5}^P(\Lambda^4(T^*N))$ such that*

$$(176) \quad (d + d_N^*)(\psi_N - (h^{-1})^*(\psi_C) - \omega_4) = 0$$

on the subset $h((2R, \infty) \times \Sigma)$ of N .

Proof Let ϑ be a smooth bounded function on N such that

$$(177) \quad \vartheta(x) = \begin{cases} 0 & \text{for } x \in L, \\ 1 & \text{for } x \in h((2R, \infty) \times \Sigma). \end{cases}$$

By Corollary 6.2, we know that $\nu = d\alpha$ for some 3-form α on $h((R, \infty) \times \Sigma)$. Now the form $d_N^*d(\vartheta\alpha)$ is a well defined 3-form on all of N which is coexact. It is identically equal to zero on the compact subset L of N and equals $d_N^*\nu = d_N^*d\alpha$ on the subset $h((2R, \infty) \times \Sigma)$ of N . By item ii) of Remark 4.21, we know that $d_N^*(\vartheta\nu)$ is in $L_{0,-7+e}^P(\Lambda^{\text{odd}}(T^*N))$ for any $e > 0$. Consider the operator

$$(178) \quad (d + d_N^*)_{1,\lambda}^{\text{even},P}: L_{1,\lambda}^P(\Lambda^{\text{even}}(T^*N)) \rightarrow L_{0,\lambda-1}^P(\Lambda^{\text{odd}}(T^*N)).$$

We claim that for any $\lambda > -6 + e$ which is *not* a critical rate, we can solve the equation

$$(179) \quad (d + d_N^*)_{1,\lambda}^P(\omega) = d_N^*d(\vartheta\alpha),$$

where $\omega = \sum_{k=0}^3 \omega_{2k}$ is an even-degree mixed form. Note that this makes sense because since $d_N^*d(\vartheta\alpha)$ is in $L_{0,-7+e}^P(\Lambda^{\text{odd}}(T^*N))$, it follows from item i) of Remark 4.21 that $d_N^*d(\vartheta\alpha)$ is in $L_{0,\lambda-1}^P(\Lambda^{\text{odd}}(T^*N))$ for any $\lambda > -6 + e$. To know that (179) is solvable for such λ , we need to use Theorem 4.28. If λ is not critical then it will be

solvable if and only if $d_N^*d(\vartheta\alpha)$ is L^2 -orthogonal to the kernel of the formal adjoint of (178), which by (145) and Theorem 4.26 is the kernel of the map

$$(d + d_N^*)_{m+1, -\lambda-6}^{\text{odd}, q}: L_{m+1, -\lambda-6}^q(\Lambda^{\text{odd}}(T^*N)) \rightarrow L_{m, -\lambda-7}^q(\Lambda^{\text{even}}(T^*N)).$$

Suppose $\beta = \sum_{k=0}^3 \beta_{2k+1}$ is in the kernel of $(d + d_N^*)_{m+1, -\lambda-6}^q$. Then in particular $d\beta_3 + d_N^*\beta_5 = 0$. Then using Corollary 4.23 twice, we see that

$$\begin{aligned} \langle \beta, d_N^*d(\vartheta\alpha) \rangle_{L^2} &= \langle \beta_3, d_N^*d(\vartheta\alpha) \rangle_{L^2} = \langle d\beta_3, d(\vartheta\alpha) \rangle_{L^2} \\ &= -\langle d_N^*\beta_5, d(\vartheta\alpha) \rangle_{L^2} = -\langle \beta_5, dd(\vartheta\alpha) \rangle_{L^2} = 0, \end{aligned}$$

and thus Equation (179) has a solution $\omega = \sum_{k=0}^3 \omega_{2k}$ in $L_{1, \lambda}^p(\Lambda^{\text{even}}(T^*N))$. This ω satisfies

$$(180) \quad d\omega_0 + d_N^*\omega_2 = 0, \quad d\omega_2 + d_N^*\omega_4 = d_N^*d(\vartheta\alpha), \quad d\omega_4 + d_N^*\omega_6 = 0, \quad d\omega_6 = 0.$$

Every term in (180) lies in $L_{0, \lambda-1}^p(\Lambda^{\text{odd}}(T^*N))$. Therefore if $\lambda - 1 < -\frac{7}{2}$, we can use Corollary 4.23 again to integrate by parts and conclude that

$$d\omega_0 = d_N^*\omega_2 = 0, \quad d\omega_2 = d_N^*\omega_4 - d_N^*d(\vartheta\alpha) = 0, \quad d\omega_4 = d_N^*\omega_6 = 0, \quad d\omega_6 = 0.$$

Hence in particular ω_4 is closed and $(d + d_N^*)(\omega_4) = d_N^*d(\vartheta\alpha)$ if $\lambda \in (-6 + \epsilon, 1 - \frac{7}{2})$. If we choose λ to be any noncritical rate less than -5 , we have $\omega_4 \in L_{1, -5}^p(\Lambda^4(T^*N))$. Then by (175) and (177) we see that

$$(d + d_N^*)(\psi_N - (h^{-1})^*(\psi_C) - \omega_4) = 0$$

on the region $h((2R, \infty) \times \Sigma)$. □

Remark 6.4 The elliptic regularity result in Theorem 4.25, combined with the smoothness of ψ_C and ψ_N , now tell us that ω_4 is smooth. Therefore we have that

$$(181) \quad |\nabla_N^j \omega_4|_{g_N} = O(r^{-5-j}) \quad \text{for all } j \geq 0.$$

We can finally prove Theorem 3.6.

Proof of Theorem 3.6. From Lemma 6.3 we know that $(\psi_N - (h^{-1})^*(\psi_C) - \omega_4)$ is in $\ker_{\infty}(d + d_N^*)_{-3+\delta}$ for any $\delta > 0$. Let $\lambda_1 = -4$ and $\lambda_2 = -3$. Since $\nu \leq -3$, we have $\nu < \lambda_1 - \lambda_2 = -1$, and thus we can apply Corollary 4.31 to obtain

$$(182) \quad h^*(\psi_N) - \psi_C - h^*(\omega_4) = \sum_{j=1}^N v_j + v',$$

where v_j is in $\mathcal{K}(\alpha_j)_{d+d_C^*}$ for α_j a critical rate in the interval $[-4, -3]$, and $|v'|_{g_C} = O(r^{-4-\delta})$. Now *a priori*, the v_j 's are even-degree mixed forms. However, since the left hand side of (182) consists of only 4-forms, we know that the v_j 's and v' are also pure 4-forms. Thus each v_j is of the form $v_j = \sum_{i=1}^m (\log(r))^i v_{j,i}$ where $v_{j,i}$ is a homogeneous 4-form of order α_j on the cone. From the calculation in the proof of Proposition 2.21, it follows that the leading coefficient $v_{j,m}$ is a 4-form, homogeneous of order -4 , and closed and coclosed. But then Corollary 2.17 says that $v_{j,m} = 0$ (and hence $v_j = 0$) for any $\alpha_j \in (-4, -3)$. So the only critical rates which can occur in (182) are $\alpha_1 = -4$ and $\alpha_2 = -3$. Proposition 2.22 and Corollary 2.19 tell us that $\alpha_1 = \eta$, a homogeneous 4-form of order -4 , harmonic with respect to the cone metric. Also, Propositions 2.21 and 2.20 tell us that $\alpha_2 = dr \wedge (*_{\Sigma}\xi)$ for some homogeneous 3-form $*_{\Sigma}\xi$ of order -3 , harmonic with respect to the cone metric. (We choose to write it as $*_{\Sigma}\xi$ for convenience.) Equation (24) says we can write $dr \wedge (*_{\Sigma}\xi) = -*_C\xi$. Now ξ is homogeneous of order -3 and harmonic, so ξ lies in Ω_{27}^3 with respect to φ_C , by Proposition 2.23. Therefore we have shown that (182) becomes

$$h^*(\psi_N) = \psi_C + \eta - *_C\xi + v' + h^*(\omega_4),$$

where $v' + h^*(\omega_4)$ is closed. We also know that $|\nabla_C^j(v' + h^*(\omega_4))|_{g_C} = O(r^{-4-\delta-j})$ as $r \rightarrow \infty$, since $|\nabla_C^j v'|_{g_C} = O(r^{-4-\delta-j})$ and $|\nabla_C^j h^*(\omega_4)|_{g_C} = O(r^{-5-j})$, which follows immediately from (181) and (57). Hence we can apply Lemma 2.11 to say that $v' + h^*(\omega_4) = d\theta$, for some 3-form θ on $(2R, \infty) \times \Sigma$ satisfying $|\nabla_C^j \theta|_{g_C} = O(r^{v'+1-j})$, for any $v' \geq -4 - \delta$. This proves Equation (172) and half of (173).

Now we apply the operator Θ^{-1} to Equation (172). We obtain

$$\begin{aligned} h^*(\varphi_N) &= \Theta^{-1}(h^*(\psi_N)) = \Theta^{-1}(\psi_C - *_C\xi + \eta + d\theta) \\ &= \varphi_C + \xi + J_{\varphi_C}(\eta + d\theta) + G_{\varphi_C}(\eta + d\theta) \end{aligned}$$

using equations (9) and (11) and the fact that ξ is in Ω_{27}^3 with respect to φ_C . Now the remainder term $J_{\varphi_C}(\eta + d\theta) + G_{\varphi_C}(\eta + d\theta) = h^*(\varphi_N) - \varphi_C - \xi$ is clearly smooth and closed. Hence (12) and (10), along with the estimates on η and $d\theta$ above tell us that

$$|\nabla_C^j(J_{\varphi_C}(\eta + d\theta) + G_{\varphi_C}(\eta + d\theta))|_{g_C} = O(r^{-4-j}) \quad \forall j \geq 0,$$

and therefore we can again apply Lemma 2.11 to conclude that $h^*(\varphi_N) - \varphi_C - \xi = d\zeta$ for some 2-form ζ on $(2R, \infty) \times \Sigma$ satisfying $|\nabla_C^j \zeta|_{g_C} = O(r^{v'+1-j})$, for any $v' \geq -4$. This proves Equation (171) and the other half of (173). It is clear that $[\xi] = [h^*(\varphi_N)] = \Phi(N)$, and $[\eta] = [h^*(\psi_N)] = \Psi(N)$ follows from the fact that $-*_C\xi = dr \wedge *_{\Sigma}\xi = d(r *_{\Sigma}\xi)$ is exact. □

Remark 6.5 To prove [Theorem 3.6](#), which is really just a result about G_2 cones, we transferred the problem to an analytic question on the AC manifold to use the machinery of [Section 4.2](#), and then transferred back to the cone. In principle, we could have avoided that by working directly on the cone itself, but then we would have needed to introduce analytic results for noncompact manifolds which had asymptotic ends as well as isolated conical singularities (which can be done), but we preferred to avoid doing so.

Remark 6.6 If we did not impose the gauge-fixing condition, the $O(r^{-7})$ expression in [\(174\)](#) would only be $O(r^{-4})$. Then the 4-form ω_4 from [Lemma 6.3](#) would be *at best* $O(r^{-3+e})$, which would be too large to ignore and the proof of [Theorem 3.6](#) would break down. Therefore the gauge-fixing condition is *necessary* to be able to prove results like [\(171\)](#) and [\(172\)](#).

References

- [1] **M Atiyah, E Witten**, *M–theory dynamics on a manifold of G_2 holonomy*, Adv. Theor. Math. Phys. 6 (2002) 1–106 [MR1992874](#)
- [2] **T Aubin**, *Some nonlinear problems in Riemannian geometry*, Springer Monogr. in Math., Springer, Berlin (1998) [MR1636569](#)
- [3] **C Bär**, *Real Killing spinors and holonomy*, Comm. Math. Phys. 154 (1993) 509–521 [MR1224089](#)
- [4] **R Bartnik**, *The mass of an asymptotically flat manifold*, Comm. Pure Appl. Math. 39 (1986) 661–693 [MR849427](#)
- [5] **A L Besse**, *Einstein manifolds*, Ergebnisse der Math. und ihrer Grenzgebiete (3) 10, Springer, Berlin (1987) [MR867684](#)
- [6] **O Biquard**, *Fibrés paraboliques stables et connexions singulières plates*, Bull. Soc. Math. France 119 (1991) 231–257 [MR1116847](#)
- [7] **R L Bryant**, *Some remarks on G_2 –structures*, from: “Proceedings of Gökova Geometry-Topology Conference 2005”, Gökova Geometry/Topology Conference (GGT), Gökova (2006) 75–109 [MR2282011](#)
- [8] **R L Bryant, S M Salamon**, *On the construction of some complete metrics with exceptional holonomy*, Duke Math. J. 58 (1989) 829–850 [MR1016448](#)
- [9] **J-B Butruille**, *Classification des variétés approximativement kähleriennes homogènes*, Ann. Global Anal. Geom. 27 (2005) 201–225 [MR2158165](#)
- [10] **Y-M Chan**, *Simultaneous desingularizations of Calabi–Yau and special Lagrangian 3–folds with conical singularities. I. The gluing construction*, to appear in Ann. Global Anal. Geom.

- [11] **Y-M Chan**, *Desingularizations of Calabi–Yau 3–folds with a conical singularity*, Q. J. Math. 57 (2006) 151–181 [MR2237517](#)
- [12] **Y-M Chan**, *Desingularizations of Calabi–Yau 3–folds with a conical singularity. II. The obstructed case*, Q. J. Math. 60 (2009) 1–44
- [13] **M Fernández**, **A Gray**, *Riemannian manifolds with structure group G_2* , Ann. Mat. Pura Appl. (4) 132 (1982) 19–45 (1983) [MR696037](#)
- [14] **A Gray**, *Riemannian manifolds with geodesic symmetries of order 3*, J. Differential Geometry 7 (1972) 343–369 [MR0331281](#)
- [15] **A Gray**, *The structure of nearly Kähler manifolds*, Math. Ann. 223 (1976) 233–248 [MR0417965](#)
- [16] **R Harvey**, **H B Lawson, Jr**, *Calibrated geometries*, Acta Math. 148 (1982) 47–157 [MR666108](#)
- [17] **D Joyce**, *Compact Riemannian 7–manifolds with holonomy G_2 . I, II*, J. Differential Geom. 43 (1996) 291–328, 329–375 [MR1424428](#)
- [18] **D Joyce**, *Compact manifolds with special holonomy*, Oxford Math. Monogr., Oxford Univ. Press (2000) [MR1787733](#)
- [19] **D Joyce**, *Special Lagrangian submanifolds with isolated conical singularities. V. Survey and applications*, J. Differential Geom. 63 (2003) 279–347 [MR2015549](#)
- [20] **D Joyce**, *Special Lagrangian submanifolds with isolated conical singularities. I. Regularity*, Ann. Global Anal. Geom. 25 (2004) 201–251 [MR2053761](#)
- [21] **D Joyce**, *Special Lagrangian submanifolds with isolated conical singularities. II. Moduli spaces*, Ann. Global Anal. Geom. 25 (2004) 301–352 [MR2054572](#)
- [22] **D Joyce**, *Special Lagrangian submanifolds with isolated conical singularities. III. Desingularization, the unobstructed case*, Ann. Global Anal. Geom. 26 (2004) 1–58 [MR2054578](#)
- [23] **D Joyce**, *Special Lagrangian submanifolds with isolated conical singularities. IV. Desingularization, obstructions and families*, Ann. Global Anal. Geom. 26 (2004) 117–174 [MR2070685](#)
- [24] **S Karigiannis**, *Moduli spaces of G_2 manifolds with isolated conical singularities*, in preparation
- [25] **S Karigiannis**, *Some notes on G_2 and Spin(7) geometry*, from: “Proceedings of the 2007 International Conference on Geometric Analysis (to appear)” [arXiv: math.DG/0608618](#)
- [26] **S Karigiannis**, *Deformations of G_2 and Spin(7) structures*, Canad. J. Math. 57 (2005) 1012–1055 [MR2164593](#)
- [27] **A Kovalev**, *Twisted connected sums and special Riemannian holonomy*, J. Reine Angew. Math. 565 (2003) 125–160 [MR2024648](#)

- [28] **R Lockhart**, *Fredholm, Hodge and Liouville theorems on noncompact manifolds*, Trans. Amer. Math. Soc. 301 (1987) 1–35 [MR879560](#)
- [29] **R Lockhart, R C McOwen**, *Elliptic differential operators on noncompact manifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 12 (1985) 409–447 [MR837256](#)
- [30] **JD Lotay**, *Desingularization of coassociative 4–folds with conical singularities*, to appear in Geom. Funct. Anal.
- [31] **JD Lotay**, *Calibrated submanifolds and the exceptional geometries*, PhD thesis, University of Oxford (2005) Available at <http://people.maths.ox.ac.uk/~joyce/theses/theses.html>
- [32] **JD Lotay**, *Coassociative 4–folds with conical singularities*, Comm. Anal. Geom. 15 (2007) 891–946 [MR2403190](#)
- [33] **S Marshall**, *Deformations of special Lagrangian submanifolds*, PhD thesis, University of Oxford (2002) Available at <http://people.maths.ox.ac.uk/~joyce/theses/theses.html>
- [34] **V G Maz'ja, B A Plamenevskii**, *Estimates in L_p and in Hölder classes, and the Miranda–Agmon maximum principle for the solutions of elliptic boundary value problems in domains with singular points on the boundary*, Math. Nachr. 81 (1978) 25–82 [MR0492821](#)
- [35] **J Nordström**, *Deformations of asymptotically cylindrical G_2 –manifolds*, Math. Proc. Cambridge Philos. Soc. 145 (2008) 311–348 [MR2442130](#)
- [36] **T Pacini**, *Deformations of asymptotically conical special Lagrangian submanifolds*, Pacific J. Math. 215 (2004) 151–181 [MR2060497](#)
- [37] **R Reyes Carrión, S Salamon**, *A survey of nearly Kähler manifolds*, Gac. R. Soc. Mat. Esp. 2 (1999) 40–49 [MR1707644](#)

Mathematical Institute, University of Oxford
24–29 St Giles', Oxford OX1 3LB, United Kingdom

Department of Pure Mathematics, University of Waterloo
200 University Avenue West, Waterloo, Ontario N2L 3G1, Canada

karigiannis@math.uwaterloo.ca

<http://www.math.uwaterloo.ca/~karigiannis>

Proposed: Simon Donaldson

Received: 23 July 2008

Seconded: Ron Fintushel and Ron Stern

Revised: 23 October 2008