

Quasisymmetric nonparametrization and spaces associated with the Whitehead continuum

JUHA HEINONEN

JANG-MEI WU

The decomposition space \mathbf{R}^3/Wh associated with the Whitehead continuum Wh is not a manifold, but the product $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ is homeomorphic to \mathbf{R}^{3+m} for any $m \geq 1$ (known since the 1960's). We study the quasisymmetric structure on $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ and show that the space $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ may be equipped with a metric resembling \mathbf{R}^{3+m} geometrically and measure theoretically—it is linearly locally contractible and Ahlfors $(3+m)$ -regular—nevertheless the resulting space does not admit a quasisymmetric parametrization by \mathbf{R}^{3+m} .

30C65; 57N16, 57N45

In memory of Juha Heinonen, an extraordinary mathematician and friend, who passed away after the mathematical work in this paper was completed. – J-M W

1 Introduction

To understand the underlying structure of a metric space, one seeks a parametrization of a special type. The classical Riemann mapping theorem is an example. The problem of characterizing metric n -spheres that are bi-Lipschitz equivalent to the standard sphere \mathbf{S}^n , or quasisymmetrically equivalent to \mathbf{S}^n , has received considerable attention in recent years. Despite a large number of results (for example by Bonk and Kleiner [6], David and Semmes [10], Heinonen and Koskela [18], Laakso [22] and Semmes [29; 31; 32]), the question is still far from being understood.

There exist finite 5-dimensional polyhedra that are homeomorphic to the standard sphere \mathbf{S}^5 but not bi-Lipschitz equivalent to \mathbf{S}^5 —an observation of Siebenmann and Sullivan [34] based on deep work of Cannon [7] and Edwards [12; 11], which asserts that the double suspension $\Sigma^2 H^3$ of a 3-dimensional homology sphere H^3 with a nontrivial fundamental group is homeomorphic to the standard sphere \mathbf{S}^5 . The double suspension $\Sigma^2 H^3$ can be considered as the join $\mathbf{S}^1 * H^3$, where H^3 is triangulated and $\Sigma^2 H^3$ is equipped with a natural barycenter metric. The complement of the suspension circle \mathbf{S}^1 in $\Sigma^2 H^3$ is not simply connected; therefore every homeomorphism f

from $\Sigma^2 H^3$ onto \mathbf{S}^5 must map \mathbf{S}^1 onto a curve $f(\mathbf{S}^1)$ whose complement in \mathbf{S}^5 is not simply connected. Consequently $f(\mathbf{S}^1)$ has positive 3-dimensional Hausdorff measure. Hence f can not be Hölder continuous of order greater than $1/3$ and is not bi-Lipschitz. It was then asked by Siebenmann and Sullivan [34] *whether there exists a quasisymmetric homeomorphism between $\Sigma^2 H^3$ and \mathbf{S}^5* , a question which seems inaccessible at the moment.

We study a problem in a similar spirit and, in some sense, of one dimension lower. Let Wh denote a Whitehead continuum in \mathbf{R}^3 (see Whitehead [39]) and \mathbf{R}^3/Wh be the decomposition space associated with Wh obtained by collapsing Wh to a point q while leaving all other points unchanged. The resulting space \mathbf{R}^3/Wh endowed with the quotient topology is not a manifold at q , therefore is not homeomorphic to \mathbf{R}^3 ; however the product space $(\mathbf{R}^3/Wh) \times \mathbf{R}^1$ is homeomorphic to \mathbf{R}^4 (see Bing [4; 5]). The complement of $\{q\} \times \mathbf{R}^1$ in $(\mathbf{R}^3/Wh) \times \mathbf{R}^1$ is not simply connected at $\{q\} \times \mathbf{R}^1$. Therefore any homeomorphism f from $(\mathbf{R}^3/Wh) \times \mathbf{R}^1$ onto \mathbf{R}^4 maps $\{q\} \times \mathbf{R}^1$ to an infinite arc, which is wild in \mathbf{R}^4 and necessarily has positive 2-dimensional Hausdorff measure. Hence f can not be Hölder continuous of order greater than $1/2$.

Semmes [31, page 206] constructed a manifold \tilde{M} modeled on \mathbf{R}^3/Wh and showed that \tilde{M} can be realized as a linearly locally contractible (resembling \mathbf{R}^3 geometrically) and Ahlfors 3-regular (resembling \mathbf{R}^3 measure theoretically) subset of \mathbf{R}^4 ; on the other hand \tilde{M} is not quasisymmetrically equivalent to \mathbf{R}^3 . Heinonen and Semmes [20] asked the following question: *When $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ is equipped with a linearly locally contractible and Ahlfors $(3+m)$ -regular metric, does $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ then admit a quasisymmetric parametrization by \mathbf{R}^{3+m} for some $m \geq 1$?* We answer this question in the negative using essentially a product of the Semmes metric with the Euclidean metric.

1.1 Theorem *For any $m \geq 1$, the metric space $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$, though homeomorphic to \mathbf{R}^{3+m} , can be equipped with a metric that is linearly locally contractible and Ahlfors $(3+m)$ -regular so that the resulting space does not admit a quasisymmetric parametrization by \mathbf{R}^{3+m} .*

The deeper reason for the nonexistence of the quasisymmetric parametrization in [Theorem 1.1](#) is the fact that the meridians of the solid tori used in construction of the Whitehead continuum Wh are trivial in homology of the complement of Wh , but nontrivial in homotopy.

We now proceed to give the formal definitions.

A homeomorphism $f: X \rightarrow Y$ between two metric spaces is said to be *bi-Lipschitz* if there is a constant $C \geq 1$ so that

$$C^{-1}|x - y| \leq |f(x) - f(y)| \leq C|x - y|$$

for all x, y in X . A homeomorphism $f: X \rightarrow Y$ between two metric spaces is said to be *quasisymmetric* if there is a homeomorphism $\eta: [0, \infty) \rightarrow [0, \infty)$ so that

$$|x - a| \leq t|x - b| \text{ implies } |f(x) - f(a)| \leq \eta(t)|f(x) - f(b)|$$

for all triples of points x, a, b in X and for all $t > 0$ (see Tukia and Väisälä [35]). Quasisymmetric maps distort relative distance by a finite amount and distort distance roughly the same in all directions. They are, in some sense, appropriate generalizations of conformal mappings for metric spaces.

A metric space is said to be *linearly locally contractible* if there is a constant $C \geq 1$ so that each metric ball of radius $0 < r < C^{-1}$ in the space can be contracted to a point inside the ball of same center with radius Cr . Linear local contractibility is a geometric condition; it is necessarily satisfied if a metric space is quasisymmetrically equivalent to \mathbf{R}^n .

A metric space X is said to be *Ahlfors Q -regular* if there is a Borel measure μ such that

$$(1.2) \quad C^{-1}r^Q \leq \mu(B_r) \leq Cr^Q$$

for some constants $Q > 0$ and $C \geq 1$, and for all closed balls B_r of radius $0 < r \leq \text{diam } X$. In this case, X has Hausdorff dimension Q and (1.2) holds when μ is replaced with the Hausdorff Q -measure of X (possibly with a different constant C). When Q is an integer, the space X resembles \mathbf{R}^Q measure theoretically but may be totally disconnected. A metric space may be quasisymmetrically equivalent to \mathbf{S}^Q but not Q -regular; the von Koch snowflake curve is an example.

A theorem of Semmes [30] states that if a linearly locally contractible topological Q -manifold is Ahlfors Q -regular, then it satisfies a $(1, 1)$ -Poincaré inequality, hence it has nice analytical properties.

Tukia and Väisälä [35] have shown that a metric doubling space (X, d) homeomorphic to \mathbf{S}^1 is quasisymmetrically equivalent to \mathbf{S}^1 if and only if X satisfies the Ahlfors 2-point condition [35].

Bonk and Kleiner [6] have given a necessary and sufficient condition for a doubling metric space to be quasisymmetrically equivalent to \mathbf{S}^2 . With this they proved the following theorem: *A linearly locally contractible metric 2-sphere is quasisymmetrically equivalent to \mathbf{S}^2 if it is Ahlfors 2-regular.*

Semmes [31] showed that the theorem of Bonk and Kleiner fails in dimension 3 for a geometrical realization of the decomposition space associated with the Bing double in \mathbf{R}^3 , and for the manifold \tilde{M} mentioned earlier. Theorem 1.1 shows that these natural conditions for quasimetric parametrization are also insufficient in dimension 4 and higher.

Acknowledgements Detailed comments of Mario Bonk, Pekka Pankka and the referee on the original manuscript are greatly appreciated. They have clarified the mathematics and improved the exposition.

JH was supported in part by NSF grants DMS-0353549 and DMS-0652915. J-MW was supported in part by NSF grants DMS-0400810 and DMS-0653088 and by the Frederick W and Lois B Gehring Professorship at the University of Michigan where the research was initiated. J-MW thanks the Department of Mathematics at Michigan for the hospitality.

2 Whitehead continuum

In 1935, JHC Whitehead [39] discovered a contractible noncompact 3-manifold in \mathbf{S}^3 that is not homeomorphic to \mathbf{R}^3 . The complement of this manifold is now called a Whitehead continuum. Topological properties and pictures accompanying the construction of a Whitehead continuum can be found in Daverman [9], Kirby [21] and Semmes [33].

Given a set E in $\{x_1 = 0, x_2 \geq 0\} \subset \mathbf{R}^3$, with a slight abuse of notation, denote by $\mathbf{S}^1 \times E$ the solid of revolution $\{(x_1, x_2, x_3) : (0, (x_1^2 + x_2^2)^{1/2}, x_3) \in E\}$ in \mathbf{R}^3 . Let B_0^2 (resp. ∂B_0^2) denote the closed disk (resp. the circle) in $\{x_1 = 0\} \subset \mathbf{R}^3$ having center $(0, 1, 0)$ and radius $\frac{1}{2}$.

A simple closed curve α on $\mathbf{S}^1 \times \partial B_0^2$ is a *meridian* (resp. a *longitude*) of the torus $\mathbf{S}^1 \times B_0^2$ if there exists a homeomorphism from ∂B_0^2 (resp. $\mathbf{S}^1 \times \{(0, \frac{1}{2}, 0)\}$) onto α which is homotopic in $\mathbf{S}^1 \times \partial B_0^2$ to the identity map on ∂B_0^2 (resp. $\mathbf{S}^1 \times \{(0, \frac{1}{2}, 0)\}$).

Let T_0 be a smooth compact solid torus embedded in \mathbf{R}^3 , identified with $\mathbf{S}^1 \times B_0^2$ for convenience. Let T_1 be a second compact solid torus smoothly embedded in the interior of T_0 and positioned in such a way that T_1 and any meridian of T_0 form a Whitehead link (see Kirby [21] and Rourke and Sanderson [27]). Note that T_1 is contractible in T_0 , but the contraction of T_1 can not take place without crossing itself. In fact, T_1 is knotted in T_0 but is not knotted in \mathbf{R}^3 . Let ϕ be a diffeomorphism defined in a neighborhood of T_0 that maps T_0 onto T_1 . Define inductively for $k \geq 2$,

$$T_k = \phi(T_{k-1}).$$

The intersection

$$Wh = \bigcap_{k=0}^{\infty} T_k$$

is a Whitehead continuum. The continuum Wh is one-dimensional and is contractible inside each $T_k, k \geq 0$. Identify \mathbf{S}^3 with $\mathbf{R}^3 \cup \{\infty\}$ in topology. The Whitehead manifold $\mathbf{S}^3 \setminus Wh$ is simply connected and in fact contractible [21].

The Whitehead link is symmetric. For $k \geq 1$, there is a homeomorphism of \mathbf{S}^3 that interchanges a meridian of T_k with a core of T_{k+1} (such as $\phi^{(k+1)}(\mathbf{S}^1 \times \{(0, 1, 0)\})$). However the homeomorphism reversing the link can not be confined to T_0 (see Kirby [21, page 83]). Therefore there are loops, namely the meridians of T_k , lying arbitrarily close to the Whitehead continuum Wh that can not be contracted in $\mathbf{S}^3 \setminus Wh$ to a point while staying near Wh . In other words the Whitehead manifold $\mathbf{S}^3 \setminus Wh$ is not simply connected at Wh , so $\mathbf{S}^3 \setminus Wh$ is not homeomorphic to \mathbf{R}^3 .

The decomposition space \mathbf{R}^3 / Wh is defined by collapsing Wh to a single point, called q , while keeping all other points unchanged, then endowed with the quotient topology. The resulting space is not a manifold at the quotient point q by the previous discussion, hence it is not homeomorphic to \mathbf{R}^3 .

Taking the product with \mathbf{R}^m ($m \geq 1$) however destroys all nonmanifold points; as a consequence $(\mathbf{R}^3 / Wh) \times \mathbf{R}^m$ is homeomorphic to \mathbf{R}^{3+m} (by Andrews and Rubin [3], also credited to Arnold Shapiro by Bing [4; 5]). See Daverman [9] and Kirby [21] for more discussions about products of decompositions with a line.

Let $\pi: \mathbf{R}^3 \rightarrow \mathbf{R}^3 / Wh$ be the quotient map defined by $\pi(Wh) = q$ and $\pi(x) = x$ if $x \notin Wh$; for simplicity denote by

$$\tilde{E} = \pi(E)$$

the image of a set E under π . Let

$$\mathcal{L} = \{q\} \times \mathbf{R}^m.$$

A curve γ on $\partial \tilde{T}_k$ is called a meridian (resp. a longitude) of \tilde{T}_k if $\pi^{-1}\gamma$ is a meridian (resp. a longitude) of T_k . The noncontractibility of the meridians of T_k in the Whitehead manifold near Wh implies the following.

2.1 Proposition *Meridians of \tilde{T}_k are not contractible in $\tilde{T}_0 \setminus \{q\}$, and meridians of $\tilde{T}_k \times \{z\}$ are not contractible in $(\tilde{T}_0 \times \mathbf{R}^m) \setminus \mathcal{L}$, for any $m \geq 1, k \geq 1$ and $z \in \mathbf{R}^m$.*

On the other hand, a meridian of \tilde{T}_k bounds a surface (a disk with a handle) in $\tilde{T}_k \setminus \{q\}$. The difference of the roles of a meridian in homotopy and in homology is important in our proof.

3 The metric

We define a metric on the decomposition space \mathbf{R}^3/Wh by adapting that of Semmes in [31]. The metric to be defined has a self-similarity property and sizes of \tilde{T}_k decrease geometrically towards the point q . Semmes embedded \mathbf{R}^3/Wh in \mathbf{R}^4 which is more visual; we consider $(\mathbf{R}^3/Wh)\setminus\{q\}$ as an abstract manifold on which the metric can be explicitly described.

Fix a number $c_0 \in (0, \frac{1}{20})$. Let $P_0 = (0, 1, 0)$, and $A = \{x \in \{0\} \times \mathbf{R}^2 : \frac{1}{2} - 2c_0 < |x - P_0| \leq \frac{1}{2} - c_0\}$, $A' = \{x \in \{0\} \times \mathbf{R}^2 : \frac{1}{2} - 2c_0 < |x - P_0| < \frac{1}{2} + 2c_0\}$ and $A'' = \{x \in \{0\} \times \mathbf{R}^2 : \frac{1}{2} - 3c_0 < |x - P_0| < \frac{1}{2} + 3c_0\}$ be three annuli in $\{0\} \times \mathbf{R}^2$. Let

$$U_0 = \mathbf{S}^1 \times A, \quad V_0 = \mathbf{S}^1 \times A' \quad \text{and} \quad W_0 = \mathbf{S}^1 \times A''$$

be the solids obtained by revolving A, A' and A'' about the x_3 -axis. Note that V_0 and W_0 are neighborhoods of ∂T_0 and that $U_0 \subset T_0 \cap V_0 \subset V_0 \subset W_0$. Recall that ϕ is a diffeomorphism defined in a neighborhood of T_0 that maps T_0 onto T_1 . Assume that c_0 has been chosen small enough so that W_0 is contained in the neighborhood of T_0 on which ϕ is defined and that $\phi(T_0 \cup W_0) \subset T_0 \setminus W_0$, thus T_1 is contained in $T_0 \setminus W_0$. Define inductively for $k \geq 1$,

$$(3.1) \quad U_k = \phi(U_{k-1}), \quad V_k = \phi(V_{k-1}) \quad \text{and} \quad W_k = \phi(W_{k-1})$$

and define for $k \geq 0$, closed regions between V_k and V_{k+1}

$$\Theta_k = T_k \setminus (T_{k+1} \cup V_k \cup V_{k+1}),$$

and solid tori

$$t_k = T_k \setminus (V_k \setminus U_k).$$

Note that

$$T_{k+1} \subset t_k \subset T_k.$$

Fix a parameter $a \in (0, 1)$. Define a metric d_a on the Whitehead continuum $\mathbf{R}^3 \setminus Wh$ as follows. Initially, define a matrix $(\lambda_{ij}) = \text{Id}$ on $(\mathbf{R}^3 \setminus T_0) \cup W_0$ and define for $k \geq 1$,

$$(\lambda_{ij}) = a^{2k} (D\phi^{(-k)})^T D\phi^{(-k)} \text{ in } W_k$$

where $\phi^{(-k)}$ is the k -th iterate of ϕ^{-1} . Later, adjust (λ_{ij}) in $\bigcup_{k=0}^\infty (W_k \setminus V_k)$ and extend (λ_{ij}) to $\mathbf{R}^3 \setminus Wh$ so as to achieve the smoothness, the positive definiteness of (λ_{ij}) and the self-similarity (3.3) below, and to preserve the previously defined values on $(\mathbf{R}^3 \setminus T_0) \cup \bigcup_{k=0}^\infty V_k$. This can be accomplished by first extending (λ_{ij}) continuously to $\Theta_0 \setminus (W_0 \cup W_1)$ with the aid of partition of unity, next smoothing (λ_{ij}) in Θ_0 by convolution with a kernel that varies with the Euclidean distance $\text{dist}(x, \partial\Theta_0)$, and finally redefining (λ_{ij}) in $\bigcup_{k=1}^\infty \Theta_k$ so that for each $k \geq 1$,

$$(\lambda_{ij}) = a^{2k} (D\phi^{(-k)})^T (\lambda_{ij}(\phi^{(-k)})) D\phi^{(-k)} \text{ in } \Theta_k.$$

The metric d_a on $\mathbf{R}^3 \setminus Wh$ associated with the matrix (λ_{ij}) is defined by

$$d_a(x, y) = \inf\{\text{length } \gamma : \gamma \text{ continuously differentiable curve joining } x \text{ and } y\},$$

where

$$(3.2) \quad \text{length } \gamma = \int_{\gamma} \sqrt{\sum \lambda_{ij} dx_i dx_j}.$$

The metric thus defined is self-similar with respect to Wh in the following sense:

$$(3.3) \quad d_a(x, y) = a^k \cdot d_a(\phi^{(-k)}x, \phi^{(-k)}y) \quad \text{for all } x \in T_{k'} \text{ and } y \in T_{k''}$$

and for all $k', k'' \geq k \geq \kappa \geq 0$ for some κ depending only on ϕ and c_0 .

The metric on the decomposition space \mathbf{R}^3/Wh , again called d_a , is the pushforward metric induced by the homeomorphism $\pi: \mathbf{R}^3 \setminus Wh \rightarrow (\mathbf{R}^3/Wh) \setminus \{q\}$ followed by the continuous extension to the point q .

Denote by d_e be the Euclidean metric on \mathbf{R}^m for any $m \geq 1$. Let δ_a be the metric on the product space $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ defined by

$$(3.4) \quad \delta_a((x, x'), (y, y')) = d_a(x, y) + d_e(x', y')$$

for all $(x, x'), (y, y') \in (\mathbf{R}^3/Wh) \times \mathbf{R}^m$ and any $m \geq 1$.

Here, as well as in the future, we use $c, c_1, c_2, \dots, C, C_1, C_2, \dots, C', C'', \dots$ to denote positive constants depending at most on m, ϕ, c_0 and a , in particular, independent of k . We use $r \lesssim s, r \gtrsim s, r \simeq s$ to mean that the ratio $\frac{r}{s}$ is bounded above, or bounded below, or bounded above and below by this type of constants. Values of c and C need not be the same at each occurrence.

Note then

$$(3.5) \quad \text{diam}_{d_a}(\tilde{T}_k) \simeq a^k,$$

$$(3.6) \quad \text{dist}_{d_a}(\tilde{t}_k, \partial\tilde{T}_k) = \text{dist}_{d_a}(\tilde{U}_k, \partial\tilde{T}_k) = c_0 a^k,$$

$$(3.7) \quad \text{dist}_{d_a}(q, \partial\tilde{T}_k) \simeq \text{dist}_{d_a}(\partial\tilde{T}_k, \partial\tilde{T}_{k+1}) \simeq \text{dist}_{d_a}(\tilde{V}_k, \tilde{V}_{k+1}) \simeq a^k.$$

We now give a concrete reformulation of [Theorem 1.1](#).

3.8 Theorem *The metric space $((\mathbf{R}^3/Wh) \times \mathbf{R}^m, \delta_a)$ is linearly locally contractible and Ahlfors $(3+m)$ -regular. However there is no quasisymmetric homeomorphism between $((\mathbf{R}^3/Wh) \times \mathbf{R}^m, \delta_a)$ and \mathbf{R}^{3+m} , when $a > 2^{-(3+m)/2m}$ and $m \geq 1$.*

The restriction on a in the theorem is a consequence of the modulus estimates. It is unclear whether the theorem remains true for the whole range of $a \in (0, 1)$.

The linear local contractibility and the Ahlfors regularity in [Theorem 3.8](#) follow from [\(3.3\)–\(3.7\)](#) and a lifting-twisting-shrinking procedure used in proving the topological equivalence between $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ and \mathbf{R}^{3+m} . By [\(3.3\)](#), [\(3.5\)](#), [\(3.6\)](#) and [\(3.7\)](#), there exists a large constant $C > 1$ such that for each ball $B_{d_a}(x, r)$ in $(\mathbf{R}^3/Wh, d_a)$, one of the following holds:

- (i) $x = q$.
- (ii) $x \in \tilde{T}_k \setminus \tilde{T}_{k+1}$ for some $k \geq -1$ and $q \in B_{d_a}(x, Cr)$; in this case $B_{d_a}(x, r) \subset B_{d_a}(q, C'r) \subset B_{d_a}(x, C''r) \subset B_{d_a}(q, C'''r)$.
- (iii) $x \in \tilde{T}_k \setminus \tilde{T}_{k+1}$ for some $k \geq -1$ and $q \notin B_{d_a}(x, Cr)$; in this case $B_{d_a}(x, r) \subset \tilde{T}_{k-1} \setminus \tilde{T}_{k+2}$ and there is an open set U homeomorphic to a Euclidean ball so that

$$B_{d_a}(x, r) \subset U \subset B_{d_a}(x, Cr).$$

Here, with a slight abuse of notation, \tilde{T}_{-1} denotes \mathbf{R}^3/Wh . By [\(3.4\)](#), there exists $C > 1$ so that

$$B_{\delta_a}((q, (0, 0, \dots, 0)), C^{-1}a^k) \subset \tilde{T}_k \times [-a^k, a^k]^m \subset B_{\delta_a}((q, (0, 0, \dots, 0)), Ca^k).$$

We first check that $\tilde{T}_1 \times [-1, 1]^m$ is contractible inside $\tilde{T}_0 \times [-4, 4]^m$. It is known that the quotient map $\pi \times \text{id}: \mathbf{R}^{3+m} \rightarrow (\mathbf{R}^3/Wh) \times \mathbf{R}^m$ can be uniformly approximated by homeomorphisms (see Daverman [\[9\]](#), pages 81–84). Choose $g: \mathbf{R}^{3+m} \rightarrow (\mathbf{R}^3/Wh) \times \mathbf{R}^m$, a homeomorphism satisfying

$$\text{dist}_{\delta_a}(g(x, z), (\pi \times \text{id})(x, z)) < c_0/30 \quad \text{on } \mathbf{R}^{3+m}.$$

Then $g^{-1}(\tilde{T}_1 \times [-1, 1]^m)$, a subset of $(T_1 \cup V_1) \times [-2, 2]^m$, is contractible inside $t_0 \times [-3, 3]^m$. Hence $\tilde{T}_1 \times [-1, 1]^m$ is contractible in $g(t_0 \times [-3, 3]^m) \subset \tilde{T}_0 \times [-4, 4]^m$. The linear local contractibility of $((\mathbf{R}^3/Wh) \times \mathbf{R}^m, \delta_a)$ can be seen by scaling and examining the projection of a given ball $B_{\delta_a}(x, r)$ into \mathbf{R}^3/Wh for the three alternative cases.

Define a measure μ on $\mathbf{R}^3 = Wh \cup (\mathbf{R}^3 \setminus Wh)$ as follows: $\mu(Wh) = 0$, $\mu|_{(\mathbf{R}^3 \setminus T_1)}$ is the Lebesgue measure, and

$$\mu(E) = a^{3k} \mu(\phi^{(-k)}(E))$$

if $E \subset T_k \setminus T_{k+1}$ and $k \geq 1$. The measure on the decomposition space $(\mathbf{R}^3/Wh, d_a)$, again called μ , is defined to be the pushforward of the measure μ on \mathbf{R}^3 induced by

the quotient map $\pi: \mathbf{R}^3 \rightarrow \mathbf{R}^3/Wh$. Note from (3.3) that

$$\mu(\tilde{T}_k \setminus \tilde{T}_{k+1}) \simeq a^{3k} \quad \text{for } k \geq 0,$$

and that $\mu(B_{d_a}(q, r)) \simeq r^3$; and note from (3.3) and the smoothness of (λ_{ij}) that $\mu(B_{d_a}(x, r)) \simeq r^3$ if $B_{d_a}(x, r) \subset \tilde{T}_{k-1} \setminus \tilde{T}_{k+2}$ and $k \geq 0$. The Ahlfors 3–regularity of $(\mathbf{R}^3/Wh, d_a, \mu)$ follows by checking the three alternative cases. The product space $((\mathbf{R}^3/Wh) \times \mathbf{R}^m, \delta_a, \mu \times dx)$ is Ahlfors $(3+m)$ –regular equipped with the product of μ with the Lebesgue measure dx in \mathbf{R}^m .

4 Modulus of surface families

Let f be a homeomorphism from $((\mathbf{R}^3/Wh) \times \mathbf{R}^m, \delta_a)$ onto \mathbf{R}^{3+m} . What is the obstruction for quasisymmetry? We give a heuristic explanation for $m = 1$. Proposition 2.1 implies that $B \setminus \mathcal{L}$ is not simply connected for any small ball B in $(\mathbf{R}^3/Wh) \times \mathbf{R}^1$ that intersects \mathcal{L} . Hence $f(B) \setminus f(\mathcal{L})$ is not simply connected and $f(\mathcal{L})$ is wild in \mathbf{R}^4 . A theorem of Martio, Rickman and Väisälä (Theorem 9.1 in the Appendix) then implies that the 2–dimensional Hausdorff measure

$$\mathcal{H}^2(f(B) \cap f(\mathcal{L})) > 0,$$

for any ball B in $(\mathbf{R}^3/Wh) \times \mathbf{R}^1$ that intersects \mathcal{L} . One would then expect that f maps all line segments of the form $\{z\} \times (t, t + s)$ in $(\mathbf{R}^3/Wh) \times \mathbf{R}^1$ near $\{q\} \times \mathbf{R}^1$ to curves of much greater length. A comparison of moduli of suitable curve families in the domain and their images would prevent f from being quasisymmetric. Nevertheless we are unable to provide quantitative estimates for the lengths of $f(\{z\} \times (t, t + s))$; so we shall work with the modulus of surface families instead.

The modulus of curve families has been an important tool in geometric function theory since the 1920’s, first used by Grötzsch and Teichmüller and later extensively studied by Beurling and Ahlfors [1]. Extending the concept of modulus from curve families to l –dimensional surface families in \mathbf{R}^n was due to Fuglede [15]. We need the notion of modulus for surface families in the setting of metric spaces. We shall not aim for full generality though.

Let (X, d) be a linearly locally contractible topological n –manifold that is also Ahlfors n –regular. Let $1 \leq l \leq n - 1$, p a real number ≥ 1 and \mathcal{S} be a family of topological l –manifolds in X . A nonnegative Borel function $\rho: X \rightarrow [0, \infty]$ is called *admissible* for \mathcal{S} if

$$\int_{\Gamma} \rho(x) d\mathcal{H}_d^l(x) \geq 1$$

for every $\Gamma \in \mathcal{S}$ that has positive Hausdorff l -measure $\mathcal{H}_d^l(\Gamma)$ and is locally finite with respect to \mathcal{H}_d^l . The p -modulus of \mathcal{S} is defined by

$$\text{Mod}_p \mathcal{S} = \inf \int_X \rho^p(x) d\mathcal{H}_d^n(x),$$

where the infimum is taken over all admissible $\rho: X \rightarrow [0, \infty]$. Here and later, we denote by \mathcal{H}_d^i the i -th dimensional Hausdorff measure with respect to the metric d . We omit the subscript d if the metric is the Euclidean one.

The most important modulus from the point of view of (quasi)conformal geometry is the n/l -modulus $\text{Mod}_{n/l}$. It is conformally invariant and hence called the *conformal modulus*. The conformal modulus of curve families is quasi-invariant under quasimetric maps. This was proved by Gehring [16] for the Euclidean n -spaces and by Heinonen and Koskela [19] for very general metric spaces.

We consider a sequence of $(1+m)$ -dimensional surface families \mathcal{S}_k in $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ defined as follows. First we select the curve family

$$\mathcal{C} = \{\mathbf{S}^1 \times \{x\} : x \in A\}$$

in $T_0 \subset \mathbf{R}^3$, where A is the planar annulus defined in Section 3. Let

$$\mathcal{C}_k = \{\pi\phi^{(k)}(\mathbf{S}^1 \times \{x\}) : x \in A\} \quad \text{for } k \geq 0,$$

be the associated curve families in the decomposition space \mathbf{R}^3/Wh , then define the surface families

$$\mathcal{S}_k = \{\gamma \times [0, 1]^m : \gamma \in \mathcal{C}_k\} \quad \text{for } k \geq 0,$$

in $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ by taking products.

We prove a one-sided inequality for the conformal moduli, which is adequate for our purpose. Constants depending on f as well as m, ϕ, c_0 and a , shall be denoted by $C(f)$.

4.1 Proposition *Let f be a quasimetric homeomorphism from $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ onto \mathbf{R}^{3+m} . Then there exists a constant $C(f) > 0$ such that*

$$(4.2) \quad \text{Mod}_{(3+m)/(1+m)} \mathcal{S}_k \leq C(f) \text{Mod}_{(3+m)/(1+m)} f(\mathcal{S}_k) \quad \text{for all } k \geq 0.$$

Proof Since the matrix (λ_{ij}) is smooth in $\mathbf{R}^3 \setminus Wh$, the quasimetric map f is in $W_{\text{loc}}^{1,r}(((\mathbf{R}^3/Wh) \times \mathbf{R}^m) \setminus \mathcal{L})$ for some $r > 3 + m$; this can be seen by following the proof of the theorem of Gehring [17] on the higher integrability of the derivatives of quasiconformal maps in Euclidean spaces. Hence f is differentiable $\mathcal{H}_{\delta_a}^{3+m}$ -almost everywhere in $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$. Set, for $x \in A$,

$$\Gamma_x = \pi\phi^{(k)}(\mathbf{S}^1 \times \{x\}) \times [0, 1]^m.$$

In Euclidean spaces, quasiconformal mappings are absolutely continuous on almost all lines. Adapting the standard proof of this fact [36, Theorem 31.2], we may conclude that for \mathcal{H}^2 -almost every $x \in A$, $f|_{\Gamma_x}$ maps sets of $\mathcal{H}^{1+m}_{\delta_a}$ -measure zero to sets of \mathcal{H}^{1+m} -measure zero. We may then conclude that for any nonnegative Borel function ρ on \mathbf{R}^{3+m} and for \mathcal{H}^2 -almost every $x \in A$,

$$(4.3) \quad \int_{\Gamma_x} \rho(f(x)) |Df(x)|^{1+m} d\mathcal{H}^{1+m}_{\delta_a}(x) \geq \int_{f(\Gamma_x)} \rho(y) d\mathcal{H}^{1+m}(y),$$

by adapting the standard argument for quasiconformal maps [36, Theorem 5.3]. Quasisymmetry implies that

$$\begin{aligned} & \int_{(\mathbf{R}^3/Wh) \times \mathbf{R}^m} (\rho(f(x)) |Df(x)|^{1+m})^{(3+m)/(1+m)} d\mathcal{H}^{3+m}_{\delta_a}(x) \\ & \leq C(f) \int_{(\mathbf{R}^3/Wh) \times \mathbf{R}^m} \rho(f(x))^{(3+m)/(1+m)} |\det Df(x)| d\mathcal{H}^{3+m}_{\delta_a}(x) \\ & \leq C(f) \int_{\mathbf{R}^{3+m}} \rho(y)^{(3+m)/(1+m)} d\mathcal{H}^{3+m}(y). \end{aligned}$$

From the fact that the union of these surfaces has $\mathcal{H}^{3+m}_{\delta_a}$ -measure zero, it follows that the exceptional surfaces for which (4.3) fails have vanishing conformal modulus. Then (4.2) can be obtained by following the standard argument for modulus of curve families (see Theorem 32.3 in [36] or Theorem II.2.4 in [26]). \square

We need the following estimates.

4.4 Proposition *Let f be a homeomorphism from $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ onto \mathbf{R}^{3+m} . Then for each $k \geq 0$ and each $\Gamma \in \mathcal{S}_k$, the $(1+m)$ -dimensional Hausdorff measures in $((\mathbf{R}^3/Wh) \times \mathbf{R}^m, \delta_a)$ and \mathbf{R}^{3+m} satisfy*

- (1) $\mathcal{H}^{1+m}_{\delta_a}(\Gamma) \simeq a^k$,
- (2) $\mathcal{H}^{1+m}(f(\Gamma)) \geq C(f)2^k$

for some positive constant $C(f)$.

4.5 Proposition *Let f be a homeomorphism from $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ onto \mathbf{R}^{3+m} . Then for each $k \geq 0$, the conformal moduli of surface families satisfy*

- (1) $\text{Mod}_{(3+m)/(1+m)} \mathcal{S}_k \simeq (a^k)^{2m/(1+m)}$,
- (2) $\text{Mod}_{(3+m)/(1+m)} f(\mathcal{S}_k) \leq C(f)(2^{-k})^{(3+m)/(1+m)}$

for some positive constant $C(f)$.

5 Proof of Theorem 1.1

Theorem 1.1 follows from **Theorem 3.8**. The last sentence of **Theorem 3.8**, the only statement yet to be proved, follows from **Proposition 4.1** and **Proposition 4.5**. The proof of **Proposition 4.5** is given first assuming the validity of **Proposition 4.4**. The proof of **Proposition 4.4** is given next assuming the validity of an estimate (5.1) on the intersection numbers. The proof of (5.1) shall occupy the remaining sections.

Proof of Proposition 4.5 Assume the validity of **Proposition 4.4**. Fix $k \geq 0$ and let ρ be a nonnegative Borel function on $(\mathbf{R}^3 / Wh) \times \mathbf{R}^m$ admissible for the surface family \mathcal{S}_k , thus

$$\int_{\Gamma} \rho d\mathcal{H}_{\delta_a}^{1+m} \geq 1$$

for every $\Gamma \in \mathcal{S}_k$. Therefore

$$\begin{aligned} & \int_{\tilde{U}_k \times [0,1]^m} \rho d\mathcal{H}_{\delta_a}^{3+m} \\ & \simeq \int_{\mathbf{S}^1 \times A} \left(\int_{[0,1]^m} \rho(\pi\phi^{(k)}(s,x), z) d\mathcal{H}^m(z) \right) a^{3k} d\mathcal{H}^3(s,x) \\ & \simeq \int_A \left(\int_{\pi\phi^{(k)}(\mathbf{S}^1 \times \{x\})} \left(\int_{[0,1]^m} \rho(\xi, z) d\mathcal{H}^m(z) \right) d\mathcal{H}_{\delta_a}^1(\xi) \right) a^{2k} d\mathcal{H}^2(x) \\ & \simeq \int_A \left(\int_{\pi\phi^{(k)}(\mathbf{S}^1 \times \{x\}) \times [0,1]^m} \rho(\xi, z) d\mathcal{H}_{\delta_a}^{1+m}(\xi, z) \right) a^{2k} d\mathcal{H}^2(x) \gtrsim a^{2k}. \end{aligned}$$

On the other hand,

$$\begin{aligned} & \int_{\tilde{U}_k \times [0,1]^m} \rho d\mathcal{H}_{\delta_a}^{3+m} \\ & \leq \left(\int_{\tilde{U}_k \times [0,1]^m} \rho^{(3+m)/(1+m)} d\mathcal{H}_{\delta_a}^{3+m} \right)^{(1+m)/(3+m)} (\mathcal{H}_{\delta_a}^{3+m}(\tilde{U}_k \times [0,1]^m))^{2/(3+m)} \\ & \leq C \left(\int_{\tilde{U}_k \times [0,1]^m} \rho^{(3+m)/(1+m)} d\mathcal{H}_{\delta_a}^{3+m} \right)^{(1+m)/(3+m)} a^{6k/(3+m)}. \end{aligned}$$

This implies

$$\int_{\tilde{U}_k \times [0,1]^m} \rho^{(3+m)/(1+m)} d\mathcal{H}_{\delta_a}^{3+m} \gtrsim (a^k)^{2m/(1+m)}.$$

Now set $\rho_0 = a^{-k}$ in $\tilde{U}_k \times [0,1]^m$ and $= 0$ outside $\tilde{U}_k \times [0,1]^m$. Note from **Proposition 4.4(1)** that $C\rho_0$ is admissible for the surface family \mathcal{S}_k for some constant $C > 0$

depending at most on m, ϕ, c_0 and a , and in particular independent of k . It is easy to check that

$$\int_{(\mathbf{R}^3/Wh) \times \mathbf{R}^m} \rho_0^{(3+m)/(1+m)} d\mathcal{H}_{\delta_a}^{3+m} \simeq (a^{-k})^{(3+m)/(1+m)} (a^k)^3 \simeq (a^k)^{2m/(1+m)}.$$

Therefore $\text{Mod}_{(3+m)/(1+m)} \mathcal{S}_k \simeq (a^k)^{2m/(1+m)}$.

To show part (2), set $\rho_1 = 2^{-k}$ in $f(\tilde{U}_k \times [0, 1]^m)$ and $\rho_1 = 0$ in $\mathbf{R}^{3+m} \setminus f(\tilde{U}_k \times [0, 1]^m)$. Note from Proposition 4.4 (2) that $C(f)\rho_1$ is admissible for $f(\mathcal{S}_k)$ for some $C(f) > 0$. Then

$$\int_{\mathbf{R}^{3+m}} \rho_1^{(3+m)/(1+m)} d\mathcal{H}^{3+m} \leq C(f)\mathcal{H}^{3+m}(f(\tilde{T}_0 \times [0, 1]^m))(2^{-k})^{(3+m)/(1+m)}.$$

Hence $\text{Mod}_{(3+m)/(1+m)} f(\mathcal{S}_k) \leq C(f)(2^{-k})^{(3+m)/(1+m)}$.

This completes the proof. □

Proof of Proposition 4.4 Part (1) follows immediately from the definition of the metric δ_a .

To prove part (2), we first construct a bi-Lipschitz homeomorphism χ of \mathbf{R}^{3+m} with some specific properties, we then slice surfaces in the collection $\bigcup_{k=k_0}^\infty (\chi \circ f(\mathcal{S}_k))$ by a certain stack of standard 2-simplices $\{D_j\}$ in \mathbf{R}^{3+m} with the parameter j ranging over an $(1+m)$ -dimensional cube J in \mathbf{R}^{3+m} orthogonal to the simplices $\{D_j\}$. Finally we estimate the number $N(k, \Gamma, j)$ of points in the intersection $D_j \cap (\chi \circ f(\Gamma))$ from below by

$$(5.1) \quad N(k, \Gamma, j) \geq C(f)2^k \quad \text{for all } k > k_0, \Gamma \in \mathcal{S}_k \text{ and } j \in J,$$

for some positive integer k_0 and some positive constant $C(f)$ depending on f .

Assume that (5.1) is valid for the moment. Given $k > k_0$ and a surface Γ in \mathcal{S}_k , let ζ be the projection map from $(\bigcup_{j \in J} D_j) \cap (\chi \circ f(\Gamma))$ to J defined by

$$\zeta(x) = j \quad \text{if } x \in D_j \cap (\chi \circ f(\Gamma)).$$

So ζ is 1-Lipschitz. It follows from (5.1) and a coarse estimate (see Federer [13, Theorem 2.10.25]) that

$$\mathcal{H}^{1+m}(\chi \circ f(\Gamma)) \geq \int_J N(k, \Gamma, j) d\mathcal{H}^{1+m}(j) \geq C(f)\mathcal{H}^{1+m}(J)2^k.$$

Since χ is bi-Lipschitz, part (2) has been proved assuming (5.1). □

To complete the proof of [Theorem 1.1](#), it remains to establish [\(5.1\)](#). In [Section 6](#) we define χ and locate the stack of two-simplices used for slicing, and in [Section 8](#) we prove the lower estimate [\(5.1\)](#).

6 A stack of two-simplices

Denote by $B_d(x, r)$ the closed ball having center x and radius r in the metric space (X, d) . We omit the subscript when the metric is Euclidean.

Let 0 be the origin of \mathbf{R}^m and Q be the point $(q, 0)$ in $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$. Recall from [\(3.5\)](#), [\(3.6\)](#) and [\(3.7\)](#) that $q \in \tilde{T}_k \subset \tilde{T}_k \cup \tilde{V}_k \subset B_{d_a}(q, C a^k) \subset \mathbf{R}^3/Wh$ and that

$$Q \in (\tilde{T}_k \cup \tilde{V}_k) \times [-a^k, a^k]^m \subset B_{\delta_a}(Q, C_1 a^k),$$

for some constants $C, C_1 > 1$. Set

$$v_k = \text{diam } f(B_{\delta_a}(Q, C_1 a^k));$$

choose and fix an integer $k_0 \geq 2$ so that

$$(6.1) \quad B(f(Q), 3v_{k_0}) \subset f(\tilde{T}_1 \times \mathbf{R}^m).$$

The 2-simplices shall be chosen to lie in $B(f(Q), 3v_{k_0})$.

Set $\alpha = \pi\phi^{(k_0)}(\partial B_0^2) \times \{0\}$,

a meridian of $\tilde{T}_{k_0} \times \{0\}$, where B_0^2 is the planar disk in \mathbf{R}^3 defined in the beginning of [Section 2](#). Let

$$(6.2) \quad \mathfrak{A} = \left\{ x \in \mathbf{R}^3/Wh : \text{dist}_{\delta_a}((x, 0), \alpha) < \frac{c_0 a^{k_0}}{30} \right\} \times \left(-\frac{c_0 a^{k_0}}{30}, \frac{c_0 a^{k_0}}{30} \right)^m$$

be a tubular neighborhood of α not intersecting $\tilde{T}_{k_0} \times \mathbf{R}^m$, where c_0 is the constant chosen in [Section 3](#).

Let $c_1 = \text{dist}(f(\alpha), f(\partial\mathfrak{A}))/10$. It follows from a PL-approximation theorem of Moise ([Theorem 9.2](#) in the [Appendix](#)) that there exists an embedding $\theta: \alpha \hookrightarrow \mathbf{R}^{3+m}$ such that $l \equiv \theta(\alpha)$ is a piecewise linear topological 1-sphere in $f(\mathfrak{A})$, and

$$d(f(x), \theta(x)) < c_1 \text{ for all } x \in \alpha.$$

The embedding θ used for defining l shall play no role in the future. We observe that l and $f(\alpha)$ are homotopic in $f(\mathfrak{A})$ and that

$$(6.3) \quad \begin{aligned} l \subset f(\mathfrak{A}) \subset f((\tilde{T}_{k_0} \cup \tilde{V}_{k_0}) \times [-a^{k_0}, a^{k_0}]^m) \\ \subset f(B_{\delta_a}(Q, C_1 a^{k_0})) \subset B(f(Q), v_{k_0}). \end{aligned}$$

Here, as well as in the future, two curves β and β' are said to be homotopic in X (or β is homotopic to β' in X) if the continuous maps defining the curves are homotopic in the target space X .

Since $3 + m \geq 4$, the PL 1–sphere l is unknotted in \mathbf{R}^{3+m} and there is a global bi-Lipschitz homeomorphism χ of \mathbf{R}^{3+m} that maps l to the boundary of a standard 2–simplex D' contained in $B(Y, 2 \text{ diam } l)$, $\chi(l) = \partial D'$, while fixes all points outside $B(Y, 2 \text{ diam } l)$ for some point $Y \in l$. (See [Theorem 9.4](#) in the [Appendix](#).) Note from (6.3) that

$$(6.4) \quad \chi = \text{id} \text{ in } \mathbf{R}^{3+m} \setminus B(f(Q), 3\nu_{k_0}).$$

Choose and fix a $(1+m)$ –dimensional cube J in \mathbf{R}^{3+m} which is orthogonal to the 2–simplex D' and is small enough such that all translates $j + \partial D'$ ($j \in J$) are contained in $\chi \circ f(\mathfrak{A})$ and are homotopic to $\chi \circ f(\alpha)$ in $\chi \circ f(\mathfrak{A})$. The translated standard 2–simplices $D_j \equiv j + D'$, $j \in J$ enclosed by $j + \partial D'$ shall be used for slicing surfaces in the collection $\bigcup_{k=k_0}^{\infty} (\chi \circ f(\mathcal{S}_k))$.

Since $\partial D_j = j + \partial D' \subset \chi \circ f(\mathfrak{A}) \subset \chi(B(f(Q), \nu_{k_0})) \subset B(f(Q), 3\nu_{k_0})$, it follows that

$$D_j \subset B(f(Q), 3\nu_{k_0}).$$

From (6.1) and (6.4), it follows that

$$f^{-1}\chi^{-1}(D_j) \subset \tilde{T}_1 \times \mathbf{R}^m, \text{ for all } j \in J.$$

Since $f^{-1}\chi^{-1}(\partial D_j)$ and α are homotopic in \mathfrak{A} , α is a meridian of $\tilde{T}_{k_0} \times \{0\}$ and $\mathfrak{A} \cap \mathcal{L} = \emptyset$, the curve $f^{-1}\chi^{-1}(\partial D_j)$ is not contractible in $(\tilde{T}_0 \times \mathbf{R}^m) \setminus \mathcal{L}$ by [Proposition 2.1](#). Therefore

$$f^{-1}\chi^{-1}(D_j) \cap \mathcal{L} \neq \emptyset, \text{ for all } j \in J.$$

6.5 Remark For simplicity and with a slight abuse of notation, we continue to use the notation f for the composition map $\chi \circ f$. Under this convention, for every $j \in J$,

$$(6.6) \quad \partial D_j \subset f(\mathfrak{A});$$

$f^{-1}(\partial D_j)$ and α are homotopic in \mathfrak{A} ; and

$$(6.7) \quad f^{-1}(D_j) \cap \mathcal{L} \neq \emptyset, D_j \cap f(\mathcal{L}) \neq \emptyset \text{ and } f^{-1}(D_j) \subset \tilde{T}_1 \times [-C_2, C_2]^m$$

for some $C_2 > 1$.

7 Interior-essential components

The tori T_k ($k \geq 0$) used in the Whitehead construction can be arranged topologically so that the meridian ∂B_0^2 on ∂T_0 bounds a 2–cell in T_0 which intersects every T_k ($k \geq 1$) in 2^k mutually disjoint 2–cells enclosed by meridians of T_k . The lower estimate (5.1) is suggested by this idealized setting followed by a homological argument of Freedman and Skora [14]. In this section, we prove a version of their Lemma 2.5 for mappings instead of embeddings.

We adopt some statements from [9, pages 73–74] in locating the essential part of a continuous image of a 2–cell inside an n –manifold. Let M be an n –manifold with boundary and ω a 2–manifold with boundary. A map $\Phi: (\omega, \partial) \rightarrow (M, \partial)$, meaning $\Phi(\omega) \subset M$ and $\Phi(\partial\omega) \subset \partial M$, is said to be *interior-inessential* if there exists a map $\Phi': \omega \rightarrow \partial M$ satisfying $\Phi'|\partial\omega = \Phi|\partial\omega$; it is said to be *interior-essential* if no such Φ' exists. In case ω is a closed 2–cell, Φ is interior-essential if and only if $\Phi(\partial\omega)$ is homotopically nontrivial in ∂M , but trivial in M .

Suppose that ω is a compact, connected 2–manifold in a 2–cell D . Denote by D_ω the unique 2–cell in D containing ω with $\partial D_\omega \subset \partial\omega$. A map $\Phi: (\omega, \partial) \rightarrow (M, \partial)$ is said to be *virtually interior-essential* if Φ extends to an interior-essential $\Phi'': D_\omega \rightarrow M$ satisfying $\Phi''(D_\omega \setminus \omega) \subset \partial M$.

In case M is a solid torus $T = \mathbf{S}^1 \times B^2$, an interior-essential map $\Phi: (\omega, \partial) \rightarrow (T, \partial)$ necessarily maps one of the boundary components of ω to a homotopically nontrivial loop on ∂T . A map $\Phi: (\omega, \partial) \rightarrow (T, \partial)$ is virtually interior-essential if and only if Φ maps the outermost component of $\partial\omega$ to a nonzero multiple of a meridian on ∂T while mapping all other components of $\partial\omega$ to trivial loops there. In other words Φ , in fact the extension $\Phi''|D_\omega$, represents a nonzero multiple of a generator of the relative homology group $H_2(T, \partial; \mathbb{Z}) \cong \mathbb{Z}$. Since a meridian and a core of a torus are linked, $\Phi(\omega)$ must intersect every core of the torus T .

Here and later, a closed curve on ∂T is said to be a *multiple of a meridian* if it is homotopic, in ∂T , to n times a meridian of T for some integer n . A boundary component of a compact 2–manifold ω in a 2–cell D is said to be *outermost* if it is not contained in the 2–cell in D enclosed by any other component of $\partial\omega$.

Recall that the relative homology group

$$H_2(\mathbf{S}^1 \times B^2, \partial; \mathbb{Z}) \cong \mathbb{Z}$$

is generated by the equivalence class $[g]$ with $g: (\Delta, \partial\Delta) \rightarrow (\mathbf{S}^1 \times B^2, \partial(\mathbf{S}^1 \times B^2))$ being continuous and of degree ± 1 on $\partial\Delta$, where Δ is a standard 2–simplex.

The following observation on meridians and longitudes is needed.

7.1 Lemma *Let T_k ($k \geq 0$) be the solid tori in the Whitehead construction. Then the meridians of T_k are not contractible in $\mathbf{S}^3 \setminus \text{Int}(T_k)$ hence not in $T_0 \setminus \text{Int}(T_k)$, and the longitudes of T_k are not contractible in $T_0 \setminus \text{Int}(T_k)$.*

Proof The noncontractibility of the meridians follows from the fact that a meridian (resp. a longitude) of the torus T is a longitude (resp. a meridian) of the torus $\mathbf{S}^3 \setminus \text{Int}(T)$.

Since a core of T_k and a core of $\mathbf{S}^3 \setminus \text{Int}(T_{k-1})$ form a Whitehead link for $k \geq 1$, there is a homeomorphism h of \mathbf{S}^3 interchanging these two cores [21, page 83]. In fact, h can be chosen to map T_k and $\mathbf{S}^3 \setminus \text{Int}(T_{k-1})$ onto $\mathbf{S}^3 \setminus \text{Int}(T_{k-1})$ and T_k respectively, thus to map T_{k-1} onto $\mathbf{S}^3 \setminus \text{Int}(T_k)$. This implies that $\mathbf{S}^3 \setminus \text{Int}(T_{k-1})$ is embedded in $\mathbf{S}^3 \setminus \text{Int}(T_k)$ in same way that T_k is embedded in T_{k-1} for all $k \geq 1$. From this and the fact mentioned in the last paragraph, the statement on the noncontractibility of the longitudes follows. \square

We now prove an analogue of Lemma 2.5 in Freedman and Skora [14] for mappings; it was proved there for embeddings and for the Bing double.

7.2 Lemma *Let T_0 and T_1 be the first two tori in the Whitehead construction, ω be a compact, connected 2-manifold in a 2-cell D , and $\Phi: (\omega, \partial) \rightarrow (T_0, \partial)$ be a virtually interior-essential map. Suppose that $\Phi(\omega)$ meets $\partial T_0 \cup \partial T_1$ transversally. Then $\Phi^{-1}(T_1)$ contains at least two components ω^1, ω^2 on which the maps $\Phi|_{\omega^i}: \omega^i \rightarrow T_1$ ($i = 1, 2$) are virtually interior-essential.*

Proof Assume that Φ has been extended to the 2-cell D_ω , which maps ∂D_ω to a nonzero multiple of a meridian of T_0 and maps $D_\omega \setminus \omega$ into ∂T_0 .

We first check that $[\Phi|_{\Phi^{-1}(T_1)}, \partial] = 0$ in $H_2(T_1, \partial; \mathbb{Z})$, an analogue of an assertion of Freedman and Skora about the Bing double [14, page 82]. Choose a smooth embedding $g: (\Delta, \partial\Delta, \Delta \setminus \partial\Delta) \hookrightarrow (T_0, \partial T_0, T_0 \setminus \partial T_0)$ in such a way that $g(\Delta)$ meets ∂T_1 transversally, $g^{-1}(T_1)$ consists of exactly two 2-cells Δ_i ($i = 1, 2$), $[g, \partial]$ generates $H_2(T_0, \partial; \mathbb{Z})$, and $g|_{\Delta_1}$ and $g|_{\Delta_2}$ represent generators of $H_2(T_1, \partial; \mathbb{Z})$ of opposite signs. Hence

$$[g|_{\Delta_1}, \partial] + [g|_{\Delta_2}, \partial] = 0 \text{ in } H_2(T_1, \partial; \mathbb{Z}).$$

Identify the 2-cells D_ω, Δ_1 and Δ_2 with the standard 2-simplex Δ with proper orientations. Since

$$[\Phi, \partial] = \pm n[g, \partial] \text{ in } H_2(T_0, \partial; \mathbb{Z}),$$

$$(7.3) \quad \Phi \mp n \cdot g + \tau = \partial\sigma,$$

for some 3-chain σ in T_0 and some 2-chain τ in ∂T_0 . After subdivision, all chains in (7.3) are sums of singular simplices in T_1 or $T_0 \setminus \text{Int}(T_1)$. Since the group $H_2(T_0, T_0 \setminus \text{Int}(T_1); \mathbb{Z})$ is isomorphic to $H_2(T_1, \partial; \mathbb{Z})$ by excision, $[\Phi] = [\Phi|\Phi^{-1}(T_1), \partial]$, $[g] = [g|\Delta_1, \partial] + [g|\Delta_2, \partial] = 0$, $[\partial\sigma] = [\partial\sigma'] = 0$ for a 3-chain σ' in T_1 , and $[\tau] = 0$ in $H_2(T_1, \partial; \mathbb{Z})$. Therefore $[\Phi|\Phi^{-1}(T_1), \partial] = 0$ in $H_2(T_1, \partial; \mathbb{Z})$.

By transversality and the Schönflies Theorem, $\Phi^{-1}(T_1)$ is a compact 2-manifold in D_ω and $\Phi^{-1}(\partial T_1)$ is a 1-manifold with no boundary. Moreover $\Phi^{-1}(T_1)$ is nonempty and contains some component on which Φ is interior-essential. Otherwise, $\Phi|\Phi^{-1}(T_1)$ could be redefined in the interior-inessential components to have images in ∂T_1 ; consequently, $\Phi(D_\omega) \subset T_0 \setminus \text{Int}(T_1)$. This would imply that $\Phi(\partial D_\omega)$, a nonzero multiple of a meridian of T_0 , is contractible in $T_0 \setminus \text{Int}(T_1)$.

Among the components of $\Phi^{-1}(T_1)$ on which Φ is interior-essential, (call the entire collection of such components Ω), choose an *innermost* component ω^1 ; ie, D_{ω^1} contains no element in $\Omega \setminus \{\omega^1\}$. We claim that $\Phi|\omega^1$ is *virtually interior-essential*. To verify the claim, note that the boundary of ω^1 consists of a special outermost component ∂D_{ω^1} and possibly some other components β 's whose interior contains no component of $\Phi^{-1}(T_1)$ on which Φ is interior-essential. Modify Φ on the interior-inessential components inside β , so that it maps the 2-cell D_β in D_ω enclosed by β into $T_0 \setminus \text{Int}(T_1)$. This implies that $\Phi(\beta)$ is contractible in $T_0 \setminus \text{Int}(T_1)$. Neither the meridians of T_1 nor the longitudes of T_1 are contractible in $T_0 \setminus \text{Int}(T_1)$ by Lemma 7.1; $\Phi(\beta)$ must be homotopically trivial in ∂T_1 . Assume as we may that $\Phi(D_\beta) \subset \partial T_1$. Since $\Phi|\omega^1$ is interior-essential, $\Phi(\partial D_{\omega^1})$ must be nontrivial in ∂T_1 . Since $\Phi(\partial D_{\omega^1}) \subset \Phi(D_{\omega^1}) \subset T_0$ is contractible in T_0 , it has to be a nonzero multiple, say n_1 , of a meridian of T_1 . This proves the claim.

Assume as we may that $\omega^1 = D_{\omega^1}$. Therefore $\Phi|\omega^1$ represents a nonzero multiple of a generator of the relative homology group $H_2(T_1, \partial; \mathbb{Z})$. Since $[\Phi|\Phi^{-1}(T_1), \partial] = 0$ in $H_2(T_1, \partial; \mathbb{Z})$, $\Omega \setminus \{\omega^1\}$ is nonempty. Suppose that $\omega^2 (\neq \omega^1)$ is another innermost component of Ω , then $\Phi|\omega^2: \omega^2 \rightarrow T_1$ must also be *virtually interior-essential*.

Suppose ω^1 is the only innermost component of Ω . We shall prove $[\Phi|\Phi^{-1}(T_1), \partial] \neq 0$. This leads to a contradiction, thus the validity of the lemma.

The assumption implies that the minimal 2-cells $D_{\omega'}$ in D_ω containing ω' ($\omega' \in \Omega$) are nested and contain ω^1 in their intersection. In other words, the elements of Ω , $\omega^1, \omega^2, \dots, \omega^N$, can be ordered so that $\omega^1 = D_{\omega^1} \subset D_{\omega^2} \subset \dots \subset D_{\omega^N} \subset D_\omega$.

We shall alter Φ in such a way, while not changing the homology class $[\Phi|\Phi^{-1}(T_1), \partial]$ in $H_2(T_1, \partial; \mathbb{Z})$, that $\Phi^{-1}(T_1)$ has only one interior-essential component, namely D_{ω^N} . Since $\Phi(\partial D_\omega)$ is noncontractible in $T_0 \setminus \text{Int}(T_1)$, $\Phi(\partial D_{\omega^N})$ must be homotopically nontrivial in ∂T_1 . It follows then $[\Phi|\Phi^{-1}(T_1), \partial] = [\Phi|D_{\omega^N}, \partial] \neq 0$.

The boundary of ω^2 consists an outermost component ∂D_{ω^2} and a special component b that encloses a 2-cell D_b in D_ω which contains ω^1 in its interior, and possibly some other components β 's whose interiors contain no elements of Ω . Since D_β does not contain any element of Ω , $\Phi(\beta)$ is trivial in ∂T_1 . After altering Φ in the interior of each D_β , we may assume that $\bigcup_\beta \Phi(D_\beta) \subset \partial T_1$ and that Φ maps the annular region in D_ω bounded by b and ∂D_{ω^2} into T_1 . The 2-cell D_b consists of D_{ω^1} and an annular region \mathcal{A} in D_ω bounded by ∂D_{ω^1} and b . The annulus \mathcal{A} contains no elements of Ω due to the ordering. Modifications can be made on the interior-inessential components in \mathcal{A} , so that $\Phi|_{\mathcal{A}}$ maps \mathcal{A} into $T_0 \setminus \text{Int}(T_1)$. The annulus \mathcal{A} provides a homotopy between $\Phi|_b$ and $\Phi|\partial D_{\omega^1}$ in $T_0 \setminus \text{Int}(T_1)$. By homotopy invariance, both $\Phi(b)$ and $\Phi(\partial D_{\omega^1})$ are n_1 times a meridian of T_1 . So Φ can again be altered in $\mathcal{A} \setminus \Phi^{-1}(T_1)$ such that $\Phi(\mathcal{A}) \subset \partial T_1$. After these changes, D_{ω^2} becomes a component of $\Phi^{-1}(T_1)$ while the homological relation $[\Phi|\omega^1, \partial] + [\Phi|\omega^2, \partial] = [\Phi|D_{\omega^2}, \partial]$ remains.

Continuing inductively if $N > 2$, Φ can be altered so that D_{ω^N} becomes the only interior-essential component of $\Phi^{-1}(T_1)$.

As noted earlier $[\Phi|\Phi^{-1}(T_1), \partial] = [\Phi|D_{\omega^N}, \partial] \neq 0$, which gives the desired contradiction. This completes the proof of the lemma. □

8 A lower estimate on the intersection numbers

In this section, we apply [Lemma 7.2](#) to obtain the estimate (5.1).

8.1 PL–approximation

Fix a parameter $j \in J$; write D for the simplex D_j used for slicing, and

$$F = f^{-1}|_D.$$

Let $\Pi: (\mathbf{R}^3/Wh) \times \mathbf{R}^m \rightarrow \mathbf{R}^3/Wh$ be the projection map

$$\Pi(x, z) = x.$$

The goal of this subsection is to construct a single-valued version of $\pi^{-1}\Pi F|_D: D \rightarrow \mathbf{R}^3$ whose image intersects the boundaries of the tori transversally.

For the purpose of applying a PL–approximation ([Theorem 9.5](#) in the [Appendix](#)), the manifold $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ is equipped with a PL–structure and a metric, induced from \mathbf{R}^{3+m} by a fixed homeomorphism (unrelated to f) $h: (\mathbf{R}^3/Wh) \times \mathbf{R}^m \rightarrow \mathbf{R}^{3+m}$,

which is $\pi^{-1} \times \text{id}$ outside $\tilde{T}_{K+1} \times \mathbf{R}^m$ and is diffeomorphic off \mathcal{L} . Fix a large integer $K > k_0$ and let

$$(8.2) \quad \epsilon = c_0 a^{K+2} / 30.$$

From [Theorem 9.5](#) it follows that for any $\epsilon' > 0$, $F|_D$ can be ϵ' -approximated by a locally flat PL embedding $L: D \hookrightarrow (\mathbf{R}^3/Wh) \times \mathbf{R}^m$. When ϵ' is sufficiently small, the approximation satisfies

$$(8.3) \quad d_{\delta_a}(F(x), L(x)) < \epsilon, \quad \text{for all } x \in D \setminus f(\tilde{T}_{K+1} \times \mathbf{R}^m),$$

with respect to the original metric. Adjusting L to one in general position with respect to the boundaries of the tori, we may assume that $L(D)$ and $(\partial\tilde{t}_{k_0} \cup \bigcup_{k=k_0}^K \partial\tilde{T}_{k+1}) \times \mathbf{R}^m$ meet transversally and their intersection is a 1-manifold having no boundary. Adjusting L again, we may further assume that the projection Π embeds each simplex in $L(D) \setminus (\tilde{T}_{K+1} \times \mathbf{R}^m)$, that $\Pi L(D)$ and $\partial\tilde{t}_{k_0} \cup \bigcup_{k=k_0}^K \partial\tilde{T}_{k+1}$ meet transversally and that $(\Pi L)^{-1}(\partial\tilde{t}_{k_0} \cup \bigcup_{k=k_0}^K \partial\tilde{T}_{k+1})$ is a 1-manifold in D having no boundary. See Rourke and Sanderson [\[27, Chapter 5\]](#), Rushing [\[28, page 33\]](#) and [Theorem 9.6](#) in the [Appendix](#).

Note from [\(6.2\)](#), [\(6.6\)](#), [\(8.2\)](#) and [\(8.3\)](#) that $L(\partial D)$ is homotopic to the meridian α of $\tilde{T}_{k_0} \times \{0\}$ in $\tilde{V}_{k_0} \times \mathbf{R}^m$. By [Proposition 2.1](#),

$$(8.4) \quad L(\partial D) = \partial L(D) \text{ is not contractible in } (\tilde{T}_0 \times \mathbf{R}^m) \setminus \mathcal{L};$$

and by [\(6.7\)](#), [\(8.2\)](#) and [\(8.3\)](#),

$$(8.5) \quad q \in \Pi L(D), L(D) \cap \mathcal{L} \neq \emptyset \quad \text{and} \quad L(D) \subset (\tilde{T}_1 \cup \tilde{V}_1) \times [-C_2 - 1, C_2 + 1]^m.$$

The function $\pi^{-1}\Pi L$ is not single-valued on the set $L^{-1}(\mathcal{L}) \cap D$; we modify $\pi^{-1}\Pi L$ near $L^{-1}(\mathcal{L})$ as follows. Let $\Lambda_1, \Lambda_2, \dots, \Lambda_N$ be the 2-cells in D enclosed by the outermost components of $L^{-1}(\partial(\tilde{T}_K \times \mathbf{R}^m))$. Then the 2-cells $\Lambda_1, \Lambda_2, \dots, \Lambda_N$ are mutually disjoint,

$$L^{-1}(\mathcal{L}) \subset \bigcup_1^N \Lambda_n, \quad \text{and} \quad L\left(D \setminus \bigcup_1^N \Lambda_n\right) \cap (\tilde{T}_K \times \mathbf{R}^m) = \emptyset.$$

Note however $L(\bigcup_1^N \Lambda_n)$ need not be contained in $\tilde{T}_K \times \mathbf{R}^m$. Since the homeomorphism $h: (\mathbf{R}^3/Wh) \times \mathbf{R}^m \rightarrow \mathbf{R}^{3+m}$ providing the PL-structure on $(\mathbf{R}^3/Wh) \times \mathbf{R}^m$ is $\pi^{-1} \times \text{id}$ outside $\tilde{T}_{K+1} \times \mathbf{R}^m$, the function $\Phi: D \rightarrow \mathbf{R}^3$ defined by

$$\Phi = \begin{cases} \pi^{-1}\Pi L & \text{in } D \setminus \bigcup_1^N \Lambda_n, \\ \Pi' h L & \text{in } \bigcup_1^N \Lambda_n, \end{cases}$$

is continuous on D . Here $\Pi': \mathbf{R}^3 \times \mathbf{R}^m \rightarrow \mathbf{R}^3$ is the projection with $\Pi'(x, z) = x$. From the discussion leading to (8.4) and (8.5), it follows that

$$\begin{aligned} \Phi(\partial D) &= \pi^{-1}\Pi L(\partial D) \subset V_{k_0} \quad \text{and} \quad \Phi(D) \subset T_1 \cup V_1 \subset T_0, \\ \Phi(\partial D) &\text{ is not contractible in } T_0 \setminus Wh, \\ (8.6) \quad \Phi(D) \cap t_{k_0} &\neq \emptyset \quad \text{and} \quad \Phi(D) \cap T_k \neq \emptyset \text{ for all } k \geq k_0 + 1. \end{aligned}$$

Define also a single-valued version of $\pi^{-1}\Pi F$:

$$\Psi = \begin{cases} \pi^{-1}\Pi F & \text{in } D \setminus \bigcup_1^N \Lambda_n, \\ \Pi' h F & \text{in } \bigcup_1^N \Lambda_n. \end{cases}$$

The ϵ -closeness (8.3) of F and L implies that $\Psi|(D \setminus \bigcup_1^N \Lambda_n)$ and $\Phi|(D \setminus \bigcup_1^N \Lambda_n)$ are ϵ -close with respect to the metric d_a on $\mathbf{R}^3 \setminus Wh$.

8.7 Intersection

We now apply Lemma 7.2 to torus pairs $(t_{k_0}, T_{k_0+1}), (T_{k_0+1}, T_{k_0+2}), \dots, (T_{K-1}, T_K)$. Notice that given any such pair, there is a homeomorphism of \mathbf{S}^3 that maps the first torus in the pair onto T_0 and the second one onto T_1 .

To begin, let ω_{k_0} be an innermost component of $\Phi^{-1}(t_{k_0})$ on which Φ is interior-essential. By (8.6), ω_{k_0} must exist and is nonempty. Following the argument in Lemma 7.2 for the statement that $\Phi|\omega^1$ is virtually interior-essential, we may conclude that $\Phi: (\omega_{k_0}, \partial) \rightarrow (t_{k_0}, \partial)$ is virtually interior-essential; here the fact that neither a meridian of t_{k_0} nor a longitude of t_{k_0} is contractible in $T_0 \setminus \text{Int}(t_{k_0})$ is needed.

Applying Lemma 7.2 to the torus pair (t_{k_0}, T_{k_0+1}) , we conclude $\Phi^{-1}(T_{k_0+1})$ contains at least two components $\omega_{k_0+1}^1$ and $\omega_{k_0+1}^2$ on which the maps $\Phi|\omega_{k_0+1}^i: \omega_{k_0+1}^i \rightarrow T_{k_0+1}$ ($i = 1, 2$) are virtually interior-essential. Apply Lemma 7.2 again to each $\omega_{k_0+1}^i$ ($i = 1, 2$) to obtain at least four components $\omega_{k_0+2}^i$ ($1 \leq i \leq 4$) of $\Phi^{-1}(T_{k_0+2})$ on which the maps $\Phi|\omega_{k_0+2}^i: \omega_{k_0+2}^i \rightarrow T_{k_0+2}$ ($1 \leq i \leq 4$) are virtually interior-essential. Continuing inductively, we get for each k , $k_0 + 1 \leq k \leq K - 1$, at least 2^{k-k_0} components ω_k^i ($1 \leq i \leq 2^{k-k_0}$) of $\Phi^{-1}(T_k)$ on which Φ is virtually interior-essential. In fact, the proof of Lemma 7.2 also shows that for a fixed k , the minimal 2-cells $D\omega_k^i$ in D containing ω_k^i ($1 \leq i \leq 2^{k-k_0}$) are mutually disjoint, and that each $\Phi(\partial D\omega_k^i)$ is a nonzero multiple of a meridian of T_k .

Fix an ω_k^i with $k_0 + 1 \leq k \leq K - 1$ and $1 \leq i \leq 2^{k-k_0}$. For any γ in the curve family \mathcal{C}_k defined in Section 4, $\pi^{-1}\gamma$ (a core of T_k) and $\Phi(\partial D\omega_k^i)$ are linked. So $\pi^{-1}\gamma$ intersects $\Phi(D\omega_k^i)$, in fact intersects $\Phi(\omega_k^i)$.

For each $\gamma \in \mathcal{C}_k$, the core $\pi^{-1}\gamma$ intersects $\Psi(\omega_k^i)$ also. This can be seen as follows. Since $\pi^{-1}\gamma \subset U_k \subset t_k$ and $\Phi(\partial D_{\omega_k^i}) = \pi^{-1}\Pi L(\partial D_{\omega_k^i}) \subset \partial T_k$, we have

$$30\epsilon = c_0 a^{K+2} \leq \text{dist}_{d_a}(t_k, \partial T_k) \leq \text{dist}_{d_a}(\pi^{-1}\gamma, \Phi(\partial D_{\omega_k^i})).$$

From the ϵ -closeness of Ψ and Φ , it follows that $\Psi(\omega_k^i)$ is contained in $V_k \cup T_k$, a neighborhood of T_k , and that $\Psi|_{\partial\omega_k^i}$ and $\Phi|_{\partial\omega_k^i}$ are homotopic in $V_k \setminus U_k$, a neighborhood of ∂T_k outside t_k . So $\pi^{-1}\gamma$ and $\Psi(\partial D_{\omega_k^i})$ are also linked. Therefore $\pi^{-1}\gamma$ intersects $\Psi(D_{\omega_k^i})$, in fact intersects $\Psi(\omega_k^i)$.

The intersection of $\pi^{-1}\gamma$ and $\Psi(\omega_k^i)$ necessarily takes place at points in U_k , which is outside the torus T_K . Therefore $\Psi = \pi^{-1}\Pi F$ at the points of intersection. So γ intersects $\Pi F(\omega_k^i)$ for each $\gamma \in \mathcal{C}_k$. Consequently, Γ intersects $F(\omega_k^i)$ for each $\Gamma \in \mathcal{S}_k$.

This together with [Remark 6.5](#) shows that for each k , $k_0 + 1 \leq k \leq K - 1$,

$$N(k, \Gamma, j) \geq 2^{k-k_0} \quad \text{for all } \Gamma \in \mathcal{S}_k \text{ and } j \in J.$$

Since K can be arbitrarily large, estimate [\(5.1\)](#) follows. This completes the proof of [Theorem 1.1](#).

9 Appendix

We state some topological facts needed in our proof.

The first theorem is about the size of the complement of a non-simply connected set proved by Martio, Rickman and Väisälä [\[23\]](#).

9.1 Theorem *Suppose that G is a simply connected domain in \mathbf{R}^n ($n \geq 3$) and A is a closed subset of G . If $G \setminus A$ is not simply connected, then A must have positive $(n-2)$ -dimensional Hausdorff measure.*

Next is a theorem about PL-approximation of embeddings of one-dimensional complexes due to Moise [\[25, page 46\]](#).

9.2 Theorem *Let K^1 be a 1-dimensional complex, $f: |K^1| \hookrightarrow \mathbf{R}^2$ an embedding, and $\psi > 0$ a continuous function on $|K^1|$. Then there exists a piecewise linear embedding $\theta: |K^1| \hookrightarrow \mathbf{R}^2$ such that*

- (1) $|\theta(x) - f(x)| < \psi(x)$ for all $x \in |K^1|$ (ψ -approximation of f),
- (2) $\theta(v) = f(v)$ for all vertices v of K^1 .

9.3 Remark The theorem remains true if the target space is \mathbf{R}^n for any $n \geq 2$. The proof is essentially unchanged.

The following theorem on unknotting spheres is due to Zeeman [40]; the second sentence in the statement below can be deduced from the proof of Theorem 5.6 in Rourke and Sanderson [27] and the propositions mentioned therein.

9.4 Theorem Any piecewise linear topological 1–sphere $l \subset \mathbf{S}^n$ is unknotted in \mathbf{S}^n when $n \geq 4$. More precisely, there is a global PL–homeomorphism L of \mathbf{S}^n with the following properties:

- (1) L maps l onto the boundary of a standard 2–simplex, in $B(Y, 2 \operatorname{diam} l)$ for some point $Y \in l$.
- (2) $L \equiv \operatorname{id}$ in $\mathbf{S}^n \setminus B(Y, 2 \operatorname{diam} l)$.

The following PL–approximation of topological embeddings of k –cells into a manifold M^n is due to Ancel and Cannon [2] when codimension is 1, due to Miller [24] and Černavskii [8] when codimension is 3 or higher and due to Venema [37; 38] when codimension is 2. We need it for $k = 2$ and $n \geq 4$.

9.5 Theorem Let $h: I^k \rightarrow M^n$ be a topological embedding of the closed k –cell I^k into a piecewise linear n –manifold M^n and $n - k \geq 1$. Then for every $\epsilon > 0$ there exists a locally flat piecewise linear embedding $L: I^k \rightarrow M^n$ such that $d(h(x), L(x)) \leq \epsilon$ for each $x \in I^k$.

The following theorem on transversality is well-known (see for example Daverman [9, page 69]). When the maps are piecewise linear the proof is simpler.

9.6 Theorem Suppose M^n is an n –manifold, Σ is an $(n - 1)$ –manifold embedded in M as a closed and bicollared subset, $F: B^2 \rightarrow M$ is a continuous map, U is an open subset of M containing Σ , and $\epsilon > 0$. Then there exists a map $L: B^2 \rightarrow M$ satisfying

- (1) $L|(B^2 - F^{-1}U) = F|(B^2 - F^{-1}U)$,
- (2) $d(L, F) < \epsilon$,
- (3) $L^{-1}\Sigma$ is a 1–manifold in B^2 with $\partial(L^{-1}\Sigma) = \partial B^2 \cap L^{-1}\Sigma$,
- (4) for any point $x \in L^{-1}\Sigma$, $L(B^2)$ pierces Σ at $L(x)$, ie, for each neighborhood W of x in B^2 , $L(W)$ touches both sides of the bicollar on Σ near $L(x)$.

We say $L(B^2)$ and Σ meet transversally if the map $L: B^2 \rightarrow M$ satisfies (3) and (4).

References

- [1] **L Ahlfors, A Beurling**, *Conformal invariants and function-theoretic null-sets*, Acta Math. 83 (1950) 101–129 [MR0036841](#)
- [2] **F D Ancel, J W Cannon**, *The locally flat approximation of cell-like embedding relations*, Ann. of Math. (2) 109 (1979) 61–86 [MR519353](#)
- [3] **J J Andrews, L Rubin**, *Some spaces whose product with E^1 is E^4* , Bull. Amer. Math. Soc. 71 (1965) 675–677 [MR0176454](#)
- [4] **R H Bing**, *The cartesian product of a certain nonmanifold and a line is E^4* , Ann. of Math. (2) 70 (1959) 399–412 [MR0107228](#)
- [5] **R H Bing**, *A set is a 3 cell if its cartesian product with an arc is a 4 cell*, Proc. Amer. Math. Soc. 12 (1961) 13–19 [MR0123303](#)
- [6] **M Bonk, B Kleiner**, *Quasisymmetric parametrizations of two-dimensional metric spheres*, Invent. Math. 150 (2002) 127–183 [MR1930885](#)
- [7] **J W Cannon**, *The recognition problem: what is a topological manifold?*, Bull. Amer. Math. Soc. 84 (1978) 832–866 [MR0494113](#)
- [8] **A V Černavskii**, *Piece-wise linear approximation of embeddings of manifolds in codimensions greater than two. A supplement to the article “Piece-wise linear approximation of embeddings of cells and spheres in codimensions greater than two”*, Mat. Sb. (N.S.) 82 (124) (1970) 499–500 [MR0275438](#)
- [9] **R J Daverman**, *Decompositions of manifolds*, Pure and Applied Math. 124, Academic Press, Orlando, FL (1986) [MR872468](#)
- [10] **G David, S Semmes**, *Quantitative rectifiability and Lipschitz mappings*, Trans. Amer. Math. Soc. 337 (1993) 855–889 [MR1132876](#)
- [11] **R D Edwards**, *Suspensions of homology spheres* [arXiv:math/0610573](#)
- [12] **R D Edwards**, *The topology of manifolds and cell-like maps*, from: “Proceedings of the International Congress of Mathematicians (Helsinki, 1978)”, Acad. Sci. Fennica, Helsinki (1980) 111–127 [MR562601](#)
- [13] **H Federer**, *Geometric measure theory*, Grundlehren der math. Wissenschaften 153, Springer, New York (1969) [MR0257325](#)
- [14] **M H Freedman, R Skora**, *Strange actions of groups on spheres*, J. Differential Geom. 25 (1987) 75–98 [MR873456](#)
- [15] **B Fuglede**, *Extremal length and functional completion*, Acta Math. 98 (1957) 171–219 [MR0097720](#)
- [16] **F W Gehring**, *Rings and quasiconformal mappings in space*, Trans. Amer. Math. Soc. 103 (1962) 353–393 [MR0139735](#)
- [17] **F W Gehring**, *The L^p -integrability of the partial derivatives of a quasiconformal mapping*, Acta Math. 130 (1973) 265–277 [MR0402038](#)

- [18] **J Heinonen, P Koskela**, *Definitions of quasiconformality*, Invent. Math. 120 (1995) 61–79 [MR1323982](#)
- [19] **J Heinonen, P Koskela**, *Quasiconformal maps in metric spaces with controlled geometry*, Acta Math. 181 (1998) 1–61 [MR1654771](#)
- [20] **J Heinonen, S Semmes**, *Thirty-three yes or no questions about mappings, measures, and metrics*, Conform. Geom. Dyn. 1 (1997) 1–12 [MR1452413](#)
- [21] **R C Kirby**, *The topology of 4-manifolds*, Lecture Notes in Math. 1374, Springer, Berlin (1989) [MR1001966](#)
- [22] **T J Laakso**, *Plane with A_∞ -weighted metric not bi-Lipschitz embeddable to \mathbb{R}^N* , Bull. London Math. Soc. 34 (2002) 667–676 [MR1924353](#)
- [23] **O Martio, S Rickman, J Väisälä**, *Topological and metric properties of quasiregular mappings*, Ann. Acad. Sci. Fenn. Ser. A I (1971) 31 [MR0299782](#)
- [24] **R T Miller**, *Approximating codimension 3 embeddings*, Ann. of Math. (2) 95 (1972) 406–416 [MR0307246](#)
- [25] **E E Moise**, *Geometric topology in dimensions 2 and 3*, Graduate Texts in Math. 47, Springer, New York (1977) [MR0488059](#)
- [26] **S Rickman**, *Quasiregular mappings*, Ergebnisse der Math. und ihrer Grenzgebiete (3) 26, Springer, Berlin (1993) [MR1238941](#)
- [27] **C P Rourke, B J Sanderson**, *Introduction to piecewise-linear topology*, Springer Study Edition (1982) [MR665919](#)
- [28] **T B Rushing**, *Topological embeddings*, Pure and Applied Math. 52, Academic Press, New York (1973) [MR0348752](#)
- [29] **S Semmes**, *Chord-arc surfaces with small constant. II. Good parameterizations*, Adv. Math. 88 (1991) 170–199 [MR1120612](#)
- [30] **S Semmes**, *Finding curves on general spaces through quantitative topology, with applications to Sobolev and Poincaré inequalities*, Selecta Math. (N.S.) 2 (1996) 155–295 [MR1414889](#)
- [31] **S Semmes**, *Good metric spaces without good parameterizations*, Rev. Mat. Iberoamericana 12 (1996) 187–275 [MR1387590](#)
- [32] **S Semmes**, *On the nonexistence of bi-Lipschitz parameterizations and geometric problems about A_∞ -weights*, Rev. Mat. Iberoamericana 12 (1996) 337–410 [MR1402671](#)
- [33] **S Semmes**, *Some topics concerning homeomorphic parameterizations*, Publ. Mat. 45 (2001) 3–67 [MR1829576](#)
- [34] **L Siebenmann, D Sullivan**, *On complexes that are Lipschitz manifolds*, from: “Geometric topology (Proc. Georgia Topology Conf., Athens, Ga., 1977)”, (J C Cantrell, editor), Academic Press, New York (1979) 503–525 [MR537747](#)

- [35] **P Tukia, J Väisälä**, *Quasisymmetric embeddings of metric spaces*, Ann. Acad. Sci. Fenn. Ser. A I Math. 5 (1980) 97–114 [MR595180](#)
- [36] **J Väisälä**, *Lectures on n -dimensional quasiconformal mappings*, Lecture Notes in Math. 229, Springer, Berlin (1971) [MR0454009](#)
- [37] **G A Venema**, *Approximating disks in 4-space*, Michigan Math. J. 25 (1978) 19–27 [MR497879](#)
- [38] **G A Venema**, *Approximating codimension two embeddings of cells*, Pacific J. Math. 126 (1987) 165–195 [MR868611](#)
- [39] **J H C Whitehead**, *A certain open manifold whose group is unity*, Quart. J. Math. Oxford (2) 6 (1935) 268–279
- [40] **E C Zeeman**, *Unknotting spheres*, Ann. of Math. (2) 72 (1960) 350–361 [MR0117738](#)

Department of Mathematics, University of Michigan
530 Church Street, Ann Arbor, MI 48109-1043, USA

Department of Mathematics, University of Illinois
1409 West Green Street, Urbana, IL 61822, USA

wu@math.uiuc.edu

Proposed: Colin Rourke

Received: 5 March 2008

Seconded: Tobias Colding, David Gabai

Revised: 28 October 2009