

## POSITIVELY INVARIANT SETS FOR FUNCTIONAL DIFFERENTIAL EQUATIONS WITH INFINITE DELAY

Dedicated to Professor Taro Yoshizawa on his sixtieth birthday

KENSUKE SAWANO

(Received July 31, 1979)

**1. Introduction.** For ordinary differential equations, many authors have discussed necessary and sufficient conditions for a closed set in the  $n$ -dimensional Euclidean space  $R^n$  to be positively invariant. Yorke [11] has discussed this problem by using a non-Lipschitzian Liapunov function which is lower-semicontinuous. For an autonomous system, Brezis [1] obtained a result under the assumption that the right hand side of the system is locally Lipschitzian, and his proof depends essentially on this assumption. Crandall [2] obtained a similar result by applying the method of polygonal approximations. For a nonautonomous system, Hartman [5] also considered an approximation which is different from the one considered in [2].

The purpose of this article is to discuss the same question for functional differential equations with infinite delay. Seifert [10] also discussed this question under the assumption that a closed set is convex. In Section 2, we introduce an abstract phase space  $B$  which satisfies some general hypotheses slightly different from those considered in [4]. We consider a subset  $\Omega$  in  $R \times R^n$  such that the cross section  $\Omega_t = \{y \in R^n; (t, y) \in \Omega\}$  is convex for all  $t \in R$  and that the cross section  $\Omega_t$  satisfies a continuity condition in the sense of Hausdorff metric. We discuss the properties of  $\Omega$  which play an important role in Section 3. In Section 3, we state the main theorem. We give the necessary and sufficient condition that, for any initial value  $(\sigma, \phi)$  in  $R \times B$  such that  $\phi(t - \sigma) \in \Omega_t$  for all  $t \leq \sigma$ , there exists at least one solution  $x(t)$  through  $(\sigma, \phi)$  which is defined on its right maximal interval of existence and satisfies  $(t, x(t)) \in \Omega$  there. Special approximate solutions are needed to prove the theorem. The construction of the solutions, although analogous to the one in [5], is much more complicated for functional differential equations. The proof of the theorem is given in Section 4. The case where the delay is finite has been considered in [7] and [8] by a different approach.

**2. Preliminaries.** Let  $R^n$  be an  $n$ -dimensional real linear vector space, and let  $R = R^1$ . We denote by  $B$  a real linear vector space of functions mapping  $(-\infty, 0]$  into  $R^n$  with a semi-norm  $|\cdot|$ . No confusion will occur if we use the same symbol  $|\cdot|$  to denote the norm in  $R^n$ . For elements  $\phi$  and  $\psi$  in  $B$ ,  $\phi = \psi$  means that  $\phi(\theta) = \psi(\theta)$  for all  $\theta$  in  $(-\infty, 0]$ . Then the quotient space  $B^* = B/|\cdot|$  is a normed linear space with the norm naturally induced by the semi-norm. The topology of  $B$  is defined by the semi-norm, that is, a family  $\{V(\phi, \varepsilon); \phi \in B, \varepsilon > 0\}$  is an open base, where  $V(\phi, \varepsilon) = \{\psi \in B; |\phi - \psi| < \varepsilon\}$ .  $B$  with this topology is a pseudo-metric space.

For any  $\phi$  in  $B$  and any  $\beta \geq 0$ , let  $\phi^\beta$  be the restriction of  $\phi$  to the interval  $(-\infty, -\beta]$ . This is a function mapping  $(-\infty, -\beta]$  into  $R^n$ . Denote the space of such functions  $\phi^\beta$  by  $B^\beta$  and define a semi-norm  $|\cdot|_\beta$  in  $B$  by

$$|\eta|_\beta = \inf \{|\psi|; \psi \in B, \psi^\beta = \eta\}, \quad \eta \in B^\beta.$$

If we let  $|\phi|_\beta = |\phi^\beta|_\beta$  for  $\phi \in B$ , then  $|\cdot|_\beta$  is also a semi-norm in  $B$ .

For an  $R^n$ -valued function  $x$  defined on  $(-\infty, \sigma)$ , we define the function  $x_t$  for each  $t \in (-\infty, \sigma)$  by the relation  $x_t(\theta) = x(t + \theta)$ ,  $-\infty < \theta \leq 0$ .

Let  $D$  be an open set in  $R \times B$  and let  $f: D \rightarrow R^n$  be a given continuous function. A functional differential equation on  $D$  is the relation

$$(1) \quad x'(t) = f(t, x_t),$$

where  $x'(t)$  stands for the right hand derivative of  $x(t)$ . For  $(\sigma, \phi)$  in  $D$ , an  $R^n$ -valued function  $x$  defined on  $(-\infty, \sigma + A)$  with  $0 < A \leq \infty$  is said to be a solution of (1) through  $(\sigma, \phi)$  if  $x_\sigma = \phi$  and if  $x$  is continuously differentiable and satisfies (1) for all  $t \in [\sigma, \sigma + A)$ .

We make the following hypotheses on the space  $B$ .

(B1) For an  $A > 0$ , let  $x: (-\infty, A) \rightarrow R^n$  be a function such that  $x_0$  is in  $B$  and  $x$  is continuous on  $[0, A)$ . Then  $x_t$  is in  $B$  for all  $t$  in  $[0, A)$  and  $x_t$  is continuous in  $t$ .

(B2) There is a continuous function  $K(\beta) > 0$  such that

$$|\phi| \leq K(\beta) \sup_{-\beta \leq \theta \leq 0} |\phi(\theta)| + |\phi|_\beta$$

for all  $\phi$  in  $B$  and for all  $\beta$  in  $[0, \infty)$ .

Under the hypotheses (B1) and (B2), there exists a solution of (1) through  $(\sigma, \phi)$  in  $D$ . This was proved by Kaminogo [4].

For  $(\sigma, \phi)$  in  $D$ , let  $Q(\sigma, \phi)$  be the collection of  $(T, x)$ , where  $T > 0$  and  $x$  is a solution of (1) through  $(\sigma, \phi)$  defined on  $(-\infty, \sigma + T)$ . We introduce a partial order  $\leq$  in  $Q(\sigma, \phi)$  in the following way. For ele-

ments  $(T^1, x^1)$  and  $(T^2, x^2)$  in  $Q(\sigma, \phi)$ , we write  $(T^1, x^1) \leq (T^2, x^2)$  when  $T^1 \leq T^2$  and the restriction of  $x^2$  to the interval  $(-\infty, \sigma + T^1)$  is equal to  $x^1$ . Then Zorn's lemma implies the existence of a maximal element  $(T, x)$  in  $Q(\sigma, \phi)$ , and  $x$  is called a right maximal solution of (1) through  $(\sigma, \phi)$  and the interval  $(-\infty, \sigma + T)$  is called the right maximal interval of existence of  $x$ .

Under the hypotheses (B1) and (B2), we have the following.

**LEMMA 1.** *For any  $\phi$  in  $B$  and constants  $A > 0, L > 0$ , let  $F'_A(\phi)$  be a set of functions  $u: (-\infty, A] \rightarrow R^n$  such that  $u_0 = \phi$  and  $|u(t) - u(s)| \leq L|t - s|$  on  $[0, A]$ . Then the set  $\Gamma = \{u_i; u \in F'_A(\phi), t \in [0, A]\}$  is compact in  $B$ .*

For the proof, see Lemma 2.1 of Hale and Kato [4], though the phase space considered in [4] is slightly different from ours.

Let  $\Omega$  be a set in  $R \times R^n$  such that the cross section  $\Omega_t = \{y \in R^n; (t, y) \in \Omega\}$  is nonempty for all  $t \in R$ . Assume that  $\Omega$  satisfies the following continuity condition (C).

(C) For any  $\epsilon > 0$  and any  $t \in R$ , there is a  $\delta = \delta(\epsilon, t) > 0$  such that if  $|t - s| < \delta$ , then

$$\inf \{r > 0; U(\Omega_t, r) \supset \Omega_s \text{ and } U(\Omega_s, r) \supset \Omega_t\} < \epsilon,$$

where  $U(\Omega_t, r)$  is an  $r$ -neighborhood of  $\Omega_t$ .

**LEMMA 2.** *If  $\Omega_t$  is a closed set in  $R^n$  for any  $t \in R$  and the condition (C) is satisfied, then  $\Omega$  is a closed set in  $R \times R^n$ .*

**PROOF.** If the conclusion is false, then there is a sequence  $\{(t_k, y_k)\}$  in  $\Omega$  such that  $(t_k, y_k) \rightarrow (t_0, y_0) \notin \Omega$  as  $k \rightarrow \infty$ . Since  $y_0 \notin \Omega_{t_0}$  and  $\Omega_{t_0}$  is closed, we see that  $U(y_0, \epsilon_0) \cap \Omega_{t_0}$  is empty for some  $\epsilon_0 > 0$ . On the other hand, if  $k$  is large,  $U(y_k, \epsilon_0/3)$  contains a point  $z_k \in \Omega_{t_0}$  since the condition (C) implies that  $\Omega_{t_k} \subset U(\Omega_{t_0}, \epsilon_0/3)$  for sufficiently large  $k$ . Moreover,  $|y_k - y_0| < \epsilon_0/3$  if  $k$  is large. Thus for sufficiently large  $k$ , we have

$$|y_0 - z_k| \leq |y_0 - y_k| + |y_k - z_k| < \epsilon_0/3 + \epsilon_0/3 < \epsilon_0,$$

a contradiction to the emptiness of  $U(y_0, \epsilon_0) \cap \Omega_{t_0}$ , and we are done.

From now on, let  $|y| = (\sum_{i=1}^n y_i^2)^{1/2}$  for  $y = (y_1, \dots, y_n)$  in  $R^n$ .

**LEMMA 3.** *Suppose that  $\Omega_t$  is closed convex for all  $t \in R$  and that the condition (C) is satisfied. For a continuous function  $p(t): [\sigma, \infty) \rightarrow R^n$ , let  $d(p(t), \Omega_t) = \inf \{|p(t) - y|; y \in \Omega_t\}$ . Then there is a continuous function  $g(t): [\sigma, \infty) \rightarrow R^n$  such that  $g(t) \in \Omega_t$  and  $d(p(t), \Omega_t) = |p(t) - g(t)|$ .*

PROOF. Since  $\Omega_t$  is closed, there exists a  $g(t) \in \Omega_t$  with  $d(p(t), \Omega_t) = |p(t) - g(t)|$  for each  $t \in [\sigma, \infty)$ . We show that  $g(t)$  is uniquely determined for each  $t$ . Otherwise, there would exist a  $z \in \Omega_s$  for some  $s \in [\sigma, \infty)$  such that  $z \neq g(s)$  and  $|p(s) - z| = d(p(s), \Omega_s) = |p(s) - g(s)|$ . Set  $d(p(s), \Omega_s) = r$  and let  $S(p(s), r)$  denote the sphere in  $R^n$  with radius  $r$  and center  $p(s)$ . Then  $g(s)$  and  $z$  belong not only to  $\Omega_s$  but also to  $S(p(s), r)$ . Since  $\Omega_s$  is convex, the segment  $\lambda g(s) + (1 - \lambda)z$  with  $0 \leq \lambda \leq 1$  belongs to  $\Omega_s$ . We see immediately that  $|p(s) - \{\lambda g(s) + (1 - \lambda)z\}| < r$  for  $0 < \lambda < 1$ , which contradicts  $d(p(s), \Omega_s) = r$ .

Next the continuity of  $d(p(t), \Omega_t)$  in  $t$  will be proved. For any  $t, s \in [\sigma, \infty)$ , we have

$$(2) \quad |d(p(t), \Omega_t) - d(p(s), \Omega_s)| \\ \leq |d(p(t), \Omega_t) - d(p(t), \Omega_s)| + |d(p(t), \Omega_s) - d(p(s), \Omega_s)|.$$

For any  $\varepsilon > 0$  and any fixed  $t$  in  $[\sigma, \infty)$ , there exists a  $\delta_1 = \delta_1(t, \varepsilon) > 0$  such that if  $|t - s| < \delta_1$ , then

$$(3) \quad |d(p(t), \Omega_s) - d(p(s), \Omega_s)| < \varepsilon/2,$$

because we have  $|d(p(t), \Omega_s) - d(p(s), \Omega_s)| \leq |p(t) - p(s)|$ . Let  $d(p(t), \Omega_s) = |p(t) - u^s|$  for  $u^s \in \Omega_s$ . Then, by the condition (C), there exists a  $\delta_2 = \delta_2(t, \varepsilon) > 0$  such that if  $|t - s| < \delta_2$ , then  $U(u^s, \varepsilon/2)$  contains a point  $v^s$  in  $\Omega_t$  and  $U(g(t), \varepsilon/2)$  contains a point  $w^s$  in  $\Omega_s$ . Therefore we have

$$d(p(t), \Omega_t) \leq |p(t) - v^s| \leq |p(t) - u^s| + |u^s - v^s| \leq d(p(t), \Omega_s) + \varepsilon/2$$

and

$$d(p(t), \Omega_s) \leq |p(t) - w^s| \leq |p(t) - g(t)| + |g(t) - w^s| \leq d(p(t), \Omega_t) + \varepsilon/2,$$

which then imply that if  $|t - s| < \delta_2$ , we have

$$(4) \quad |d(p(t), \Omega_t) - d(p(t), \Omega_s)| \leq \varepsilon/2.$$

Combining (3) and (4), the right hand side of (2) is less than  $\varepsilon$  if  $|t - s| < \delta$ , where  $\delta = \min\{\delta_1, \delta_2\}$ . Thus  $d(p(t), \Omega_t)$  is continuous in  $t$ .

Finally we show that  $g(t)$  is continuous. Suppose that  $g(t)$  is not continuous at  $t = t_0 \geq \delta$ . Then there exists an  $\varepsilon_0 > 0$  and a sequence  $\{t_k\}$  such that  $t_k \rightarrow t_0$  as  $k \rightarrow \infty$  and that  $|g(t_k) - g(t_0)| \geq \varepsilon_0$  for all  $k = 1, 2, \dots$ . Since  $p(t)$  and  $d(p(t), \Omega_t)$  are continuous in  $t$ , the sequence  $\{g(t_k)\}$  is bounded, and hence we may assume that the sequence is convergent. Set  $\lim_{k \rightarrow \infty} g(t_k) = z_0$ . Then  $z_0 \in \Omega_{t_0}$  by Lemma 2. Moreover, since  $d(p(t), \Omega_t) = |p(t) - g(t)|$  and  $p(t)$  are continuous in  $t$ , we have

$$|p(t_0) - z_0| = \lim_{k \rightarrow \infty} |p(t_k) - g(t_k)| = \lim_{k \rightarrow \infty} d(p(t_k), \Omega_{t_k}) = d(p(t_0), \Omega_{t_0}).$$

Thus  $z_o = g(t_o)$  because of the uniqueness of  $g(t)$ . On the other hand,  $|g(t_k) - g(t_o)| \geq \epsilon_o$  implies  $|g(t_o) - z_o| \geq \epsilon_o$ , which contradicts  $z_o = g(t_o)$ . This proves that  $g(t)$  is continuous and completes the proof.

**3. The main result.** Consider a system

$$(5) \quad x'(t) = f(t, x_t),$$

where  $f: R \times B \rightarrow R^n$  is a continuous function.

**THEOREM.** Assume that  $\Omega_t$  is closed convex for all  $t \in R$  and the condition (C) is satisfied. Then the following two statements are equivalent:

(i) For any  $(\sigma, \phi) \in R \times B$  with  $\phi(t - \sigma) \in \Omega_t$  for all  $t \leq \sigma$ , there exists at least one solution  $x$  of (5) through  $(\sigma, \phi)$  defined on its right maximal interval of existence and satisfying  $(t, x(t)) \in \Omega$  on the interval.

(ii) For any  $(\sigma, \phi) \in R \times B$  with  $\phi(t - \sigma) \in \Omega_t$  for all  $t \leq \sigma$ , it holds that

$$\lim_{h \rightarrow 0^+} d(\phi(0) + hf(\sigma, \phi), \Omega_{\sigma+h})/h = 0.$$

We prove this theorem in the next section. In the rest of this section, we consider special approximate solutions under the condition (ii).

Let  $(\sigma, \phi) \in R \times B$  be such that  $\phi(t - \sigma) \in \Omega_t$  for all  $t \leq \sigma$ . Since  $f$  is continuous at  $(\sigma, \phi)$ , there are positive constants  $r, A$  and  $\delta$  such that  $|f| \leq r$  on  $[\sigma, \sigma + A] \times V(\phi, \delta)$ . Let  $L = \max \{K(\beta); 0 \leq \beta \leq A\} > 0$ . Define  $\tilde{\phi}$  by

$$\tilde{\phi}(t) = \begin{cases} \phi(t - \sigma), & t \leq \sigma, \\ \phi(0), & t \geq \sigma. \end{cases}$$

Then  $\tilde{\phi}_t$  belongs to  $B$  for all  $t \geq \sigma$  by the hypothesis (B1) and  $\tilde{\phi}_\sigma = \phi$ . Furthermore, by the hypothesis (B1), there is an  $\alpha = \alpha(\sigma, \phi)$  with  $0 < \alpha \leq A$  such that

$$(6) \quad 3Lr\alpha + |\tilde{\phi}_t - \phi| < \delta \quad \text{for all } t \in [\sigma, \sigma + \alpha].$$

The set  $W$  defined by

$$W = \{(t, u_t); \sigma \leq t \leq \sigma + \alpha, u_\sigma = \phi \text{ and } |u(t) - u(s)| \leq 2r|t - s| \text{ on } [\sigma, \sigma + \alpha]\}$$

is compact in  $R \times B$  by Lemma 1.

Let  $\epsilon, 0 < \epsilon < r$ , be given. Since  $W$  is compact, there is an  $\eta(\epsilon, W) > 0$  such that

$$(7) \quad |f(t, \phi^1) - f(t, \phi^2)| < \epsilon$$

if  $(t, \phi^1) \in W$  and  $|\phi^1 - \phi^2| < \eta(\varepsilon, W)$ , where we can assume that

$$(8) \quad \eta(\varepsilon, W) < Lr\alpha.$$

Now consider the set  $Q_\varepsilon(\sigma, \phi)$  which consists of all  $(T, x)$ , where  $0 < T \leq \alpha$  and  $x$  is a function mapping  $(-\infty, \sigma + T]$  into  $R^n$  with the following properties:

(I)  $x_\sigma = \phi$ ,  $x(\sigma + T) \in \Omega_{\sigma+T}$  and  $d(x(t), \Omega_t) < \eta(\varepsilon, W)L^{-1}$  for all  $t \in [\sigma, \sigma + T]$ .

(II)  $|x(t) - x(t')| \leq 2r|t - t'|$  on  $[\sigma, \sigma + T]$ .

(III)  $|\dot{x}(t) - f(t, x_t)| \leq 3\varepsilon$  for almost all  $t \in [\sigma, \sigma + T]$ , where  $\dot{x}(t)$  is the derivative of  $x(t)$ .

(IV) Every subinterval of  $[\sigma, \sigma + T]$  of length  $\varepsilon$  contains a point  $s$  such that  $(s, x(s)) \in \Omega$ .

LEMMA 4. *The set  $Q_\varepsilon(\sigma, \phi)$  is nonempty for any small  $\varepsilon > 0$ .*

PROOF. By Lemma 3, there is a continuous mapping  $g: [\sigma, \infty) \rightarrow R^n$  such that  $d(\phi(0) + hf(\sigma, \phi), \Omega_{\sigma+h}) = |\phi(0) + hf(\sigma, \phi) - g(\sigma + h)|$  and  $g(\sigma + h) \in \Omega_{\sigma+h}$  for all  $h \geq 0$ . For  $S$  with  $0 < S \leq \varepsilon$ , define a function  $y$  by

$$y(t) = \begin{cases} \phi(t - \sigma), & t \leq \sigma, \\ \phi(0) + \{(g(\sigma + S) - \phi(0))/S\}(t - \sigma), & \sigma < t \leq \sigma + S. \end{cases}$$

We show that  $(S, y)$  belongs to  $Q_\varepsilon(\sigma, \phi)$  if  $S$  is sufficiently small.

The condition (ii) implies that there is a  $\delta_1$  with  $0 < \delta_1 \leq \varepsilon$  such that

$$(9) \quad |(g(\sigma + h) - \phi(0))/h - f(\sigma, \phi)| < \varepsilon$$

for all  $h \in (0, \delta_1]$ . Hence if  $S \leq \delta_1$ , we have

$$(10) \quad \begin{aligned} |y(t) - y(t')| &= |(g(\sigma + S) - \phi(0))/S||t - t'| \\ &\leq (|f(\sigma, \phi)| + \varepsilon)|t - t'| \leq 2r|t - t'| \end{aligned}$$

on  $[\sigma, \sigma + S]$ . Then by the hypothesis (B2), we have  $|y_t - \phi| \leq |y_t - \tilde{\phi}_t| + |\tilde{\phi}_t - \phi| \leq 2rL(t - \sigma) + |\tilde{\phi}_t - \phi|$  for all  $t \in [\sigma, \sigma + S]$ . Hence the continuity of  $f$  implies that there is a  $\delta_2$  with  $0 < \delta_2 \leq \delta_1$  such that  $|f(\sigma, \phi) - f(t, y_t)| < \varepsilon$  for all  $t \in [\sigma, \sigma + S]$  if  $S \leq \delta_2$ . From this and (9), it follows that

$$(11) \quad \begin{aligned} |\dot{y}(t) - f(t, y_t)| &\leq |(g(\sigma + S) - \phi(0))/S - f(\sigma, \phi)| + |f(\sigma, \phi) - f(t, y_t)| \\ &\leq 2\varepsilon \end{aligned}$$

for all  $t \in [\sigma, \sigma + S]$  if  $S \leq \delta_2$ .

Since  $g(\sigma) = \phi(0) = y(\sigma)$  and  $y(t)$  satisfies (10) for  $S \leq \delta_2$ , there is a  $\delta_3$  with  $0 < \delta_3 \leq \delta_2$  such that  $|g(t) - y(t)| < \eta(\varepsilon, W)L^{-1}$  on  $[\sigma, \sigma + S]$  if  $S \leq \delta_3$ . Therefore we have

$$(12) \quad d(y(t), \Omega_t) \leq |y(t) - g(t)| < \eta(\varepsilon, W)L^{-1}$$

for all  $t \in [\sigma, \sigma + S]$  if  $S \leq \delta_3$ . From (10), (11) and (12), it follows that  $y(t)$  satisfies (I), (II) and (III) if  $S = \delta_3$ . The condition (IV) is also satisfied because  $0 < \delta_3 \leq \varepsilon$ . This completes the proof.

LEMMA 5. *There is an element  $(\alpha, x)$  in  $Q_\varepsilon(\sigma, \phi)$  for any small  $\varepsilon > 0$ .*

PROOF. Introduce a partial order  $\leq$  in  $Q_\varepsilon(\sigma, \phi)$  as follows. For elements  $(T^1, x^1)$  and  $(T^2, x^2)$  in  $Q_\varepsilon(\sigma, \phi)$ , we write  $(T^1, x^1) \leq (T^2, x^2)$  when  $T^1 \leq T^2$  and the restriction of  $x^2$  to the interval  $(-\infty, \sigma + T^1]$  is equal to  $x^1$ . First, we show that there is a maximal element.  $Q_\varepsilon(\sigma, \phi)$  is non-empty by Lemma 4. Let  $E = \{(T^\lambda, x^\lambda); \lambda \in A\}$  be any totally ordered set in  $Q_\varepsilon(\sigma, \phi)$ . Set  $J = \sup \{T^\lambda; \lambda \in A\}$ . If  $(T^\lambda, x^\lambda) \leq (T^\mu, x^\mu)$  for  $\lambda, \mu \in A$ , we see that

$$|x^\lambda(\sigma + T^\lambda) - x^\mu(\sigma + T^\mu)| = |x^\mu(\sigma + T^\lambda) - x^\mu(\sigma + T^\mu)| \leq 2r|T^\lambda - T^\mu|$$

by the condition (II). Hence  $\lim_{T^\lambda \rightarrow J} x^\lambda(\sigma + T^\lambda) = p$  exists, and  $p \in \Omega_{\sigma+J}$  by Lemma 2. Define  $x^*(t)$  by

$$x^*(t) = \begin{cases} x^\lambda(t), & t \leq \sigma + T^\lambda, \lambda \in A, \\ p, & t = \sigma + J. \end{cases}$$

Then  $(J, x^*)$  is in  $Q_\varepsilon(\sigma, \phi)$  and is the supremum of  $E$ . Therefore there is a maximal element  $(T, x)$  in  $Q_\varepsilon(\sigma, \phi)$  by Zorn's lemma.

Next, we prove that  $T = \alpha$  for the maximal element  $(T, x)$  obtained above. Suppose that  $T < \alpha$ . By Lemma 3, there is a continuous mapping  $g_1: [\sigma, \sigma + T] \rightarrow R^n$  such that  $d(x(t), \Omega_t) = |x(t) - g_1(t)|$  and  $g_1(t) \in \Omega_t$  for all  $t \in [\sigma, \sigma + T]$ . Let  $\xi: (-\infty, \sigma + T] \rightarrow R^n$  be a function such that  $\xi_\sigma = \phi$  and  $\xi(t) = g_1(t)$  on  $[\sigma, \sigma + T]$ . Then  $\xi_t \in B$  for all  $t \in [\sigma, \sigma + T]$  by the hypothesis (B1). Recall that  $|x(t) - \xi(t)| = |x(t) - g_1(t)| < \eta(\varepsilon, W)L^{-1}$  on  $[\sigma, \sigma + T]$  by (I). Since  $x(t)$  satisfies (I) and (II), it follows from the hypothesis (B2) and (6), (8) that

$$\begin{aligned} |\xi_{\sigma+T} - \phi| &\leq |\xi_{\sigma+T} - \tilde{\phi}_{\sigma+T}| + |\tilde{\phi}_{\sigma+T} - \phi| \\ &\leq L \sup_{-T \leq \theta \leq 0} |\xi(\sigma + T + \theta) - \phi(0)| + |\tilde{\phi}_{\sigma+T} - \phi| \\ &\leq L \sup_{-T \leq \theta \leq 0} \{|\xi(\sigma + T + \theta) - x(\sigma + T + \theta)| + |x(\sigma + T + \theta) - \phi(0)|\} \\ &\quad + |\tilde{\phi}_{\sigma+T} - \phi| \\ &\leq L\{\eta(\varepsilon, W)L^{-1} + 2rT\} + |\tilde{\phi}_{\sigma+T} - \phi| \leq 3Lr\alpha + |\tilde{\phi}_{\sigma+T} - \phi| < \delta. \end{aligned}$$

Therefore we have

$$(13) \quad |f(\sigma + T, \xi_{\sigma+T})| \leq r.$$

Since  $\xi_{\sigma+T}(t - \sigma - T) \in \Omega_t$  for all  $t \leq \sigma + T$  and  $\xi(\sigma + T) = x(\sigma + T)$ , we have

$$(14) \quad \lim_{h \rightarrow 0^+} d(x(\sigma + T) + hf(\sigma + T, \xi_{\sigma+T}), \Omega_{\sigma+T+h})/h = 0$$

by the condition (ii). Again by Lemma 3, there is a continuous function  $g_2(t): [\sigma + T, \infty) \rightarrow R^n$  such that

$$\begin{aligned} d(x(\sigma + T) + hf(\sigma + T, \xi_{\sigma+T}), \Omega_{\sigma+T+h}) \\ = |x(\sigma + T) + hf(\sigma + T, \xi_{\sigma+T}) - g_2(\sigma + T + h)| \end{aligned}$$

and  $g_2(\sigma + T + h) \in \Omega_{\sigma+T+h}$  for all  $h \geq 0$ . Then by (14), there is a  $\delta_1$  with  $0 < \delta_1 \leq \varepsilon$  such that

$$(15) \quad |f(\sigma + T, \xi_{\sigma+T}) - \{g_2(\sigma + T + h) - x(\sigma + T)\}/h| \leq \varepsilon$$

for all  $h \in (0, \delta_1]$ .

Let  $S$  be a constant such that  $0 < S < \alpha - T$  and  $S \leq \delta_1$ , and define a function  $y$  by

$$y(t) = \begin{cases} x(t), & t \leq \sigma + T, \\ x(\sigma + T) + \{(g_2(\sigma + T + S) - x(\sigma + T))/S\}(t - \sigma - T), & \sigma + T \leq t \leq \sigma + T + S. \end{cases}$$

We show that  $(T + S, y)$  belongs to  $Q_\varepsilon(\sigma, \phi)$  if  $S$  is sufficiently small. Since  $y(t) = x(t)$  for  $t \leq \sigma + T$ , it is sufficient to consider the case  $t \geq \sigma + T$ . By (13) and (15) and as in the proof of Lemma 4, we can find a  $\delta_2$  with  $0 < \delta_2 \leq \delta_1$  such that  $y(t)$  satisfies (I), (II) and (IV) for  $S \leq \delta_2$ .

To show (III), define another function  $z$  by  $z(t) = \xi(t)$  on  $(-\infty, \sigma + T]$  and  $z(t) = y(t)$  on  $[\sigma + T, \sigma + T + S]$ , where  $S = \delta_2$ . Then  $y_\sigma = z_\sigma = \phi$  and  $\sup \{|z(t) - y(t)|; t \in [\sigma, \sigma + T + S]\} = \sup \{|x(t) - \xi(t)|; t \in [\sigma, \sigma + T]\} < \eta(\varepsilon, W)L^{-1}$ , and hence

$$|y_t - z_t| \leq L \sup_{-(t-\sigma) \leq \theta \leq 0} |y(t + \theta) - z(t + \theta)| < L\eta(\varepsilon, W)L^{-1} = \eta(\varepsilon, W)$$

for all  $t \in [\sigma, \sigma + T + S]$  by the hypothesis (B2).  $(t, y_t)$  belongs to the compact set  $W$  for all  $t \in [\sigma, \sigma + T + S]$  since  $y(t)$  satisfies (II) on  $[\sigma, \sigma + T + S]$ . Thus we have

$$(16) \quad |f(t, y_t) - f(t, z_t)| < \varepsilon \quad \text{on } [\sigma, \sigma + T + S]$$

by (7). Since  $z(t)$  is  $2r$ -Lipschitzian in  $t \in [\sigma + T, \sigma + T + S]$ , we see that  $|z_t - \xi_{\sigma+T}| = |z_t - z_{\sigma+T}|$  is small if  $t - (\sigma + T) > 0$  is sufficiently small by the hypotheses (B1) and (B2). Therefore the continuity of  $f$  implies that there is a  $\delta_3$  with  $0 < \delta_3 \leq \delta_2$  such that

$$(17) \quad |f(t, z_t) - f(\sigma + T, \xi_{\sigma+T})| < \varepsilon$$

for all  $t \in [\sigma + T, \sigma + T + S]$  if  $S \leq \delta_3$ . Let  $S = \delta_3$ . Then it follows from (15), (16) and (17) that

$$\begin{aligned} |\dot{y}(t) - f(t, y_t)| &\leq |\{g_2(\sigma + T + S) - x(\sigma + T)\}/S - f(\sigma + T, \xi_{\sigma+T})| \\ &\quad + |f(\sigma + T, \xi_{\sigma+T}) - f(t, z_t)| + |f(t, z_t) - f(t, y_t)| \\ &\leq 3\epsilon \end{aligned}$$

for all  $t \in [\sigma + T, \sigma + T + S]$ .

Consequently, we obtain an element  $(T + S, y)$  in  $Q_\epsilon(\sigma, \phi)$  such that  $(T, x) \leq (T + S, y)$  and  $(T, x) \neq (T + S, y)$ , which contradicts the maximality of  $(T, x)$ . Thus  $T$  should be equal to  $\alpha$ , and we are done.

**4. The proof of the theorem.** It is easily proved that (i) implies (ii), and so we prove the converse.

Let  $\{\epsilon_k\}$  be a sequence such that  $\epsilon_k > 0$  and  $\epsilon_k \rightarrow 0$  as  $k \rightarrow \infty$ . Let  $(\sigma, \phi) \in R \times B$  be such that  $\phi(t - \sigma) \in \Omega_t$  for all  $t \leq \sigma$ . By Lemma 5, there exists an  $\alpha > 0$  such that the set  $Q_{\epsilon_k}(\sigma, \phi)$  contains an element  $(\alpha, x^k)$  for each  $k$ . Since the sequence of the functions  $\{x^k(t)\}$  is uniformly bounded and equicontinuous on  $[\sigma, \sigma + \alpha]$ , we may assume that the sequence converges uniformly to a continuous function  $x(t)$  on  $[\sigma, \sigma + \alpha]$  as  $k \rightarrow \infty$ . Let  $x(t) = \phi(t - \sigma)$  for  $t \leq \sigma$ . Then  $x_t$  belongs to  $B$  for all  $t \in [\sigma, \sigma + \alpha]$  by the hypothesis (B1). Also,  $|x_t^k - x_t| \rightarrow 0$  as  $k \rightarrow \infty$  for all  $t \in [\sigma, \sigma + \alpha]$  by the hypothesis (B2). Since  $(t, x_t^k)$  belongs to the compact set  $W$  for all  $t \in [\sigma, \sigma + \alpha]$  by (II), we have  $|f(t, x_t^k)| \leq C$  for all  $t \in [\sigma, \sigma + \alpha]$  and all  $k$ , where  $C$  is a constant. Hence, applying Lebesgue's dominant convergence theorem, we see by (II) and (III) that

$$\begin{aligned} x(t) &= \lim_{k \rightarrow \infty} x^k(t) = \lim_{k \rightarrow \infty} \left\{ \phi(0) + \int_\sigma^t \dot{x}^k(s) ds \right\} \\ &= \phi(0) + \lim_{k \rightarrow \infty} \left\{ \int_\sigma^t f(s, x_s^k) ds + \int_\sigma^t [\dot{x}^k(s) - f(s, x_s^k)] ds \right\} \\ &= \phi(0) + \int_\sigma^t f(s, x_s) ds \end{aligned}$$

for all  $t \in [\sigma, \sigma + \alpha]$ . Thus  $x(t)$  is a solution of (5) through  $(\sigma, \phi)$ .

By (IV), for each  $t \in [\sigma, \sigma + \alpha]$  and  $k$ , there is a point  $s^k \in [\sigma, \sigma + \alpha]$  such that  $|t - s^k| \leq \epsilon_k$  and  $(s^k, x^k(s^k)) \in \Omega$ . Then, by (II), we have  $|x(t) - x^k(s^k)| \leq |x(t) - x^k(t)| + |x^k(t) - x^k(s^k)| \leq |x(t) - x^k(t)| + 2r\epsilon_k$ , which implies  $\lim_{k \rightarrow \infty} (s^k, x^k(s^k)) = (t, x(t))$ , and hence  $(t, x(t)) \in \Omega$  by Lemma 2. Consequently,  $x(t)$  is the solution of (5) through  $(\sigma, \phi)$  such that  $(t, x(t)) \in \Omega$  on  $(-\infty, \sigma + \alpha]$ .

Now let  $Q(\sigma, \phi, \Omega)$  be the set defined by

$$Q(\sigma, \phi, \Omega) = \{(T, y) \in Q(\sigma, \phi); (t, y(t)) \in \Omega \text{ on } (-\infty, \sigma + T)\}.$$

Then  $Q(\sigma, \phi, \Omega)$  is nonempty because  $(\alpha, x) \in Q(\sigma, \phi, \Omega)$ . Introducing the same partial order in  $Q(\sigma, \phi, \Omega)$  as in  $Q(\sigma, \phi)$ , we obtain a maximal element  $(T, y)$  in  $Q(\sigma, \phi, \Omega)$  by Zorn's lemma. We show that the  $(T, y)$  is also a maximal element in  $Q(\sigma, \phi)$ . Otherwise,  $y$  can be extended up to  $t = \sigma + T$ , and then  $(t, y(t)) \in \Omega$  for all  $t \leq \sigma + T$  by Lemma 2. Clearly,  $y_{\sigma+T}$  belongs to  $B$  by the hypothesis (B1). Therefore, by applying the condition (ii) to  $(\sigma + T, y_{\sigma+T})$  and by the same argument as above we obtain an element  $(\alpha', z)$  in  $Q(\sigma + T, y_{\sigma+T}, \Omega)$ . Then  $(T + \alpha', z)$  is in  $Q(\sigma, \phi, \Omega)$ ,  $(T + \alpha', z) \geq (T, y)$  and  $(T + \alpha', z) \neq (T, y)$ . This contradicts the maximality of  $(T, y)$  in  $Q(\sigma, \phi, \Omega)$ . Thus  $(T, y)$  is in  $Q(\sigma, \phi, \Omega)$  and is the maximal element in  $Q(\sigma, \phi)$ , that is,  $y$  is the solution of (5) through  $(\sigma, \phi)$  defined on its right maximal interval of existence  $(-\infty, \sigma + T)$  and satisfying  $(t, y(t)) \in \Omega$  there.

## REFERENCES

- [1] H. BREZIS, On a characterization of flow-invariant sets, *Comm. Pure Appl. Math.* 23 (1970), 261-263.
- [2] M. G. CRANDALL, A generalization of Peano's existence theorem and flow invariance, *Proc. Amer. Math. Soc.* 36 (1972), 151-155.
- [3] J. K. HALE, Dynamical systems and stability, *J. Math. Anal. Appl.* 26 (1969), 39-59.
- [4] J. K. HALE AND J. KATO, Phase space for retarded differential equations with infinite delay, *Funkcial. Ekvac.* 21 (1978), 11-41.
- [5] P. HARTMAN, On invariant sets and on a theorem of Wazewski, *Proc. Amer. Math. Soc.* 32 (1972), 511-520.
- [6] T. KAMINOGO, Kneser's property and boundary value problem for some retarded functional differential equations, *Tohoku. Math. J.* 30 (1978), 471-486.
- [7] V. LAKSHMIKANTHAM, S. LEELA AND V. MOAURO, Existence and uniqueness of solutions of delay differential equations on a closed subset of a Banach spaces, *Nonlinear Analysis* 2 (1978), 311-327.
- [8] S. LEELA AND V. MOAURO, Existence of solutions in a closed set for delay differential equations in Banach spaces, *Nonlinear Analysis* 2 (1978), 47-58.
- [9] R. H. MARTIN, Approximation and existence of solutions to ordinary differential equations in Banach spaces, *Funkcial. Ekvac.* 16 (1973), 195-211.
- [10] G. SEIFERT, Positively invariant closed sets for systems of delay differential equations, *J. Differential Equations* 22 (1976), 292-304.
- [11] J. A. YORKE, Differential inequalities and non-Lipschitz scalar functions, *Math. System Theory* 4 (1970), 140-153.

MATHEMATICAL INSTITUTE  
TÔHOKU UNIVERSITY  
SENDAI, 980 JAPAN