

THE CHARACTERIZATION OF RIEMANNIAN METRIC ARISING FROM PHASE TRANSITION PROBLEMS

MAO-SHENG CHANG, SHU-CHENG LEE AND CHIEN-CHANG YEN

(Received February 26, 2008, revised February 12, 2009)

Abstract. We present one property of the Riemannian metric which is derived from the positive power of potential functions. Then this property is applied to the study of the Γ -convergence of energy functionals which are associated with the Euler-Lagrange p -Laplacian equation.

1. Introduction. The singular perturbation nonconvex functionals,

$$(1) \quad E_\varepsilon(v) = \int_{\Omega} \left[\frac{1}{\varepsilon} W(v(x)) + \varepsilon |\nabla v(x)|^2 \right] dx$$

with a nonnegative double-wells potential W , were studied by Modica and Mortola [22] in 1977. In 1987, Modica [21] applied the Γ -convergence theory to solve the minimal interface problem in the Van der Waals-Cahn-Hilliard theory of phase transitions. Suppose that $W(v) = 0$ if and only if $v \in \{a, b\}$. It was shown in [21, 22, 24] that the Γ -limit of $\{E_\varepsilon\}_{\varepsilon>0}$ is of the form

$$E_0(v) = \begin{cases} \beta \text{Per}_{\Omega}(A) & \text{if } v \in T, \\ +\infty & \text{otherwise,} \end{cases}$$

where $T = \{v \in \text{BV}(\Omega); v = \chi_A a + (1 - \chi_A)b, A \subset \Omega \text{ with } 0 < |A| < |\Omega|\}$ and

$$(2) \quad \beta = 2 \int_a^b W^{1/2}(s) ds.$$

The notation $\text{BV}(\Omega)$ and $\text{Per}_{\Omega}(A)$ stand for the space of functions of bounded variation and the perimeter of A in Ω , respectively. Let $p > 1$ and q the exponent conjugate to p , that is, $1/p + 1/q = 1$, and let K be a positive integer in this whole article. Let a_i ($i = 1, \dots, K$) be given points in \mathbf{R}^N and $\mathcal{A} = \{a_i; i = 1, \dots, K\}$. We denote by $C_{\text{pw}}^1(I)$ the set of piecewise C^1 curves $f: I \subset \mathbf{R} \rightarrow \mathbf{R}^N$, and we define

$$T_{F,L} \equiv \{g \in C_{\text{pw}}^1([-L, L]) \mid g(-L) = a_i, \text{ and } g(L) = a_j \text{ for some } i \neq j\},$$

$$(3) \quad T_F \equiv \bigcup_{L>0} T_{F,L},$$

2000 *Mathematics Subject Classification.* Primary 49J45.

Key words and phrases. Riemannian metric, Γ -convergence, functions of bounded variations.

and

$$(4) \quad S_F \equiv \left\{ \int_{-L}^L \left(\frac{W(g(s))}{q} + \frac{|g'(s)|^p}{p} \right) ds; \quad g \in T_F \right\}.$$

Barroso and Fonseca [5], Fonseca and Tartar [16] generalize the result of Γ -convergence to the vector-valued case for which $v : \Omega \rightarrow \mathbf{R}^N$, $\Omega \subset \mathbf{R}^n$ and the equation (2) is replaced by $\beta = \inf S_F$ for $p = 2$ and $K = 2$. Baldo [4] in 1990 established the Γ -limit of $\{F_\varepsilon\}_{\varepsilon>0}$ for multiple-well potential. Suppose that $W(v) = 0$ if and only if $v \in \{a_1, \dots, a_K\}$. The Γ -limit of $\{E_\varepsilon\}_{\varepsilon>0}$ is essentially of the form

$$E_0(v) = \begin{cases} \sum_{i,j=1}^K d(a_i, a_j) H^{n-1}(\partial^* S_i \cap \partial^* S_j \cap \Omega) & \text{if } v \in T, \\ +\infty & \text{otherwise,} \end{cases}$$

where $T = \{v \in \text{BV}(\Omega; \mathbf{R}^N); v = \sum_{i=1}^K \chi_{S_i} a_i, \text{ with } S_i \subset \Omega \text{ and } |\Omega - \bigcup_{i=1}^K S_i| = 0, \text{ and } S_i \cap S_j = \emptyset, i \neq j\}$, $d(a_i, a_j) = \inf S(a_i, a_j)$ and

$$S(a_i, a_j) = \left\{ \int_{-L}^L [W(g(s)) + |g'(s)|^2] ds; g \in C_{\text{pw}}^1([-L, L]), g(-L) = a_i, g(L) = a_j \right\}.$$

For more generalization of this case, see Ambrosio [1] and Sternberg [24].

The development of this type of Γ -convergence for gradient vector fields, i.e., $\{E_\varepsilon(\nabla v)\}_{\varepsilon>0}$ was studied by Jin and Kohn [18], and Conti, Fonseca and Leoni [9]. There have been many works on the related subjects, for instance, Fonseca and Mantegazza [15], Fonseca [14], Ambrosio, De Lellis and Mantegazza [2], Aviles and Giga [3], and DeSimone, Kohn, Müller and Otto [11].

In this paper, we consider a family of functionals $\{F_\varepsilon\}_{\varepsilon>0}$ given by

$$(5) \quad F_\varepsilon(v) = \begin{cases} \int_{\Omega} \left[\frac{1}{\varepsilon} \frac{W(v(x))}{q} + \varepsilon^{p-1} \frac{|\nabla v(x)|^p}{p} \right] dx & \text{if } v \in \mathcal{W}^{1,p}(\Omega; \mathbf{R}^N), \\ +\infty & \text{otherwise.} \end{cases}$$

The Euler-Lagrange equation of these functionals is the following p -Laplacian equation,

$$(6) \quad \varepsilon^p \Delta_p v - \frac{W'(v)}{q} = 0 \quad \text{in } \Omega,$$

where $\Delta_p v = \text{div}(|\nabla v|^{p-2} \nabla v)$. The functionals defined by Equation (1) could be regarded as a special case of the functionals defined by Equation (5) for $p = 2$. The minimizer and Γ -convergence for scalar case with a special potential $W(v) = v^2(1-v)^2/4$ has been studied in [8]. It follows from the important role of constants $\inf S(a_i, a_j)$ in the study of Γ -convergence of functional $\{E_\varepsilon\}_{\varepsilon>0}$ that give us a motivation to find properties of the Riemannian metric which is derived from the positive power of potential functions. In this paper, we present one property of this Riemannian metric and apply it to the study of the Γ -convergence of functionals $\{F_\varepsilon\}_{\varepsilon>0}$ for which the potential is of double-well type.

Define $\mathcal{A} = \{a_1, \dots, a_K\}$ and let us introduce two hypothesis on the potential W below.

(H1) The potential $W \in \mathcal{W}_{\text{loc}}^{1,\infty}(\mathbf{R}^N; \mathbf{R})$ is a nonnegative function such that $W(x) = 0$ if and only if $x = a_i$ for some $i = 1, \dots, K$.

(H2) Let $d = \min\{\text{dist}(a_i, a_j); i \neq j, \text{ and } i, j = 1, \dots, K\}$. There exist $\alpha, \delta \in \mathbf{R}$ with $0 < \alpha \leq 1$ and $0 < \delta < d/2$ such that, if $x \in \mathcal{A}$ and $y \in \mathbf{R}^N \setminus \mathcal{A}$ with $|y - x| \leq \delta$, then $\alpha|y - x|^p \leq W(y) \leq \alpha^{-1}|y - x|^p$.

Furthermore, let us define two sets of functions and two sets of real numbers. They are

$$(7) \quad T_\infty \equiv \{g \in C_{\text{pw}}^1(\mathbf{R}); g(-\infty) = a_i, \text{ and } g(\infty) = a_j \text{ for some } i \neq j\},$$

$$(8) \quad T_\star \equiv \{g \in C_{\text{pw}}^1([-1, 1]); g(-1) = a_i, \text{ and } g(1) = a_j \text{ for some } i \neq j\},$$

and

$$(9) \quad S_\infty \equiv \left\{ \int_{-\infty}^{\infty} \left(\frac{W(g(s))}{q} + \frac{|g'(s)|^p}{p} \right) ds; g \in T_\infty \right\},$$

$$(10) \quad S_\star \equiv \left\{ \int_{-1}^1 \left(W(g(s))^{1/q} \cdot |g'(s)| \right) ds; g \in T_\star \right\}.$$

The following theorem states one property of our Riemannian metric. We will prove it in the next section.

THEOREM 1. *Let S_∞ , S_F and S_\star be the set of values defined by (9), (4) and (10), respectively. Then*

$$\inf S_\infty = \inf S_F = \inf S_\star.$$

We will apply Theorem 1 to the Γ -convergence theory for the case of two wells, that is $K = 2$, with $a_1 = a$ and $a_2 = b$. Let Ω be an open bounded strongly Lipschitz domain of \mathbf{R}^n . The sets of states are defined as

$$(11) \quad A = \{x \in \Omega; v(x) = a\}, \quad B = \{x \in \Omega; v(x) = b\}.$$

The role of $F_0(v)$ is played by the limit problem

$$(12) \quad F_0(v) = \begin{cases} (\inf S_\star) \text{Per}_\Omega(A) & \text{if } v \in \text{BV}(\Omega; \mathbf{R}^N) \text{ with } v(x) \in \{a, b\} \text{ a.e.} \\ +\infty & \text{otherwise.} \end{cases}$$

We will show that the functional F_0 is the $\Gamma(L^1(\Omega; \mathbf{R}^N))$ -limit of $\{F_\varepsilon\}_{\varepsilon>0}$ (denoted by $F_0 = \Gamma(L^1(\Omega; \mathbf{R}^N))$ -limit $\{F_\varepsilon\}_{\varepsilon>0}$). This means by definition that for each $v \in L^1(\Omega; \mathbf{R}^N)$,

- (i) if v_ε converges to v in $L^1(\Omega; \mathbf{R}^N)$, then $F_0(v) \leq \liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(v_\varepsilon)$,
- (ii) there exists a family $\{v_\varepsilon\}_{\varepsilon>0}$ in $L^1(\Omega; \mathbf{R}^N)$ such that v_ε converges to v in $L^1(\Omega; \mathbf{R}^N)$ and $\limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(v_\varepsilon) \leq F_0(v)$.

The Γ -convergence theorem below will be proved in Section 3.

THEOREM 2. *In addition to the assumptions (H1) and (H2), assume that*

(H3) *there exist $\alpha, C > 0$ such that $W(v) \geq \alpha|v|$ whenever $|v| > C$.*

Then the functional F_0 defined in (12) is the $\Gamma(L^1(\Omega; \mathbf{R}^N))$ -limit of $\{F_\varepsilon\}_{\varepsilon>0}$ defined by (5).

Let a, b and m be vectors in \mathbf{R}^N with $m = \theta a + (1 - \theta)b$ for some $\theta \in (0, 1)$. Define the spaces V_m and V as follows.

$$V_m \equiv \left\{ u \in W^{1,p}(\Omega; \mathbf{R}^N); \int_{\Omega} u(x) dx = |\Omega| m \right\},$$

$$V \equiv \left\{ u \in \text{BV}(\Omega; \mathbf{R}^N); \quad u(x) \in \{a, b\} \text{ a.e. and } \int_{\Omega} u(x) dx = |\Omega| m \right\}.$$

We would like to give some remarks on our works for more applications and further studies.

Remark 1. For each $\varepsilon > 0$, the existence of a minimizer u_{ε} of F_{ε} on V_m can be obtained by virtue of the direct methods in the calculus of variations.

Remark 2. Using the definition of Γ -convergence, we observe that if (i) $F_0 = \Gamma(L^1(\Omega; \mathbf{R}^N)\text{-}\liminf\{F_{\varepsilon}\}_{\varepsilon>0}$, (ii) u_{ε} is a minimizer of F_{ε} on V_m for each $\varepsilon > 0$, and (iii) u_{ε} converges to u_0 in $L^1(\Omega; \mathbf{R}^N)$ as ε goes to 0^+ , then

$$\lim_{\varepsilon \rightarrow 0^+} F_{\varepsilon}(u_{\varepsilon}) = F_0(u_0)$$

and u_0 is a minimizer of F_0 on V .

Remark 3. Another consequence is that Γ -convergence is stable under continuous perturbations: If $\{F_{\varepsilon}\}_{\varepsilon>0}$ Γ -converges to F_0 in $L^1(\Omega; \mathbf{R}^N)$ and the functional G is continuous with respect to the $L^1(\Omega; \mathbf{R}^N)$ -topology, then $\{F_{\varepsilon} + G\}_{\varepsilon>0}$ Γ -converges to $F_0 + G$ in $L^1(\Omega; \mathbf{R}^N)$.

Remark 4. Theorem 1 is useful for characterizing the behavior of the minimizers of Γ -limit F_0 of $\{F_{\varepsilon}\}_{\varepsilon>0}$ with multiple potential wells. We anticipate further developments in this direction in the future.

2. Riemannian metric. We divide the proof of Theorem 1 into Lemmas 3, 4 and 5 in this section.

LEMMA 3. *Let S_{∞} and S_F be the sets defined in Equations (9) and (4), respectively. Then $\inf S_{\infty} = \inf S_F$.*

PROOF. We first prove that $\inf S_{\infty} \leq \inf S_F$ by showing

$$(13) \quad S_F \subset S_{\infty}.$$

For any $\alpha \in S_F$, Equation (4) implies that there exists a function $g_L \in T_F$ such that

$$\alpha = \int_{-L}^L \left(\frac{W(g_L(s))}{q} + \frac{|g'_L(s)|^p}{p} \right) ds.$$

The function $g_L \in T_F$ can be extended to $g_{\infty} \in T_{\infty}$ by defining

$$(14) \quad g_{\infty}(s) \equiv \begin{cases} g_L(-L) & \text{if } s < -L, \\ g_L(s) & \text{if } -L \leq s \leq L, \\ g_L(L) & \text{if } s > L. \end{cases}$$

The function g_∞ belongs to T_∞ by Equations (3) and (7). Since $W(g_\infty(s)) = 0$ and $|g'_\infty(s)| = 0$ for $s \in \mathbf{R} \setminus [-L, L]$, we evaluate

$$\alpha = \int_{-L}^L \left(\frac{W(g_L(s))}{q} + \frac{|g'_L(s)|^p}{p} \right) ds = \int_{-\infty}^{\infty} \left(\frac{W(g_\infty(s))}{q} + \frac{|g'_\infty(s)|^p}{p} \right) ds.$$

It follows that α is in S_∞ , and we get (13).

Next we show $\inf S_F \leq \inf S_\infty$. Now, we assume that α is in S_∞ . By Equation (9), there is a piecewise C^1 curve $g : \mathbf{R} \rightarrow \mathbf{R}^N$ with $g(-\infty) = a = a_i$ and $g(\infty) = b = a_j$ for some $i \neq j$ such that

$$(15) \quad \alpha = \int_{-\infty}^{\infty} \left(\frac{W(g(s))}{q} + \frac{|g'(s)|^p}{p} \right) ds < +\infty.$$

Condition (H2) implies that there exist $0 < \delta < d/2$ and $L > 0$ such that

$$(16) \quad |g(s) - a| < \delta \quad \text{for all } s \in (-\infty, -L) \quad \text{and} \quad |g(s) - b| < \delta \quad \text{for all } s \in (L, \infty).$$

Without loss of generality, we may assume $g(s) \in \mathbf{R}^N \setminus \mathcal{A}$ for $s \in \mathbf{R} \setminus [-L, L]$. By (H2) again, we see that

$$(17) \quad \alpha |g(s) - a|^p \leq W(g(s)) \leq \frac{1}{\alpha} |g(s) - a|^p \quad \text{for } s \in (-\infty, -L).$$

Using Equations (15) and (17) gives us

$$\int_{-\infty}^{-L} |g(s) - a|^p ds \leq \frac{1}{\alpha} \int_{-\infty}^{-L} W(g(s)) ds < +\infty.$$

It follows that

$$(18) \quad \int_{-\infty}^L |g(s) - a|^p ds < +\infty$$

by $\int_{-L}^L |g(s) - a|^p ds < +\infty$. The identity $|(g(s) - a)'| = |g'(s)|$ and Equation (17) imply

$$(19) \quad \int_{-\infty}^L |(g(s) - a)'|^p ds < +\infty.$$

From Equations (18) and (19), the function $g - a$ is in $\mathcal{W}^{1,p}((-\infty, L); \mathbf{R}^N)$. Similarly, the function $g - b$ is in $\mathcal{W}^{1,p}((-L, \infty); \mathbf{R}^N)$. It is easy to observe that for all $\tilde{L} > 0$, $g - a$ is in $\mathcal{W}^{1,p}((-\infty, \tilde{L}); \mathbf{R}^N)$ and $g - b$ is in $\mathcal{W}^{1,p}((-\tilde{L}, \infty); \mathbf{R}^N)$.

By the theorem on partition of unity [23], there exists a function $\gamma : \mathbf{R} \rightarrow \mathbf{R}$ in C_c^∞ with $0 \leq \gamma \leq 1$, $\gamma(s) = 1$ on $[-1, 1]$ and $\gamma(s) = 0$ on $\mathbf{R} \setminus (-2, 2)$. Let us define a sequence of functions $\{g_k\}_{k=1}^\infty$ by

$$(20) \quad g_k(s) \equiv \begin{cases} \gamma(s/k)g(s) + (1 - \gamma(s/k))a & \text{if } s < 0, \\ g(s) & \text{if } s = 0, \\ \gamma(s/k)g(s) + (1 - \gamma(s/k))b & \text{if } s > 0. \end{cases}$$

Note that for $k > L$, we have

$$(21) \quad g_k(s) = \begin{cases} a & \text{if } s < -2k, \\ g(s) & \text{if } -k \leq s \leq k, \\ b & \text{if } s > 2k, \end{cases}$$

and hence $g_k \in T_\infty$. We claim that

$$(22) \quad \lim_{k \rightarrow \infty} \int_{-2k}^{2k} \left(\frac{W(g_k(s))}{q} + \frac{|g'_k(s)|^p}{p} \right) ds = \int_{-\infty}^{\infty} \left(\frac{W(g(s))}{q} + \frac{|g'(s)|^p}{p} \right) ds.$$

We divide the proof into the two parts

$$\lim_{k \rightarrow \infty} \int_{-2k}^{2k} W(g_k(s)) ds = \int_{-\infty}^{\infty} W(g(s)) ds \quad \text{and} \quad \lim_{k \rightarrow \infty} \int_{-2k}^{2k} |g'_k(s)|^p ds = \int_{-\infty}^{\infty} |g'(s)|^p ds.$$

It is equivalent to show that

$$\lim_{k \rightarrow \infty} \int_{-\infty}^{\infty} W(g_k(s)) ds = \int_{-\infty}^{\infty} W(g(s)) ds \quad \text{and} \quad \lim_{k \rightarrow \infty} \int_{-\infty}^{\infty} |g'_k(s)|^p ds = \int_{-\infty}^{\infty} |g'(s)|^p ds,$$

since $W(g_k(s)) = 0$ and $g'_k(s) = 0$ for all $|s| \geq 2k$ and $k \in \mathcal{N}$ by Equation (21).

Let $L > 0$ be given. By Equation (20), we have $g_k(s) - a = \gamma(s/k)(g(s) - a)$ for $s < 0$. By the inequality $0 \leq \gamma \leq 1$ and Condition (H2), we have

$$(23) \quad W(g_k(s)) \leq \frac{1}{\alpha} |g_k(s) - a|^p \leq \frac{1}{\alpha} |g(s) - a|^p \quad \text{on } (-\infty, -L),$$

for each $k \in \mathcal{N}$. By the continuity of the potential W and the convergence $g_k \rightarrow g$ on \mathbf{R} as $k \rightarrow \infty$, we have

$$(24) \quad W \circ g_k \rightarrow W \circ g \quad \text{on } \mathbf{R} \text{ as } k \rightarrow \infty.$$

Applying the Lebesgue Dominated Convergence Theorem and by Equations (24) and (23), we get

$$(25) \quad \lim_{k \rightarrow \infty} \int_{-\infty}^{-L} W(g_k(s)) ds = \int_{-\infty}^{-L} W(g(s)) ds.$$

A similar argument gives us

$$(26) \quad \lim_{k \rightarrow \infty} \int_L^{\infty} W(g_k(s)) ds = \int_L^{\infty} W(g(s)) ds.$$

If $k > L$, by Equation (21), we get $g_k(s) = g(s)$ for $-L \leq s \leq L$. Hence we have

$$(27) \quad \int_{-L}^L W(g_k(s)) ds = \int_{-L}^L W(g(s)) ds.$$

By combining Equations (25), (26) and (27), we get the equality

$$\lim_{k \rightarrow \infty} \int_{-\infty}^{\infty} W(g_k(s)) ds = \int_{-\infty}^{\infty} W(g(s)) ds.$$

Since $\gamma \in C_c^\infty(\mathbf{R})$, there exists a constant M such that $|\gamma'(s)| \leq M$ for all $s \in \mathbf{R}$. A simple calculation shows that, if $s < 0$, then $g'_k(s) = (1/k)\gamma'(s/k)(g(s) - a)^T + \gamma(s/k)g'(s)$, and $|g'_k(s)| \leq M|g(s) - a| + |g'(s)|$ on $(-\infty, 0)$. This also implies $|g'_k(s)|^p \leq 2^p \max\{M^p|g(s) -$

$a|P, |g'(s)|^P\}$ on $(-\infty, 0)$. It is easy to show g'_k converges to g' on \mathbf{R} as k go to ∞ . By the fact that $g - a$ is in $\mathcal{W}^{1,p}(\mathbf{R}; \mathbf{R}^N)$ and by Lebesgue's Dominated Convergence Theorem, we conclude that

$$\lim_{k \rightarrow \infty} \int_{-\infty}^0 |g'_k(s)|^p ds = \int_{-\infty}^0 |g'(s)|^p ds.$$

Similary, $\lim_{k \rightarrow \infty} \int_0^\infty |g'_k(s)|^p ds = \int_0^\infty |g'(s)|^p ds$. Hence

$$\lim_{k \rightarrow \infty} \int_{-\infty}^\infty |g'_k(s)|^p ds = \int_{-\infty}^\infty |g'(s)|^p ds.$$

Thus Equation (22) holds and the proof of Lemma 3 is completed. \square

LEMMA 4. *Let S_F and S_\star be the sets defined in Equations (4) and (10), respectively. Then $\inf S_\star \leq \inf S_F$.*

PROOF. Let $\alpha \in S_F$. By Equation (4), there exists a function $g_L \in T_F$ such that

$$\alpha = \int_{-L}^L \left(\frac{W(g_L(s))}{q} + \frac{|g'_L(s)|^p}{p} \right) ds.$$

Using the technique of rescaling set, we define $g_{L,1}(s) = g_L(sL)$ for $s \in [-1, 1]$. Applying the change of variables and Young's inequality, we have

$$\int_{-1}^1 (W(g_{L,1}(s)))^{1/q} \cdot |g'_{L,1}(s)| ds = \int_{-L}^L W(g_L(s))^{1/q} \cdot |g'_L(s)| ds,$$

and

$$\int_{-L}^L W(g_L(s))^{1/q} \cdot |g'_L(s)| ds \leq \int_{-L}^L \left(\frac{W(g_L(s))}{q} + \frac{|g'_L(s)|^p}{p} \right) ds.$$

Define $\tilde{\alpha} \equiv \int_{-1}^1 (W(g_{L,1}(s)))^{1/q} \cdot |g'_{L,1}(s)| ds$. We have shown that for each $\alpha \in S_F$, there exists a number $\tilde{\alpha} \in S_\star$ such that $\tilde{\alpha} \leq \alpha$. Therefore, $\inf S_\star \leq \inf S_F$. \square

LEMMA 5. *Let S_∞ and S_\star be the sets defined in Equations (9) and (10), respectively. Then $\inf S_\infty \leq \inf S_\star$.*

PROOF. Let $\alpha \in S_\star$. Then there is a function $g_\star \in T_\star$ such that

$$(28) \quad \alpha = \int_{-1}^1 (W(g_\star(s)))^{1/q} \cdot |g'_\star(s)| ds.$$

We change the variable to the arclength $\tau(s)$, $\tau(s) \equiv \int_{-1}^s |g'_\star(t)| dt$ for $-1 \leq s \leq 1$ and denote $L = \tau(1)$. Since $\tau'(s) = |g'_\star(s)| > 0$ on $(-1, 1)$, the function τ is monotone increasing, and its inverse function $\tau^{-1} : [0, L] \rightarrow [-1, 1]$ exists. Furthermore, $(\tau^{-1})'(s) = (|g'_\star(\tau^{-1}(s))|)^{-1} > 0$ on $(0, L)$. We define $g_L(s) = g_\star(\tau^{-1}(s))$ for $0 \leq s \leq L$. Then

$$(29) \quad g'_L(s) = \frac{g'_\star(\tau^{-1}(s))}{|g'_\star(\tau^{-1}(s))|}, \quad \text{and} \quad |g'_L(s)| = 1 \quad \text{for } s \in [0, L].$$

Using the variable of arclength to rewrite Equation (28) as

$$\alpha = \int_0^L (W(g_\star(\tau^{-1}(s)))^{1/q} \cdot |g'_\star(\tau^{-1}(s))| |(\tau^{-1})'(s)| ds.$$

Since $|g'_\star(\tau^{-1}(s))| |\tau^{-1}(s)| = |g'_L(s)|$, it further implies

$$(30) \quad \alpha = \int_0^L (W(g_L(s)))^{1/q} \cdot |g'_L(s)| ds.$$

For the function g_L , if there is a function $g_\infty \in T_\infty$ such that

$$\int_{-\infty}^{\infty} \left(\frac{W(g_\infty(s))}{q} + \frac{|g'_\infty(s)|^p}{p} \right) ds = \int_0^L (W(g_L(s)))^{1/q} \cdot |g'_L(s)| ds = \alpha,$$

then $\alpha \in S_\infty$. Moreover, $S_\star \subset S_\infty$ and this will show that Lemma 5 holds. The connection between the two functions g_L and g_∞ is contained in the two equalities

$$(31) \quad \int_{-\infty}^{\infty} \left(\frac{W(g_\infty(s))}{q} + \frac{|g'_\infty(s)|^p}{p} \right) ds = \int_{-\infty}^{\infty} (W(g_\infty(s)))^{1/q} \cdot |g'_\infty(s)| ds$$

and

$$(32) \quad \int_{-\infty}^{\infty} (W(g_\infty(s)))^{1/q} \cdot |g'_\infty(s)| ds = \int_0^L (W(g_L(s)))^{1/q} \cdot |g'_L(s)| ds.$$

The condition for Equation (31) means that the equality holds in Young's inequality, that is,

$$(33) \quad |g'_\infty(t)| = W(g_\infty(t))^{1/p} \quad \text{for all } t \in \mathbf{R}.$$

The condition for Equation (32) is that there is a monotone increasing function $h \in C_{pw}^1(\mathbf{R})$ such that

$$(34) \quad g_\infty(t) = g_L(h(t)) \quad \text{for all } t \in \mathbf{R},$$

$$(35) \quad \lim_{t \rightarrow -\infty} h(t) = 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} h(t) = L.$$

Since h is increasing, Equation (33) is reduced to

$$(36) \quad h'(t) = |h'(t)| = W(g_L(h(t)))^{1/p} \quad \text{for all } t \in \mathbf{R}$$

by Equations (29) and (34). The above discussion reduces the problem to the existence of an increasing function h which satisfies the ordinary differential equation $y' = W(g_L(y))^{1/p}$ with initial value $y(0) = L/2$. Since W is in $\mathcal{W}_{loc}^{1,\infty}(\mathbf{R}^N; \mathbf{R})$ and W is locally Lipschitz on \mathbf{R}^N , it implies that $W(g_L(y))^{1/p}$ is in $\mathcal{W}_{loc}^{1,\infty}((0, L))$. Thanks to Picard's Existence and Uniqueness Theorem [17], we obtain the existence of a unique continuously differentiable function $y : I \rightarrow \mathbf{R}$ with $0 \in I$ satisfying the differential equation and the initial condition, and also by the Continuation Theorem [17], we extend the interval I into the maximum interval of existence, say (t_0, t_1) with $I \subset (t_0, t_1)$, and

$$(37) \quad \lim_{t \rightarrow t_0^+} y(t) = 0, \quad \lim_{t \rightarrow t_1^-} y(t) = L.$$

Define

$$(38) \quad h(t) \equiv \begin{cases} 0 & \text{if } t \leq t_0, \\ y(t) & \text{if } t_0 < t < t_1, \\ L & \text{if } t \geq t_1. \end{cases}$$

The function $h \in C_{\text{pw}}^1(\mathbf{R})$ is as desired and it also satisfies Equations (36) and (35). By Equations (34), (37) and (38), it follows that g_∞ is in $C_{\text{pw}}^1(\mathbf{R})$,

$$g_\infty(-\infty) \equiv \lim_{t \rightarrow -\infty} g_\infty(t) = g_L(0) = g_\star(-1) \in A_i,$$

and

$$g_\infty(\infty) \equiv \lim_{t \rightarrow \infty} g_\infty(t) = g_L(L) = g_\star(1) \in A_j.$$

Hence g_∞ is in T_∞ and satisfies Equations (31) and (32). As a consequence, S_\star is included in S_∞ and Lemma 5 is proved. \square

3. Γ -convergence theory. We first investigate the structure of a limit function.

LEMMA 6. *If the functionals F_ε are defined by Equation (5) with $\liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(v_\varepsilon) < \infty$ and functions v_ε converge to v in $L^1(\Omega; \mathbf{R}^N)$, then the values of the function $v(x)$ belong to $\{a, b\}$ a.e. Furthermore, the function $v(x)$ is equal to $\chi_A(x)a + \chi_B(x)b$, where the set A and B are defined by (11).*

PROOF. Since v_ε converges to v in $L^1(\Omega; \mathbf{R}^N)$, there is a subsequence $\{v_{\varepsilon_j}\}$ which converges to v a.e. It follows that $W \circ v_{\varepsilon_j}$ converges to $W \circ v$ a.e. By Fatou's lemma, we have

$$0 \leq \int_{\Omega} W(v(x)) dx \leq \liminf_{j \rightarrow \infty} \int_{\Omega} W(v_{\varepsilon_j}(x)) dx \leq \liminf_{j \rightarrow \infty} \int_{\Omega} \varepsilon_j q F_{\varepsilon_j}(v_{\varepsilon_j}) = 0.$$

The last equality holds due to the assumption. The integral of the nonnegative potential function W over Ω vanishes, and hence $W(v(x)) = 0$. That implies $v(x)$ is in $\{a, b\}$ a.e., and this lemma is proved. \square

It follows from (H3) that we can choose $r_1 > |a - b|/2 = r_0$, and $C = \max\{W^{1/q}(v); |v - (a + b)/2| \leq r_1\}$, such that

$$\int_{r_0}^{r_1} \left[\inf_{|v - (a+b)/2| = r} W^{1/q}(v) \right] dr > \frac{1}{2} \inf S_\star.$$

Define $W^\star(v) \equiv \min\{W^{1/q}(v), C\}$, $T(\xi) = \{\gamma \in C_{\text{pw}}^1([-1, 1]); \gamma(-1) = a, \gamma(1) = \xi\}$, and

$$(39) \quad \Phi(\xi) = \inf \left\{ \int_{-1}^1 W^\star(\gamma(s)) |\gamma'(s)| ds; \gamma \in T(\xi) \right\}.$$

The technique of truncation has been used to guarantee the Lipschitz continuity of the function Φ on \mathbf{R}^N . A similar argument in Fonseca and Tartar [16] establishes the properties of the function Φ in the following lemma.

LEMMA 7. Let Φ be the function defined by (39).

- (i) The function $\Phi : \mathbf{R}^N \rightarrow \mathbf{R}$ is Lipschitz continuous on \mathbf{R}^N .
- (ii) If $v \in \mathcal{W}^{1,p}(\Omega; \mathbf{R}^N)$, then $\Phi \circ v \in \mathcal{W}^{1,p}(\Omega; \mathbf{R}^N)$ and

$$|\nabla(\Phi \circ v)(x)| \leq W^{1/q}(v(x)) \cdot |\nabla v(x)| \quad \text{for a.e. } x \in \Omega.$$

- (iii) $\inf S_\star = \Phi(b)$.

Let us recall the following lemma.

LEMMA 8 (cf. [13, 20], Lower semicontinuous of variation measures). Suppose that $\Omega \subset \mathbf{R}^n$ is open. Suppose f_k is in $BV(\Omega)$ for each k in N and f_k converges to f in $L^1(\Omega)$. Then $\int_\Omega |\nabla f(x)| dx \leq \liminf_{k \rightarrow \infty} \int_\Omega |\nabla f_k(x)| dx$.

Now, we will show the theorem below.

THEOREM 9. Let functionals F_ε and F_0 be defined by Equations (5) and (12), respectively. If v_ε converges to v in $L^1(\Omega; \mathbf{R}^N)$, then $F_0(v) \leq \liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(v_\varepsilon)$.

PROOF. We first claim that the limit function v is a BV-function on Ω . If v_ε converges to v in $L^1(\Omega; \mathbf{R}^N)$, then $\liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(v_\varepsilon)$ is finite and $\Phi \circ v_\varepsilon$ converges to $\Phi \circ v$ in $L^1(\Omega)$. It implies that v_ε is in $\mathcal{W}^{1,p}(\Omega; \mathbf{R}^N)$. By the fact that if u is in $\mathcal{W}^{1,p}(\Omega; \mathbf{R}^N)$ then $\Phi \circ u$ is in $\mathcal{W}^{1,p}(\Omega; \mathbf{R}^N)$ and also in $BV(\Omega)$, we have $\Phi \circ v_\varepsilon \in BV(\Omega)$. Applying Lemma 8 to $\Phi \circ v_\varepsilon$ and $\Phi \circ v$, we have

$$\int_\Omega |\nabla(\Phi \circ v)(x)| dx \leq \liminf_{\varepsilon \rightarrow 0^+} \int_\Omega |\nabla(\Phi \circ v_\varepsilon)(x)| dx.$$

Using condition (ii) in Lemma 7 and Young's inequality to (5), we get

$$\int_\Omega |\nabla(\Phi \circ v)(x)| dx \leq \liminf_{\varepsilon \rightarrow 0^+} \int_\Omega W^{1/q}(v_\varepsilon(x)) |\nabla v_\varepsilon(x)| dx$$

and

$$\int_\Omega W^{1/q}(v_\varepsilon(x)) |\nabla v_\varepsilon(x)| dx \leq F_\varepsilon(v_\varepsilon).$$

It follows that

$$(40) \quad \int_\Omega |\nabla(\Phi \circ v)(x)| dx \leq \liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(v_\varepsilon),$$

and $\Phi \circ v$ is in $BV(\Omega)$. By (39), $\Phi(a)$ is zero and $\Phi(b)$ is equal to $\inf S_\star$. Hence we obtain $(\Phi \circ v)(x) = \Phi(v(x)) = \Phi(\chi_A a + \chi_B b) = \Phi(b) \chi_B$. Thus, χ_B is in $BV(\Omega)$ and $\chi_A = \chi_{\Omega \setminus B}$ is in $BV(\Omega)$. Furthermore, the function v is in $BV(\Omega)$ since $v = \chi_A a + \chi_B b$. We have proved the claim. The next step is to show the identity

$$(41) \quad \int_\Omega |\nabla(\Phi \circ v)(x)| dx = F_0(v).$$

Since $(\Phi \circ v)(x) = \Phi(b) \chi_B$, we have

$$\int_\Omega |\nabla(\Phi \circ v)(x)| dx = \int_\Omega \Phi(b) |\nabla \chi_B| dx = \Phi(b) \text{Per}_\Omega(B) = \Phi(b) \cdot \text{Per}_\Omega(A).$$

From the equality $\Phi(b) = \inf S_\star$ again, it follows that

$$(42) \quad \int_{\Omega} |\nabla(\Phi \circ v)(x)| dx = (\inf S_\star) \text{Per}_{\Omega}(A).$$

Combining Lemma 7 with Equation (12), we get Equation (41). Theorem 9 follows from Equations (41), (42) and (40). \square

THEOREM 10. *Let $v \in \mathcal{W}^{1,p}(\Omega; \mathbf{R}^N)$. Then there exists a family $\{v_\varepsilon\}_{\varepsilon>0}$ in $L^1(\Omega; \mathbf{R}^N)$ which converges to v in $L^1(\Omega; \mathbf{R}^N)$ and $\limsup_{\varepsilon \rightarrow 0^+} F_\varepsilon(v_\varepsilon)$ is less than or equal to $F_0(v)$.*

Before the proof of Theorem 10, let us recall two lemmas.

LEMMA 11 ([4, 21]). *Let $\Omega \subset \mathbf{R}^n$ be open, and A be a polygonal domain in \mathbf{R}^n with the compact boundary ∂A and $H^{n-1}(\partial A \cap \partial \Omega) = 0$. Then the following two statements hold:*

(i) *There exists a constant $\eta > 0$ such that the function $h(x)$ defined by*

$$h(x) \equiv \begin{cases} -d(x, \partial A) & \text{if } x \in A, \\ d(x, \partial A) & \text{if } x \notin A \end{cases}$$

is Lipschitz continuous on $D_\eta \equiv \{x \in \mathbf{R}^n; |h(x)| < \eta\}$ and $|\nabla h(x)| = 1$ a.e. on D_η .

(ii) *If $S_t \equiv \{x \in \mathbf{R}^n; h(x) = t\}$, then $\lim_{t \rightarrow 0} H^{n-1}(S_t \cap \Omega) = H^{n-1}(\partial A \cap \Omega)$.*

LEMMA 12 (cf. [13, 20], Coarea formula). *Suppose that $f : \mathbf{R} \rightarrow \mathbf{R}$ is a measurable function and that $h : \mathbf{R}^n \rightarrow \mathbf{R}$ is a Lipschitz function. Then*

$$\int_{\Omega} f(h(x)) |\nabla h(x)| dx = \int_{-\infty}^{\infty} f(t) H^{n-1}(\{x \in \Omega; h(x) = t\}) dt$$

for each measurable subset Ω of \mathbf{R}^n . In particular, we have

$$\int_{\Omega} |\nabla h(x)| dx = \int_{-\infty}^{\infty} H^{n-1}(\{x \in \Omega; h(x) = t\}) dt$$

for each measurable subset Ω of \mathbf{R}^n .

PROOF OF THEOREM 10. The strategy for the proof is to observe the form of the limit function v , $v = \chi_A a + (1 - \chi_A)b$, then to construct a sequence of functions $v_\varepsilon^g \in L^1(\Omega; \mathbf{R}^N)$ which converges to v in $L^1(\Omega; \mathbf{R}^N)$ starting from a given function g in T_F . Moreover, this sequence of functions satisfies

$$(43) \quad \lim_{\varepsilon \rightarrow 0} F_\varepsilon(v_\varepsilon^g) = \text{Per}_{\Omega}(A) \int_{-L}^L \left(\frac{W(g(s))}{q} + \frac{|g'(s)|^p}{p} \right) ds.$$

Finally, we apply the above result to construct a sequence which we want.

By the assumption, $v \in \mathcal{W}^{1,p}(\Omega; \mathbf{R}^N)$ and Lemma 6, v must be of the form

$$(44) \quad v = \chi_A a + (1 - \chi_A)b \quad \text{a.e. in } \Omega.$$

Without loss of generality, we assume that $\partial A \cap \Omega \in C^2$ and $H^{n-1}(\partial A \cap \partial \Omega) = 0$. We define the function $h : \Omega \rightarrow \mathbf{R}$ by

$$(45) \quad h(x) \equiv \begin{cases} -d(x, \partial A) & \text{if } x \in A, \\ d(x, \partial A) & \text{if } x \notin A. \end{cases}$$

Now, let $g \in T_F$, where T_F was defined by (3). That is, there is $L > 0$ such that $g(-L) = a$ and $g(L) = b$. For this function g , we define a sequence of functions $v_\varepsilon^g : \Omega \rightarrow \mathbf{R}^N$ by

$$(46) \quad v_\varepsilon^g(x) = \begin{cases} a & \text{if } h(x) \leq -\varepsilon L, \\ g_\varepsilon(h(x)) & \text{if } -\varepsilon L < h(x) < \varepsilon L, \\ b & \text{if } \varepsilon L \leq h(x), \end{cases}$$

where $g_\varepsilon(s) = g(s/\varepsilon)$ for all s in $[-L, L]$. For the simplification of notation, we define $f : \mathbf{R} \rightarrow \mathbf{R}$ by

$$f(t) = \begin{cases} |g_\varepsilon(t) - a| & \text{if } -\varepsilon L < t \leq 0, \\ |g_\varepsilon(t) - b| & \text{if } 0 < t < \varepsilon L, \\ 0 & \text{otherwise.} \end{cases}$$

Then $|v_\varepsilon^g - v|$ is equal to $f \circ h$ on Ω . By Lemma 11, we have

$$\int_\Omega |v_\varepsilon^g - v|(x) dx = \int_\Omega f \circ h(x) dx = \int_\Omega f(h(x)) |\nabla h|(x) dx,$$

and by Lemma 12, we have

$$\begin{aligned} \int_\Omega |v_\varepsilon^g - v|(x) dx &= \int_{-\infty}^{\infty} f(t) \cdot H^{n-1}(\{x \in \Omega; h(x) = t\}) dt \\ &= \int_{-\varepsilon L}^0 |g_\varepsilon(t) - a| \cdot H^{n-1}(\{x \in \Omega; h(x) = t\}) dt \\ &\quad + \int_0^{\varepsilon L} |g_\varepsilon(t) - b| \cdot H^{n-1}(\{x \in \Omega; h(x) = t\}) dt. \end{aligned}$$

Using the change of variables for the last term in the last equality, we get the equality

$$\begin{aligned} &\int_0^{\varepsilon L} |g_\varepsilon(t) - b| \cdot H^{n-1}(\{x \in \Omega; h(x) = t\}) dt \\ &= \varepsilon \int_0^L |g(t) - b| \cdot H^{n-1}(\{x \in \Omega; h(x) = \varepsilon t\}) dt. \end{aligned}$$

By Lemma 11 again, $\lim_{\tau \rightarrow 0} H^{n-1}(\{x \in \Omega; h(x) = \tau\}) = H^{n-1}(\partial A \cap \Omega) = \text{Per}_\Omega(A)$. Taking $\tau = \varepsilon t$ for each $\varepsilon > 0$ with $|\varepsilon L| < \delta$, we have $H^{n-1}(\{x \in \Omega; h(x) = \varepsilon t\}) < \text{Per}_\Omega(A) + 1$ for all $-L \leq t \leq L$. Therefore, we have

$$\varepsilon \int_0^L |g(t) - b| \cdot H^{n-1}(\{x \in \Omega; h(x) = \varepsilon t\}) dt \leq \varepsilon L (\text{Per}_\Omega(A) + 1) \|g - b\|_\infty.$$

Similary, we have

$$\varepsilon \int_{-L}^0 |g(t) - a| \cdot H^{n-1}(\{x \in \Omega; h(x) = \varepsilon t\}) dt \leq \varepsilon L (\text{Per}_\Omega(A) + 1) \|g - a\|_\infty.$$

The above argument gives us the estimate

$$\int_{\Omega} |v_{\varepsilon}^g - v|(x) dx \leq \varepsilon L(\text{Per}_{\Omega}(A) + 1)[\|g - a\|_{\infty} + \|g - b\|_{\infty}],$$

provided $\varepsilon > 0$ small enough. Furthermore, we know $\lim_{\varepsilon \rightarrow 0} \int_{\Omega} |v_{\varepsilon}^g - v|(x) dx$ is equal to zero, and the convergence of v_{ε}^g to v in $L^1(\Omega; \mathbf{R}^N)$ is proved. Next step is to show Equation (43).

By Equation (5), we evaluate

$$\begin{aligned} F_{\varepsilon}(v_{\varepsilon}^g) &= \int_{\Omega} \left[\frac{1}{\varepsilon} \frac{W(v_{\varepsilon}^g(x))}{q} + \varepsilon^{p-1} \frac{|\nabla v_{\varepsilon}^g|^p(x)}{p} \right] dx \\ &= \int_{\Omega} \left[\frac{1}{\varepsilon} \frac{W(g_{\varepsilon}(h(x)))}{q} + \varepsilon^{p-1} \frac{|\nabla v_{\varepsilon}^g(h(x))|^p(x) |\nabla h(x)|}{p} \right] dx \\ &= \int_{\Omega} \left[\frac{1}{\varepsilon} \frac{W(g_{\varepsilon}(h(x)))}{q} + \varepsilon^{p-1} \frac{|\nabla v_{\varepsilon}^g(h(x))|^p(x)}{p} \right] |\nabla h(x)| dx. \end{aligned}$$

The last equality follows from $|\nabla h| = 1$ a.e. (cf. Lemma 11) whenever $\varepsilon > 0$ small enough. By using Lemmas 11, 12 and technique of change of variables to the last term, we rewrite $F_{\varepsilon}(v_{\varepsilon}^g)$ as

$$F_{\varepsilon}(v_{\varepsilon}^g) = \int_{-L}^L \left(\frac{W \circ g}{q} + \frac{|g'|^p}{p} \right)(t) H^{n-1}(\{x \in \Omega; h(x) = \varepsilon t\}) dt.$$

Therefore,

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} F_{\varepsilon}(v_{\varepsilon}^g) &= \lim_{\varepsilon \rightarrow 0} \int_{-L}^L \left(\frac{W \circ g}{q} + \frac{|g'|^p}{p} \right)(t) H^{n-1}(\{x \in \Omega; h(x) = \varepsilon t\}) dt \\ &= \int_{-L}^L \left(\frac{W \circ g}{q} + \frac{|g'|^p}{p} \right)(t) \lim_{\varepsilon \rightarrow 0} H^{n-1}(\{x \in \Omega; h(x) = \varepsilon t\}) dt \\ &= \text{Per}_{\Omega}(A) \int_{-L}^L \left(\frac{W(g(t))}{q} + \frac{|g'(t)|^p}{p} \right) dt. \end{aligned}$$

Let $\tilde{\varepsilon}$ be a positive number. Let us denote $\tilde{S}_F = \text{Per}_{\Omega}(A)S_F$, $\tilde{S}_{\infty} = \text{Per}_{\Omega}(A)S_{\infty}$ and $\tilde{S}_{\star} = \text{Per}_{\Omega}(A)S_{\star}$. By Theorem 1, we have $\inf \tilde{S}_F = \inf \tilde{S}_{\infty} = \inf \tilde{S}_{\star} = F_0(v)$. Since $F_0(v) = \inf \tilde{S}_F$, it follows that there exists a number $\eta = \eta(\tilde{\varepsilon})$ in \tilde{S}_F and g_{η} in T_F such that $\eta < F_0(v) + \tilde{\varepsilon}$ and

$$\eta = \text{Per}_{\Omega}(A) \int_{-L}^L \left[\frac{W(g_{\eta}(s))}{q} + \frac{|g'_{\eta}(s)|^p}{p} \right] ds \quad \text{for some } L > 0.$$

Furthermore, for $\eta \in \tilde{S}_F$, there exists a family $\{v_{\tau}^{g_{\eta}}\}_{\tau > 0}$ in $L^1(\Omega; \mathbf{R}^N)$ such that $v_{\tau}^{g_{\eta}}$ converges to v in $L^1(\Omega; \mathbf{R}^N)$ and $F_{\tau}(v_{\tau}^{g_{\eta}})$ converges to η as τ goes to 0^+ . We get that for each $\tilde{\varepsilon} > 0$, there is a $\eta < F_0(v) + \tilde{\varepsilon}$ and $v_{\tau_{\eta}}^{g_{\eta}}$ in $L^1(\Omega; \mathbf{R}^N)$ such that $\|v_{\tau_{\eta}}^{g_{\eta}} - v\|_{L^1(\Omega; \mathbf{R}^N)} < \tilde{\varepsilon}$ and $F_{\tau_{\eta}}(v_{\tau_{\eta}}^{g_{\eta}}) < F_0(v) + \tilde{\varepsilon}$. Let $v_{\tilde{\varepsilon}} = v_{\tau_{\eta}}^{g_{\eta}}$. The sequence $\{v_{\tilde{\varepsilon}}\}_{\tilde{\varepsilon} > 0}$ has the desired property and Theorem 10 is proved. \square

Acknowledgment. The authors are grateful to the referees for carefully reading the manuscript and giving us many valuable advices and comments.

REFERENCES

- [1] L. AMBROSIO, Metric space valued functions of bounded variation, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* 17 (1990), 439–478.
- [2] L. AMBROSIO, C. DE LELLIS AND C. MANTEGAZZA, Line energies for gradient vector fields in the plane, *Calc. Var. Partial Differential Equations* 9 (1999), 327–355.
- [3] P. AVILES AND Y. GIGA, On lower semicontinuity of a defect energy obtained by a singular limit of the Ginzburg-Landau type energy for gradient fields, *Proc. Roy. Soc. Edinburgh Sect. A* 129 (1999), 1–17.
- [4] S. BALDO, Minimal interface criterion for phase transitions in mixtures of Cahn-Hilliard fluids, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 7 (1990), 67–90.
- [5] A. C. BARROSO AND I. FONSECA, Anisotropic singular perturbations—the vectorial case, *Proc. Roy. Soc. Edinburgh Sect. A* 124 (1994), 527–571.
- [6] G. BOUCHITTÉ, Singular perturbations of variational problems arising from a two-phase transition model, *Appl. Math. Optim.* 21 (1990), 289–314.
- [7] A. BRAIDES, Γ -convergence for beginners, *Oxford Lecture Ser. Math. Appl.* 22, Oxford University Press, Oxford, 2002.
- [8] M. S. CHANG, S. C. LEE AND C. C. YEN, Minimizers and Γ -convergence of energy functionals derived from p -Laplacian equation, to appear in *Taiwanese Journal of Mathematics*, 2009.
- [9] S. CONTI, I. FONSECA AND G. LEONI, A Γ -convergence result for the two-gradient theory of phase transitions, *Comm. Pure Appl. Math.* 55 (2002), 857–936.
- [10] G. DAL MASO, An introduction to Γ -convergence, Birkhäuser Boston Inc., Boston, MA, 1993.
- [11] A. DESIMONE, R. V. KOHN, S. MÜLLER AND F. OTTO, A compactness result in the gradient theory of phase transitions, *Proc. Roy. Soc. Edinburgh Sect. A* 131 (2001), 833–844.
- [12] L. C. EVANS, *Partial differential equations*, American Mathematical Society, Providence, 1998.
- [13] L. C. EVANS AND R. F. GARIEPY, *Measure theory and fine properties of functions*, CRC Press, Boca Raton, FL, 1992.
- [14] I. FONSECA, Phase transitions of elastic solid materials, *Arch. Ration. Mech. Anal.* 107 (1989), 195–223.
- [15] I. FONSECA AND C. MANTEGAZZA, Second order singular perturbation models for phase transitions, *SIAM J. Math. Anal.* 31 (2000), 1121–1143.
- [16] I. FONSECA AND L. TARTAR, The gradient theory of phase transitions for systems with two potential wells, *Proc. Roy. Soc. Edinburgh Sect. A* 111 (1989), 89–102.
- [17] P. HARTMAN, *Ordinary differential equations*, John Wiley & Sons, Inc., New York-London-Sydney, 1964.
- [18] W. JIN AND R. V. KOHN, Singular perturbation and the energy of folds, *J. Nonlinear Sci.* 10 (2000), 355–390.
- [19] R. V. KOHN AND P. STERNBERG, Local minimizers and singular perturbations, *Proc. Roy. Soc. Edinburgh Sect. A* 111 (1989), 69–84.
- [20] F. H. LIN AND X. P. YANG, *Geometric measure theory—an introduction*, Science Press, Beijing, International Press, Boston, MA, 2002.
- [21] L. MODICA, The gradient theory of phase transitions and the minimal interface criterion, *Arch. Ration. Mech. Anal.* 98 (1987), 123–142.
- [22] L. MODICA AND S. MORTOLA, Un esempio di Γ -convergenza, *Boll. Un. Mat. Ital. B* (5) 14 (1977), 285–299.
- [23] J. R. MUNKRES, *Analysis on manifolds*, Addison-Wesley Publishing Company, Advanced Book Program, Redwood City, CA, 1991.
- [24] P. STERNBERG, The effect of a singular perturbation on non-convex variational problems, *Arch. Ration. Mech. Anal.* 101 (1988), 209–260.

DEPARTMENT OF MATHEMATICS
FU JEN CATHOLIC UNIVERSITY
TAIPEI
TAIWAN

E-mail addresses: mschang@math.fju.edu.tw
sclee@math.fju.edu.tw
yen@math.fju.edu.tw