

FEKETE-SZEGÖ PROBLEM FOR A CLASS DEFINED BY AN INTEGRAL OPERATOR

AKSHAYA KUMAR MISHRA AND PRIYABRAT GOCHHAYAT¹

Abstract

By making use of an integral operator due to Noor, a new subclass of analytic functions, denoted by $k - \mathcal{UCV}_n$ ($n \in \mathbf{N}_0 := \{0, 1, 2, \dots\}; 0 \leq k < \infty$); is introduced. For this class the Fekete-Szegö problem is completely settled. The results obtained here also give the Fekete-Szegö inequalities for the classes of k -uniformly convex functions and k -parabolic starlike functions.

1. Introduction and definitions

Let \mathcal{A} denote the family of functions analytic in the *open* unit disk

$$\mathcal{U} := \{z : z \in \mathbf{C}, |z| < 1\}$$

and let \mathcal{A}_0 be the class of functions f in \mathcal{A} satisfying the normalization condition $f(0) = f'(0) - 1 = 0$. Thus, the functions in \mathcal{A}_0 are given by the power series

$$(1.1) \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (z \in \mathcal{U}).$$

Let $\mathcal{D}^n : \mathcal{A}_0 \rightarrow \mathcal{A}_0$ be the operator defined by

$$\begin{aligned} \mathcal{D}^n f(z) &= \frac{z}{(1-z)^{n+1}} * f(z) \quad n > -1 \\ &= z + \sum_{k=2}^{\infty} \binom{k+n-1}{k-1} a_k z^k, \quad (f \in \mathcal{A}_0, n \in \mathbf{N}_0 := \{0, 1, 2, \dots\}), \end{aligned}$$

where '*' denotes the convolution or Hadamard product. We note that $\mathcal{D}^0 f = f$, $\mathcal{D}^1 f = zf'$ and $\mathcal{D}^n f = \frac{z(z^{n-1} f)^{(n)}}{n!}$. The operator $\mathcal{D}^n f$ is called the

2000 *Mathematics Subject Classification*: Primary 30C45.

Keywords and Phrases: Integral operator, k -uniformly convex function, elliptic integral, Fekete-Szegö problem, Gauss hypergeometric series.

¹(Corresponding author) The present research work of the second author was supported by Council of Scientific and Industrial Research, New Delhi, Government of India, under Senior Research Fellowship grant. Sanction letter no. 9/297/(73)/2008—EMR-I.

Received October 21, 2008; revised December 8, 2009.

Ruscheweyh derivative of n^{th} order of f . Analogous to $\mathcal{D}^n f$, Noor [13] defined an integral operator $\mathcal{I}_n : \mathcal{A}_0 \rightarrow \mathcal{A}_0$ as follows:

Write $f_n(z) = z/(1-z)^{n+1}$ and let f_n^\dagger be defined by the relation

$$(1.2) \quad f_n(z) * f_n^\dagger(z) = \frac{z}{(1-z)^2}.$$

For $f \in \mathcal{A}_0$ let

$$(1.3) \quad \mathcal{I}_n f(z) = f_n^\dagger(z) * f(z).$$

Note that $\mathcal{I}_0 f(z) = zf'(z)$ and $\mathcal{I}_1 f(z) = f(z)$. The operator $\mathcal{I}_n f$ defined by (1.3) is called n^{th} order Noor integral operator of f . For details see (cf. [13], [14], also see [15]).

For any real or complex numbers a, b, c other than $0, -1, -2, \dots$, the Gauss hypergeometric series is defined by

$$(1.4) \quad {}_2F_1(a, b; c; z) = 1 + \frac{ab}{c}z + \frac{a(a+1)b(b+1)}{c(c+1)} \frac{z^2}{2!} + \dots$$

We can write (1.3) as

$$(1.5) \quad \begin{aligned} \mathcal{I}_n f(z) &= [z {}_2F_1(2, 1; n+1; z)] * f(z) \\ &= \left[z + \sum_{k=2}^{\infty} \frac{k!}{(n+1) \cdots (n+k-1)} z^k \right] * f(z). \end{aligned}$$

Let \mathcal{S} , \mathcal{S}^* , \mathcal{CV} , \mathcal{UCV} and \mathcal{SP} denote, respectively the subclasses of \mathcal{A}_0 consisting of functions which are *univalent*, *starlike*, *convex* (cf. [4]), *uniformly convex* (cf. [6]) and *parabolic starlike* (cf. [16])) in \mathcal{U} . For fixed k ($0 \leq k < \infty$), the function $f \in \mathcal{A}_0$ is said to be in $k - \mathcal{UCV}$; the class of *k -uniformly convex functions* in \mathcal{U} , if the image of every circular arc γ contained in \mathcal{U} , with centre ξ where $|\xi| \leq k$, is a convex arc. This interesting unification of the concepts of *convex functions* and *uniformly convex functions* is due to [8]. The class $k - \mathcal{SP}$, consisting of *k -parabolic starlike functions*, is defined from $k - \mathcal{UCV}$ via the *Alexander's transform* (see [9]) i.e.

$$(1.6) \quad f \in k - \mathcal{UCV} \Leftrightarrow g \in k - \mathcal{SP}, \quad \text{where } g(z) = zf'(z) \quad (z \in \mathcal{U}).$$

The one variable characterization theorem (cf. [8]) of the the class $k - \mathcal{UCV}$ gives that $f \in k - \mathcal{UCV}$ (respectively $f \in k - \mathcal{SP}$) if and only if the values of

$$p(z) = 1 + \frac{zf''(z)}{f'(z)} \left(\text{respectively } \frac{zf'(z)}{f(z)} \right) \quad (z \in \mathcal{U})$$

lie in the conic region Ω_k in the w -plane, where

$$\Omega_k := \{w = u + iv \in \mathbf{C} : u^2 > k^2(u-1)^2 + k^2v^2; 0 \leq k < \infty\}.$$

For details of the geometric description of Ω_k see [8, 9]. Taking cue from (1.6), we now define a new subclass of analytic functions by using the linear operator \mathcal{I}_n .

DEFINITION 1. The function $f \in \mathcal{A}_0$ is said to be in the class $k - \mathcal{UCV}_n$ ($0 \leq k < \infty; n \in \mathbf{N}_0$) if $\mathcal{I}_n f \in k - \mathcal{SP}$. Or equivalently

$$\Re \left(\frac{z(\mathcal{I}_n f)'(z)}{(\mathcal{I}_n f)(z)} \right) > k \left| \frac{z(\mathcal{I}_n f)'(z)}{(\mathcal{I}_n f)(z)} - 1 \right| \quad (z \in \mathcal{U}).$$

Note that the class $k - \mathcal{UCV}_n$ unifies many subclasses of \mathcal{A}_0 related to \mathcal{S} . In particular, for $k = 0, n = 0 : 0 - \mathcal{UCV}_0 := \mathcal{CV}$, the class of univalent convex functions (see [4]); for $k = 1, n = 0 : 1 - \mathcal{UCV}_0 := \mathcal{UCV}$, the class of uniformly convex functions (see [6]); for $k = 0, n = 1 : 0 - \mathcal{UCV}_1 := \mathcal{S}^*$, the class of univalent starlike functions (see [4]); for $k = 1, n = 1 : 1 - \mathcal{UCV}_1 := \mathcal{SP}$, the class of parabolic starlike functions (see [16]); for $k \neq 0, n = 0 : k - \mathcal{UCV}_0 := k - \mathcal{UCV}$ (see [8]); for $k \neq 0, n = 1 : k - \mathcal{UCV}_1 := k - \mathcal{SP}$ (see [8]).

It is well known (cf. [4]) that for $f \in \mathcal{S}$ and given by (1.1), the sharp inequality $|a_3 - a_2^2| \leq 1$ holds. Fekete-Szegö, in their seminal work [5], found sharp bounds for $|\mu a_2^2 - a_3|$, ($\mu \in \mathbf{R}, 0 \leq \mu < 1$), for $f \in \mathcal{S}$. Thus, for any family \mathcal{F} of functions in \mathcal{A}_0 , the problem of finding sharp estimates on the nonlinear functional $|\mu a_2^2 - a_3|$, ($\mu \in \mathbf{R}$ or $\mu \in \mathbf{C}$) is popularly known as the Fekete-Szegö problem for \mathcal{F} . Recently the present authors [12] settled this problem for the classes $k - \mathcal{UCV}$, $k - \mathcal{SP}$ and some related classes defined using fractional calculus. One may also see [7, 10, 11, 17, 18] for some interesting results on this topic. In the present paper the Fekete-Szegö problem for the class $k - \mathcal{UCV}_n$ ($0 \leq k < \infty; n \in \mathbf{N}_0$) is settled completely. For $k = 1, n = 0$ and $n = 0; n = 1$, the result on Fekete-Szegö inequalities obtained here include, respectively, earlier results of Ma and Minda [11] on \mathcal{UCV} and recent results of the authors on $k - \mathcal{UCV}$ and $k - \mathcal{SP}$ in [12]. For values of $n > 1$ our results provide new information.

In the present investigation we also need the following definitions and notations, for the presentation of our results.

The *Jacobi elliptic integral* (or normal elliptic integral) of first kind (cf. [1], [2], also see [19, p. 50]) is defined by

$$\mathcal{F}(\omega, t) = \int_0^\omega \frac{dx}{\sqrt{(1-x^2)(1-t^2x^2)}} \quad (0 < t < 1).$$

The function $\mathcal{F}(1, t) := \mathcal{K}(t)$ is called the complete elliptic integral of the first kind. Changing to the variable $t' = \sqrt{1-t^2}$, $t \in (0, 1)$, we write $\mathcal{K}'(t) := \mathcal{K}(t')$. It should be emphasized here that the symbol ' (prime) does not stand for derivative. The following properties of $\mathcal{K}(t)$ and $\mathcal{K}'(t)$ are well known (cf. [7]).

$$\lim_{t \rightarrow 0^+} \mathcal{K}(t) = \frac{\pi}{2} \quad \lim_{t \rightarrow 1^-} \mathcal{K}(t) = \infty.$$

Moreover the function

$$v(t) = \frac{\pi}{2} \frac{\mathcal{K}'(t)}{\mathcal{K}(t)}, \quad (t \in (0, 1))$$

strictly decreases from ∞ to 0 as t moves from 0 to 1. Therefore every positive number k can be expressed as

$$(1.7) \quad k = \cosh(v(t))$$

for some unique $t \in (0, 1)$.

Define the function \mathcal{G} on \mathcal{U} by

$$(1.8) \quad \mathcal{G}(z) = [z_2 F_1(n+1, 1; 2; z)] * \left\{ z \exp\left(\int_0^z \frac{q_k(\zeta) - 1}{\zeta} d\zeta\right) \right\}, \quad (z \in \mathcal{U}),$$

where q_k is the function defined in Lemma 1 (below) and write the function $\psi(z, \theta, \eta)$ in $k - \mathcal{UCV}_n$ by

$$(1.9) \quad \psi(z, \theta, \eta) = [z_2 F_1(n+1, 1; 2; z)] * z \exp\left(\int_0^z \left[q_k\left(\frac{e^{i\theta}\zeta(\zeta + \eta)}{1 + \eta\zeta}\right) - 1\right] \frac{d\zeta}{\zeta}\right) \quad (0 \leq \theta \leq 2\pi; 0 \leq \eta \leq 1).$$

Note that $\psi(z, 0, 1) = \mathcal{G}(z)$ defined by (1.8) and $\psi(z, \theta, 0)$ is an odd function.

2. Preliminary lemmas

We need the following results in our investigation:

LEMMA 1 [7]. *Let $k \in [0, \infty)$ be fixed and q_k be the Riemann map of \mathcal{U} on to Ω_k , satisfying $q_k(0) = 1$ and $q'_k(0) > 0$. If*

$$(2.1) \quad q_k(z) = 1 + Q_1 z + Q_2 z^2 + \dots, \quad (z \in \mathcal{U})$$

then

$$Q_1 = \begin{cases} \frac{2A^2}{1 - k^2}; & 0 < k < 1, \\ \frac{8}{\pi^2}; & k = 1, \\ \frac{\pi^2}{4(k^2 - 1)\mathcal{K}^2(t)\sqrt{t}(1+t)}; & k > 1, \end{cases}$$

$$Q_2 = \begin{cases} \frac{(A^2 + 2)}{3} Q_1; & 0 < k < 1, \\ \frac{2}{3} Q_1; & k = 1, \\ \frac{(4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2)}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} Q_1; & k > 1, \end{cases}$$

where

$$(2.2) \quad A = \frac{2}{\pi} \arccos k,$$

and $\mathcal{K}(t)$ is the complete elliptic integral of first kind.

LEMMA 2 [10]. *Let h be analytic, with $h(0) = 1$, $\Re(h(z)) > 0$ and given by the series*

$$(2.3) \quad h(z) = 1 + c_1 z + c_2 z^2 + \dots \quad (z \in \mathcal{U}).$$

Then

$$(2.4) \quad |c_n| \leq 2 \quad (n \in \mathbb{N}),$$

$$(2.5) \quad |c_2 - c_1^2| \leq 2 \quad \text{and} \quad \left| c_2 - \frac{1}{2} c_1^2 \right| \leq 2 - \frac{|c_1|^2}{2}.$$

3. Fekete-Szegö inequalities

The following calculations shall be used in each of the proofs of Theorems 1, 2 and 3.

By Definition 1 there exists a function $w \in \mathcal{A}$ satisfying the conditions of the Schwarz lemma such that

$$(3.1) \quad \frac{z(\mathcal{I}_n f(z))'}{(\mathcal{I}_n f(z))} = q_k(w(z)) \quad (z \in \mathcal{U}),$$

where q_k is the function defined as in Lemma 1.

Let the function p_1 be defined by

$$p_1(z) = \frac{1 + w(z)}{1 - w(z)} = 1 + c_1 z + c_2 z^2 + \dots, \quad (z \in \mathcal{U}),$$

so that $\Re(p_1(z)) > 0$. This gives

$$w(z) = \frac{c_1}{2} z + \frac{1}{2} \left(c_2 - \frac{c_1^2}{2} \right) z^2 + \dots.$$

Substituting this in the series (2.1) we get

$$\begin{aligned} (3.2) \quad q_k(w(z)) &= 1 + Q_1 \left\{ \frac{c_1}{2} z + \frac{1}{2} \left(c_2 - \frac{c_1^2}{2} \right) z^2 + \dots \right\} \\ &\quad + Q_2 \left\{ \frac{c_1}{2} z + \frac{1}{2} \left(c_2 - \frac{c_1^2}{2} \right) z^2 + \dots \right\}^2 + \dots \\ &= 1 + \frac{Q_1 c_1}{2} z + \left\{ \frac{1}{2} \left(c_2 - \frac{c_1^2}{2} \right) Q_1 + \frac{1}{4} c_1^2 Q_2 \right\} z^2 + \dots, \end{aligned}$$

Using the expansion (1.5) in (3.1) and equating coefficients we find that

$$(3.3) \quad a_2 = \frac{(n+1)Q_1 c_1}{4}$$

and

$$(3.4) \quad a_3 = \frac{(n+1)(n+2)}{12} \left[\frac{Q_1^2 c_1^2}{4} + \frac{Q_1}{2} \left(c_2 - \frac{c_1^2}{2} \right) + \frac{Q_2 c_1^2}{4} \right].$$

We have the following:

THEOREM 1. *Let the function f given by (1.1) be in the class $k - \mathcal{UCV}_n$ ($0 \leq k < 1$). Then*

$$(3.5) \quad |\mu a_2^2 - a_3| \leq \begin{cases} \frac{(n+1)(n+2)A^2}{6(1-k^2)} \left(\frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} - \frac{(2+A^2)}{3} - \frac{2A^2}{(1-k^2)} \right); & \mu \geq \alpha_1, \\ \frac{(n+1)(n+2)A^2}{6(1-k^2)}; & \alpha_2 \leq \mu \leq \alpha_1, \\ \frac{(n+1)(n+2)A^2}{6(1-k^2)} \left(\frac{2A^2}{(1-k^2)} + \frac{(2+A^2)}{3} - \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} \right); & \mu \leq \alpha_2, \end{cases}$$

where the constant A is given by (2.2),

$$(3.6) \quad \alpha_1 := \alpha_1(k) = \frac{(n+2)}{(n+1)} \left(\frac{(5+A^2)(1-k^2)}{18A^2} + \frac{1}{3} \right)$$

and $\alpha_2 := \alpha_2(k) = \frac{(n+2)}{(n+1)} \left(\frac{1}{3} - \frac{(1-A^2)(1-k^2)}{18A^2} \right).$

Each of the estimates in (3.5) is sharp.

Proof. Putting the value of Q_1 and Q_2 for $0 \leq k < 1$ from Lemma 1 in (3.3) and (3.4) we get

$$a_2 = \frac{(n+1)A^2}{2(1-k^2)} c_1$$

and

$$a_3 = \frac{(n+1)(n+2)A^2}{12(1-k^2)} \left\{ c_2 - \frac{1}{6} \left((1-A^2) - \frac{6A^2}{1-k^2} \right) c_1^2 \right\}.$$

Therefore

$$\begin{aligned}
 |\mu a_2^2 - a_3| &= \left| \frac{(n+1)^2 A^4 c_1^2}{4(1-k^2)^2} \mu - \frac{(n+1)(n+2)A^2}{12(1-k^2)} \left\{ c_2 \right. \right. \\
 &\quad \left. \left. - \frac{1}{6} \left((1-A^2) - \frac{6A^2}{1-k^2} \right) c_1^2 \right\} \right| \\
 &= \left| \left\{ \frac{(n+1)^2 A^4}{4(1-k^2)^2} \mu + \frac{(n+1)(n+2)A^2}{72(1-k^2)} \left((1-A^2) \right. \right. \right. \\
 &\quad \left. \left. \left. - \frac{6A^2}{(1-k^2)} \right) c_1^2 - \frac{(n+1)(n+2)A^2}{12(1-k^2)} c_2 \right\} \right| \\
 &= \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left| \left\{ \frac{6(n+1)A^2 \mu}{(n+2)(1-k^2)} \right. \right. \\
 &\quad \left. \left. + \frac{1}{3} \left((1-A^2) - \frac{6A^2}{(1-k^2)} \right) c_1^2 - 2c_2 \right\} \right| \\
 (3.7) \quad &= \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left| \left\{ \frac{6(n+1)A^2 \mu}{(n+2)(1-k^2)} \right. \right. \\
 &\quad \left. \left. + \frac{1-A^2}{3} - \frac{2A^2}{(1-k^2)} - 2 \right\} c_1^2 + 2c_1^2 - 2c_2 \right| \\
 (3.8) \quad &\leq \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left\{ \left| \frac{6(n+1)A^2 \mu}{(n+2)(1-k^2)} \right. \right. \\
 &\quad \left. \left. - \frac{5+A^2}{3} - \frac{2A^2}{(1-k^2)} \right| |c_1^2| + 2|c_1^2 - c_2| \right\}.
 \end{aligned}$$

Now we note that

$$\begin{aligned}
 (3.9) \quad &\frac{6(n+1)A^2 \mu}{(n+2)(1-k^2)} - \frac{5+A^2}{3} - \frac{2A^2}{(1-k^2)} \geq 0 \\
 \text{provided } \mu &\geq \frac{(n+2)}{(n+1)} \left(\frac{(5+A^2)(1-k^2)}{18A^2} + \frac{1}{3} \right) = \alpha_1.
 \end{aligned}$$

Thus if $\mu \geq \alpha_1$, the expression inside the first modulus symbol on the right hand side of (3.8) is non negative. An application of Lemma 2 yields,

$$\begin{aligned}
 (3.10) \quad |\mu a_2^2 - a_3| &\leq \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left\{ \left(\frac{6(n+1)A^2 \mu}{(n+2)(1-k^2)} \right. \right. \\
 &\quad \left. \left. - \frac{5+A^2}{3} - \frac{2A^2}{(1-k^2)} \right) 4 + 4 \right\}
 \end{aligned}$$

$$\leq \frac{(n+1)(n+2)A^2}{6(1-k^2)} \left\{ \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} - \frac{2+A^2}{3} - \frac{2A^2}{(1-k^2)} \right\}.$$

which is the first part of assertion (3.5).

Rewriting (3.7)

$$(3.11) \quad |\mu a_2^2 - a_3| = \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left| -2c_2 - \left\{ \frac{2A^2}{(1-k^2)} - \frac{1-A^2}{3} - \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} \right\} c_1^2 \right| \\ = \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left| \left\{ \frac{2A^2}{(1-k^2)} - \frac{1-A^2}{3} - \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} \right\} c_1^2 + 2c_2 \right|.$$

Now if $\mu \leq \alpha_2$, where α_2 is given by (3.6), then

$$|\mu a_2^2 - a_3| \leq \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left\{ \left(\frac{2A^2}{(1-k^2)} - \frac{1-A^2}{3} - \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} \right) |c_1^2| + 2|c_2| \right\}$$

Applying Lemma 2, we get

$$(3.12) \quad |\mu a_2^2 - a_3| \leq \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left\{ \left(\frac{2A^2}{(1-k^2)} - \frac{1-A^2}{3} - \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} \right) 4 + 4 \right\} \\ = \frac{(n+1)(n+2)A^2}{6(1-k^2)} \left(\frac{2A^2}{(1-k^2)} + \frac{2+A^2}{3} - \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} \right)$$

which is the third part of the assertion (3.5).

Lastly from (3.11), write

$$(3.13) \quad |\mu a_2^2 - a_3| = \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left| 2c_2 + \left\{ \frac{2A^2}{(1-k^2)} - \frac{1-A^2}{3} - \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} \right\} c_1^2 \right|$$

$$\begin{aligned} &\leq \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left\{ 2 \left| c_2 - \frac{1}{2}c_1^2 \right| \right. \\ &\quad \left. + \left| \frac{2A^2}{(1-k^2)} + \frac{2+A^2}{3} - \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} \right| |c_1^2| \right\}. \end{aligned}$$

Observe that $\alpha_2 < \mu < \alpha_1$ gives

$$(3.14) \quad \left| \frac{2A^2}{(1-k^2)} + \frac{2+A^2}{3} - \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} \right| \leq 1.$$

Therefore an application of Lemma 2 in (3.13) gives

$$(3.15) \quad |\mu a_2^2 - a_3| \leq \frac{(n+1)(n+2)A^2}{6(1-k^2)}$$

which is the second part of assertion (3.5).

We next discuss the sharpness of (3.5).

If $\mu > \alpha_1$, equality holds in (3.5) if and only if equality holds in (3.10). This happens if and only if $|c_1| = 2$ and $|c_1^2 - c_2| = 2$. Thus $w(z) = z$. Equivalently the extremal function is $\mathcal{G}(z)$ defined by (1.8) or one of its rotations.

If $\mu < \alpha_2$ then equality holds in (3.12) if and only if $c_1^2 = -4$ and $c_2 = -2$ in (3.7) if and only if $c_1 = 2e^{i\pi/2}$ or $c_1 = 2e^{i3\pi/2}$ which also gives $c_2 = -2$. Thus $w(z) = e^{i\theta}z$ where $\theta = \frac{\pi}{2}$ or $\theta = \frac{3\pi}{2}$ and the extremal function is $\psi(z, \theta, 1)$ or one of its rotations.

If $\mu = \alpha_2$, the equality holds if and only if $|c_2| = 2$. Equivalently, we have

$$p_1(z) = \frac{1+\eta}{2} \left(\frac{1+z}{1-z} \right) + \frac{1-\eta}{2} \left(\frac{1-z}{1+z} \right) \quad (0 < \eta < 1; z \in \mathcal{U}).$$

Thus the extremal function f is $\psi(z, 0, \eta)$ or one of its rotations.

Similarly if $\mu = \alpha_1$, is equivalent to

$$\frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} - \frac{5+A^2}{3} - \frac{2A^2}{(1-k^2)} = 0.$$

Therefore equality holds true in (3.10) if and only if $|c_1^2 - c_2| = 2$ in (3.8). This happens if and only if

$$\frac{1}{p_1(z)} = \frac{1+\eta}{2} \left(\frac{1+z}{1-z} \right) + \frac{1-\eta}{2} \left(\frac{1-z}{1+z} \right) \quad (0 < \eta < 1; z \in \mathcal{U}).$$

Thus the extremal function is $\psi(z, \pi, \eta)$ or one of its rotations.

Lastly if $\alpha_2 < \mu < \alpha_1$, then equality holds true if $|c_1| = 0$ and $|c_2| = 2$. Equivalently we have

$$p_1(z) = \frac{1+\eta z^2}{1-\eta z^2} \quad (0 \leq \eta \leq 1; z \in \mathcal{U}).$$

Thus the function f is $\psi(z, 0, 0)$ or one of its rotation. The proof of Theorem 1 is complete. ■

Taking $n = 0$ and $n = 1$ in Theorem 1, we get the Fekete-Szegö inequalities for the classes $k - \mathcal{UCV}$ and $k - \mathcal{SP}$ ($0 \leq k < 1$) respectively, obtained earlier by the authors [12].

COROLLARY 1. *Let the function f given by (1.1) be in the class $k - \mathcal{UCV}$ ($0 \leq k < 1$). Then*

$$|\mu a_2^2 - a_3| \leq \begin{cases} \frac{2A^2}{3(1-k^2)} \left(\frac{3A^2\mu}{2(1-k^2)} - \frac{(7-k^2)A^2}{6(1-k^2)} - \frac{1}{3} \right); & \mu \geq \alpha_1^*, \\ \frac{A^2}{3(1-k^2)}; & \alpha_2^* \leq \mu \leq \alpha_1^*, \\ \frac{2A^2}{3(1-k^2)} \left(\frac{1}{3} + \frac{(7-k^2)A^2}{6(1-k^2)} - \frac{3A^2\mu}{2(1-k^2)} \right); & \mu \leq \alpha_2^*, \end{cases}$$

where the constant A is given by (2.2),

$$\alpha_1^* = \frac{5(1-k^2)}{9A^2} + \frac{7-k^2}{9} \quad \text{and} \quad \alpha_2^* = \frac{7-k^2}{9} - \frac{1-k^2}{9A^2}.$$

The result is best possible.

COROLLARY 2. *Let the function f given by (1.1) be in the class $k - \mathcal{SP}$ ($0 \leq k < 1$). Then*

$$|\mu a_2^2 - a_3| \leq \begin{cases} \frac{2A^2}{(1-k^2)} \left(\frac{2A^2\mu}{(1-k^2)} - \frac{(7-k^2)A^2}{6(1-k^2)} - \frac{1}{3} \right); & \mu \geq \alpha_1^{**}, \\ \frac{A^2}{(1-k^2)}; & \alpha_2^{**} \leq \mu \leq \alpha_1^{**}, \\ \frac{2A^2}{(1-k^2)} \left(\frac{1}{3} + \frac{(7-k^2)A^2}{6(1-k^2)} - \frac{2A^2\mu}{(1-k^2)} \right); & \mu \leq \alpha_2^{**}, \end{cases}$$

where the constant A is given by (2.2),

$$\alpha_1^{**} = \frac{5(1-k^2)}{12A^2} + \frac{7-k^2}{12} \quad \text{and} \quad \alpha_2^{**} = \frac{(7-k^2)}{12} - \frac{1-k^2}{12A^2}.$$

The result is best possible.

THEOREM 2. *Let the function f given by (1.1) be in the class $k - \mathcal{UCV}_n$ ($k = 1, n \in \mathbb{N}$). Then*

$$(3.16) \quad |\mu a_2^2 - a_3| \leq \begin{cases} \frac{4(n+1)(n+2)}{3\pi^2} \left(\frac{12(n+1)\mu}{(n+2)\pi^2} - \frac{1}{3} - \frac{4}{\pi^2} \right); & \mu \geq \beta_1, \\ \frac{2(n+1)(n+2)}{3\pi^2}; & \beta_2 \leq \mu \leq \beta_1, \\ \frac{4(n+1)(n+2)}{3\pi^2} \left(\frac{4}{\pi^2} + \frac{1}{3} - \frac{12(n+1)\mu}{(n+2)\pi^2} \right); & \mu \leq \beta_2, \end{cases}$$

where,

$$\beta_1 := \beta_1(k) = \frac{(n+2)}{(n+1)} \left(\frac{5\pi^2}{72} + \frac{1}{3} \right) \quad \text{and} \quad \beta_2 := \beta_2(k) = \frac{(n+2)}{(n+1)} \left(\frac{1}{3} - \frac{\pi^2}{72} \right).$$

Each of the estimates in (3.16) is sharp.

Proof. We put the values of Q_1 and Q_2 for $k = 1$ given in (3.5) and (3.6). By following the lines of proof of Theorem 1 the result follows. ■

Taking $n = 0$ in Theorem 2, we get the Fekete-Szegö inequalities for the class \mathcal{UCV} obtained earlier by Ma and Minda [11]. Similarly the choice $n = 1$ yields a result for the class \mathcal{SP} obtained recently by the present authors [12], which is as follows:

COROLLARY 3. *Let the function f given by (1.1) be in the class \mathcal{SP} . Then*

$$|\mu a_2^2 - a_3| \leq \begin{cases} \frac{8}{\pi^2} \left(\frac{8\mu}{\pi^2} - \frac{1}{3} - \frac{4}{\pi^2} \right); & \mu \geq \beta_1^*, \\ \frac{4}{\pi^2}; & \beta_2^* \leq \mu \leq \beta_1^*, \\ \frac{8}{\pi^2} \left(\frac{4}{\pi^2} + \frac{1}{3} - \frac{8\mu}{\pi^2} \right); & \mu \leq \beta_2^*, \end{cases}$$

where,

$$\beta_1^* = \frac{3}{2} \left(\frac{5\pi^2}{72} + \frac{1}{3} \right) \quad \text{and} \quad \beta_2^* = \frac{3}{2} \left(\frac{1}{3} - \frac{\pi^2}{72} \right).$$

The result is best possible.

THEOREM 3. *Let the function f given by (1.1) be in the class $k - \mathcal{UCV}_n$ ($k > 1$). Then*

$$(3.17) \quad |\mu a_2^2 - a_3| \leq \begin{cases} \frac{(n+1)(n+2)Q_1}{12} \left(\frac{3(n+1)\mu}{(n+2)} Q_1 - Q_1 \right. \\ \left. - \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right); & \mu \geq \gamma_1, \\ \frac{(n+1)(n+2)Q_1}{12}; & \gamma_2 \leq \mu \leq \gamma_1, \\ \frac{(n+1)(n+2)Q_1}{12} \left(Q_1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right. \\ \left. - \frac{3(n+1)\mu}{(n+2)} Q_1 \right); & \mu \leq \gamma_2, \end{cases}$$

where $\mathcal{K}(t)$ is the complete elliptic integral of the first kind, Q_1 is given in (2.1),

$$\gamma_1 := \gamma_1(k) = \frac{(n+2)}{3Q_1(n+1)} \left(1 + Q_1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right)$$

and

$$\gamma_2 := \gamma_2(k) = \frac{(n+2)}{3Q_1(n+1)} \left(Q_1 - 1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right).$$

Each of the estimates in (3.17) is sharp.

Proof. We follow the lines of proof of Theorem 1 and give here only the essential steps. Putting the value of Q_2 for $k > 1$ from Lemma 1 in (3.4) we get

$$a_2 = \frac{Q_1(n+1)}{4} c_1,$$

$$a_3 = \frac{Q_1(n+1)(n+2)}{12} \left[\frac{Q_1 c_1^2}{4} + \frac{1}{2} \left(c_2 - \frac{c_1^2}{2} \right) + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \frac{c_1^2}{4} \right].$$

Therefore

$$(3.18) \quad |\mu a_2 - a_3| = \frac{(n+1)(n+2)Q_1}{48} \left| \left\{ \frac{3Q_1(n+1)\mu}{(n+2)} - \left(Q_1 - 1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right) \right\} c_1^2 - 2c_2 \right|$$

$$(3.19) \quad \leq \frac{(n+1)(n+2)Q_1}{48} \left\{ \left| \frac{3Q_1(n+1)\mu}{(n+2)} - 1 - Q_1 - \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right| |c_1^2| + 2|c_1^2 - c_2| \right\}.$$

Let

$$(3.20) \quad \mu \geq \frac{(n+2)}{3Q_1(n+1)} \left(1 + Q_1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right) = \gamma_1 \quad (\text{say})$$

Observe that if $\mu \geq \gamma_1$, then the expression inside the first modulus of right-hand side (3.19) is non negative. Thus by applying Lemma 2 we get

$$(3.21) \quad |\mu a_2 - a_3| \leq \frac{(n+1)(n+2)Q_1}{12} \left(\frac{3Q_1(n+1)\mu}{(n+2)} - Q_1 - \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right)$$

which is the first part of the assertion (3.17).

Rewriting (3.18)

$$\begin{aligned}
 |μa_2^2 - a_3| &= \frac{(n+1)(n+2)Q_1}{48} \left| -2c_2 - \left\{ Q_1 - 1 \right. \right. \\
 &\quad \left. \left. + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} - \frac{3Q_1(n+1)\mu}{(n+2)} \right\} c_1^2 \right| \\
 (3.22) \quad &= \frac{(n+1)(n+2)Q_1}{48} \left| 2c_2 + \left\{ Q_1 - 1 \right. \right. \\
 &\quad \left. \left. + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} - \frac{3Q_1(n+1)\mu}{(n+2)} \right\} c_1^2 \right| \\
 (3.23) \quad &\leq \frac{(n+1)(n+2)Q_1}{48} \left\{ \left| Q_1 - 1 \right. \right. \\
 &\quad \left. \left. + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} - \frac{3Q_1(n+1)\mu}{(n+2)} \right| |c_1^2| + 2|c_2| \right\}
 \end{aligned}$$

Let,

$$(3.24) \quad \mu \leq \frac{(n+2)}{3Q_1(n+1)} \left(Q_1 - 1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right) = \gamma_2 \quad \text{say}$$

If $\mu \leq \gamma_2$, then the expression inside the first modulus of right hand side of (3.23) is non negative. Thus by applying Lemma 2 we get

$$\begin{aligned}
 (3.25) \quad |μa_2^2 - a_3| &= \frac{(n+1)(n+2)Q_1}{12} \left(Q_1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right. \\
 &\quad \left. - \frac{3Q_1(n+1)\mu}{(n+2)} \right).
 \end{aligned}$$

Which is the third part of assertion (3.17). Lastly (3.22) gives

$$\begin{aligned}
 (3.26) \quad |μa_2^2 - a_3| &= \frac{(n+1)(n+2)Q_1}{48} \left| 2 \left(c_2 - \frac{1}{2}c_1^2 \right) \right. \\
 &\quad \left. + \left(Q_1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} - \frac{3Q_1(n+1)\mu}{(n+2)} \right) c_1^2 \right| \\
 &\leq \frac{(n+1)(n+2)Q_1}{48} \left\{ 2 \left| c_2 - \frac{1}{2}c_1^2 \right| \right. \\
 &\quad \left. + \left| Q_1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} - \frac{3Q_1(n+1)\mu}{(n+2)} \right| |c_1^2| \right\}.
 \end{aligned}$$

Notably $\gamma_2 < \mu < \gamma_1$ gives

$$\left| Q_1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} - \frac{3Q_1(n+1)\mu}{(n+2)} \right| \leq 1.$$

Therefore, if $\gamma_2 < \mu < \gamma_1$, an application of Lemma 2 in (3.26) yields

$$\begin{aligned} |\mu a_2^2 - a_3| &\leq \frac{(n+1)(n+2)Q_1}{48} \left\{ 2 \left| c_2 - \frac{1}{2}c_1^2 \right| + |c_1^2| \right\} \\ &\leq \frac{(n+1)(n+2)Q_1}{12}. \end{aligned}$$

This is the middle part of the assertions in (3.17). The sharpness of each estimate in (3.17) can be established as in Theorem 1. The proof of Theorem 3 is complete. \blacksquare

Taking $n = 0$ and $n = 1$ in Theorem 3, we get the Fekete-Szegö inequalities for the classes $k - \mathcal{UCV}$ and $k - \mathcal{SP}$ respectively obtained recently by the present authors [12]. We list them here for the sake of completeness.

COROLLARY 4. *Let the function f given by (1.1) be in the class $k - \mathcal{UCV}$ ($k > 1$). Then*

$$|\mu a_2^2 - a_3| \leq \begin{cases} \frac{Q_1}{6} \left(\frac{3Q_1\mu}{2} - Q_1 - \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right); & \mu \geq \gamma_1^*, \\ \frac{Q_1}{6}; & \gamma_2^* \leq \mu \leq \gamma_1^*, \\ \frac{Q_1}{6} \left(Q_1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} - \frac{3Q_1\mu}{2} \right); & \mu \leq \gamma_2^*, \end{cases}$$

where $\mathcal{K}(t)$ is the complete elliptic integral of the first kind, Q_1 is given in (2.1),

$$\gamma_1^* = \frac{2}{3Q_1} \left(1 + Q_1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right)$$

and

$$\gamma_2^* = \frac{2}{3Q_1} \left(Q_1 - 1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right).$$

Each of the estimates is sharp.

COROLLARY 5. *Let the function f given by (1.1) be in the class $k - \mathcal{SP}$ ($k > 1$). Then*

$$|\mu a_2^2 - a_3| \leq \begin{cases} \frac{Q_1}{2} \left(2Q_1\mu - Q_1 - \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right); & \mu \geq \gamma_1^{**}, \\ \frac{Q_1}{2}; & \gamma_2^{**} \leq \mu \leq \gamma_1^{**}, \\ \frac{Q_1}{2} \left(Q_1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} - 2Q_1\mu \right); & \mu \leq \gamma_2^{**}, \end{cases}$$

where $\mathcal{K}(t)$ is the complete elliptic integral of the first kind, Q_1 is given in (2.1),

$$\gamma_1^{**} = \frac{1}{2Q_1} \left(1 + Q_1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right)$$

and

$$\gamma_2^{**} = \frac{1}{2Q_1} \left(Q_1 - 1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right).$$

Each of the estimates is sharp.

4. Remarks on main results

In this section we discuss some improvements of the middle inequalities in (3.5), (3.16) and (3.17) respectively.

Remark 1. The second part of assertion in (3.5) can be improved as follows:

$$(4.1) \quad \begin{aligned} |\mu a_2^2 - a_3| &+ \left\{ \mu - \frac{(n+2)}{(n+1)} \left(\frac{1}{3} - \frac{(1-A^2)(1-k^2)}{18A^2} \right) \right\} |a_2^2| \\ &\leq \frac{(n+1)(n+2)A^2}{6(1-k^2)}; \quad \alpha_2 \leq \mu \leq \alpha_3, \end{aligned}$$

and

$$(4.2) \quad \begin{aligned} |\mu a_2^2 - a_3| &+ \left\{ \frac{(n+2)}{(n+1)} \left(\frac{1}{3} + \frac{(5+A^2)(1-k^2)}{18A^2} \right) - \mu \right\} |a_2^2| \\ &\leq \frac{(n+1)(n+2)A^2}{6(1-k^2)}; \quad \alpha_3 \leq \mu \leq \alpha_1, \end{aligned}$$

where α_3 is given by

$$(4.3) \quad \alpha_3 := \frac{n+2}{n+1} \left(\frac{1}{3} + \frac{(2+A^2)(1-k^2)}{18A^2} \right).$$

Proof. Suppose $\alpha_2 \leq \mu \leq \alpha_3^*$. We continue with the estimate in (3.13) and write

$$\begin{aligned}
& |\mu a_2^2 - a_3| + (\mu - \alpha_2)|a_2|^2 \\
&= |\mu a_2^2 - a_3| + \left\{ \mu - \frac{(n+2)}{(n+1)} \left(\frac{1}{3} - \frac{(1-A^2)(1-k^2)}{18A^2} \right) \right\} |a_2^2| \\
&\leq \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left\{ 2 \left| c_2 - \frac{1}{2} c_1^2 \right| + \left| \frac{2A^2}{(1-k^2)} + \frac{2+A^2}{3} - \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} \right| |c_1^2| \right\} \\
&\quad + \left\{ \mu - \frac{(n+2)}{(n+1)} \left(\frac{1}{3} - \frac{(1-A^2)(1-k^2)}{18A^2} \right) \right\} \left| \frac{(n+1)^2 A^4 c_1^2}{4(1-k^2)^2} \right| \\
&\leq \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left\{ 2 \left| c_2 - \frac{1}{2} c_1^2 \right| + \left(\frac{2A^2}{(1-k^2)} + \frac{2+A^2}{3} - \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} \right) |c_1^2| \right. \\
&\quad \left. + \left\{ \frac{6(n+1)A^2\mu}{(n+2)(1-k^2)} - \frac{6A^2}{(1-k^2)} \left(\frac{1}{3} - \frac{(1-A^2)(1-k^2)}{18A^2} \right) \right\} |c_1^2| \right\} \\
&\leq \frac{(n+1)(n+2)A^2}{24(1-k^2)} \left\{ 2 \left| c_2 - \frac{1}{2} c_1^2 \right| + |c_1^2| \right\} \\
&= \frac{(n+1)(n+2)A^2}{6(1-k^2)}.
\end{aligned}$$

We get (4.1). On the other hand suppose $\alpha_3 \leq \mu \leq \alpha_1$. In this case we estimate $|\mu a_2^2 - a_3| + (\alpha_1 - \mu)|a_2|^2$. Now following the lines of proof for (4.1) with obvious changes we get (4.2).

The proof of next two remarks are same as in Remark 1. Therefore we omit details.

Remark 2. The second part of assertion in (3.16) can be improved as follows:

$$(4.4) \quad |\mu a_2^2 - a_3| + \left\{ \mu - \frac{n+2}{n+1} \left(\frac{1}{3} - \frac{\pi^2}{72} \right) \right\} |a_2^2| \leq \frac{2(n+1)(n+2)}{3\pi^2}; \quad \beta_2 \leq \mu \leq \beta_3,$$

and

$$(4.5) \quad |\mu a_2^2 - a_3| + \left\{ \frac{n+2}{n+1} \left(\frac{5\pi^2}{72} + \frac{1}{3} \right) - \mu \right\} |a_2^2| \leq \frac{2(n+1)(n+2)}{3\pi^2}; \quad \beta_3 \leq \mu \leq \beta_1,$$

where β_3 is given by

$$(4.6) \quad \beta_3 := \frac{n+2}{n+1} \left(\frac{1}{3} + \frac{\pi^2}{36} \right).$$

Remark 3. The second part of assertion in (3.17) can be improved as follows:

$$(4.7) \quad |\mu a_2^2 - a_3| + \left\{ \mu - \frac{(n+2)}{3Q_1(n+1)} \left(Q_1 - 1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right) \right\} \times |a_2^2| \\ \leq \frac{(n+1)(n+2)Q_1}{12}; \quad \gamma_2 \leq \mu \leq \gamma_3,$$

and

$$(4.8) \quad |\mu a_2^2 - a_3| + \left\{ \frac{(n+2)}{3Q_1(n+1)} \left(Q_1 + 1 + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{24\mathcal{K}^2(t)\sqrt{t}(1+t)} \right) - \mu \right\} \times |a_2^2| \\ \leq \frac{(n+1)(n+2)Q_1}{12}; \quad \gamma_3 \leq \mu \leq \gamma_1,$$

where γ_3 is given by

$$(4.9) \quad \gamma_3 := \frac{n+2}{n+1} \left(\frac{1}{3} + \frac{4\mathcal{K}^2(t)(t^2 + 6t + 1) - \pi^2}{72Q_1\mathcal{K}^2(t)\sqrt{t}(1+t)} \right).$$

5. An alternate method

Let $\mathcal{P}(q_k)$ denote the subclass of \mathcal{A} , consisting of functions $h(z) \prec q_k(z)$ in \mathcal{U} , where $q_k(z)$ is as in Lemma 1 and \prec denotes subordination. We need the following result:

LEMMA 3 [7]. Let $k \in [0, \infty)$ be fixed and the function $h \in \mathcal{P}(q_k)$ be given by the series expansion

$$(5.1) \quad h(z) = 1 + b_1 z + b_2 z^2 + \dots, \quad (z \in \mathcal{U})$$

then

$$(5.2) \quad |b_2 - ub_1^2| \leq \begin{cases} Q_1(k) - uQ_1^2(k); & u \leq 0, \\ Q_1(k); & u \in (0, 1], \\ Q_1(k) + (u-1)Q_1^2(k); & u \geq 1, \end{cases}$$

where $Q_1(k)$ is as in Lemma 1. If $0 < u < 1$ then equality holds in (5.2) if $h(z) = q_k(z^2)$.

Now replacing $q_k(w(z))$ by $h(z)$ in the equation (3.1) we get

$$a_2 = \left(\frac{n+1}{2} \right) b_1 \\ a_3 = \frac{(n+1)(n+2)}{12} (b_1^2 + b_2)$$

and

$$\mu a_2^2 - a_3 = \frac{(n+1)(n+2)}{12} \left[\left\{ \frac{3(n+1)\mu}{(n+2)} - 1 \right\} b_1^2 - b_2 \right]$$

An application of Lemma 3 with,

$$u = 3 \left(\frac{n+1}{n+2} \right) \mu - 1$$

yields

$$(5.3) \quad |\mu a_2^2 - a_3| = \frac{(n+1)(n+2)}{12} Q_1(k) \quad \text{if } \frac{(n+2)}{3(n+1)} < \mu \leq \frac{2}{3} \left(\frac{n+2}{n+1} \right)$$

Putting the values of $Q_1(k)$ from Lemma 1 for $0 \leq k < 1$, $k = 1$ and $k > 1$ the estimate (5.3) simplifies to the middle inequalities in (3.5), (3.16) and (3.17) respectively. However, following the lines of proof Theorem 3.1 in [7] it can be shown that

$$\alpha_1(k), \beta_1(k), \gamma_1(k) > \frac{2}{3} \left(\frac{n+2}{n+1} \right)$$

and

$$\alpha_2(k), \beta_2(k), \gamma_2(k) < \frac{1}{3} \left(\frac{n+2}{n+1} \right)$$

for every $0 \leq k < \infty$ where $\alpha_1(k), \alpha_2(k); \beta_1(k), \beta_2(k)$ and $\gamma_1(k), \gamma_2(k)$ are defined as in Theorems 3.1, 3.2 and 3.3 respectively. Thus the methods of proof adopted for Theorems 3.1, 3.2 and 3.3 yield larger interval for μ than the interval obtained for the estimates of $|\mu a_2^2 - a_3|$ in (5.3).

REFERENCES

- [1] N. I. AHIEZER, Elements of theory of elliptic functions (in Russian), Moscow, 1970.
- [2] G. D. ANDERSON, M. K. VAMANAMURTHY AND M. K. VUORINEN, Conformal invariants, inequalities and quasiconformal maps, Wiley-Interscience, 1997.
- [3] B. C. CARLSON AND D. B. SHAFFER, Starlike and pre-starlike hypergeometric functions, SIAM J. Math. Anal. **15** (1984), 737–745.
- [4] P. L. DUREN, Univalent Functions, Grundlehren der mathematischen Wissenschaften **259**, Bd., Springer-Verlag, New York, Berlin, Heidelberg, Tokyo, 1983.
- [5] M. FEKETE AND G. SZEGÖ, Eine bemerkung über ungerade schlichte funktionen, J. London Math. Soc. **8** (1933), 85–89.
- [6] A. W. GOODMAN, On uniformly convex functions, Ann. Polon. Math. **56** (1991), 87–92.
- [7] S. KANAS, Coefficient estimates in subclasses of the Carathéory class related to conical domains, Acta Math. Univ. Comenian. **74** (2005), 149–161.
- [8] S. KANAS AND A. WISNIOWSKA, Conic regions and k -uniform convexity, J. Comput. Appl. Math. **105** (1999), 327–336.

- [9] S. KANAS AND A. WISNIEWSKA, Conic domains and starlike functions, *Rev. Roumaine Math. Pures Appl.* **45** (2000), 647–657.
- [10] W. KOEPEL, On the Fekete-Szegö problem for close to convex functions, *Proc. Amer. Math. Soc.* **101**(1) (1987), 89–95.
- [11] W. MA AND D. MINDA, Uniformly convex functions I and II, *Ann. Polon. Math.* **57**(2) (1992), 165–175, *Ann. Polon. Math.* **58**(3) (1993), 275–285.
- [12] A. K. MISHRA AND P. GOCHHAYAT, Fekete-Szegö problem for k -uniformly convex functions and for a class defined by the Owa-Srivastava operator, *J. Math. Anal. Appl.* **347** (2008), 563–572.
- [13] K. I. NOOR, On new class of integral operator, *J. Natur. Geom.* **16** (1999), 71–80.
- [14] K. I. NOOR AND M. A. NOOR, On integral operators, *J. Math. Anal. Appl.* **238** (1999), 341–352.
- [15] K. I. NOOR AND M. A. NOOR, On certain classes of analytic functions defined by Noor integral operator, *J. Math. Anal. Appl.* **281** (2003), 244–252.
- [16] F. RØNNING, Uniformly convex functions and a corresponding class of starlike functions, *Proc. Amer. Math. Soc.* **118** (1993), 189–196.
- [17] H. M. SRIVASTAVA, A. K. MISHRA AND M. K. DAS, A nested class of analytic functions defined by fractional calculus, *Commun. Appl. Anal.* **2** (1998), 321–332.
- [18] H. M. SRIVASTAVA AND A. K. MISHRA, Applications of fractional calculus to parabolic starlike and uniformly convex functions, *Comput. Math. Appl.* **39**(3/4) (2000), 57–69.
- [19] H. M. SRIVASTAVA AND S. OWA (EDS.), Current topics in analytic function theory, World Scientific Publishing Company, Singapore, New Jersey, London and Hong Kong, 1992.

Akshaya Kumar Mishra
 DEPARTMENT OF MATHEMATICS
 BERHAMPUR UNIVERSITY
 BHANJA BIHAR
 BERHAMPUR-760007, ORISSA
 INDIA
 E-mail: akshayam2001@yahoo.co.in

Priyabrat Gochhayat
 DEPARTMENT OF MATHEMATICS
 SAMBALPUR UNIVERSITY
 JYOTI VIHAR
 SAMBALPUR-768019, ORISSA
 INDIA
 E-mail: pb_gochhayat@yahoo.com