

180. Further Characterizations for the Jacobson Radical of a Ring^{*)}

By Ferenc Andor SZÁSZ

(Comm. by Kinjirô KUNUGI, M. J. A., Feb. 12, 1971)

Ring will mean in this note always an associative ring. For the fundamental notations, used here, we refer the reader e.g. to N. Divinsky [2] and N. Jacobson [3]. Various characterizations for the Jacobson radical F of a ring A are given (cf. N. Jacobson [3], manuscript of author's book [4] or survey [5]). Among others, author's paper [8] asserts, that any quasiprimitive ideal coincides with a primitive ideal of the ring, the Jacobson radical being the intersection of all quasiprimitive ideals (cf. author's papers [6] and [7]). Four new characterizations for the Brown-McCoy radical, for the other well used concrete radical of rings, can be found in author's earlier paper [11].

Let A be a ring. Then the equalities $(a)_r = (b)_r$ for the principal right ideals $(x)_r = Tx + xA$ ($a, b, x \in A$ and T is the ring of the rational integers), of A , define an equivalence relation $a \equiv b$ in the set of the elements of the ring, such that \equiv is suitable for yielding of some characterizations of the Jacobson radical. Obviously the relation \equiv is a left congruence of the multiplicative semigroup of the ring.

Here we mention only without proof our results, obtained spring 1968, whose details [10] will be published with their proofs later:

Theorem 1. *The Jacobson radical F of a ring A coincides with the subset B of those elements b of A , such the equivalence relation $a \equiv a + ac$ for any $a \in A$ and for any element c of the principal right ideal $(b)_r$, generated by $b \in B$, of A , holds.*

Theorem 2. *F coincides with the subset D of those elements d of A , such the equivalence relation $a \equiv a + adb$ for any $a, b \in A$ holds.*

Remark. On the basis of these statements, F can be considered, as a "right-sided antisimple" (twosided) radical (cf. V. A. Andrunakievitch [1]).

References

- [1] V. A. Andrunakievitch: Antisimple and strong idempotent rings. *Izvestiya Akad. Nauk SSSR, Mat. Ser.*, **21**, 125-144 (1957) (in Russian).
- [2] N. Divinsky: *Rings and Radicals*. London (1965).
- [3] N. Jacobson: *Structure of Rings* (2 edition). Providence (1964).
- [4] F. Szász: *Radikale der Ringe*. Budapest, Akadémiai Kiadó (to appear).

^{*)} Dedicated to Professor N. Jacobson.

- [5] F. Szász: On radicals of rings. *Matematikai Lapok*, I: **19** (3-4), 259-301 (1968), II: **20**(1-2), 99-116 (1969), III: **20**(3-4), 311-346 (1969) (in Hungarian, survey without proofs).
- [6] —: Lösung eines Problems bezüglich einer Charakterisierung des Jacobsonischen Radikals. *Acta Math. Acad. Sci. Hungar.*, **18**, 261-272 (1967).
- [7] —: Eine Charakterisierung des Jacobsonischen Radikals eines Ringes. *Bull. Acad. Polon. Sci. Classe Troisième*, **15**, 53-56 (1967).
- [8] —: The sharpening of a result concerning the primitive ideals of an associative ring. *Proc. Amer. Math. Soc.*, **18**, 910-912 (1967).
- [9] —: Die Lösung eines Problems bezüglich des Durchschnittes zweier modularer Rechtsideale in einem Ring. *Acta Math. Acad. Sci. Hungar.*, **20**(1-2), 211-216 (1969).
- [10] —: Äquivalenzrelation für Charakterisierung des Jacobsonischen Radikals eines Ringes. *Acta Math. Acad. Sci. Hungar.*, **22** (to appear).
- [11] —: An observation on the Brown-McCoy radical. *Proc. Japan Acad.*, **37**, 413-416 (1961).