

uniquely as a product of a finite number of prime differential ideals. The consistency of the axioms is easily shown. If we define differentiation in the ring $C(x)$, obtained by adjoining x to the field of the rational numbers, in *any* way so as to leave it closed, it may be shown that Axioms I–IV are always satisfied. In $C(x, y)$ differentiation *may* be defined in such way that the statement as above still holds. This is of interest because every ordinary ideal in $C(x, y)$ may not be expressed uniquely as a product of a finite number of prime ideals.

HOFSTRA COLLEGE

ON THE ITERATION OF LINEAR HOMOGENEOUS TRANSFORMATIONS

ARNOLD DRESDEN

1. Statement of problem. The question which this note tries to answer is that of determining under what conditions on the matrix (a_{ij}) , $(i, j = 1, \dots, n)$, the n -fold multiple sequence of complex numbers $x_k, x'_k, \dots, x_k^{(m)}, \dots$ ($k = 1, 2, \dots, n$) obtained by iteration of the linear homogeneous transformation $x'_k = a_{kj}x_j$ will converge for every initial set $x = (x_1, x_2, \dots, x_n)$. Convergence is to be understood in the sense that there exists a set X_1, X_2, \dots, X_n such that, for $k = 1, 2, \dots, n$, $x_k^{(m)} \rightarrow X_k$, as $m \rightarrow \infty$.

2. Jordan normal form. We begin by recalling that a matrix $A = (a_{ij})$ with complex elements is similar to its Jordan normal form J_0 . This means that there exists a unimodular matrix P , such that $A = P^{-1}J_0P$ and $J_0 = PAP^{-1}$, where J_0 is the direct sum of Jordan matrices J_1, \dots, J_N . To each elementary divisor $(\lambda - \lambda_\rho)^{e_\rho}$ of the characteristic matrix $\lambda I - A$ ($\rho = 1, 2, \dots, N$) and $e_1 + e_2 + \dots + e_N = n$, corresponds a Jordan matrix J_ρ . If $e_\rho > 1$, then J_ρ has zero elements everywhere, except in the principal diagonal, all of whose elements are λ_ρ , and in the diagonal immediately below the principal diagonal, all of whose elements are 1.¹ If $e_\rho = 1$, then J_ρ consists of the single element λ_ρ .

It follows that any integral power of J_0 is the direct sum of the same powers of the Jordan matrices J_ρ . Let us now denote by J an arbitrary Jordan matrix of order $n > 1$,

Received by the editors August 25, 1941.

¹ See, for example, MacDuffee, *Introduction to Abstract Algebra*, p. 241.

$$J = \begin{pmatrix} \lambda & 0 & 0 & 0 & \cdots & \cdots & \cdots & 0 \\ 1 & \lambda & 0 & 0 & \cdots & \cdots & \cdots & 0 \\ 0 & 1 & \lambda & 0 & \cdots & \cdots & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdots & \cdots & \cdot & \cdot \\ 0 & \cdot & \cdot & \cdot & \cdots & \cdots & 0 & \lambda & 0 \\ 0 & \cdot & \cdot & \cdot & \cdots & \cdots & 0 & 1 & \lambda \end{pmatrix}.$$

It is then easily seen that J^2 is a matrix of order n , whose first column is $\lambda^2, 2\lambda, 1, 0, \dots, 0$, and whose second column is obtained from the first column by shifting the elements one place downward, as far as the length permits, and placing 0 in the first place; the third column is obtained from the second in similar manner; the last column is $0\ 0\ \cdots\ 0\ \lambda^2$. A simple induction enables us to prove that, for any positive integer h , the matrix J^h is of order n , that its first column consists of the successive terms in the binomial expansion of $(\lambda + 1)^h$, followed by zeros, provided $n \geq h + 1$ and otherwise of as many of these terms as the length of the column permits, that each of its columns after the first is obtained from the one to the left of it by shifting the elements one place downward and filling the open position above with a zero, and that its last column is $0\ 0\ 0\ \cdots\ 0\ \lambda^h$. If J is a Jordan matrix of order 1, consisting of the single element λ then J^n consists of the single element λ^n .

3. Reduction of problem; solution. Returning now to our linear homogeneous transformation, we write it in the form

$$(1) \quad x' = xA,$$

where x and x' stand for the row-vectors (x_1, x_2, \dots, x_n) and $(x'_1, x'_2, \dots, x'_n)$, respectively. By means of the unimodular matrix P , we map the space of the row-vectors x on the space of the row-vectors y : $x = yP, \quad y = xP^{-1}$. This mapping carries the transformation (1) into a transformation in the y -space:

$$(2) \quad y' = x'P^{-1} = xAP^{-1} = yPAP^{-1} = yJ_0.$$

It follows from the mapping that the vectors $x^{(m)}$ will converge, in the sense described in §1, if and only if the vectors $y^{(m)}$ converge. But $y^{(m)} = y^{(m-1)}J_0 = \cdots = y_0J_0^m = y(J_1^m \dot{+} J_2^m \dot{+} \cdots \dot{+} J_N^m)$. We obtain therefore the following form for the m th iterate of the transformation (2) in terms of the roots λ_ρ of the characteristic equation $|a_{ij} - \lambda\delta_{ij}| = 0$ and the exponents e_ρ of the corresponding elementary divisors $(\lambda - \lambda_\rho)^{e_\rho}$ of $\lambda I - A$, provided every $e_\rho > 1$.

$$\begin{aligned} y_1^{(m)} &= \lambda_1^m y_1 + C_{m,1}\lambda_1^{m-1} y_2 + \cdots + C_{m,e_1-1}\lambda_1^{m-e_1+1} y_{e_1}, \\ y_2^{(m)} &= \lambda_1^m y_2 + C_{m,1}\lambda_1^{m-1} y_3 + \cdots + C_{m,e_1-2}\lambda_1^{m-e_1+2} y_{e_1}, \end{aligned}$$

