

# SPACES OF MEASURABLE TRANSFORMATIONS<sup>1</sup>

BY ROBERT J. AUMANN

Communicated by Paul R. Halmos, February 27, 1960

By a *space* we shall mean a measurable space, i.e. an abstract set together with a  $\sigma$ -ring of subsets, called *measurable sets*, whose union is the whole space. The *structure* of a space will be the  $\sigma$ -ring of its measurable subsets. A *measurable transformation* from one space to another is a mapping such that the inverse image of every measurable set is measurable.

Let  $X$  and  $Y$  be spaces,  $F$  a set of measurable transformations from  $X$  into  $Y$ , and  $\phi_F: F \times X \rightarrow Y$  the natural mapping defined by  $\phi_F(f, x) = f(x)$ . A structure  $R$  on  $F$  will be called *admissible* if  $\phi_F$ , considered as a mapping from the product space  $(F, R) \times X$  into  $Y$ , is a measurable transformation.<sup>2</sup> It may not be possible to define an admissible structure on  $F$ ; if it is,  $F$  itself will also be called *admissible*. We are concerned with the problem of characterizing, for given  $X$  and  $Y$ , the admissible sets  $F$  and the admissible structures  $R$  on the admissible sets.

The following three theorems may be established fairly easily:

**THEOREM A.** *A set consisting of a single measurable transformation is admissible.*

**THEOREM B.** *A subset of an admissible set is admissible. Indeed, if  $G \subset F$ ,  $R$  is an admissible structure on  $F$ , and  $R_G$  is the subspace structure on  $G$  induced<sup>3</sup> by  $R$ , then  $R_G$  is admissible on  $G$ .*

**THEOREM C.** *The union of denumerably many admissible sets is admissible. Indeed, if  $F = \bigcup_{i=1}^{\infty} F_i$  and  $R_1, R_2, \dots$  are admissible structures on  $F_1, F_2, \dots$  respectively, then the structure  $R$  on  $F$  generated by the members of all the  $R_i$  is admissible on  $F$ .*

Much more can be said if  $X$  and  $Y$  are assumed to be *separable*, i.e. to have countably generated structures.<sup>4</sup> To state our theorems in this case we first define the concept of Banach class, closely related to that of Baire class. Let  $\mathfrak{A}$  be an arbitrary class of meas-

<sup>1</sup> The author is much indebted to Professor P. R. Halmos, who suggested a number of significant improvements in the complete version of this note.

<sup>2</sup>  $(F, R)$  is the space whose underlying abstract set is  $F$  and whose structure is  $R$ .

<sup>3</sup>  $R_G$  consists of all intersections of  $G$  with members of  $R$ .

<sup>4</sup> The term is used by analogy with its topological use. We will also use the term "separable structure," meaning a countably generated structure.

urable subsets of  $X$ . For each denumerable ordinal number  $\alpha \geq 1$ , we define classes  $P_\alpha(\mathfrak{A})$  and  $Q_\alpha(\mathfrak{A})$  inductively as follows:  $Q_1(\mathfrak{A})$  consists of all denumerable unions of members of  $\mathfrak{A}$ , and  $P_1(\mathfrak{A})$  consists of all complements of members of  $Q_1(\mathfrak{A})$ ; supposing  $Q_\beta(\mathfrak{A})$  and  $P_\beta(\mathfrak{A})$  to have been defined for all  $\beta < \alpha$ , we define  $Q_\alpha(\mathfrak{A}) = Q_1(\bigcup_{\beta < \alpha} P_\beta(\mathfrak{A}))$  and  $P_\alpha(\mathfrak{A}) = P_1(\bigcup_{\beta < \alpha} P_\beta(\mathfrak{A}))$ .  $Q_\alpha(\mathfrak{A}) \cup P_\alpha(\mathfrak{A})$  is the set of all subsets of  $X$  which can be "reached from  $\mathfrak{A}$ " by performing at most  $\alpha$  operations, where each operation consists of forming a denumerable union and a complement. If  $\mathfrak{A}$  generates the structure of  $X$ , then the union (over  $\alpha$ ) of all the  $Q_\alpha(\mathfrak{A})$  (or of the  $P_\alpha(\mathfrak{A})$ ) is the set of all measurable subsets of  $X$ . If  $\mathfrak{B}$  is a class of measurable subsets of  $Y$  and  $\alpha \geq 0$  is a denumerable ordinal number, then we define  $L_\alpha(\mathfrak{A}, \mathfrak{B})$  to be the set of all functions  $f: X \rightarrow Y$  such that for all  $B \in Q_1(\mathfrak{B})$ ,  $f^{-1}(B) \in Q_{\alpha+1}(\mathfrak{A})$ . If  $X$  and  $Y$  are separable and  $\mathfrak{A}$  and  $\mathfrak{B}$  are denumerable generating sets for their respective structures, then the union (over  $\alpha$ ) of all the  $L_\alpha(\mathfrak{A}, \mathfrak{B})$  is the set of all measurable transformations from  $X$  into  $Y$ . It will be denoted  $Y^X$ . In this case  $L_\alpha(\mathfrak{A}, \mathfrak{B})$  is called the *Banach class*<sup>5</sup> of order  $\alpha$  for  $(\mathfrak{A}, \mathfrak{B})$ . A subset  $F$  of  $Y^X$  is said to be of *bounded Banach class* if there is an  $\alpha$  and denumerable generating sets  $\mathfrak{A}, \mathfrak{B}$  such that  $F \subset L_\alpha(\mathfrak{A}, \mathfrak{B})$ . It is important to note that the definition of bounded Banach class is independent of the choice of  $\mathfrak{A}$  and  $\mathfrak{B}$ , i.e. that if  $F \subset L_\alpha(\mathfrak{A}, \mathfrak{B})$ , then for any other generating pair  $\mathfrak{A}', \mathfrak{B}'$ , there is an  $\alpha'$  such that  $F \subset L_{\alpha'}(\mathfrak{A}', \mathfrak{B}')$ . If  $X$  and  $Y$  are separable metric spaces and  $Y$  is pathwise connected, then the Banach classes coincide with the Baire classes (for appropriate choice of  $\mathfrak{A}$  and  $\mathfrak{B}$ ).

**THEOREM D.** *If  $X$  and  $Y$  are separable, then  $F$  is admissible if and only if it is of bounded Banach class.*

**THEOREM E.** *If  $X$  and  $Y$  are separable, then every admissible subset of  $Y^X$  has a separable admissible structure.*

A space  $Z$  and its structure are called *regular* if for all  $x, y \in Z$ , there is a measurable set in  $Z$  containing  $x$  but not  $y$ . It is known (cf. [2]) that a space is separable and regular if and only if it is isomorphic<sup>6</sup> to a subspace of  $I$ , where  $I$  denotes the unit interval  $[0, 1]$  with the usual Borel structure.

**THEOREM F.** *If  $X$  and  $Y$  are separable and regular, then every admissible subset of  $Y^X$  has a separable and regular admissible structure.*

<sup>5</sup> Because of the work that Banach [1] did in characterizing these classes.

<sup>6</sup> Two spaces are said to be *isomorphic* if there is a 1-1 correspondence between them that preserves measurability (in both directions).

The *natural* admissible structure on a given admissible set  $F$  is defined to be the smallest admissible structure on  $F$ , if it exists. Alternatively, it may be defined to be the intersection of all the admissible structures on  $F$ , in case this is admissible. Not every admissible set need have a natural admissible structure; the counterexample is due to P. R. Halmos.

If  $a \in X$  and  $B \subset Y$ , define  $F(a, B) = \{f: f \in F, f(a) \in B\}$ . It is not hard to prove that if  $B$  is measurable and  $a$  is arbitrary, then every admissible structure on  $F$  must contain  $F(a, B)$ . A "converse" would be that the structure generated by the  $F(a, B)$  is admissible, and it would follow that it is also natural.

**THEOREM G.** *If  $X$  and  $Y$  are separable metric spaces and  $F$  contains continuous functions only, then  $F$  has a natural admissible structure, which is generated by the set of all  $F(a, B)$ , where  $B$  is measurable and  $a$  is arbitrary.*

We now give some applications. A space is said to have the *discrete* structure if every subset is measurable. Let  $J$  be the space consisting of 0 and 1 only, and  $K$  the space of all positive integers, both with the discrete structure. If  $X$  is an arbitrary space, then  $X^J$  and  $X^K$  are both admissible, and possess natural admissible structures which make them isomorphic to  $X \times X$  and  $\times_{i=1}^{\infty} X_i$  respectively, where the  $X_i$  are copies of  $X$ . In particular,  $J^K$  is admissible and has a natural admissible structure which makes it isomorphic to  $I$ . These results are relatively trivial or at least easily derivable from known results.

The situation changes when we pass to exponent spaces with non-discrete structures. For example,  $J^I$  may be considered the set of all measurable subsets of  $I$ . It is not itself admissible. The set of all open subsets of  $I$  is admissible, as is the set of all closed subsets, the set of all  $G_\delta$ , etc. In general, a subset  $F$  of  $J^I$  is admissible if and only if all members of  $F$  can be constructed from the open subsets of  $I$  by taking denumerable unions and intersections at most  $\alpha$  times, where  $\alpha$  is an arbitrary denumerable ordinal number (which is fixed for given  $F$ , but may differ for different  $F$ ). I do not know whether or not every admissible subset of  $J^I$  has a natural admissible structure, but if  $F$  is admissible, then we may endow it with an admissible structure in such a way so that it will be isomorphic to a subset of  $I$ .

$I^I$  is not admissible. The set of all continuous functions from  $I$  into  $I$  is admissible; more generally, a necessary and sufficient condition that a subset  $F$  of  $I^I$  be admissible is that there exist a denumerable ordinal number  $\alpha$  such that all members of  $F$  are of Baire class  $\alpha$

at most. The set  $H$  of all continuous functions from  $I$  into  $I$  has a natural admissible structure; it is the Borel structure of  $H$  when considered as a metric space (in the uniform convergence topology). Again, I do not know whether or not every admissible subset of  $I^I$  has a natural admissible structure, but if  $F$  is admissible, we may endow it with an admissible structure in such a way so that it will be isomorphic to a subset of  $I$ .

The above theory may be applied to give a generalization of Kuhn's theorem [3] about optimal behavior strategies in games of perfect recall, to games in which there may be a continuum of alternatives at some of the moves.

A fuller account of the theory outlined above, together with proofs, will be published elsewhere.

#### REFERENCES

1. S. Banach, *Über analytisch darstellbare Operationen in abstrakten Räumen*, Fund. Math. vol. 17 (1931) pp. 283–295.
2. G. W. Mackey, *Borel structures in groups and their duals*, Trans. Amer. Math. Soc. vol. 85 (1957) pp. 134–165.
3. H. W. Kuhn, *Extensive games and the problem of information*, Contributions to the Theory of Games II, Princeton University Press, 1953, pp. 245–266.

HEBREW UNIVERSITY,  
JERUSALEM, ISRAEL