

Stability Theory for Solitary-Wave Solutions of Scalar Field Equations

Daniel B. Henry^{1*}, J. Fernando Perez^{2**}, and Walter F. Wreszinski^{2**}

1 Instituto de Matemática, Universidade de São Paulo, São Paulo, Brazil

2 Instituto de Física, Universidade de São Paulo, São Paulo, Brazil

Abstract. We prove stability and instability theorems for solitary-wave solutions of classical scalar field equations.

0. Introduction

In this paper, we study the stability of special travelling-wave (“solitary-wave”, [1]) solutions of classical scalar field equations of the form

$$\square\phi + U'(\phi) = 0. \quad (0-1)$$

This problem has attracted much attention recently in the physics literature ([2]), in part because classical solutions may be recovered from suitable expectation values of quantum fields in the classical limit ([3]).

Apart from the main motivation, which is to provide a simple and clear mathematical theory of stability for classical field equations, there is also a three-fold physical motivation. Firstly, most of the discussion in the physics literature ([2]), which is heuristically correct, relies on the linear theory. It may be shown, however, using methods of the present paper, that the latter is not applicable, because the linearized operator (on the natural Hilbert space, after proper “subtraction” of the zero mode) is skew-adjoint, a reflection of the fact that the mechanism of stability in these theories is dispersive, not dissipative (see also the discussion in [6] for K-dV equation). Secondly, the existing rigorous nonlinear stability theories ([4], generalized and corrected in [5], and [6]) are in principle applicable only to a class of equations (such as the K-dV equation) which may be treated either by inverse scattering theory ([6]), or which possess more than one scalar conservation law ([4], [5]), and are, therefore, unsuitable to describe, for instance, the stability of “kinks” of the nonlinear Klein-Gordon equation ([2]). Thirdly, and perhaps most importantly, the heuristic discussion disregards the somewhat delicate technical problems posed

* Partial financial support by FAPESP and FINEP.

** Partial financial support by CNPq.

by the zero-mode, which is always present due to translation invariance ([2]): this is also the reason why stability of the solitary-waves is a *form-stability* ([4], [6]).

The plan of the paper is as follows. In Sect. 1 we describe the relevant class of equations and solutions, state and prove our stability result (Theorem 1), and in Sect. 2 we prove a general instability result (Theorem 2). In Sect. 3 we provide a brief discussion of applications of Theorems 1 and 2 and prove, in particular, instability of the solitary-waves in higher dimensions which have been constructed by Parenti *et al.* ([13]) and Strauss ([17]). The Derrick–Strauss theorem ([7],[8]) is thereby revisited.

1. Stability

Let ϕ be a bounded static (time-independent) solution of (0-1), i.e., satisfying

$$\phi''(x) = U'(\phi(x)), \quad x \in \mathbb{R}. \tag{1-1}$$

We further assume

$$\phi'(x) > 0, \quad \forall x \in \mathbb{R}, \quad \phi'(x) \rightarrow 0, \tag{1-2}$$

$x \rightarrow \pm \infty$

and that there exist constants $-\infty < a_- < a_+ < \infty$ such that

$$\phi(x) \rightarrow a_- \quad \text{and} \quad \phi(x) \rightarrow a_+ \tag{1-3}$$

$x \rightarrow -\infty$ $x \rightarrow +\infty$

Since $U(\phi(x)) = \frac{1}{2}\phi'(x)^2 + \text{const}$, it follows that $U(a_-) = U(a_+)$. We normalize the energy of the “vacua” $\phi(x) \equiv a_+$ and $\phi(x) \equiv a_-$ to zero and assume

$$U(a_-) = U(a_+) = 0 \quad \text{and} \quad U(x) \geq 0, \quad x \in \mathbb{R}. \tag{1-4}$$

We also assume that

$$E_0 \equiv \int_{-\infty}^{\infty} (\frac{1}{2}\phi'^2 + U(\phi)) < \infty. \tag{1-5}$$

This means that ϕ has finite energy relative to the vacua. This point and the restriction to one space dimension are related to the Derrick–Strauss theorem ([7], [8]), see the discussion in Sect. 3. On U we further impose the following condition:

$$U \in C^2 \text{ in a neighbourhood of } [a_-, a_+] \quad \text{and} \quad U''(a_{\pm}) > 0. \tag{1-6}$$

We note that by the first of (1-2) we are describing “lumps” ([2]). “Antilumps” ($\phi'(x) < 0, x \in \mathbb{R}$) may be handled by reflection $x \rightarrow -x$.

Let $H^1_{loc}(\mathbb{R}) \equiv \left\{ \psi : \mathbb{R} \rightarrow \mathbb{R} : \int_{\Omega} dx [\psi'(x)^2 + \psi(x)^2] < \infty \text{ for every bounded region } \Omega \in \mathbb{R} \right\},$

$$H^1(\mathbb{R}) \equiv \left\{ \psi : \mathbb{R} \rightarrow \mathbb{R} : \int_{-\infty}^{\infty} dx [\psi'(x)^2 + \psi(x)^2] < \infty \right\}.$$

$H^1_{loc}(\mathbb{R})$ is the natural space for solutions of (0-1) ([12]), and we shall therefore assume $\phi \in H^1_{loc}(\mathbb{R})$. Such a ϕ defines (under the previous assumptions) a Hilbert sector in the language of ([12]), and it is natural to inquire upon stability within a sector, which means that “perturbed” solutions are required to satisfy $\begin{pmatrix} \psi - \phi \\ \psi_t \end{pmatrix} \in H^1(\mathbb{R}) \oplus L_2(\mathbb{R})$. Due to translation invariance, the following is (for $q > 0$) a natural distance function (see also [4]):

$$d_q(\psi)^2 \equiv \min_{-\infty < c < \infty} \int_{-\infty}^{\infty} dx [(\psi'(x) - \phi'(x+c))^2 + q(\psi(c) - \psi(x+c))^2]$$

We may now state our stability theorem:

Theorem 1. *There exist positive constants r, q and k such that if $u \in H^1_{loc}(\mathbb{R})$ satisfies*

$$(u_x(\cdot, t_0), u_t(\cdot, t_0)) \in L_2(\mathbb{R}) \times L_2(\mathbb{R})$$

and is a solution of (0-1) satisfying

$$d_q(u(\cdot, t_0)) < r$$

at some time t_0 , having energy

$$E = \int_{-\infty}^{\infty} dx [\frac{1}{2}u_t^2 + \frac{1}{2}u_x^2 + U(u)] < E_0 + kr^2,$$

then the solution exists for all t and has

$$d_q(u(\cdot, t)) \leq \sqrt{\frac{1}{k}(E - E_0)} < r$$

for all t .

The theorem is a consequence of the following

Proposition 1. *There exist positive constants q, k and r such that any solution $\psi \in H^1_{loc}(\mathbb{R})$ of (0-1) with $d_q(\psi(\cdot, t_0)) \leq r$ satisfies*

$$\int_{-\infty}^{\infty} dx [\frac{1}{2}\psi'(x)^2 + U(\psi(x))] \geq E_0 + kd_q(\psi)^2.$$

Proof. For clarity we divide the proof into a succession of steps:

1) $\phi'(x) = 0[e^{-\lambda_{\pm}|x|}]$ as $x \rightarrow \pm \infty$, $\lambda_{\pm} = \sqrt{U''(a_{\pm})} > 0$, since $(\phi, \phi') = (a_+, 0)$ or $(a_-, 0)$ are saddle points of the equilibrium equation (1-1).

2) The potential energy $V(\psi) = \int_{-\infty}^{\infty} dx (\frac{1}{2}\psi'^2 + U(\psi))$ has ϕ as a critical point:

$$(V'(\phi), \psi) = \int_{-\infty}^{\infty} dx (\phi' \psi' + U'(\phi)\psi) = 0 \text{ for all } \psi \in H^1(\mathbb{R})$$

and

$$(\psi, V''(\phi)\psi) = \int_{-\infty}^{\infty} dx (\psi'^2 + U''(\phi)\psi^2) = \int_{-\infty}^{\infty} dx (\psi A_{\phi} \psi),$$

where $A_\phi \equiv -d^2/dx^2 + U''(\phi(x))$ is a self-adjoint operator in $L_2(\mathbb{R})$, bounded below, with essential spectrum $\sigma_e(A_\phi) = [b, \infty]$, $b = \min(U''(a_+), U''(a_-)) > 0$ [[9], Theorem 16, pg. 1448 and Theorem 4, pg. 1438]. In particular, $0 \notin \sigma_e(A_\phi)$. If $\tau_c: \psi \rightarrow \psi(\cdot + c)$ is the translation operator in $L_2(\mathbb{R})$ then $\tau_c^{-1} A_{\tau_c \phi} \tau_c = A_\phi$, so $\sigma(A_{\tau_c \phi}) = \sigma(A_\phi)$ for all c .

3) $\phi'' = U'(\phi)$ so $\phi''' = U''(\phi)\phi'$, i.e. $A_\phi \phi' = 0$. By hypothesis $E_0 < \infty$, hence $\phi' \in L_2(\mathbb{R})$. Since $\phi' > 0$, 0 is the smallest eigenvalue of A_ϕ and it is simple, by standard methods.

4) Let $\beta > 0$ denote the first positive eigenvalue of A_ϕ , or b , whichever is smaller.

Then, for $\psi \in H^1(\mathbb{R})$, $\int_{-\infty}^{\infty} dx \psi \phi' = 0$ implies

$$(\psi, V''(\phi)\psi) = \int_{-\infty}^{\infty} dx (\psi'^2 + U''(\phi)\psi^2) \geq \beta \int_{-\infty}^{\infty} dx \psi^2.$$

The same estimate holds (with the same β) when ϕ is replaced by $\Phi(\cdot + c)$.

5) Suppose $q \geq U''(\phi(x))$ for all x and $\int_{-\infty}^{\infty} dx \psi \phi'(q - U''(\phi)) = 0$; then

$$(\psi, V''(\phi)\psi) > \frac{\beta}{(1 + K_q)^2} \int_{-\infty}^{\infty} dx \psi^2,$$

where

$$K_q \equiv \frac{\|\phi'\|_{L_2} \|\phi'(q - U''(\phi))\|_{L_2}}{\int_{-\infty}^{\infty} dx \phi'^2 (q - U''(\phi))}.$$

To see this, set $\psi = \alpha \phi' + \theta$, $\alpha = \text{constant}$, $\int_{-\infty}^{\infty} dx \phi' \theta = 0$. Then

$$(\psi, V''(\phi)\psi) = (\theta, V''(\phi)\theta) \geq \beta \int_{-\infty}^{\infty} dx \theta^2$$

and

$$0 = \alpha \int dx \phi'^2 (q - U''(\phi)) + \int dx \theta \phi' (q - U''(\phi)),$$

so

$$\|\psi\|_{L_2} \leq |\alpha| \|\phi'\|_{L_2} + \|\theta\|_{L_2} \leq (K_q + 1) \|\theta\|_{L_2}.$$

6) $q \int_{-\infty}^{\infty} dx (u(x) - \phi(x + c))^2 \rightarrow \infty$ as $c \rightarrow \pm \infty$, so there exists c with $d_q(u)^2 =$

$$\int_{-\infty}^{\infty} dx [(u'(x) - \phi'(x + c))^2 + q(u(x) - \phi(x + c))^2].$$

The minimum may be achieved at several values of c , but any one will serve, and we shall assume $c = 0$ for simplicity.

The derivative with respect to c must vanish at $c = 0$ and

$$\begin{aligned} 0 &= \int_{-\infty}^{\infty} dx [(u' - \phi')\phi'' + q(u - \phi)\phi'] = \int_{-\infty}^{\infty} dx (u - \phi)[q\phi' - \phi''] \\ &= \int_{-\infty}^{\infty} dx (u - \phi)\phi'(q - U''(\phi)), \end{aligned}$$

so with $\psi = u - \phi$ in step 5),

$$((u - \phi), V''(\phi)(u - \phi)) = (\psi, V''(\phi)\psi) \geq \frac{\beta}{(1 + K_q)^2} \int_{-\infty}^{\infty} dx \psi^2.$$

The same estimate holds with ϕ replaced by any translate $\phi(\cdot + c)$.

7) Let $\psi \in H^1(\mathbb{R})$. By an inequality of Sobolev type,

$$\|\psi\|_{L^\infty}^2 \leq \frac{2}{\sqrt{q}} \int_{-\infty}^{\infty} dx (\psi'^2 + q\psi^2).$$

Suppose ψ as in step 5). Choose $\varepsilon > 0$ so that $u''(a + s) > u''(a) - (\beta/2)(1 + K_q)^2$ for $a_- \leq a \leq a_+$ and $-\varepsilon \leq s \leq \varepsilon$. If $(\int dx (\psi'^2 + q\psi^2))^{1/2} \leq \frac{\varepsilon q^{1/4}}{\sqrt{2}} \equiv r$, it follows that

$\|\psi\|_{L^\infty} \leq \varepsilon$ and

$$\begin{aligned} V(\phi + \psi) - V(\phi) &= \int_{-\infty}^{\infty} dx \left\{ \frac{1}{2} \psi'^2 + U(\phi + \psi) - U(\phi) - U'(\phi)\psi \right\} \\ &\geq \int_{-\infty}^{\infty} dx \left\{ \frac{1}{2} \psi'^2 + \frac{1}{2} \left(U''(\phi) - \frac{\beta}{2(1 + K_q)^2} \right) \psi^2 \right\} \\ &\geq \frac{1}{2} \left[(\psi, V''(\phi)\psi) - \frac{\beta}{2(1 + K_q)^2} \int_{-\infty}^{\infty} dx \psi^2 \right] \\ &\geq \frac{\beta}{4(1 + K_q)^2} \int_{-\infty}^{\infty} dx \psi^2. \end{aligned}$$

Also $U(\phi + \psi) - U(\phi) - U'(\phi)\psi \geq -\frac{1}{2} B_1 \psi^2$, where

$$\begin{aligned} -B &= \inf_x U''(\phi(x)), \\ B_1 &= B + \frac{\beta}{2(1 + K_q)^2}. \end{aligned}$$

So

$$\begin{aligned} V(\phi + \psi) - V(\phi) &\geq \max \left\{ \frac{\beta}{4(1 + K_q)^2} \|\psi\|_{L^2}^2, \right. \\ &\quad \left. \frac{1}{2} \left(\int_{-\infty}^{\infty} dx (\psi'^2 + q\psi^2) - (B_1 + q) \|\psi\|_{L^2}^2 \right) \right\} \\ &\geq \frac{\delta}{2} \int_{-\infty}^{\infty} dx (\psi'^2 + q\psi^2) + \left\{ -\frac{1}{2} \delta (B_1 + q) + (1 - \delta) \frac{\beta}{4(1 + K_q)^2} \right\} \int_{-\infty}^{\infty} dx \psi^2 \end{aligned}$$

for any $0 \leq \delta \leq 1$. Choose δ such as to make

$$-\frac{1}{2} \delta (B_1 + q) + (1 - \delta) \frac{\beta}{4(1 + K_q)^2} = 0.$$

so as to get

$$V(\phi + \psi) - V(\phi) \geq K \int_{-\infty}^{\infty} dx(\psi'^2 + q\psi^2)$$

$$K = \frac{\beta/4}{\beta + (1 + K_q)^2(B_1 + q)}.$$

By (6), (7) and $V(\phi(\cdot + c)) = V(\phi)$ we obtain finally

$$d_q(u) \leq r \Rightarrow V(u) \geq V(\phi) + Kd_q(u)^2.$$

In particular, $E(u) \geq E_0 + Kd_q(u)^2$.

Proof of Theorem 1. Suppose $d(u(\cdot, 0)) < r$, $E = \int_{-\infty}^{\infty} dx(\frac{1}{2}u_t^2) + V(u) < V(\phi) + Kr^2$ at $t = 0$, and $u_{tt} - u_{xx} + U'(u) = 0$ for $t > 0$ (the equation is reversible, $t \rightarrow -t$, so we need only consider $t > 0$). There exists a unique mild solution with $\{u_x(\cdot, t), u_t(\cdot, t)\} \in L_2 \times L_2$ continuous in t and with constant energy E which exists on some maximal interval $0 \leq t < t_\infty$. If $t_\infty < +\infty$, $\{u_x(\cdot, t), u_t(\cdot, t)\}$ cannot converge in $L_2 \times L_2$ as $t \rightarrow t_\infty$ — so $\|U'(u(\cdot, t))\|_{L_2}$ must be unbounded as $t \rightarrow t_\infty$. But on $0 \leq t < t_\infty$, $V(\phi) + Kr^2 > E \geq V(\phi) + Kd(u)^2$ so $d(u(\cdot, t)) < r$. Thus for each t there is a $c = c(t)$ so that

$$\int_{-\infty}^{\infty} dx[(u_x - \phi'(x + c))^2 + q(u - \phi(x + c))^2] < r^2.$$

So

$$|U'(u(x, t))| \leq |U'(\phi(x + c))| + \text{const } |u(x, t) - \phi(x + c)|$$

for all $x \in \mathbb{R}$, so that $\|U'(u(\cdot, t))\|_{L_2}$ is uniformly bounded as $t \rightarrow t_\infty$. Hence $t_\infty = +\infty$ and the solutions exist for all $t \geq 0$ — and similarly for all $t \leq 0$.

Remark. If the initial values have $\{u_{xx}(\cdot, 0), u_{xt}(\cdot, 0)\} \in L_2 \times L_2$ as well, then $t \rightarrow \{u_x(\cdot, t), u_t(\cdot, t)\} \in L_2 \times L_2$ is continuously differentiable for all t and we have a strict solution (see, for example, [10], Th. VIII, 3.2).

2. Instability

The following theorem is featured along the lines of reference [16].

Theorem 2. Let X be a Banach space, $U \in X$ an open set containing 0, suppose $T : U \rightarrow X$ has $T(0) = 0$, and for some $p > 1$ and continuous linear L with spectral radius $r(L) > 1$.

$$\|T(x) - Lx\| = 0(\|x\|^p) \quad \text{as } x \rightarrow 0.$$

Then 0 is unstable as a fixed point of T .

In fact, we may estimate the direction in which points move away from 0 under successive applications of T . Choose any positive integer m and any μ , $0 < \mu < 1/\sqrt{2}$,




Fig. 1.

and define the cone

$$\Sigma = \Sigma(m, \mu) = \{v \in X : \|L^m v\| \leq \mu r^m \|v\|\},$$

where $r = r(L)$. If $0 < q < (1/\sqrt{2} - \mu)/(\mu + r^{-m}\|L^m\|)$ there exists $a_q > 0$ such that: given any a in $0 < a \leq a_q$, arbitrarily small $\epsilon_0 > 0$ and arbitrarily large $N_0 > 0$, there exist $N > N_0$ and $x \in X$ such that $\|x\| \leq \epsilon_0$, $\|T^n(x)\| \leq a$ for $0 \leq n \leq N$ and $\text{dist}(T^n(x), \Sigma) \geq qa$. In particular, $\|T^N(x)\| \geq qa$.

Remark. If Γ is a C^1 curve of fixed points of T with $0 \in \Gamma$, the tangent of Γ at 0 will be in $N(L - 1) \in \Sigma(M, 1/2)$ if $r^m \geq 2$. Choosing a small, so Γ is close to its tangent, we may conclude there exist arbitrarily small x such that, for some N , $\|T^n(x)\| \leq a$ when $0 \leq n \leq N$ but $\text{dist}(T^N(x), \Gamma \cap B_a) \geq \frac{1}{2}qa$. (Here B_a is the ball of radius a about 0.) Thus the points $\{T^n(x), n \geq 0\}$ not only move away from 0, but also away from Γ : the curve Γ is unstable.

Proof. Since $r = r(L) > 1$, $p > 1$, we may choose η in $0 < \eta < r^p - r$, and then choose K so $\|L^n\| \leq K(r + \eta)^n$ for all $n \geq 0$. There exist $a_0 > 0$ and b , so $\|x\| \leq a_0$ implies $x \in U$ and $\|T(x) - Lx\| \leq b\|x\|^p$. Given μ, m, q as above, choose $\delta > 0$ and $0 < a_q \leq a_0$ so small that

$$\Delta \equiv \frac{bK}{r^p - r - \eta} a_q^{p-1} < 1 \quad \text{and}$$

$$q \leq \frac{1 - \Delta}{\sqrt{2} + \delta} \cdot C_\delta - \Delta, \quad C_\delta = (1 - 3\delta - \mu(\sqrt{2} + \delta))/(\mu + r^{-M}\|L^M\|).$$

Choose $\lambda \in \sigma(L)$ with $|\lambda| = r$, say $\lambda = re^{i\theta}$, and choose $N \geq N_0$ so $r^N \geq a_0/\epsilon_0$ and $|e^{i(N+m)\theta} - 1| < \delta$. Since λ is in the boundary of the spectrum, it is an approximate eigenvalue and there exist $\zeta \in X + iX$ (the complexification of X) with $\|\zeta\| \geq 1$ but $\|L\zeta - \lambda\zeta\|$ arbitrarily small. Choose $\zeta = \xi + i\eta(\xi, \eta \text{ in } X)$ so $\|\xi\| = 1 \geq \|\eta\|$ and for $0 \leq n \leq N + m$ $\|\text{Re}(L^n \zeta - \lambda^n \zeta)\| = \|L^n \xi - r^n(\cos n\theta \xi - \sin n\theta \eta)\| \leq \delta r^n$. Note $\|L^n \xi\| \leq (\sqrt{2} + \delta)r^n$ and $L^{N+m} \xi \geq r^{N+m}(1 - 3\delta)$. We prove $\text{dist}(L^N \xi, \Sigma) \geq C_\delta r^N$, by contradiction. If $v \in \Sigma$ and $\|L^N \xi - v\| < C_\delta r^N$, then $\|v\| < (C_\delta + \sqrt{2} + \delta)r^N$ and $C_\delta r^N \|L^m\|$

$> \|L^m\| \|L^n \xi - v\| \geq \|L^{N+m} \xi - L^m v\| > r^{N+m}(1 - 3\delta) - \mu r^{m+N}(C_\delta + \sqrt{2} + \delta) = C_\delta r^N \|L^m\|$, a contradiction.

Now let $0 < a \leq a_q$, $R = (\sqrt{2} + \delta)/(1 - \Delta)$, and $\varepsilon = a/Rr^N$. Note $a \leq a_0$ and $\varepsilon \leq a_0/r^N \leq \varepsilon_0$. Let $x_0 = \varepsilon \xi$, $x_{n+1} = T(x_n)$ for $n \geq 0$; then

$$X_n = L^n x_0 + \sum_{k=0}^{n-1} L^{n-1-k}(T(x_k) - Lx_k).$$

Suppose $n \leq N$ and $\|x_k\| \leq \varepsilon Rr^k$ for $0 \leq k \leq n - 1$; this is certainly true for $n = 1$. It follows that

$$\|X_n - L^n x_0\| \leq \sum_{k=0}^{n-1} K(r + \eta)^{n-1-k} b(\varepsilon Rr^k)^p \leq \frac{bK(\varepsilon Rr^n)^p}{r^p - r - \eta} \leq \Delta \varepsilon Rr^n,$$

so $\|x_n\| \leq (\sqrt{2} + \delta + \Delta R)\varepsilon r^n = \varepsilon Rr^n$. By induction, $\|x_n\| \leq \varepsilon Rr^n \leq a$ for all $n \leq N$, and $\|x_N - L^N x_0\| \leq \Delta \varepsilon Rr^N$. Finally $\text{dist}(L^N x_0, \Sigma) = \varepsilon \text{dist}(L^N \xi, \Sigma) \geq C_\delta \varepsilon r^N$, so $\text{dist}(x_N, \Sigma) \geq (C_\delta/R - \Delta)\varepsilon Rr^N \geq qa$.

3. Applications

Solitary-wave solutions of (0-1) are of type

$$\Phi_0(x, t) = \phi_0(x - ut) \tag{3-1}$$

for suitable u . In order to apply Theorems 1 and 2 to the stability of (3-1), we first transform to a fixed-point problem by a Lorentz-transformation (which leaves (0-1) invariant)

$$x \rightarrow \xi = \frac{x - ut}{\sqrt{1 - u^2}}, \quad t \rightarrow \tau = \frac{t - ux}{\sqrt{1 - u^2}}.$$

(3-1) describes then (for $|u| < 1$) static (i.e., τ -independent) solutions of (0-1). In one space-dimension, $s = 1$, there are two types of nontrivial solutions of the form (3-1) (see Fig. 2):

a) solutions joining two distinct absolute minima a_- and a_+ of U : $\lim_{x \rightarrow -\infty} \phi(x) = a_- < a_+ = \lim_{x \rightarrow +\infty} \phi(x)$, with $U(a_-) = U(a_+)$; or solutions joining two distinct relative minima $b_- < b_+$, with $U(b_-) = U(b_+)$;

b) solutions around a relative minimum α of U : $\lim_{x \rightarrow -\infty} \phi(x) = \alpha = \lim_{x \rightarrow +\infty} \phi(x)$.

In case a), Theorem 1 applies directly yielding *form-stability* of the solitary-wave ([4], [6]).

The following condition follows from b):

c) $\text{sgn } \phi'(x)$ is not constant in $x \in \mathbb{R}$.

In view of the wealth of solutions for $s > 1$ ([13], [17]), we now state the appropriate analogous conditions for general s :

- c1) $\lim_{|x| \rightarrow \infty} \phi(x) = \alpha$ and $\alpha_1 \equiv U''(\alpha) > 0$ and $\Phi(\cdot)$ is not constant;
- c2) $[\phi(\cdot) - \alpha] \in L_2(\mathbb{R}^s)$;




Fig. 2.

In Theorem 2 $F_t(t \geq 0)$ is a (nonlinear) semigroup and T is F_t at any fixed $t > 0$. $L = DF_t(0) = \exp(tA)$, where A is the linearized operator

$$A \equiv \begin{pmatrix} 0 & 1 \\ -K & 0 \end{pmatrix}$$

on $H^1_{\alpha_1}(\mathbb{R}^s) \oplus L_2(\mathbb{R}^s)$, where

$$H^1_{\alpha_1}(\mathbb{R}^s) = \{u : \|u\|_{\delta_1}^2 \equiv (-\Delta + \alpha_1)^{1/2}u, (-\Delta + \alpha_1)^{1/2}u < \infty\}$$

and

$$K \equiv -\Delta + U''(\phi).$$

We have

$$A = \begin{pmatrix} 0 & 1 \\ -\Delta + \alpha_1 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ U''(\phi) - \alpha_1 & 0 \end{pmatrix}.$$

The first operator is skew adjoint on $D((-\Delta + \alpha_1)) \oplus D((-\Delta + \alpha_1)^{1/2})$ ([15], Theorem 1, pg. 26). The multiplication operator $(U''(\phi) - \alpha_1)$ is by c2) a relatively compact perturbation $(-\Delta + \alpha_1)$ on $L_2(\mathbb{R}^s)$, hence it is a compact operator on $H^1_{\alpha_1}(\mathbb{R}^s)$, and $\begin{pmatrix} 0 & 0 \\ U''(\phi) - \alpha_1 & 0 \end{pmatrix}$ is a compact operator on $H^1_{\alpha_1}(\mathbb{R}^s) \oplus L_2(\mathbb{R}^s)$. Therefore $|\text{Re } \sigma(A)| \leq \text{constant}$, and $L = \exp(tA)$ exists as a semigroup of bounded linear operators. We have now

Proposition 2. Under assumption (c) (for $s = 1$) or c1) and c2) (for $s > 1$), K has an eigenvalue $e_0 < 0$ and A has an eigenvalue $\sqrt{|e_0|} > 0$, so $r(L) > 1$.

Proof. It follows from c1) and c2) that $(U''(\phi) - \alpha_1)$ is a relatively compact perturbation of $(-\Delta + \alpha_1)$, as remarked above. Hence $\sigma(K) \setminus [\alpha_1, \infty]$ consists of point eigenvalues of finite multiplicities which can accumulate at most at α_1 . We have

$$K \partial_{x_i} \phi = 0, \quad i = 1, \dots, s.$$

Hence (by c3)) zero is a discrete eigenvalue, and the bottom of $\sigma(K)$ is an eigenvalue e_0 . By ([12], theorems XIII-43 and XIII-45), K is ergodic and by ([12], theorem

XIII-43) e_0 is a *simple* eigenvalue and the corresponding eigenfunction $\psi \in D(K) = D(\Delta)$ is strictly positive. But by c) (for $s = 1$) $\partial_x \phi \neq \lambda \Psi$. For $s > 1$, $\partial_{x_i} \phi \neq \lambda \Psi$ for all $i = 1, \dots, s$ because Ψ is simple and $\partial_{x_i} \phi \neq \lambda \partial_{x_j} \phi$ for some $i \neq j$ (i.e., the zero eigenvalue is degenerate). To see this, suppose $\sum_1^s \lambda_j \partial_{x_j} B \equiv 0$ for some constant $\lambda \in \mathbb{R}^s$.

This implies

$$\frac{d}{dt} \phi(x + \lambda t) = 0, \quad \phi(x) = \phi(x + \lambda t)$$

for all x, t : if $\lambda \neq 0$, let $t \rightarrow \infty$ and conclude from c1) that $\phi(x) = \alpha$ for all x . Hence, in all cases, $e_0 < 0$. Let $\beta = -|e_0|$. The vector

$$v = \begin{pmatrix} \psi \\ \sqrt{\beta} \psi \end{pmatrix} \in L_2(\mathbb{R}^s) \oplus L_2(\mathbb{R}^s)$$

is an eigenvector of A corresponding to eigenvalue $\sqrt{\beta} > 0$. But $\psi \in D(\Delta)$, hence $\psi \in H_{x_1}^1(\mathbb{R}^s)$ and $v \in H_{x_1}^1(\mathbb{R}^s) \oplus L_2(\mathbb{R}^s)$.

The above proposition proves instability for case b) (and $s = 1$) under assumption (3). There are many examples in higher dimensions ([13], [17]). For an explicit example in dimension $s = 3$, consider the following potential ([13]):

$$U(\phi) = \begin{cases} g[(\phi - c)^3 - 3(\phi - c)^4], & c \leq \phi \leq 1 + c, \\ g(\phi - c)^4, & \text{for } \phi \leq c, \\ f(\phi), & \text{for } \phi \geq 1 + c, \end{cases}$$

where $f \in C^2$, smoothly matched at $\phi = 1 + c$, with $|f''(\phi)| \leq \text{const}(1 + \phi^2)$, $g > 0$ and c is a given constant. $U \in C^2$, and $\phi = c$ is a relative minimum. The function

$$\phi(r) = c + \left[1 + \left(\frac{r}{a} \right)^2 \right]^{-1}$$

with $a^2 = 2/3g$, is a radial solution of (0-1). It satisfies c1) and c2). Proposition 2, coupled with Theorem 2, implies therefore that the above solution is unstable, as conjectured in [13]. Similarly, the (infinite series of) radial solitary-wave solutions of (0-1) constructed by Strauss in [17] are *unstable*, by the same reasoning.

The above results lead us to revisit the Derrick–Strauss theorem ([7], [8]). The latter states (in the form originally proposed in [7]) that solitary-wave solutions of scalar field theories do not exist for $s > 1$, *provided* they have finite energy relative to the absolute minimum (or minima) of U . However, solutions joining two relative minima of U (as b_- and b_+ in Fig. 1) are stable and define a Hilbert sector, although they have infinite energy relative to the absolute minima. Hence, the finite energy property does not seem to be relevant. If we accept this, there exist solutions around relative minima of U (as defined in b)) both for $s = 1$ and $s > 1$, as the previous examples show. The latter are, however, *unstable*. Stability seems therefore to be the main issue.

References

1. Lax, P. D.: *Comm. Pure Appl. Math.* **21**, 467 (1968)
2. Coleman, S.: Classical lumps and their quantum descendants. Lectures at the 1975 International School of Subnuclear Physics "Ettore Majorana" (Erice), in "New Phenomena in Subnuclear Physics," Zichichi A. (ed.). New York: Plenum Press, 1977
3. Hepp, K.: *Comm. Math. Phys.* **35**, 265 (1974)
4. Benjamin, T. B.: *Proc. R. Soc. (London)* **A328**, 153 (1972)
5. Bona, J. L.: *Proc. R. Soc. (London)* **A344**, 363 (1975)
6. Scharf, G., Wreszinski, W. F.: *Ann. Phys. (N.Y.)* **134**, 56 (1981)
7. Derrick, G.: *J. Math. Phys.* **5**, 1252 (1964)
8. Strauss, W.: Nonlinear invariant wave equations. In: *Invariant Wave Equations*, Velo, G., Wightman, A. S. (eds.), Berlin, Heidelberg, New York: Springer 1977
9. Dunford, N., Schwartz, J.: *Linear operators*, Vol. II. New York: Interscience 1963
10. Martin, R. H.: *Nonlinear operators and differential equations in Banach spaces*. New York: Wiley 1976
11. Reed, M., Simon, B.: *Methods of modern mathematical physics*, Vol. IV. New York: Academic Press 1978
12. Parenti, C., Strocchi, F., Velo, G.: In: *Invariant Wave Equations*, Velo, G., Wightman, A. S.: (eds.), Berlin, Heidelberg, New York: Springer 1977, and *Commun. Math. Phys.* **53**, 65 (1977)
13. Parenti, C., Strocchi, F., Velo, G.: *Dynamical charges and symmetries in non-linear classical field theory*. Bielefeld Preprint
14. Chernoff, P. R., Marsden, J. E.: *properties of infinite dimensional Hamiltonian systems*. Berlin, Heidelberg, New York: Springer 1974
15. Marsden, J., McCracken, M.: *The Hopf bifurcation and its applications*. Berlin, Heidelberg, New York: Springer 1976
16. Henry, Daniel B.: *Stability and instability by the linear approximation*, Proc. of the 12th. Brazilian Analysis Seminar (Soc. Bras. de Matematica, Rio de Janeiro)
17. Strauss, W. A.: *Commun. Math. Phys.* **55**, 149 (1977)

Communicated by J. Moser

Received August 28, 1981

