

GROTHENDIECK AND WITT RINGS OF HERMITIAN FORMS OVER DEDEKIND RINGS

MANFRED KNEBUSCH, ALEX ROSENBERG AND ROGER WARE

The prime ideal theory of the Grothendieck and Witt ring of non-degenerate hermitian forms over a Dedekind ring with involution is studied. The relationship of these rings to those defined over the quotient field of the Dedekind ring is also investigated.

The main goal of this paper is to extend the structure theory for Witt rings over fields of Pfister [18] and Harrison-Leicht-Lorenz ([10], [16]) to the Grothendieck ring $K(C, J)$ and the Witt ring $W(C, J)$ of a Dedekind ring C with an involution J . Since the case $J = \text{identity}$ is allowed, the Grothendieck and Witt rings of [12] are included. We shall see that the main theorems of Pfister and Harrison-Leicht-Lorenz remain true for $W(C, J)$ and that if J is the identity they are also true for $K(C, J)$. However, for $K(C, J)$ with $J \neq \text{identity}$ there is some deviation: there may be p -torsion for primes $p \neq 2$ and there may be nilpotent elements which are not torsion (Example 1.3). This fact has been overlooked in [13].

In §1 we extend some elementary results of [12, §11, §13] to the case $J \neq \text{identity}$. We conjecture that they are well known to the specialists but we did not find an appropriate reference in the literature. We show that the canonical map from $W(C, J)$ to the Witt ring $W(L, J)$ of the quotient field L of C is injective and give some information about the kernel $A(C, J)$ of the map $K(C, J) \rightarrow K(L, J)$. Since the exact determination of $A(C, J)$ is not needed for our structure theory we delay this matter to §4, where such a determination is given along the same lines as in [12, §11.2]. We then show that $W(C, J)$ is the intersection of certain subrings $W(C_i, J)$ of $W(L, J)$ which are Witt rings for abelian groups of exponent 2 in the sense of [14, Def. 3.12] and we describe the image $K'(C, J)$ of $K(C, J)$ in $K(L, J)$ in an analogous way.

We are thus led to study subrings T of an "abstract" Witt ring R for an arbitrary abelian q -group [14, Def. 3.12]. If T is the intersection of a family $\{T_\alpha\}$ of subrings of R which are also Witt rings for some abelian q -groups, the entire prime ideal theory of R remains true for T .

In §3 we show that if T is either $K(C, J)$ or $W(C, J)$ then the group of units of T is generated by $1 + \text{Nil } T$ and the rank one spaces

The second author was partially supported by NSF Grant GP-25600, and the third author was partially supported by NSF Grant GP-23915.

over (C, J) .

The results of [14] are used throughout.

1. Elementary facts about $K(C, J)$ and $W(C, J)$. (cf. [12, §11, §13.3], [9].)

In this paper C will always denote a Dedekind ring and J will be an involution on C which is allowed to be the identity. The quotient field of C will be denoted by L and J will also denote the unique extension of our involution to L . Further, F denotes the fixed field under J and A denotes the intersection $F \cap C$. Since C is integral over A ($c^2 - (J(c) + c)c + J(c)c = 0$ for all c in C), A is also a Dedekind ring and C is the integral closure of A in L . The unadorned \otimes means \otimes_c .

Throughout we use the notations of [14]. In particular, we often write \bar{x} for the value $J(x)$ of some x in L under J . A space (E, Φ) , or more briefly E , over (C, J) always means a finitely generated projective C -module E equipped with a non degenerate J -hermitian form Φ (see [14, Sec. 1], Φ is linear in the first argument and anti-linear in the second). A space E over (C, J) is called *metabolic* if E contains a direct summand V as C -module with $V = V^\perp$ (see [14, Sec. 1]). By $S(C, J)$ we denote the semiring of isometry classes of spaces over (C, J) , by $K(C, J)$ we denote the corresponding Grothendieck ring, and by $W(C, J)$ the *Witt ring* of (C, J) , i.e., the residue class ring $K(C, J)/KM(C, J)$, where $KM(C, J)$ denotes the Grothendieck group of metabolic spaces over (C, J) which is an ideal in $K(C, J)$ [14, Cor. 1.6]. We write $[E]$ for the image of the isometry class of a space E in $K(C, J)$.

For any finitely generated projective C -module U we denote by U^* the group $\text{Hom}_c(U, C)$ with C -module structure defined such that

$$\langle u, cu^* \rangle = \bar{c} \langle u, u^* \rangle$$

for all u in U , u^* in U^* , c in C . We further denote by $H(U)$ the *hyperbolic* space $U \oplus U^*$ with the hermitian form Φ characterized by $\Phi(U \times U) = \Phi(U^* \times U^*) = 0$ and $\Phi(u, u^*) = \langle u, u^* \rangle$ for u in U , u^* in U^* . The elements of $KM(C, J)$ all have the form $[H(U)] - [H(V)]$ ([14, Lemma 1.3(i)]). For the space $H(C)$ corresponding to the free C -module C we simply write H . Analogous notations will be used over (L, J) .

LEMMA 1.1. *A space E over (C, J) is metabolic if the space $L \otimes E$ over (L, J) is metabolic. Hence the canonical map from $W(C, J)$ to $W(L, J)$ induced by $E \mapsto L \otimes E$, is injective.*

Proof. We repeat the well known argument for the convenience

of the reader. Let W be a subspace of $L \otimes E$ with $W^\perp = W$ and consider E as a subset of $L \otimes E$. Then $V = W \cap E$ is a submodule of E with $V = V^\perp$ and V is also a direct summand of E since E/V is finitely generated and torsion free, hence projective [7, Prop. 4.1, p. 133]. The last assertion follows from the fact that over (L, J) a space M whose class in $W(L, J)$ is zero is metabolic ([3, §4] and in case of characteristic 2 and $J = \text{identity}$, [12, Lemma 8.2.2, p. 119] or [17]).

We denote by $\Lambda(C, J)$ the kernel of the canonical map $[E] \mapsto [L \otimes E]$ from $K(C, J)$ to $K(L, J)$. Since the natural map $W(C, J) \rightarrow W(L, J)$ is injective it is clear that $\Lambda(C, J)$ is contained in $KM(C, J)$. Hence any element x of $\Lambda(C, J)$ has the form

$$x = [H(U)] - [H(V)]$$

with $[H(U) \otimes L] = [H(V) \otimes L]$ in $K(L, J)$. Thus for some space M over (L, J) there is an isometry $M_\perp(H(U) \otimes L) \cong M_\perp(H(V) \otimes L)$ so that $\text{rank } U = \text{rank } V$. Conversely, it is evident that all such x lie in $\Lambda(C, J)$. Now there exist ideals \mathfrak{a} and \mathfrak{b} of C such that $U \cong \mathfrak{a} \oplus r \times C$ and $V \cong \mathfrak{b} \oplus r \times C$ for some $r \geq 0$ where $r \times C$ denotes the free C -module of rank r [22] or [20, Thm.1]. Hence by [14, Lemma 1.3(iii)],

$$x = [H(\mathfrak{a})] - [H(\mathfrak{b})] = [H(\mathfrak{a})] + [H(\mathfrak{b}^{-1})] - ([H(\mathfrak{b})] + [H(\mathfrak{b}^{-1})]) .$$

Recalling that

$$(*) \quad \mathfrak{a}_1 \oplus \mathfrak{a}_2 \cong C \oplus \mathfrak{a}_1 \mathfrak{a}_2$$

for any ideals $\mathfrak{a}_1, \mathfrak{a}_2$ of C and again using Lemma 1.3(iii) of [14], we obtain $x = [H(\mathfrak{a}\mathfrak{b}^{-1})] - [H]$.

We denote by $\text{Pic}(C)$ the ideal class group of C , by $\text{Pic}(C)^G$ the subgroup of all elements invariant under $G = \{1, J\}$, and by $(1+J)\text{Pic}(C)$ the subgroup of all classes $(\mathfrak{a}\bar{\mathfrak{a}})$. Our consideration about $\Lambda(C, J)$ shows that we have a surjective map

$$f: \text{Pic}(C) \longrightarrow \Lambda(C, J)$$

(cf. [12, 11.1.4, p. 136], [9, Th. 1]) which sends an ideal class (\mathfrak{a}) in $\text{Pic}(C)$ onto $[H(\mathfrak{a})] - [H]$. By (*) we see that $f(\mathfrak{a}\mathfrak{b}) = f(\mathfrak{a}) + f(\mathfrak{b})$.

PROPOSITION 1.2.

- (i) $(1 + J)\text{Pic}(C) \subset \text{Ker } f \subset \text{Pic}(C)^G$.
- (ii) If $J = \text{identity}$ then $2\Lambda(C, J) = 0$.
- (iii) $\Lambda(C, J)^2 = 0$ (cf. [9, Prop. 5]).

Proof. (i) Since for any ideal \mathfrak{a} of C the C -modules \mathfrak{a}^* and $\bar{\mathfrak{a}}^{-1}$ are isomorphic, we have $H(\mathfrak{a}) \cong H(\bar{\mathfrak{a}}^{-1})$ and hence $f(\mathfrak{a}) = f(\bar{\mathfrak{a}}^{-1})$. Thus $f(\mathfrak{a}\bar{\mathfrak{a}}) = 0$ for all ideals \mathfrak{a} . On the other hand, if \mathfrak{a} is an ideal

with $f(\mathfrak{a}) = 0$ then there exists a space E over (C, J) with $H(\mathfrak{a}) \perp E \cong H \perp E$. Comparing the Steinitz-invariants (= highest exterior powers) of both sides we see that $(\mathfrak{a}\bar{\mathfrak{a}}^{-1}) = 1$ in $\text{Pic}(C)$, hence $(\mathfrak{a}) = (\bar{\mathfrak{a}})$.

Statement (ii) is now clear since $2 \text{ Pic}(C) \subset \text{Ker } f$.

(iii) For two ideals $\mathfrak{a}, \mathfrak{b}$ we have

$$H(\mathfrak{a}) \otimes H(\mathfrak{b}) \cong H(\mathfrak{a} \otimes (\mathfrak{b} \oplus \bar{\mathfrak{b}}^{-1})) \cong H(\mathfrak{a}\mathfrak{b}) \perp H(\mathfrak{a}\bar{\mathfrak{b}}^{-1})$$

[14, Prop. 1.5 and Prop. 1.3]. From this one computes

$$\begin{aligned} f(\mathfrak{a})f(\mathfrak{b}) &= [H(\mathfrak{a}\mathfrak{b})] + [H(\mathfrak{a}\bar{\mathfrak{b}}^{-1})] - 2[H(\mathfrak{a})] - 2[H(\mathfrak{b})] + 2[H] \\ &= f(\mathfrak{a}\mathfrak{b}) + f(\mathfrak{a}\bar{\mathfrak{b}}^{-1}) - 2f(\mathfrak{a}) - 2f(\mathfrak{b}) \\ &= f(\mathfrak{b}^{-1}\bar{\mathfrak{b}}^{-1}) = 0. \end{aligned}$$

EXAMPLE 1.3. Let C be the ring of all integers of a quadratic number field with automorphism J . We have $A = \mathbb{Z}$, hence

$$(1 + J)\text{Pic}(C) = 0,$$

so that $2 \text{ Pic}(C)^G = 0$. Hence $\Lambda(C, J)$ is a torsion group whose part prime to 2 equals the part prime to 2 of $\text{Pic}(C)$. Thus we obtain many examples of Dedekind rings (C, J) with $K(C, J)$ having odd torsion.

Furthermore, there are Dedekind rings (C, J) such that $\Lambda(C, J)$ is not torsion at all. To obtain an example let Y be an elliptic curve over the complex numbers and $X = Y - p$ for some point p of Y . Since X is a non singular affine curve its affine ring is a Dedekind ring C with an involution J induced by the inverse map j of the abelian group Y with unit p . There is a canonical isomorphism $\text{Pic}(C) \cong Y$ which carries the induced action of J on $\text{Pic}(C)$ to j . Since as an abelian group Y is $S^1 \times S^1$, there are only four fixed elements in Y and Y is not torsion. This yields an example with $\Lambda(C, J)$ not torsion.

REMARK 1.4. In [9, Cor. to Th. 4] and [12, §11] it has been shown that for $J = \text{identity}$ the kernel of f is generated by $2 \text{ Pic}(C)$ and the classes of all maximal ideals \mathfrak{P} with $2 \in \mathfrak{P}$. By the same method we show in §4 that for $J \neq \text{identity}$, $\text{Ker } f$ is generated by the image of $\text{Pic}(A)$ and all maximal ideals \mathfrak{P} with $\mathfrak{P}^2 = C(\mathfrak{P} \cap A)$.

For each maximal ideal \mathfrak{p} of A we set $C_{\mathfrak{p}} = A_{\mathfrak{p}} \otimes_A C =$ the localization of C with respect to \mathfrak{p} , which is a semi-local Dedekind ring. The involution on $C_{\mathfrak{p}}$ induced by J will also be denoted by J . We consider the $C_{\mathfrak{p}}$ as subrings of L and have $C_{\mathfrak{p}} = A_{\mathfrak{p}}C$. Let $\text{Im } S(C, J)$ and $\text{Im } S(C_{\mathfrak{p}}, J)$ be the images of $S(C, J)$ and $S(C_{\mathfrak{p}}, J)$ in $S(L, J)$.

LEMMA 1.5. $\text{Im } S(C, J) = \bigcap_{\mathfrak{p}} \text{Im } S(C_{\mathfrak{p}}, J)$ where \mathfrak{p} runs through the set, $\text{Max } A$, of all maximal ideals of A .

Proof. The assertion means that a space over (L, J) , which contains a (C_p, J) -space of full rank for all p in $\text{Max } A$, also contains a (C, J) -space of full rank. This is elementary lattice theory (cf. [6, Thm. 3, p. 54]).

By Lemma 1.1 we regard $W(C, J)$ and all $W(C_p, J)$ as subrings of $W(L, J)$. Since $\text{Pic}(C_p) = 0$ for each p and hence by Proposition 1.2, $\Lambda(C_p, J) = 0$, we similarly regard the rings $K(C_p, J)$ as subrings of $K(L, J)$. We denote by $K'(C, J)$ the image of $K(C, J)$ in $K(L, J)$.

LEMMA 1.6.

- (i) $K'(C, J) = \bigcap_p K(C_p, J)$
- (ii) $W(C, J) = \bigcap_p W(C_p, J)$

where in both equations p runs through the maximal ideals of A .

Proof. Since all projective modules over C_p are free, $KM(L, J)$ and all $KM(C_p, J)$ coincide with the additive group generated by $[H]$. Evidently $KM(C, J)$ also maps onto $KM(L, J)$. Thus as subsets of $W(L, J)$ we get $W(C_p, J) = K(C_p, J)/KM(L, J)$ and

$$W(C, J) = [K(C, J)/\Lambda(C, J)]/[KM(C, J)/\Lambda(C, J)] = K'(C, J)/KM(L, J)$$

so that the two assertions of the lemma are equivalent.

We proceed to prove (ii). Let x be an element of $\bigcap W(C_p, J)$ and (V, ϕ) a space over (L, J) representing x . As for any (L, J) -space there exists a finite "exceptional set" $T \subset \text{Max } A$ such that V contains a space over (C_p, J) of full rank for all $p \notin T$. If $T \neq \emptyset$ let p_0 be a prime ideal in T . Since the class of V lies in $W(C_{p_0}, J)$ there exists a space V_0 over (L, J) containing a (C_{p_0}, J) -space of full rank and metabolic (L, J) -spaces U, U_0 such that $V \perp U \cong V_0 \perp U_0$ [14, Lemma 1.4]. Since any metabolic space over (L, J) contains (C_p, J) -spaces of full rank for all p in $\text{Max } A$ (see the definition of metabolic spaces in [14, Sec. 1]) it follows that $V \perp U$ is a space over (L, J) which represents x and has an exceptional set contained in $T - \{p_0\}$. After a finite number of such steps we obtain a space \tilde{V} representing x and containing (C_p, J) -spaces of full rank for all p . Hence by Lemma 1.5 \tilde{V} contains a (C, J) -space of full rank. Thus $\bigcap W(C_p, J) \subset W(C, J)$. The reverse inclusion is clear.

REMARK. Using the cancellation theorem for hermitian forms over fields [3, Thm. 1, p. 71], Hilfsatz 13.3.3 of [12] can be extended to the case $J \neq \text{identity}$, stating that if a space V over (L, J) contains a (C_p, J) -space of full rank then the same is true for all spaces V' with the same class in $W(L, J)$. This offers a shorter proof of Lemma 1.6(ii).

For any p in $\text{Max } A$ we have either $pC = \mathfrak{P}$ or $= \mathfrak{P}^2$ or $= \mathfrak{P}\overline{\mathfrak{P}}, \mathfrak{P} \neq$

$\bar{\mathfrak{P}}$, with \mathfrak{P} in $\text{Max } C$ [5, Th. 2, p. 42].

LEMMA 1.7. *If $\mathfrak{p}C = \mathfrak{P}\bar{\mathfrak{P}}$ with $\mathfrak{P} \neq \bar{\mathfrak{P}}$ or $\mathfrak{p}C = \mathfrak{P}^2$ then $K(C_{\mathfrak{p}}, J) = K(L, J)$ and $W(C_{\mathfrak{p}}, J) = W(L, J)$.*

Proof. We denote the norm function $x \mapsto x\bar{x}$ ($x \mapsto x^2$ if $J = \text{identity}$) by N . Since $K(L, J)$ and $W(L, J)$ are both generated by one dimensional forms it suffices to show that any coset of $F^* \text{ mod } NL^*$ contains a unit of $C_{\mathfrak{p}}$. Now, the coset of an element x of $F^* \text{ mod } NL^*$ contains a unit of $C_{\mathfrak{p}}$ if and only if the coset of x in $\hat{F}_{\mathfrak{p}}^* \text{ mod } N\hat{L}_{\mathfrak{p}}^*$ contains a unit of $\hat{C}_{\mathfrak{p}}$ ($\hat{F}_{\mathfrak{p}}, \hat{L}_{\mathfrak{p}}, \hat{C}_{\mathfrak{p}}, \dots$ denote the completions of F, L, C, \dots with respect to \mathfrak{p}). Evidently this is true if $\mathfrak{p}C = \mathfrak{P}^2$. If $\mathfrak{p}C = \mathfrak{P}\bar{\mathfrak{P}}$ with $\mathfrak{P} \neq \bar{\mathfrak{P}}$ then $\hat{L}_{\mathfrak{p}} = \hat{F}_{\mathfrak{p}} \times \hat{F}_{\mathfrak{p}}$ and the involution of $\hat{L}_{\mathfrak{p}}$ corresponds to the involution $(a, b) \mapsto (b, a)$ of $\hat{F}_{\mathfrak{p}} \times \hat{F}_{\mathfrak{p}}$ [14, Example 1.7]. Thus $N(\hat{L}_{\mathfrak{p}}^*) = \hat{F}_{\mathfrak{p}}^*$ and the assertion is also true in this case.

REMARK 1.8. If $\mathfrak{p}C = \mathfrak{P}$ then $K(C_{\mathfrak{p}}, J) \neq K(L, J)$ and $W(C_{\mathfrak{p}}, J) \neq W(L, J)$. In fact, one has an exact sequence

$$0 \longrightarrow W(C_{\mathfrak{p}}, J) \longrightarrow W(L, J) \xrightarrow{\partial_{\mathfrak{p}}} W(C/\mathfrak{P}, J) \longrightarrow 0$$

with residue class form homomorphism $\partial_{\mathfrak{p}}$ (cf. [21], [12, §13.3], [19]).

Combining Lemmas 1.6 and 1.7 we get

PROPOSITION 1.9.

- (i) $K'(C, J) = \bigcap_{\mathfrak{p}} K(C_{\mathfrak{p}}, J)$
- (ii) $W(C, J) = \bigcap_{\mathfrak{p}} W(C_{\mathfrak{p}}, J)$

where in both equations \mathfrak{p} runs through all maximal ideals of A with $\mathfrak{p}C$ a maximal ideal of C .

2. Subrings of Witt rings. Let q be a rational prime and G an abelian q -group. In [14, Def. 3.12] we have called a commutative ring R a Witt ring for G if R is a homomorphic image of the integral group ring $\mathbb{Z}[G]$ and the torsion subgroup, R_t , of R is q -primary. Let us recall some facts proved for a Witt ring R in [14]: There is only one prime ideal M_0 of R which contains q . The ideal M_0 is of finite index q (i.e., $R/M_0 \cong F_q^{(v)}$) and contains all minimal prime ideals of R . Moreover, any maximal ideal $M \neq M_0$ of R properly contains a unique prime ideal and this prime ideal is a minimal prime ideal. All the zero divisors of R lie in M_0 , the ring R is connected (i.e., has no idempotents other than 0 or 1) and $R_i \neq 0$ if and only if M_0 consists entirely of zero divisors. R is integral over \mathbb{Z} and hence R is a Jacobson ring [5, p. 67] of Krull dimension, $\dim R \leq 1$. In particular, its Jacobson radical, $\text{Rad } R$, coincides with its nil radical,

¹ F_q denotes the field of q -elements.

Nil R . In addition, $R_i = \text{Nil } R$ if and only if $Z \rightarrow R$ is injective and $R_i = R$ otherwise. In the former case the minimal prime ideals P of R are characterized by $P \cap Z = 0$ and in the latter case R is local and M_0 is the only prime ideal of R .

Let C be a Dedekind ring with involution J and quotient field L . It is pointed out in [14, Remark 3.11] that $K(L, J)$ and $W(L, J)$ are Witt rings for an abelian group of exponent two. By Lemma 1.1, $W(C, J)$ is a subring of $W(L, J)$ and by Proposition 1.2 the natural map $K(C, J) \rightarrow K(L, J)$ has a kernel $\Lambda(C, J)$ which is small in the sense then $\Lambda(C, J)^2 = 0$, and if J is the identity, in addition $2\Lambda(C, J) = 0$. We are thus led to consider subrings T of a Witt ring R for an abelian q -group.

The following Lemma follows easily from the properties of R described above, the Lying over theorem [5, Th. 1, p. 38], and [14, Lemma 2.5]. Its proof will be omitted.

LEMMA 2.1. *Let T be a subring of a Witt ring R for an abelian q -group. Then*

- (i) *T is integral over Z , T is a Jacobson ring, and T is connected.*
- (ii) *T has a unique prime ideal $M_{0,T}$ containing q and $T/M_{0,T} \cong F_q$.*
- (iii) *If $M_{0,T}$ is not the only prime ideal of T then every maximal ideal properly contains a prime ideal. In this case $Z \rightarrow T$ is injective and a prime ideal P is minimal if and only if $P \cap Z = 0$.*
- (iv) *$T = T_i$ if and only if $M_{0,T}$ is the only prime ideal of T .*
- (v) *T_i is q -primary.*
- (vi) *All zero divisors of T lie in $M_{0,T}$ and $M_{0,T}$ consists entirely of zero divisors if and only if $T_i \neq 0$.*
- (vii) *If $T_i \neq T$ then $\text{Nil } T = T_i$.*

In particular, the statements of the lemma are true for $W(C, J)$ and $K'(C, J)$. Because of the stated properties of $\Lambda(C, J)$ they also remain true for $K(C, J)$ if J is the identity. If J is not the identity (i), (ii), and (iii) remain true for $K(C, J)$ and (iv) is vacuous because $K(C, J)$ always contains Z . As shown by Example 1.3, (v), (vi), and (vii) are generally not true. However, even in this case $K(C, J)_i \subset \text{Nil } K(C, J)$ since $\Lambda(C, J)^2 = 0$ and $K'(C, J)_i \subset \text{Nil } K'(C, J)$.

REMARK 2.2. Suppose G is group of exponent two and $Z \rightarrow T$ is injective. If P is a minimal prime ideal of T there is a minimal prime ideal P' of R such that $P' \cap T = P$ [11, Ex. 1, p. 41]. Moreover, by [14, Remark 3.2], $R/P' \cong Z$ so T/P is a subring of Z and hence $T/P \cong Z$. Thus we see that any homomorphism from T to Z extends to one from R to Z and $\text{Nil } T$ is the intersection of the

kernels of the homomorphisms from T to \mathbf{Z} . Moreover, if M is a maximal ideal of T we must have $T/M \cong \mathbf{F}_p$ for some rational prime p . Thus if C is a Dedekind ring with involution J and P is a non maximal, whence minimal, prime ideal of $T = K(C, J)$ or $W(C, J)$ then $T/P \cong \mathbf{Z}$. If M is a maximal ideal of $T = K(C, J)$ or $W(C, J)$ then $T/M \cong \mathbf{F}_p$ for some rational prime p . Moreover, in these cases the ideal $M_{0,T}$ is exactly the ideal of forms of even rank.

At the beginning of this section we stated one property of Witt rings R which is not necessarily inherited by all subrings of R : a maximal ideal $M \neq M_0$ contains a unique minimal prime ideal. We obtained this property in [14] as a consequence of the following theorem: If \tilde{R} denotes the integral closure of the image $\bar{R} = 1 \otimes R$ of R in $Q \otimes R$ then \tilde{R}/\bar{R} is a q -group. Here, and until the end of §2, \otimes always denotes $\otimes_{\mathbf{Z}}$. This results from the rather evident fact, that R/pR is von Neumann regular for all $p \neq q$ (see [14]). Thus we are now looking for subrings T of R such that T/pT is von Neumann regular for all $p \neq q$. For any such subring it will be true that a maximal ideal $M \neq M_{0,T}$ contains a unique minimal prime ideal.

For any abelian group X and rational prime p we let $X[p]$ be the p -primary component of X .

LEMMA 2.3. *Let $T \subset R$ be an integral extension of commutative rings and p a rational prime such that $R[p] = 0$ and R/pR is von Neumann regular. Then T/pT is von Neumann regular if and only if $(R/T)[p] = 0$.*

Proof. By [14, Lemma 2.8], $\text{Nil}(T/pT) = 0$ if and only if $(R/T)[p] = 0$. Now by Lemma 2.7 of [14] the kernel of the map $T/pT \rightarrow R/pR$ is a nil ideal. Since R/pR is integral over T/pT , $\dim T/pT = \dim R/pR = 0$. Thus T/pT is von Neumann regular if and only if $\text{Nil}(T/pT) = 0$ [4, Ex. 16(d), p. 173], which proves the lemma.

COROLLARY 2.4. *Let R be a commutative ring and $\{T_i\}_{i \in I}$ a family of subrings such that R is integral over $T = \bigcap_{i \in I} T_i$. If for a rational prime p with $R[p] = 0$ all the rings $R/pR, T_i/pT_i$ are von Neumann regular then T/pT is also von Neumann regular.*

Proof. Consider the exact sequence

$$0 \longrightarrow R/T \longrightarrow \prod_{i \in I} R/T_i$$

of abelian groups. By Lemma 2.3, we have $(R/T_i)[p] = 0$ for all i so $(R/T)[p] = 0$. Hence Lemma 2.3 shows that T/pT is von Neumann regular.

THEOREM 2.5. *Let R be a Witt ring for an abelian q -group G and let $\{T_i\}_{i \in I}$ be a family of subrings of R such that each T_i is a Witt ring for an abelian q -group H_i . If $T = \bigcap_{i \in I} T_i$ then*

- (i) *T/pT is von Neumann regular for all rational primes $p \neq q$.*
- (ii) *If \tilde{T} denotes the integral closure of $\bar{T} = 1 \otimes T$ in $\mathbb{Q} \otimes T$ then \tilde{T}/\bar{T} is a q -group.*
- (iii) *Any maximal ideal of T distinct from $M_{0,T}$ properly contains a unique prime ideal of T which is a minimal prime ideal.*

Proof. Since R_i is q -primary, $R[p] = 0$ for all rational primes $p \neq q$. Moreover, by [14, Example 2.6] R/pR and T_i/pT_i are von Neumann regular rings. Statement (i) now follows immediately from Corollary 2.4. Statement (ii) is then a consequence of [14, Prop. 2.9]. Finally, if T contains a maximal ideal distinct from $M_{0,T}$ then by Lemma 2.1 (iv) and (vii), $\text{Nil } T = T_i$. Thus (iii) follows from (i) and [14, Cor. 2.10 and Th. 2.12].

EXAMPLES 2.6. (i) Let G be an abelian q -group and $\{H_i\}_{i \in I}$ a family of subgroups of G . Let K be an ideal of $\mathbb{Z}[G]$, $R = \mathbb{Z}[G]/K$, and for i in I , let $T_i = \mathbb{Z}[H_i]/(\mathbb{Z}[H_i] \cap K)$. If R is a Witt ring for G then each T_i has only q -torsion and so is a Witt ring for H_i . Hence $T = \bigcap T_i$ satisfies the conclusions of Theorem 2.5.

(ii) Let C be a Dedekind ring with involution J and quotient field L . Denote the fixed ring of J on C by A . By Proposition 1.9,

$$K(C, J) = \bigcap_{\mathfrak{p}} K(C_{\mathfrak{p}}, J), \quad W(C, J) = \bigcap_{\mathfrak{p}} W(C_{\mathfrak{p}}, J)$$

where \mathfrak{p} runs through all maximal ideals of A with $\mathfrak{p}C$ a maximal ideal of C . Since each $C_{\mathfrak{p}}$ is a local ring it follows that $K(C_{\mathfrak{p}}, J)$ and $W(C_{\mathfrak{p}}, J)$ are Witt rings for an abelian group of exponent two [14, Remark 3.11]. Hence the conclusions of Theorem 2.5 apply to $K(C, J)$ and $W(C, J)$. Since by Proposition 1.2(iii) the kernel $\lambda(C, J)$ of $K(C, J) \rightarrow K(L, J)$ is nilpotent, statement (iii) of Theorem 2.5 remains true for $K(C, J)$. In case J is the identity, Proposition 1.2(ii) asserts that $2\lambda(C, J) = 0$. Hence $\mathbb{Q} \otimes K(C, Id) = \mathbb{Q} \otimes K'(C, Id)$ and

$$K(C, Id)/pK(C, Id) \cong K'(C, Id)/pK'(C, Id)$$

for all odd p so that statements (i) and (ii) of Theorem 2.5 hold for $K(C, Id)$ also.

We finally remark that a shorter proof of Theorem 2.5(ii) and hence, using [14, Cor. 2.10 and Th. 2.12], also of Theorem 2.5(iii) can be given by way of

LEMMA 2.7. *Let R be a commutative ring, $\{T_i\}_{i \in I}$ a family of*

subrings and $T = \bigcap_{i \in I} T_i$. Let p be a rational prime such that $R[p] = (\tilde{T}_i/\bar{T}_i)[p] = 0$ for all i in I . Then $(\tilde{T}/\bar{T})[p] = 0$.

Proof. Since \mathbf{Q} is a flat \mathbf{Z} -module we may identify $\mathbf{Q} \otimes T_i$ and $\mathbf{Q} \otimes T$ with subrings of $\mathbf{Q} \otimes R$. Thus we regard $\bar{T}_i, \tilde{T}_i, \bar{T}$, and \tilde{T} as subrings of $\mathbf{Q} \otimes R$. Let \tilde{x} be an element of \tilde{T} with $p\tilde{x}$ in \bar{T} . Then for all i in I we have $\tilde{x} \in \tilde{T}_i$ and $p\tilde{x} \in \bar{T}_i$. Hence $\tilde{x} \in \bigcap_{i \in I} \tilde{T}_i$. There thus exist elements x_i in T_i and y in T such that $\tilde{x} = 1 \otimes x_i$ and $p\tilde{x} = 1 \otimes y$. Hence $px_i - y$ lies in R_i for all i in I . Thus there exist integers n_i with $(n_i, p) = 1$ such that $n_i px_i = n_i y$. Now there are integers a_i, b_i with $1 = a_i n_i + b_i p$. Multiplying this last relation by y yields elements z_i in T_i with $y = pz_i$ for all i in I . But since $R[p] = 0$ the relation $pz_i = pz_j$ shows $z_i = z_j$ for all i, j in I . Hence there is an element z in T with $y = pz$. Thus $p\tilde{x} = 1 \otimes pz$ and therefore $\tilde{x} = 1 \otimes z$ which shows that \tilde{x} lies in \bar{T} , i.e. $(\tilde{T}/\bar{T})[p] = 0$.

3. Units in $K(C, J)$ or $W(C, J)$. We first recall some elementary facts true for spaces over an arbitrary commutative ring C with involution J . For any space (E, Φ) over (C, J) there is an *inverse space* $(\hat{E}, \hat{\Phi})$ [3, Def. 8, p. 23] defined as follows: \hat{E} is a C -module on the additive group E with a new scalar multiplication $c \cdot x = J(c)x$ and the form $\hat{\Phi}$ on \hat{E} is defined by $\hat{\Phi}(x, y) = \Phi(y, x)$. Now assume that E has rank one. Then it is easily checked that the C -linear map $g: E \otimes \hat{E} \rightarrow C$ defined by $g(x \otimes y) = \Phi(x, y)$ is bijective and gives an isometry from $(E \otimes \hat{E}, \Phi \otimes \hat{\Phi})$ to the space (C, Ω) where $\Omega(c, d) = c\bar{d}$ for c, d in C (by a localization argument it suffices to check this for E a free space in which case the proof is straightforward). Since the space (C, Ω) has matrix (1) it is clear that the tensor product makes the set $S_1(C, J)$ of isometry classes of (C, J) -spaces of rank one into an abelian group. The subgroup of classes of free spaces of rank one can be identified with A^*/NC^* where A denotes the fixed ring of J and C^*, A^* are the groups of units of C, A .

REMARK 3.1. On the group $\text{Pic}(C)$ of projective C -modules of rank one, J induces an involution $(M) \mapsto (\hat{M})$, where again \hat{M} denotes the abelian group M with new scalar multiplication $c \cdot x = J(c)x$. Denote by ${}_{(1+J)}\text{Pic}(C)$ the subgroup of all (M) in $\text{Pic}(C)$ with $(M)(\hat{M}) = 1$. The "forgetful map" $(E, \Phi) \mapsto (E)$ from $S_1(C, J)$ to $\text{Pic}(C)$ yields an exact sequence

$$1 \longrightarrow A^*/NC^* \longrightarrow S_1(C, J) \longrightarrow {}_{(1+J)}\text{Pic}(C) \longrightarrow 1.$$

(The surjectivity on the right can easily be proved, cf. [12, §2.3]).

REMARK 3.2. Let C be an integral domain with quotient field L .

Any element of $\text{Pic}(C)$ can be represented by an invertible ideal α and $\hat{\alpha}$ is isomorphic with the ideal $\bar{\alpha} = J(\alpha)$. For any hermitian form Φ on α we have $\Phi(x, y) = dx\bar{y}$ for some d in L such that $d\alpha\bar{\alpha} \subset C$, since Φ is the restriction of an (L, J) -form on L . Note that Φ is non degenerate if and only if $d\alpha\bar{\alpha} = C$.

Now let C be connected. For any (C, J) -space (E, Φ) of rank n we define the *determinant*, $\det(E)$, of (E, Φ) to be the element of $S_1(C, J)$ represented by the space $(\bigwedge^n E, \Delta)$ with Δ defined by

$$\Delta(x_1 \wedge \cdots \wedge x_n, y_1 \wedge \cdots \wedge y_n) = \det(\Phi(x_i, y_j)) ,$$

which is again a nondegenerate space [3, §2]. The map

$$\det: S(C, J) \longrightarrow S_1(C, J)$$

is additive but it does not vanish on all hyperbolic spaces. Thus we have to modify this map in the following well known way: denote by $\mathbb{Z}/2\mathbb{Z} \cdot S_1(C, J)$ the set $\mathbb{Z}/2\mathbb{Z} \times S_1(C, J)$ with the twisted multiplication

$$(u_1, x_1)(u_2, x_2) = (u_1 + u_2, (-1)^{u_1 u_2} x_1 x_2) .$$

We have a surjective map

$$\chi: \mathbb{Z} \times S_1(C, J) \longrightarrow \mathbb{Z}/2\mathbb{Z} \cdot S_1(C, J)$$

via
$$\chi(n, x) = (n \bmod 2, (-1)^{(n(n-1)/2}) x) .$$

It is easy to verify that χ is multiplicative so that, in particular, $\mathbb{Z}/2\mathbb{Z} \cdot S_1(C, J)$ is a group. We now consider the additive map

$$(\nu, d): S(C, J) \xrightarrow{(\text{rank}, \det)} \mathbb{Z} \times S_1(C, J) \xrightarrow{\chi} \mathbb{Z}/2\mathbb{Z} \cdot S_1(C, J) ,$$

i.e. $(\nu(E), d(E)) = (n \bmod 2, (-1)^{(n(n-1)/2}) \det E)$ for any (E) in $S(C, J)$ of rank n , and call $d(E)$ the *signed determinant* of E . Since (ν, d) vanishes on all hyperbolic spaces [12, Satz 4.1.2, p. 108], it induces maps from $K(C, J)$ and $W(C, J)$ to $\mathbb{Z}/2\mathbb{Z} \cdot S_1(C, J)$ which we will also denote by (ν, d) . Since for (E) in $S_1(C, J)$ we have $d(E) = (E)$ it follows that the canonical maps $S_1(C, J) \rightarrow K(C, J)$ and $S_1(C, J) \rightarrow W(C, J)$ are injective. Thus we may regard $S_1(C, J)$ as a subgroup of the units of $K(C, J)$ or of $W(C, J)$.

THEOREM 3.3. *Let C be either a Dedekind ring or a connected semi-local ring with involution J and let T be either $K(C, J)$ or $W(C, J)$. Then for any unit x of T we have*

$$x = d(x)(1 + n)$$

with a nilpotent element n in T . In particular, the group T^ is*

generated by $S_1(C, J)$ and $1 + \text{Nil } T$.

Proof. If $T = T_t$ then by Proposition 1.2 (iii) and Lemma 2.1 (iv), $M_{0,T} = \text{Nil } T$. Since x is a unit, it must be represented by a form of odd rank. Hence $x - d(x) \in M_{0,T}$ is nilpotent, so setting

$$n = d(x)^{-1}(x - d(x))$$

we have $x = d(x)(1 + n)$ with n nilpotent.

If $T \neq T_t$ then $Z \rightarrow T$ is injective so by Proposition 1.2 (iii) and Remark 2.2, $\text{Nil } T$ is the set of y in T such that $\sigma(y) = 0$ for all ring homomorphisms $\sigma: T \rightarrow Z$. Thus we only need to check that $\sigma(x) = \sigma(d(x))$ for all σ . This has been done in [14], proof of Theorem 3.23, for C connected semi-local. Now let C be a Dedekind ring, L its quotient field, and R be $K(L, J)$ or respectively $W(L, J)$. Denote the canonical map $T \rightarrow R$ by φ . By Proposition 1.2 (iii) and Remark 2.2 any homomorphism $\sigma: T \rightarrow Z$ can be factored $\sigma = \sigma' \circ \varphi$ with some $\sigma': R \rightarrow Z$. Since $\varphi(x)$ is a unit of R we know from the preceding that $\sigma'(\varphi(x)) = \sigma'(d(\varphi(x)))$. But by the functorial properties of exterior powers clearly $d(\varphi(x)) = \varphi(d(x))$. Thus we get the desired equation

$$(\sigma' \circ \varphi)(x) = (\sigma' \circ \varphi)(d(x)) .$$

REMARK 3.4. Let q be any rational prime, $R = Z[G]/K$ a Witt ring for an abelian group G of exponent q and T be the intersection of the subrings $T_i = Z[H_i]/(Z[H_i] \cap K)$ corresponding to some family $\{H_i\}$ of subgroups of G . It can be shown that any unit of T has the form $\pm \bar{g}(1 + n)$ where \bar{g} is the image of some g in $\bigcap_i H_i$ and n is in $\text{Nil } T$, in the following cases: ($I_G =$ augmentation ideal).

- (i) $q \neq 2: K \subset I_G^2 + qZ[G]$
- (ii) $q = 2: K \subset I_G^2 + 4Z[G]$
- (iii) $q = 2:$ there exists an element w of $\bigcap H_i$ such that

$$1 + w \in K \subset I_G^2 + 4Z[G] + (1 + w)Z[G] .$$

Note that for $R = W(L, J)$, $G = F^*/NL^*$, $T = W(C, J)$ condition (ii) is violated but (iii) holds with $w = (-1)$. The assumptions (i)-(iii) are needed to construct "determinant functions" playing a role similar to that of the signed determinant in the proof of Theorem 3.2.

4. $A(C, J)$ for $J \neq$ identity. If J is the identity the kernel $A(C, J)$ of the canonical map $K(C, J) \rightarrow K(L, J)$ has been determined for any Dedekind ring in [9, Cor. to Th. 4] and [12, 11.3.5, p. 138]. In this section we give a computation of $A(C, J)$ for $J \neq$ identity. We argue along the same lines as in [12] and use well-known classical methods ([2], [15]).

Throughout this section we use the notations of §1: C is an arbitrary Dedekind ring with an involution $J \neq \text{identity}$, whose fixed ring is denoted by A . For any prime ideal \mathfrak{p} of A , we denote by $\hat{A}_{\mathfrak{p}}$ the completion of the localization $A_{\mathfrak{p}}$, and we write $\hat{C}_{\mathfrak{p}}$ for $C \otimes_A \hat{A}_{\mathfrak{p}}$, the completion of $C \otimes_A A_{\mathfrak{p}}$ with respect to \mathfrak{p} . We set $\hat{C}_{\mathfrak{p}} = L$ for $\mathfrak{p} = 0$ and for a maximal ideal \mathfrak{p} of A we write $\hat{L}_{\mathfrak{p}}$ for the completion $L \otimes_F \hat{F}_{\mathfrak{p}} = C \otimes_A \hat{F}_{\mathfrak{p}}$ of L which is the total ring of quotients of the semi-local ring $\hat{C}_{\mathfrak{p}}$. The involutions induced by J on $\hat{C}_{\mathfrak{p}}$ and $\hat{L}_{\mathfrak{p}}$ will both be written as $\hat{J}_{\mathfrak{p}}$. For any space E over (C, J) we denote by $\hat{E}_{\mathfrak{p}}$ the completed localization $E \otimes_A \hat{A}_{\mathfrak{p}}$ which is a space over $(\hat{C}_{\mathfrak{p}}, \hat{J}_{\mathfrak{p}})$. As usual we consider $\hat{E}_{\mathfrak{p}}$ as a subset of the space $E \otimes_A \hat{F}_{\mathfrak{p}}$ over $(\hat{L}_{\mathfrak{p}}, \hat{J}_{\mathfrak{p}})$ and the automorphism group $\mathfrak{U}(\hat{E}_{\mathfrak{p}})$ as a subgroup of $\mathfrak{U}(E \otimes_A \hat{F}_{\mathfrak{p}})$. For a space M over (L, J) we denote by $\hat{M}_{\mathfrak{p}}$ the completion with respect to \mathfrak{p} , which is a space over $(\hat{L}_{\mathfrak{p}}, \hat{J}_{\mathfrak{p}})$. Note that $\hat{M}_{\mathfrak{p}} = M$ if $\mathfrak{p} = 0$.

We define the *genus*, $\Gamma(E)$, of a space E over (C, J) as the set of isometry classes (E') in $S(C, J)$ such that $\hat{E}_{\mathfrak{p}} \cong \hat{E}'_{\mathfrak{p}}$ for all prime ideals \mathfrak{p} of A . If (E') lies in $\Gamma(E)$ we can find an isometry $\sigma: E \otimes L \rightarrow E' \otimes L$ and for every \mathfrak{p} in $\text{Max } A$ an element $\sigma_{\mathfrak{p}}$ in $\mathfrak{U}(E' \otimes \hat{L}_{\mathfrak{p}})$ such that

$$(4.1) \quad \hat{E}'_{\mathfrak{p}} = \sigma_{\mathfrak{p}} \sigma \hat{E}_{\mathfrak{p}} .$$

Since $\sigma_{\mathfrak{p}}$ is unitary, the determinant of $\sigma_{\mathfrak{p}}$ over $\hat{L}_{\mathfrak{p}}$ has norm 1 and hence by Hilbert's Theorem 90,

$$(4.2) \quad \det \sigma_{\mathfrak{p}} = a_{\mathfrak{p}} \bar{a}_{\mathfrak{p}}^{-1}$$

for some $a_{\mathfrak{p}}$ in $\hat{L}_{\mathfrak{p}}^*$. We use (4.2) to define for any genus Γ a map

$$\varphi: \Gamma \times \Gamma \longrightarrow \text{Div } C / \mathfrak{S}(\text{Div } C)^G$$

where $\text{Div } C$ denotes the divisor group of C , $(\text{Div } C)^G$ the subgroup left fixed by $G = \{1, J\}$ and \mathfrak{S} the group of principal divisors of C . For E, E' in Γ we define $\varphi(E, E')$ to be the image of

$$(4.3) \quad \prod_{\mathfrak{p}} \prod_{\mathfrak{P}|\mathfrak{p}} \mathfrak{P}^{\text{ord}_{\mathfrak{P}}(a_{\mathfrak{p}})} , \quad \mathfrak{P} \text{ in Max } C$$

in $\text{Div } C / \mathfrak{S}(\text{Div } C)^G$ with $a_{\mathfrak{p}}$ coming from (4.1) and (4.2) chosen arbitrarily. Here \mathfrak{p} runs only through the ideals of $\text{Max } A$ which split in C . Next, we show that $\varphi(E, E')$ is well defined, i.e., that (4.3) does not depend on the particular choices in (4.1) and (4.2). Given (4.1), the $a_{\mathfrak{p}}$ in (4.2) can only be changed by a factor in $\hat{F}_{\mathfrak{p}}^*$ which does not affect (4.3) mod (Dw) . Let $\hat{E}'_{\mathfrak{p}} = \sigma'_{\mathfrak{p}} \sigma' E$ be another system of equations of type (4.1). Then there exists some ρ in $\mathfrak{U}(E' \otimes L)$ and $\mu_{\mathfrak{p}}$ in $\mathfrak{U}(\hat{E}'_{\mathfrak{p}})$ such that

$$(4.4) \quad \sigma'_\mathfrak{p} = \mu_\mathfrak{p} \sigma_\mathfrak{p} \rho$$

for all \mathfrak{p} in $\text{Max } A$. If \mathfrak{p} splits in C then $\hat{L}_\mathfrak{p} \cong \hat{F}_\mathfrak{p} \times \hat{F}_\mathfrak{p}$ and the involution $\hat{J}_\mathfrak{p}$ is given by $\hat{J}_\mathfrak{p}(x, y) = (y, x)$ [14, Lemma 1.8]. Hence the element $\det(\mu_\mathfrak{p})$ of $\hat{C}_\mathfrak{p}^*$ of norm 1 can be written in the form $b_\mathfrak{p} \bar{b}_\mathfrak{p}^{-1}$ with $b_\mathfrak{p}$ in $\hat{C}_\mathfrak{p}^*$. Again by Hilbert's Theorem 90, there is an element c in L such that $\det \rho = c \bar{c}^{-1}$. Thus starting from (4.4), we obtain for the computation of $\varphi(E, E')$, elements $a'_\mathfrak{p} = b_\mathfrak{p} a_\mathfrak{p} c$ at split prime ideals and this yieldst he same value in (4.3) mod \mathfrak{S} . Hence it is also clear that $\varphi(E, E')$ depends only on the isometry classes of E and E' . Clearly for spaces V, V' in some other genus we have

$$(4.5) \quad \begin{aligned} \varphi(E \perp V, E' \perp V') &= \varphi(E, E') \varphi(V, V') \quad \text{and also} \\ \varphi(E, E') &= \varphi(E', E)^{-1} \end{aligned}$$

EXAMPLE 4.6. Let (E, Φ) be the hyperbolic space of rank 2 over (L, J) , i.e., $E = Ce \oplus Cf$ with $\Phi(e, e) = \Phi(f, f) = 0, \Phi(e, f) = 1$. For any α in $\text{Div } C$ we regard $H(\alpha)$ as the sublattice $\alpha e \oplus \bar{\alpha}^{-1} f$ of E . Choose for every \mathfrak{p} in $\text{Max } A$ a generator $a_\mathfrak{p}$ of the principal ideal $\alpha \hat{C}_\mathfrak{p}$. Then the automorphism $e \mapsto a_\mathfrak{p} e, f \mapsto \bar{a}_\mathfrak{p}^{-1} f$ of $\hat{E}_\mathfrak{p}$ maps $H(\hat{C})_\mathfrak{p}$ to $\hat{H}(\alpha)_\mathfrak{p}$. Thus $H(\alpha)$ and H are in the same genus and $\varphi(H, H(\alpha)) = [\alpha]$, where $[\alpha]$ denotes the class of the divisor α in $\text{Div } C / \mathfrak{S}(\text{Div } C)^G$.

We define the *SU-genus* $\Sigma(E)$ of a space E over (C, J) as the set of all isometry classes (E') in $\Gamma(E)$ such that there exists an equation (4.1) with $\sigma_\mathfrak{p}$ in $SU(\hat{E}'_\mathfrak{p})$ for all \mathfrak{p} in $\text{Max } A$, i.e., with $\det \sigma_\mathfrak{p} = 1$.

LEMMA 4.7. Let E be a (C, J) -space and assume that $\hat{E}_\mathfrak{p}$ represents a unit of $\hat{A}_\mathfrak{p}$ for every \mathfrak{p} in $\text{Max } A$. (This hypothesis is always fulfilled if \mathfrak{p} splits or if 2 does not lie in \mathfrak{p} .) Then $\Sigma(E)$ is the set of all (E') in $\Gamma(E)$ with $\varphi(E, E') = 1$.

Proof. If (E') lies in $\Sigma(E)$ it is clear that $\varphi(E, E') = 1$. Let now (E') be in $\Gamma(E)$ with $\varphi(E, E') = 1$. Without loss of generality we assume that E and E' are lattices of full rank in the same space M over (L, J) . First, for every \mathfrak{p} in $\text{Max } A$, we choose some $\sigma_\mathfrak{p}$ in $\mathfrak{U}(\hat{M}_\mathfrak{p})$ and some $a_\mathfrak{p}$ in $\hat{L}_\mathfrak{p}^*$ such that $\hat{E}'_\mathfrak{p} = \sigma_\mathfrak{p} \hat{E}_\mathfrak{p}$ with $\det \sigma_\mathfrak{p} = a_\mathfrak{p} \bar{a}_\mathfrak{p}^{-1}$. We want $\det \sigma_\mathfrak{p}$ to be a unit in $\hat{C}_\mathfrak{p}^*$. This is automatically true unless $\mathfrak{p}C = \mathfrak{P}\bar{\mathfrak{P}}$ with $\mathfrak{P} \neq \bar{\mathfrak{P}}$. But since $\varphi(E, E') = 1$ we can find a c in L^* such that $\text{ord}_\mathfrak{p}(a_\mathfrak{p} c) = \text{ord}_\mathfrak{p}(a_\mathfrak{p})$ whenever $\mathfrak{p}C = \mathfrak{P}\bar{\mathfrak{P}}, \mathfrak{P} \neq \bar{\mathfrak{P}}$. Since $J \neq \text{identity}$, M contains a one-dimensional subspace $W = Le$ with $\Phi(e, e) \neq 0$ [3, Lemma 1, p. 90] so that $M = Le \perp W^\perp$. Now let ρ in $\mathfrak{U}(M)$ be the automorphism which is the identity on W^\perp and multiplies e by $c^{-1} \bar{c}$. Setting $\tau_\mathfrak{p} = \sigma_\mathfrak{p} \rho^{-1}$ we obtain

$$\hat{E}'_\mathfrak{p} = \tau_\mathfrak{p} \rho \hat{E}_\mathfrak{p}$$

and $\det \tau_p = (\det \sigma_p)(e\bar{e}^{-1})$, which is a unit of \hat{C}_p for all p . By hypothesis every \hat{E}'_p has a subspace $\hat{C}_p e_p$ with $\Phi(e_p, e_p)$ in A_p^* . So by an argument as above we can find a μ_p in $\mathfrak{U}(\hat{E}'_p)$ with $\det \mu_p = (\det \tau_p)^{-1}$. The equations

$$\hat{E}'_p = \mu_p \tau_p \circ \hat{E}_p$$

show that (E') lies in the SU -genus of E .

We now sketch a proof of a weak version of Kneser's strong approximation theorem for the special unitary group [15]. Let (M, Φ) be any space over (L, J) of dimension ≥ 2 . Then $\mathfrak{U}(M)$ is generated by the symmetries

$$\sigma(x, l): z \longmapsto z - \Phi(z, x)l^{-1}x$$

where (x, l) runs through all pairs of $M \times L^*$ such that $l + \bar{l} = \Phi(x, x) \neq 0$ [8, p. 41]. We first indicate a proof for the following well-known fact [2, §3] since we feel that the argument in [2] is not quite clear.

LEMMA 4.8. *Given maximal ideals $\mathfrak{p}_1, \dots, \mathfrak{p}_r$ of A and pairs (x_i, l_i) in $\hat{M}_{\mathfrak{p}_i} \times \hat{L}_{\mathfrak{p}_i}^*$, $1 \leq i \leq r$, with $l_i + \bar{l}_i = \Phi(x_i, x_i) \neq 0$, there exists for every $\varepsilon > 0$ a pair (x, l) in $M \times L^*$ with $l + \bar{l} = \Phi(x, x) \neq 0$ such that $\|x - x_i\|_{\mathfrak{p}_i} < \varepsilon$, $|l - l_i|_{\mathfrak{p}_i} < \varepsilon$ for $1 \leq i \leq r$. Here $|\cdot|_{\mathfrak{p}}$ denotes a valuation belonging to \mathfrak{p} and $\|\cdot\|_{\mathfrak{p}}$ an associated norm on $\hat{M}_{\mathfrak{p}}$.*

Proof. Since $J \neq$ identity there is some m in L with $m + \bar{m} = 1$. We can find a vector x in M which is near x_i at \mathfrak{p}_i and an l' in L which is near l_i at \mathfrak{p}_i , for $1 \leq i \leq r$. Then

$$l = l' + m(\Phi(x, x) - l' - \bar{l}')$$

satisfies $l + \bar{l} = \Phi(x, x)$ and the pair (x, l) has the desired properties if the approximation of the x_i by the x and the l_i by l' is good enough.

According to [2, §3] this lemma implies that the weak approximation theorem holds for $\mathfrak{U}(M)$, i.e., the diagonal map from $\mathfrak{U}(M)$ to

$$\prod_{\mathfrak{p} \text{ in } S} \mathfrak{U}(\hat{M}_{\mathfrak{p}})$$

has dense image for any finite subset S of $\text{Max } A$.

Assume now that $[M: L] \geq 3$ and that M is isotropic, i.e., M contains some $x \neq 0$ with $\Phi(x, x) = 0$. Then $SU(M)$ is the commutator subgroup of $\mathfrak{U}(M)$ [8, Chap. II, §4, §5]. Thus it can be verified that the weak approximation theorem also holds for $SU(M)$. Again pick some m in L with $m + \bar{m} = 1$. As is shown in [1] (cf. [8, p. 61] for the orthogonal case), $SU(M)$ is generated by automorphisms

$$E(x, y): z \longmapsto z + \Phi(z, x)y - \Phi(z, y)x - m\Phi(z, x)\Phi(y, y)x$$

where (x, y) runs through all elements of $M \times M$ with $\Phi(x, x) = \Phi(x, y) = 0$. Now the simple argument in [12, Proof of Satz 11.2.8, p. 137] shows that we have strong approximation in $SU(M)$. (It also seems possible to adopt the proof of [2, Satz 18]):

THEOREM 4.9. *Let M be an isotropic space over (L, J) with² $[M: L] \geq 3$ and let $E \subset M$ be a (C, J) -lattice of full rank. Prescribe lattices E'_i of full rank in \hat{M}_{v_i} and elements σ_i in $SU(\hat{M}_{v_i})$ for finitely many ideals $\mathfrak{p}_1, \dots, \mathfrak{p}_r$ of $\text{Max } A$. Then there exists an element σ of $SU(M)$ such that $(\sigma - \sigma_i)\hat{E}_{v_i} \subset E'_{v_i}$ for $1 \leq i \leq r$ and $\sigma\hat{E}_a = \hat{E}_a$ for \mathfrak{a} in $\text{Max } A$ distinct from $\mathfrak{p}_1, \dots, \mathfrak{p}_r$.*

From this theorem a standard argument [15], [2, p. 99] establishes

COROLLARY 4.10. *If E is an isotropic (C, J) -space of rank ≥ 3 , the SU -genus $\Sigma(E)$ contains only the class (E) .*

Finally, Corollary 4.10 and Lemma 4.7 yield the desired

THEOREM 4.11. *The kernel of the surjection $f: \text{Pic}(C) \rightarrow \mathcal{A}(C, J)$, where $f(\mathfrak{a}) = [H(\mathfrak{a})] - [H]$, is the subgroup $\mathfrak{S}(\text{Div } C)^g/\mathfrak{S}$.*

Proof. For any \mathfrak{a} in $(\text{Div } C)^g$ we consider the spaces $E = H \perp (1)$ and $E' = H(\mathfrak{a}) \perp (1)$. According to (4.5) and Example 4.6, we have $\varphi(E, E') = 1$. Now, since $(H(\hat{\mathfrak{a}}))_v$ and \hat{H}_v represent all of A_v^* , Lemma 4.7 shows that E' is in the SU -genus of E and Corollary 4.10 forces $E \cong E'$. Hence $f(\mathfrak{a}) = [E'] - [E] = 0$.

Let (\mathfrak{a}) be in $\text{Ker } f$. Then there exists a (C, J) -space V such that $H \perp V \cong H(\mathfrak{a}) \perp V$. Again, (4.5) yields $\varphi(H, H(\mathfrak{a})) = 1$ and so by Example 4.6, the class of \mathfrak{a} is in $\mathfrak{S}(\text{Div } C)^g/\mathfrak{S}$.

REFERENCES

1. R. Baeza, *Unitäre Gruppen über lokalen Ringen*, Dissertation, Saarbrücken, 1970.
2. S. Böge, *Schiefermitesche Formen über Zahlkörper und Quaternionenschieferkörpern*, J. reine angew. Math., **221** (1966), 85-112.
3. N. Bourbaki, *Algèbre*, Chap. 9, Act. Sc. Ind. 1272, Hermann, Paris 1959.
4. ———, *Algèbre commutative*, Chap. 1-2, Act. Sc. Ind. 1290, Hemann, Paris 1961.
5. ———, *Algèbre commutative*, Chap. 5, Act. Sc. Ind. 1308, Hermann, Paris 1964.
6. ———, *Algèbre commutative*, Chap. 7, Act. Sc. Ind. 1314, Hermann, Paris 1965.
7. H. Cartan and S. Eilenberg, *Homological algebra*, Princeton University Press, Princeton 1956.

² The condition $[M: L] \geq 3$ has inadvertently been omitted in [2, 11.2].

8. J. Dieudonné, *La géométrie des groupes classiques*, 2nd Ed., Ergebnisse d. Math., Springer, Berlin 1963.
9. A. Fröhlich, *On the K-theory of unimodular forms over rings of algebraic integers*, Quar. J. Math., (Oxford), **22** (1971), 401-423.
10. D. K. Harrison, *Witt rings*, Lecture notes, Department of Mathematics, University of Kentucky, Lexington, Kentucky, 1970.
11. I. Kaplansky, *Commutative rings*, Allyn and Bacon, Boston, 1970.
12. M. Knebusch, *Grothendieck und Witttringe von nichtausgearteten symmetrischen Bilinearformen*, Sitzber. Heidelberg Akad. Wiss. (1969/1970), 93-157.
13. M. Knebusch, A. Rosenberg and R. Ware, *Structure of Witt rings, quotients of abelian group rings, and orderings of fields*, Bull. Amer. Math. Soc., **77** (1971), 205-210.
14. ———, *Structure of Witt rings and quotients of abelian group rings*, Amer. J. Math., **94** (1972), 119-155.
15. M. Kneser, *Klassenzahlen indefiniter quadratischer Formen in drei und mehr Veränderlichen*, Archiv Math., **7** (1956), 323-332.
16. J. Leicht und F. Lorenz, *Die Primideale des Wittschen Ringes*, Invent. Math., **10** (1970), 82-88.
17. J. Milnor, *Symmetric inner product spaces in characteristic two*, in Ann. of Math. Studies 70, Princeton University Press, Princeton, New Jersey, 1971.
18. A. Pfister, *Quadratische Formen in beliebigen Körpern*, Invent. Math., **1** (1966), 116-132.
19. W. Scharlau, *Klassifikation hermitescher Formen über lokalen Körpern*, Math. Ann., **186** (1970), 201-208.
20. J. Serre, *Modules projectifs et espaces fibrés à fibre vectorielle*, Sémin. Dubreil-Pisot, Paris 1957-58.
21. T. A. Springer, *Quadratic forms over fields with a discrete valuation*, Indag. Math., **17** (1955), 352-362.
22. E. Steinitz, *Rechteckige Systeme und Moduln in algebraischen Zahlkörpern*, Math. Ann., **71** (1912), 328-354.

Received July 21, 1971.

UNIVERSITÄT DES SAARLANDES
CORNELL UNIVERSITY
NORTHWESTERN UNIVERSITY

