

Parasar Mohanty, Department of Pure Mathematics, University of Waterloo,  
Waterloo ON, Canada N2L 3G1. email: pmohanty@math.uwaterloo.ca

Erik Talvila\*, Department of Mathematics and Statistics, University College  
of the Fraser Valley, Abbotsford BC, Canada V2S 7M8.  
email: Erik.Talvila@ucfv.ca

## A PRODUCT CONVERGENCE THEOREM FOR HENSTOCK-KURZWEIL INTEGRALS

### Abstract

Necessary and sufficient for  $\int_a^b f g_n \rightarrow \int_a^b f g$  for all Henstock–Kurzweil integrable functions  $f$  is that  $g$  be of bounded variation,  $g_n$  be uniformly bounded and of uniform bounded variation and, on each compact interval in  $(a, b)$ ,  $g_n \rightarrow g$  in measure or in the  $L^1$  norm. The same conditions are necessary and sufficient for  $\|f(g_n - g)\| \rightarrow 0$  for all Henstock–Kurzweil integrable functions  $f$ . If  $g_n \rightarrow g$  a.e., then convergence  $\|f g_n\| \rightarrow \|f g\|$  for all Henstock–Kurzweil integrable functions  $f$  is equivalent to  $\|f(g_n - g)\| \rightarrow 0$ . This extends a theorem due to Lee Peng-Yee.

Let  $-\infty \leq a < b \leq \infty$  and denote the Henstock–Kurzweil integrable functions on  $(a, b)$  by  $\mathcal{HK}$ . The Alexiewicz norm of  $f \in \mathcal{HK}$  is  $\|f\| = \sup_I |\int_I f|$  where the supremum is taken over all intervals  $I \subset (a, b)$ . If  $g$  is a real-valued function on  $[a, b]$ , we write  $V_{[a,b]}g$  for the variation of  $g$  over  $[a, b]$ , dropping the subscript when the identity of  $[a, b]$  is clear. The set of functions of normalized bounded variation,  $\mathcal{NBV}$ , consists of the functions on  $[a, b]$  that are of bounded variation, are left continuous and vanish at  $a$ . It is known that the multipliers for  $\mathcal{HK}$  are  $\mathcal{NBV}$ ; i.e.,  $f g \in \mathcal{HK}$  for all  $f \in \mathcal{HK}$  if and only if  $g$  is equivalent to a function in  $\mathcal{NBV}$ . This paper is concerned with necessary and sufficient conditions under which  $\int_a^b f g_n \rightarrow \int_a^b f g$  for all  $f \in \mathcal{HK}$ . One such set of conditions was given by Lee Peng-Yee in [2, Theorem 12.11]. If  $g$  is of bounded variation, changing  $g$  on a countable set will make it an element of  $\mathcal{NBV}$ . With this observation, a minor modification of Lee’s theorem produces the following result.

---

Key Words: Henstock–Kurzweil integral, convergence theorem, Alexiewicz norm

Mathematical Reviews subject classification: 26A39, 46E30

Received by the editors December 2, 2002

Communicated by: Peter Bullen

\*Research partially supported by the Natural Sciences and Engineering Research Council of Canada.

**Theorem 1.** [2, Theorem 12.11] *Let  $-\infty < a < b < \infty$ , let  $g_n$  and  $g$  be real-valued functions on  $[a, b]$  with  $g$  of bounded variation. In order for  $\int_a^b f g_n \rightarrow \int_a^b f g$  for all  $f \in \mathcal{HK}$  it is necessary and sufficient that*

$$\left. \begin{array}{l} \text{for each interval } (c, d) \subset (a, b), \int_c^d g_n \rightarrow \int_c^d g \text{ as } n \rightarrow \infty, \\ \text{for each } n \geq 1, g_n \text{ is equivalent to a function } h_n \in \mathcal{NBV}, \\ \text{and there is } M \in [0, \infty) \text{ such that } Vh_n \leq M \text{ for all } n \geq 1. \end{array} \right\} \quad (1)$$

We extend this theorem to unbounded intervals, show that the condition  $\int_c^d g_n \rightarrow \int_c^d g$  in (1) can be replaced by  $g_n \rightarrow g$  on each compact interval in  $(a, b)$  either in measure or in the  $L^1$  norm, and that this also lets us conclude  $\|f(g_n - g)\| \rightarrow 0$ . We also show that if  $g_n \rightarrow g$  in measure or almost everywhere, then  $\|f g_n\| \rightarrow \|f g\|$  for all  $f \in \mathcal{HK}$  if and only if  $\|f g_n - f g\| \rightarrow 0$  for all  $f \in \mathcal{HK}$ .

One might think the conditions (1) imply  $g_n \rightarrow g$  almost everywhere. This is not the case, as is illustrated by the following example [1, p. 61].

**Example 2.** Let  $g_n = \chi_{(j2^{-k}, (j+1)2^{-k}]}$  where  $0 \leq j < 2^k$  and  $n = j + 2^k$ . Note that  $\|g_n\|_\infty = 1$ ,  $g_n \in \mathcal{NBV}$ ,  $Vg_n \leq 2$ , and  $|\int_c^d g_n| \leq \|g_n\| = 2^{-k} < 2/n \rightarrow 0$ , so that (1) is satisfied with  $g = 0$ . For each  $x \in (0, 1]$  we have  $\inf_n g_n(x) = 0$ ,  $\sup_n g_n(x) = 1$ , and for no  $x \in (0, 1]$  does  $g_n(x)$  have a limit. However,  $g_n \rightarrow 0$  in measure since if  $T_n = \{x \in [0, 1] : |g_n(x)| > \epsilon\}$ , then for each  $0 < \epsilon \leq 1$ , we have  $\lambda(T_n) < 2/n \rightarrow 0$  as  $n \rightarrow \infty$  ( $\lambda$  is Lebesgue measure).

We have the following extension of Theorem 1.

**Theorem 3.** *Let  $[a, b]$  be a compact interval in  $\mathbb{R}$ , let  $g_n$  and  $g$  be real-valued functions on  $[a, b]$  with  $g$  of bounded variation. In order for  $\int_a^b f g_n \rightarrow \int_a^b f g$  for all  $f \in \mathcal{HK}$  it is necessary and sufficient that*

$$\left. \begin{array}{l} g_n \rightarrow g \text{ in measure as } n \rightarrow \infty, \\ \text{for each } n \geq 1, g_n \text{ is equivalent to a function } h_n \in \mathcal{NBV}, \\ \text{and there is } M \in [0, \infty) \text{ such that } Vh_n \leq M \text{ for all } n \geq 1. \end{array} \right\} \quad (2)$$

*If  $(a, b) \subset \mathbb{R}$  is unbounded, then change the first line of (2) by requiring  $g_n \chi_I \rightarrow g \chi_I$  in measure for each compact interval  $I \in (a, b)$ .*

**PROOF.** By working with  $g_n - g$  we can assume  $g = 0$ . First consider the case when  $(a, b)$  is a bounded interval. If  $\int_a^b f g_n \rightarrow 0$  for all  $f \in \mathcal{HK}$ , then using Theorem 1 and changing  $g_n$  on a countable set, we can assume  $g_n \in \mathcal{NBV}$ ,  $Vg_n \leq M$ ,  $\|g_n\|_\infty \leq M$  and  $\int_c^d g_n \rightarrow 0$  for each interval  $(c, d) \subset (a, b)$ . Suppose  $g_n$  does not converge to 0 in measure. Then there are  $\delta, \epsilon > 0$  and

an infinite index set  $\mathcal{J} \subset \mathbb{N}$  such that  $\lambda(S_n) > \delta$  for each  $n \in \mathcal{J}$ , where  $S_n = \{x \in (a, b) : g_n(x) > \epsilon\}$ . (Or else there is a corresponding set on which  $g_n(x) < -\epsilon$  for all  $n \in \mathcal{J}$ .) Now let  $n \in \mathcal{J}$ . Since  $g_n$  is left continuous, if  $x \in S_n$ , there is a number  $c_{n,x} > 0$  such that  $[x - c_{n,x}, x] \subset S_n$ . Hence,  $V_n := \{[c, x] : x \in S_n \text{ and } [c, x] \subset S_n\}$  is a Vitali cover of  $S_n$ . So there is a finite set of disjoint closed intervals,  $\sigma_n \subset V_n$ , with  $\lambda(S_n \setminus \cup_{I \in \sigma_n} I) < \delta/2$ . Write  $(a, b) \setminus \cup_{I \in \sigma_n} I = \cup_{I \in \tau_n} I$  where  $\tau_n$  is a set of disjoint open intervals with  $\text{card}(\tau_n) = \text{card}(\sigma_n) + 1$ . Let

$$P_n = \text{card}(\{I \in \tau_n : g_n(x) \leq \epsilon/2 \text{ for some } x \in I\}).$$

Each interval  $I \in \tau_n$  that does not have  $a$  or  $b$  as an endpoint has contiguous intervals on its left and right that are in  $\sigma_n$  (for each of which  $g_n > \epsilon$ ). The interval  $I$  then contributes more than  $(\epsilon - \epsilon/2) + (\epsilon - \epsilon/2) = \epsilon$  to the variation of  $g_n$ . If  $I$  has  $a$  as an endpoint, then since  $g_n(a) = 0$ ,  $I$  contributes more than  $\epsilon$  to the variation of  $g_n$ . If  $I$  has  $b$  as an endpoint, then  $I$  contributes more than  $\epsilon/2$  to the variation of  $g_n$ . Hence,

$$Vg_n \geq (P_n - 1)\epsilon + \epsilon/2 = (P_n - 1/2)\epsilon.$$

(This inequality is still valid if  $P_n = 1$ .) But,  $Vg_n \leq M$ ; so  $P_n \leq P$  for all  $n \in \mathcal{J}$  and some  $P \in \mathbb{N}$ . Then we have a set of intervals,  $U_n$ , formed by taking unions of intervals from  $\sigma_n$  and those intervals in  $\tau_n$  on which  $g_n > \epsilon/2$ . Now,  $\lambda(\cup_{I \in U_n} I) > \delta/2$ ,  $\text{card}(U_n) \leq P + 1$  and  $g_n > \epsilon/2$  on each interval  $I \in U_n$ . Therefore, there is an interval  $I_n \in U_n$  such that  $\lambda(I_n) > \delta/[2(P + 1)]$ . The sequence of centers of intervals  $I_n$  has a convergent subsequence. There is then an infinite index set  $\mathcal{J}' \subset \mathcal{J}$  with the property that for all  $n \in \mathcal{J}'$  we have  $g_n > \epsilon/2$  on an interval  $I \subset (a, b)$  with  $\lambda(I) > \delta/[3(P + 1)]$ . Hence,  $\limsup_{n \geq 1} \int_I g_n > \delta\epsilon/[6(P + 1)]$ . This contradicts the fact that  $\int_I g_n \rightarrow 0$ , showing that indeed  $g_n \rightarrow 0$  in measure.

Suppose (2) holds. As above, we can assume  $g_n \in \mathcal{NBV}$ ,  $Vg_n \leq M$ ,  $\|g_n\|_\infty \leq M$  and  $g_n \rightarrow 0$  in measure. Let  $\epsilon > 0$ . Define

$$T_n = \{x \in (a, b) : |g_n(x)| > \epsilon\}.$$

Then

$$\begin{aligned} \left| \int_a^b g_n \right| &\leq \int_{T_n} |g_n| + \int_{(a,b) \setminus T_n} |g_n| \\ &\leq M\lambda(T_n) + \epsilon(b - a). \end{aligned}$$

Since  $\lim \lambda(T_n) = 0$ , it now follows that  $\int_c^d g_n \rightarrow 0$  for each  $(c, d) \subset (a, b)$ . Theorem 1 now shows  $\int_a^b f g_n \rightarrow 0$  for all  $f \in \mathcal{HK}$ .

Now consider integrals on  $\mathbb{R}$ . If  $\int_{-\infty}^{\infty} f g_n \rightarrow 0$  for all  $f \in \mathcal{HK}$ , then it is necessary that  $\int_a^b f g_n \rightarrow 0$  for each compact interval  $[a, b]$ . By the current theorem,  $g_n \rightarrow g$  in measure on each  $[a, b]$ . And, it is necessary that  $\int_1^{\infty} f g_n \rightarrow 0$ . The change of variables  $x \mapsto 1/x$  now shows it is necessary that  $g_n$  be equivalent to a function that is uniformly bounded and of uniform bounded variation on  $[1, \infty]$ . Similarly with  $\int_{-\infty}^1 f g_n \rightarrow 0$ . Hence, it is necessary that  $g_n$  be uniformly bounded and of uniform bounded variation on  $\mathbb{R}$ .

Suppose (2) holds with  $g_n \rightarrow g$  in measure on each compact interval in  $\mathbb{R}$ . Write  $\int_{-\infty}^{\infty} f g_n = \int_{-\infty}^a f g_n + \int_a^b f g_n + \int_b^{\infty} f g_n$ . Use Lemma 24 in [4] to write  $|\int_{-\infty}^a f g_n| \leq \|f \chi_{(-\infty, a)}\| V_{[-\infty, a]} g_n \leq \|f \chi_{(-\infty, a)}\| M \rightarrow 0$  as  $a \rightarrow -\infty$ . We can then take a large enough interval  $[a, b] \subset \mathbb{R}$  and apply the current theorem on  $[a, b]$ . Other unbounded intervals are handled in a similar manner.  $\square$

**Remark 4.** If (2) holds, then dominated convergence shows  $\|g_n - g\|_1 \rightarrow 0$ . And, convergence in  $\|\cdot\|_1$  implies convergence in measure. Therefore, in the first statement of (2) and in the last statement of Theorem 3, ‘convergence in measure’ can be replaced with ‘convergence in  $\|\cdot\|_1$ ’. Similar remarks apply to Theorem 6.

**Remark 5.** The change of variables argument in the second last paragraph of Theorem 3 can be replaced with an appeal to the Banach–Steinhaus Theorem on unbounded intervals. See [3, Lemma 7]. A similar remark holds for the proof of Theorem 8.

The sequence of Heaviside functions  $g_n = \chi_{(n, \infty]}$  shows (2) is not necessary to have  $\int_{-\infty}^{\infty} f g_n \rightarrow 0$  for all  $f \in \mathcal{HK}$ . For then,  $\int_{-\infty}^{\infty} f g_n = \int_n^{\infty} f \rightarrow 0$ . In this case,  $g_n \in \mathcal{NBV}$  and  $V g_n = 1$ . However,  $\lambda(T_n) = \infty$  for all  $0 < \epsilon < 1$ . Note that for each compact interval  $[a, b]$  we have  $\int_a^b g_n \rightarrow 0$  and  $g_n \rightarrow 0$  in measure on  $[a, b]$ .

It is somewhat surprising that condition (2) is also necessary and sufficient to have  $\|f(g_n - g)\| \rightarrow 0$  for all  $f \in \mathcal{HK}$ .

**Theorem 6.** *Let  $[a, b]$  be a compact interval in  $\mathbb{R}$ , let  $g_n$  and  $g$  be real-valued functions on  $[a, b]$  with  $g$  of bounded variation. In order for  $\|f(g_n - g)\| \rightarrow 0$  for all  $f \in \mathcal{HK}$  it is necessary and sufficient that*

$$\left. \begin{aligned} &g_n \rightarrow g \text{ in measure as } n \rightarrow \infty, \\ &\text{for each } n \geq 1, g_n \text{ is equivalent to a function } h_n \in \mathcal{NBV}, \\ &\text{and there is } M \in [0, \infty) \text{ such that } V h_n \leq M \text{ for all } n \geq 1. \end{aligned} \right\} \quad (3)$$

*If  $(a, b) \subset \mathbb{R}$  is unbounded, then change the first line of (3) by requiring  $g_n \chi_I \rightarrow g \chi_I$  in measure for each compact interval  $I \in (a, b)$ .*

PROOF. Certainly (3) is necessary in order for  $\|f(g_n - g)\| \rightarrow 0$  for all  $f \in \mathcal{HK}$ .

If we have (3), let  $I_n$  be any sequence of intervals in  $(a, b)$ . We can again assume  $g = 0$ . Write  $\tilde{g}_n = g_n \chi_{I_n}$ . Then

$$\|\tilde{g}_n\|_\infty \leq \|g_n\|_\infty, V\tilde{g}_n \leq Vg_n + 2\|g_n\|_\infty \text{ and } \tilde{g}_n \rightarrow 0 \text{ in measure.}$$

The result now follows by applying Theorem 3 to  $f\tilde{g}_n$ .

Unbounded intervals are handled as in Theorem 3. □

By combining Theorem 3 and Theorem 6 we have the following.

**Theorem 7.** *Let  $(a, b) \subset \mathbb{R}$ . Then  $\int_a^b f g_n \rightarrow \int_a^b f g$  for all  $f \in \mathcal{HK}$  if and only if  $\|f g_n - f g\| \rightarrow 0$  for all  $f \in \mathcal{HK}$ .*

Note that  $\|f(g_n - g)\| \geq |\|f g_n\| - \|f g\||$ ; so if  $\|f(g_n - g)\| \rightarrow 0$ , then  $\|f g_n\| \rightarrow \|f g\|$ . Thus, (3) is sufficient to have  $\|f g_n\| \rightarrow \|f g\|$  for all  $f \in \mathcal{HK}$ . However, this condition is not necessary. For example, let  $[a, b] = [0, 1]$ . Define  $g_n(x) = (-1)^n$ . Then  $\|g_n\|_\infty = 1$  and  $Vg_n = 0$ . Let  $g = g_1$ . For no  $x \in [-1, 1]$  does the sequence  $g_n(x)$  converge to  $g(x)$ . For no open interval  $I \subset [0, 1]$  do we have  $\int_I (g_n - g) \rightarrow 0$ . And,  $g_n$  does not converge to  $g$  in measure. However, let  $f \in \mathcal{HK}$  with  $\|f\| > 0$ . Then  $\|f(g_n - g)\| = 0$  when  $n$  is odd and when  $n$  is even,  $\|f(g_n - g)\| = 2\|f\|$ . And yet, for all  $n$ ,  $\|f g_n\| = \|f\| = \|f g\|$ .

It is natural to ask what extra condition should be given so that  $\|f g_n\| \rightarrow \|f g\|$  will imply  $\|f g_n - f g\| \rightarrow 0$ . We have the following.

**Theorem 8.** *Let  $g_n \rightarrow g$  in measure or almost everywhere. Then  $\|f g_n\| \rightarrow \|f g\|$  for all  $f \in \mathcal{HK}$  if and only if  $\|f g_n - f g\| \rightarrow 0$  for all  $f \in \mathcal{HK}$ .*

PROOF. Let  $[a, b]$  be a compact interval. If  $\|f g_n\| \rightarrow \|f g\|$ , then  $g$  is equivalent to  $h \in \mathcal{NBV}$  [2, Theorem 12.9] and for each  $f \in \mathcal{HK}$  there is a constant  $C_f$  such that  $\|f g_n\| \leq C_f$ . By the Banach–Steinhaus Theorem [2, Theorem 12.10], each  $g_n$  is equivalent to a function  $h_n \in \mathcal{NBV}$  with  $Vh_n \leq M$  and  $\|h_n\|_\infty \leq M$ . Let  $(c, d) \subset (a, b)$ . By dominated convergence,  $\int_c^d g_n \rightarrow \int_c^d g$ . It now follows from Theorem 1 that  $\int_a^b f g_n \rightarrow \int_a^b f g$  for all  $f \in \mathcal{HK}$ . Hence, by Theorem 7,  $\|f g_n - f g\| \rightarrow 0$  for all  $f \in \mathcal{HK}$ .

Now suppose  $(a, b) = \mathbb{R}$  and  $\|f g_n\| \rightarrow \|f g\|$  for all  $f \in \mathcal{HK}$ . The change of variables  $x \mapsto 1/x$  shows the Banach–Steinhaus Theorem still holds on  $\mathbb{R}$ . We then have each  $g_n$  equivalent to  $h_n \in \mathcal{NBV}$  with  $Vh_n \leq M$  and  $\|h_n\|_\infty \leq M$ . As with the end of the proof of Theorem 3, given  $\epsilon > 0$  we can find  $c \in \mathbb{R}$  such that  $|\int_{-\infty}^c f g_n| < \epsilon$  for all  $n \geq 1$ . The other cases are similar. □

**Acknowledgment.** An anonymous referee provided reference [3] and pointed out that in place of convergence in measure we can use convergence in  $\|\cdot\|_1$  (cf. Remark 4).

## References

- [1] G. B. Folland, *Real analysis*, New York, Wiley, 1999.
- [2] P.-Y. Lee, *Lanzhou lectures on integration*, Singapore, World Scientific, 1989.
- [3] W. L. C. Sargent, *On some theorems of Hahn, Banach and Steinhaus*, J. London Math. Soc., **28** (1953), 438–451.
- [4] E. Talvila, *Henstock–Kurzweil Fourier transforms*, Illinois J. Math., **46** (2002), 1207–1226.