

M. Federson, Departamento de Matemática, Universidade de São Paulo, CP 688, São Carlos - SP, 13560-970, Brazil. email: federson@icmc.sc.usp.br

SOME PECULIARITIES OF THE HENSTOCK AND KURZWEIL INTEGRALS OF BANACH SPACE-VALUED FUNCTIONS

Abstract

Some examples, due to G. Birkhoff, are used to explore the differences and peculiarities of the Henstock and Kurzweil integrals in abstract spaces. We also include a proof, due to C. S. Hönl, of the fact that the Bochner-Lebesgue integral is equivalent to the variational Henstock-McShane integral.

1 Introduction

In 1988, Professor Stefan Schwabik came to Brazil on a visit to Professor Chaim Samuel Hönl and Professor Luciano Barbanti. On that occasion, Professor Schwabik gave a series of lectures on generalized ODE's which motivated Professor Hönl to deal with the Henstock-Kurzweil integration theory for some years. In 1993, in a course on the subject at the University of São Paulo, São Paulo, Brazil, Professor Hönl presented some examples borrowed from [1] in order to clarify the differences and peculiarities of the integrals defined by Henstock ([12]) and by Kurzweil ([19]) for Banach space-valued functions. The notes on such examples are contained here. We also include a proof, due to Hönl ([17]), of the fact that the Bochner-Lebesgue integral is equivalent to the variational Henstock-McShane integral.

2 Basic Definitions and Terminology

Let $[a, b]$ be a compact interval of the real line \mathbb{R} . A *division* of $[a, b]$ is any finite set of closed non-overlapping intervals $[t_{i-1}, t_i] \subset [a, b]$ such that $\cup_i [t_{i-1}, t_i] = [a, b]$. We write $(t_i) \in D_{[a,b]}$ in this case. When $(t_i) \in D_{[a,b]}$ and $\xi_i \in [t_{i-1}, t_i]$

Key Words: Henstock-Kurzweil integral, Bochner-Lebesgue integral, McShane integral
Mathematical Reviews subject classification: 26A39
Received by the editors July 15, 2002
Communicated by: Peter Bullen

for every i , then (ξ_i, t_i) is a *tagged division* of $[a, b]$. By $TD_{[a,b]}$ we mean the set of all tagged divisions of $[a, b]$.

A *gauge* of $[a, b]$ is any function $\delta : [a, b] \rightarrow]0, \infty[$. Given a gauge δ of $[a, b]$, we say $(\xi_i, t_i) \in TD_{[a,b]}$ is δ -*fine*, if $[t_{i-1}, t_i] \subset \{t \in [a, b]; |t - \xi_i| < \delta(\xi_i)\}$ for every i .

In what follows X denotes a Banach space.

A function $f : [a, b] \rightarrow X$ is integrable in the sense of Kurzweil or *Kurzweil integrable* (we write $f \in K([a, b], X)$) and $I = (K) \int_a^b f = (K) \int_a^b f(t) dt \in X$ is its integral if given $\varepsilon > 0$, there is a gauge δ of $[a, b]$ such that for every δ -fine $(\xi_i, t_i) \in TD_{[a,b]}$,

$$\left\| (K) \int_a^b f - \sum_i f(\xi_i)(t_i - t_{i-1}) \right\| < \varepsilon.$$

As it should be expected, the Kurzweil integral is linear and additive over non-overlapping intervals. The basic literature on this subject includes [11], [14], [20], [21], [22], [23], [26].

We use the notation “ \sim ” to indicate the indefinite integral of a function $f \in K([a, b], X)$, that is, $\tilde{f} : [a, b] \rightarrow X$ is given by $\tilde{f}(t) = (K) \int_a^t f(s) ds$ for all $t \in [a, b]$. We have $\tilde{f} \in \mathcal{C}([a, b], X)$ (see [6] for instance), where $\mathcal{C}([a, b], X)$ is the Banach space of all continuous functions $f : [a, b] \rightarrow X$ equipped with the usual supremum norm, $\|f\|_\infty$.

A function $f : [a, b] \rightarrow X$ is integrable in the sense of Henstock or *Henstock integrable* or even *variationally Henstock integrable* (we write $f \in H([a, b], X)$) if given $\varepsilon > 0$, there is a function $F : [a, b] \rightarrow X$ and a gauge δ of $[a, b]$ such that for every δ -fine $(\xi_i, t_i) \in TD_{[a,b]}$,

$$\sum_i \|F(t_i) - F(t_{i-1}) - f(\xi_i)(t_i - t_{i-1})\| < \varepsilon.$$

In this case, we write $(H) \int_a^t f = F(t) - F(a)$, $t \in [a, b]$.

Let $R([a, b], X)$ be the space of abstract Riemann integrable functions $f : [a, b] \rightarrow X$ with integral $\int_a^b f$. It is immediate that

$$H([a, b], X) \subset K([a, b], X) \text{ and } R([a, b], X) \subset K([a, b], X),$$

and the integrals coincide when they exist.

Two functions $g, f \in K([a, b], X)$ are called *equivalent*, whenever $\tilde{g}(t) = \tilde{f}(t)$ for all $t \in [a, b]$. When this is the case, $K([a, b], X)_A$ denotes the space of all equivalence classes of functions of $K([a, b], X)$ endowed with the Alexiewicz

norm

$$f \in K([a, b], X) \mapsto \|f\|_A = \|\tilde{f}\|_\infty = \sup_{t \in [a, b]} \left\| (K) \int_a^t f(s) ds \right\|.$$

In an analogous way, $H([a, b], X)_A$ denotes the space of all equivalence classes of functions of $H([a, b], X)$ endowed with the Alexiewicz norm.

If $g, f \in H([a, b], X)$ are equivalent, then $g = f$ almost everywhere in the sense of the Lebesgue measure ([7]). On the other hand, we may have $f \in R([a, b], X) \setminus H([a, b], X)$ (i.e., f belongs to $R([a, b], X)$ but not to $H([a, b], X)$) such that $\tilde{f} = 0$ but $f(t) \neq 0$ for almost every $t \in [a, b]$ (see Example 2.1). Thus $g, f \in R([a, b], X) \subset K([a, b], X)$ and f equivalent to g do not imply $g = f$ almost everywhere.

Let $I \subset \mathbb{R}$ be an arbitrary set and let E be a normed space. A family $(x_i)_{i \in I}$ of elements of E is *summable* with sum $x \in E$ (we write $\sum_{i \in I} x_i = x$) if for every $\varepsilon > 0$, there is a finite subset $F_\varepsilon \subset I$ such that for every finite subset $F \subset I$ with $F \supset F_\varepsilon$,

$$\|x - \sum_{i \in F} x_i\| < \varepsilon.$$

Let $l_2(I)$ be the set of all families $(x_i)_{i \in I}$, $x_i \in \mathbb{R}$, such that the family $(|x_i|^2)_{i \in I}$ is summable. We write

$$l_2(I) = \left\{ x = (x_i)_{i \in I}, x_i \in \mathbb{R}; \sum_{i \in I} |x_i|^2 < \infty \right\}.$$

The expression

$$\langle x, y \rangle = \sum_{i \in I} x_i y_i$$

defines an inner product and $l_2(I)$ equipped with the norm

$$\|x\|_2 = \left(\sum_{i \in I} |x_i|^2 \right)^{1/2}$$

is a Hilbert space. Moreover by the Basis Theorem $\{e_i; i \in I\}$, where

$$e_i(j) = \begin{cases} 1, & j = i \\ 0, & j \neq i \end{cases},$$

is a complete orthonormal system for $l_2(I)$. We refer to the relation

$$\|x\|_2^2 = \sum_{i \in I} |\langle x_i, e_i \rangle|^2 = \sum_{i \in I} |x_i|^2, \quad \forall x \in l_2(I),$$

as the Bessel equality.

Example 2.1. Let $[a, b]$ be non-degenerate and $X = l_2([a, b])$ be equipped with the norm

$$x \mapsto \|x\|_2 = \left(\sum_{i \in [a, b]} |x_i|^2 \right)^{1/2}.$$

Consider a function $f : [a, b] \rightarrow X$ given by $f(t) = e_t$, $t \in [a, b]$. Given $\varepsilon > 0$, there exists $\delta > 0$, with $\delta^{\frac{1}{2}} < \frac{\varepsilon}{(b-a)^{\frac{1}{2}}}$, such that for every $(\frac{\delta}{2})$ -fine $(\xi_j, t_j) \in TD_{[a, b]}$,

$$\begin{aligned} \left\| \sum_j f(\xi_j)(t_j - t_{j-1}) - 0 \right\|_2 &= \left\| \sum_j e_{\xi_j}(t_j - t_{j-1}) \right\|_2 = \left[\sum_j |t_j - t_{j-1}|^2 \right]^{\frac{1}{2}} < \\ &< \delta^{\frac{1}{2}} \left[\sum_j (t_j - t_{j-1}) \right]^{\frac{1}{2}} < \varepsilon \end{aligned}$$

where we applied the Bessel equality. Thus $f \in R([a, b], X) \subset K([a, b], X)$ and $\tilde{f} = 0$, since $\int_a^t f(s)ds = 0$ for every $t \in [a, b]$.

If $f \in H([a, b], X)$, then $(H) \int_a^t f = 0$ for every $t \in [a, b]$, since $H([a, b], X) \subset K([a, b], X)$ and $(H) \int_a^t f = (K) \int_a^t f = \int_a^t f = 0$. But

$$\sum_i \|f(\xi_i)(t_i - t_{i-1}) - 0\|_2 = b - a$$

for every $(\xi_i, t_i) \in TD_{[a, b]}$. Hence $f \notin H([a, b], X)$. \square

Let $\mathcal{L}_1([a, b], X)$ be the space of Bochner-Lebesgue integrable functions $f : [a, b] \rightarrow X$ with finite absolute Lebesgue integral, that is, $(L) \int_a^b \|f\| < \infty$. We denote by $(L) \int_a^b f$ the Bochner-Lebesgue integral of $f \in \mathcal{L}_1([a, b], X)$ (and also the Lebesgue integral of $f \in \mathcal{L}_1([a, b], \mathbb{R})$). The inclusion

$$\mathcal{L}_1([a, b], X) \subset H([a, b], X)$$

always holds (see [4], [17] or the Appendix).

In particular,

$$R([a, b], \mathbb{R}) \subset \mathcal{L}_1([a, b], \mathbb{R}) \subset H([a, b], \mathbb{R}) = K([a, b], \mathbb{R})$$

(see [23], for instance, for a proof of the equality). On the other hand, when X is a general Banach space it is possible to find a function $f : [a, b] \rightarrow X$ which is abstract Riemann integrable but not Bochner-Lebesgue integrable. Both Examples 2.1 and 3.1 in the sequel show functions $f \in R([a, b], X) \setminus H([a, b], X)$ (i.e., f belongs to $R([a, b], X)$ but not to $H([a, b], X)$). In particular, such functions belong to $R([a, b], X) \setminus \mathcal{L}_1([a, b], X)$ and also to $K([a, b], X) \setminus H([a, b], X)$.

When real-valued functions are considered only, the Lebesgue integral is equivalent to a modified version of the Kurzweil integral. The idea of slightly modifying Kurzweil's definition is due to E. J. McShane ([24], [25]). Instead of taking δ -fine tagged divisions, McShane considered what we call δ -fine *semi-tagged divisions* (ξ_i, t_i) of $[a, b]$, that is $(t_i) \in D_{[a, b]}$ and $[t_{i-1}, t_i] \subset \{t \in [a, b] : |t - \xi_i| < \delta(\xi_i)\}$ for every i . In this case, we write $(\xi_i, t_i) \in STD_{[a, b]}$. Notice that in the definition of semi-tagged divisions, it is *not* required that $\xi_i \in [t_{i-1}, t_i]$ for *any* i . In this manner, McShane's modification of the Kurzweil integral gives an elegant characterization of the Lebesgue integral through Riemann sums (see the Appendix).

Let us denote by $KMS([a, b], \mathbb{R})$ the space of real-valued Kurzweil-McShane integrable functions $f : [a, b] \rightarrow \mathbb{R}$, that is, $f \in KMS([a, b], \mathbb{R})$ is integrable in the sense of Kurzweil with the modification of McShane. Formally, $f \in KMS([a, b], \mathbb{R})$ if and only if there exists $I \in \mathbb{R}$ such that for every $\varepsilon > 0$, there is a gauge δ of $[a, b]$ such that

$$\left| I - \sum_i f(\xi_i)(t_i - t_{i-1}) \right| < \varepsilon.$$

whenever $(\xi_i, t_i) \in STD_{[a, b]}$ is δ -fine. This definition can be extended to Banach space-valued functions.

We have

$$R([a, b], \mathbb{R}) \subset \mathcal{L}_1([a, b], \mathbb{R}) = KMS([a, b], \mathbb{R}) \subset K([a, b], \mathbb{R}) = H([a, b], \mathbb{R}).$$

Furthermore, $K([a, b], \mathbb{R}) \setminus \mathcal{L}_1([a, b], \mathbb{R}) \neq \emptyset$. The next classical example exhibits an $f \in K([a, b], \mathbb{R}) \setminus \mathcal{L}_1([a, b], \mathbb{R})$.

Example 2.2. Let $F(t) = t^2 \sin(t^{-2})$ for $t \in]0, 1]$ and $F(0) = 0$. Let $f = \frac{d}{dt}F$. Because f is Riemann improper integrable, it follows that $f \in K([a, b], \mathbb{R}) = H([a, b], \mathbb{R})$, since the Kurzweil and the Henstock integrals contain their improper integrals (see [21], Cauchy Extension). However $f \notin \mathcal{L}_1([a, b], \mathbb{R})$ (see [28]).

Example 2.2 says $K([a, b], \mathbb{R}) = H([a, b], \mathbb{R})$ is not an absolute integrable space. More generally, $H([a, b], X)$ and hence $K([a, b], X)$ are non-absolute integrable spaces (see Example 3.4 and Lemma 4.3 in the Appendix).

The generalization of the Riemannian characterization of the Banach space-valued Lebesgue-type integral, namely the Bochner-Lebesgue integral, is not straightforward. In fact, Example 3.1 shows that the modification of McShane applied to the abstract Kurzweil integral can give a more general space than that of Bochner-Lebesgue. On the other hand, if McShane's idea is used to modify the variational definition of Henstock, then we obtain a Riemannian definition of the Bochner-Lebesgue integral (see [4], [17] or the Appendix). Thus, if $HMS([a, b], X)$ denotes the space of Henstock-McShane integrable functions $f : [a, b] \rightarrow X$, that is, $f \in HMS([a, b], X)$ is integrable in the sense of Henstock with the modification of McShane, then

$$HMS([a, b], X) = \mathcal{L}_1([a, b], X).$$

In addition,

$$\begin{cases} HMS([a, b], X) \subset H([a, b], X), \\ KMS([a, b], X) \subset K([a, b], X) \text{ and} \\ RMS([a, b], X) \subset R([a, b], X), \end{cases}$$

where $KMS([a, b], X)$ and $RMS([a, b], X)$ denote, respectively, the spaces of Kurzweil-McShane and Riemann-McShane integrable functions $f : [a, b] \rightarrow X$.

For other interesting results, the reader may want to consult [5].

3 Birkhoff's Examples

The first example of this section shows a Banach space-valued function which is integrable in the sense of Riemann-McShane, but not integrable in the variational sense of Henstock (and neither in the Bochner-Lebesgue sense).

Example 3.1. Let $G([a, b], X)$ be the Banach space, endowed with the usual supremum norm, $\|\cdot\|_\infty$, of all *regulated functions* $f : [a, b] \rightarrow X$ (i.e., f has discontinuities of the first kind only - see [16], p. 16). Let $X = G^-([0, 1], \mathbb{R})$, where

$$G^-([0, 1], \mathbb{R}) = \{f \in G([0, 1], \mathbb{R}); f \text{ is left continuous}\},$$

and consider the function

$$f : t \in [0, 1] \mapsto f(t) = 1_{[t, 1]} \in X,$$

where 1_A denotes the characteristic function of a set $A \subset [0, 1]$. Since f is a function of weak bounded variation (we write $f \in BW([0, 1], X)$ - see [16], p. 23) and $\phi(t) = t$, $t \in [0, 1]$, is an element of $\mathcal{C}([0, 1], \mathbb{R})$, it follows from [16], Theorem 4.6, p. 24, that the abstract Riemann-Stieltjes integral,

$\int_0^1 df \phi$, exists. Moreover, the Riemann-Stieltjes integral, $\int_0^1 f d\phi$, exists and the integration by parts formula

$$\int_0^1 f(t)dt = \int_0^1 f d\phi = f(t) \cdot t|_0^1 - \int_0^1 df \phi$$

holds (see [16], Theorem 1.3, p. 18). Hence $f \in R([0, 1], X) \subset K([0, 1], X)$. The indefinite integral $\tilde{f}(t) = \int_0^t f(r)dr$, $t \in [0, 1]$, of f is given by $\tilde{f}(t)(s) = t \wedge s = \inf \{t, s\}$, since

$$\left(\int_0^t f(r)dr \right)(s) = \left(\int_0^t 1_{[r,1]}dr \right)(s) = \int_0^t 1_{[r,1]}(s)dr = \int_0^{t \wedge s} dr = t \wedge s.$$

Hence, \tilde{f} is absolutely continuous. However \tilde{f} is nowhere differentiable as we will show later. Then the Lebesgue Theorem implies $f \notin \mathcal{L}_1([0, 1], X)$. More generally, $f \notin H([0, 1], X)$ by the Fundamental Theorem of Calculus for the Henstock integral (see [7]). Or we can prove directly that $f \notin H([0, 1], X)$, since

$$\left\| f(\xi_i)(t_i - t_{i-1}) - \int_{t_{i-1}}^{t_i} f \right\| \geq \frac{1}{2} (t_i - t_{i-1}),$$

for every $(\xi_i, t_i) \in TD_{[0,1]}$. Thus $f \in R([0, 1], X) \setminus H([0, 1], X)$ and, in particular, $f \in R([0, 1], X) \setminus \mathcal{L}_1([0, 1], X)$. Moreover, we assert that $f \in RMS([0, 1], X)$, that is, f is Riemann-McShane integrable. It is enough to show that for every $\varepsilon > 0$, there exists $\delta > 0$ such that for every δ -fine $(\xi_i, t_i) \in STD_{[0,1]}$,

$$\left\| \tilde{f}(1) - \sum_i f(\xi_i)(t_i - t_{i-1}) \right\| < \varepsilon.$$

Given $\varepsilon > 0$, let $0 < \delta < \varepsilon$ and suppose $(\xi_i, t_i) \in STD_{[0,1]}$ is δ -fine. If $\xi_i \leq s$ and $t_i < \xi_i + \delta$, then $t_i < s + \delta$ which implies $\sum_{\xi_i \leq s} (t_i - t_{i-1}) < s + \delta$ and then

$$s - \sum_{\xi_i \leq s} (t_i - t_{i-1}) < \delta. \quad (1)$$

If $\xi_j > s$ and $t_{j-1} > \xi_j - \delta$, then $t_{j-1} > s - \delta$ and therefore $\sum_{\xi_j > s} (t_j - t_{j-1}) < 1 - (s - \delta) = \sum_i (t_i - t_{i-1}) - s + \delta$. Then $0 \leq \sum_{\xi_i \leq s} (t_i - t_{i-1}) + \delta - s$ which

implies

$$s - \sum_{\xi_i \leq s} (t_i - t_{i-1}) < \delta. \quad (2)$$

By (1) and (2), we have

$$\begin{aligned} \left\| \tilde{f}(1) - \sum_i f(\xi_i)(t_i - t_{i-1}) \right\|_\infty &= \sup_{0 \leq s \leq 1} \left| \tilde{f}(1)(s) - \sum_i f(\xi_i)(s)(t_i - t_{i-1}) \right| = \\ &= \sup_{0 \leq s \leq 1} \left| s - \sum_{\xi_i \leq s} (t_i - t_{i-1}) \right| < \delta < \varepsilon \end{aligned}$$

and the assertion follows.

Now we give a proof of the fact that \tilde{f} is neither strongly nor weakly differentiable. We begin by showing that \tilde{f} is not strongly differentiable in the sense that the limit

$$\lim_{\varepsilon_1 \rightarrow 0_+, \varepsilon_2 \rightarrow 0_+} \left[\frac{\tilde{f}(t + \varepsilon_2) - \tilde{f}(t)}{\varepsilon_2} - \frac{\tilde{f}(t + \varepsilon_1) - \tilde{f}(t)}{\varepsilon_1} \right], \quad t \in [0, 1[,$$

does not exist. In an analogous way, it can be proved that the limit

$$\lim_{\varepsilon_1 \rightarrow 0_-, \varepsilon_2 \rightarrow 0_-} \left[\frac{\tilde{f}(t) - \tilde{f}(t + \varepsilon_2)}{\varepsilon_2} - \frac{\tilde{f}(t) - \tilde{f}(t + \varepsilon_1)}{\varepsilon_1} \right], \quad t \in]0, 1],$$

does not exist.

For $0 < \varepsilon_1 < \varepsilon_2$, we have

$$\begin{aligned} \left\| \frac{\tilde{f}(t + \varepsilon_2) - \tilde{f}(t)}{\varepsilon_2} - \frac{\tilde{f}(t + \varepsilon_1) - \tilde{f}(t)}{\varepsilon_1} \right\| &= \\ &= \sup_{0 \leq s \leq 1} \left| \frac{(t + \varepsilon_2) \wedge s - t \wedge s}{\varepsilon_2} - \frac{(t + \varepsilon_1) \wedge s - t \wedge s}{\varepsilon_1} \right| \\ &\geq \left| \frac{(t + \varepsilon_2) \wedge s - t \wedge s}{\varepsilon_2} - \frac{(t + \varepsilon_1) \wedge s - t \wedge s}{\varepsilon_1} \right|_{s=t+\varepsilon_1} \\ &= \left| \frac{t + \varepsilon_1 - t}{\varepsilon_2} - \frac{t + \varepsilon_1 - t}{\varepsilon_1} \right| = \left| \frac{\varepsilon_1}{\varepsilon_2} - 1 \right| \rightarrow 1, \end{aligned}$$

as we suppose, without loss of generality, that ε_1 goes faster than ε_2 to zero.

Let us show that \tilde{f} is not weakly differentiable in the following sense: if Y is a Banach space and Y' is its topological dual, then $g : [a, b] \rightarrow Y$ is *weakly*

right differentiable at a point $t \in [a, b[$ with *weak right derivative* denoted by $\frac{d^{\sigma^+}g(t)}{dt}$ whenever for every $y' \in Y'$,

$$\lim_{\varepsilon \rightarrow 0_+} \left\langle \frac{g(t+\varepsilon) - g(t)}{\varepsilon}, y' \right\rangle = \left\langle \frac{d^{\sigma^+}g(t)}{dt}, y' \right\rangle.$$

Analogously we define the *weak left derivative* of g at a point $t \in]a, b]$.

Let $BV_0([0, 1], \mathbb{R})$ be the Banach space of all functions $h : [0, 1] \rightarrow \mathbb{R}$ of bounded variation which vanish at $t = 0$ equipped with the norm given by the variation of h , $V(h)$. Then $BV_0([0, 1], \mathbb{R}) = G^-([0, 1], \mathbb{R})'$ (see [16], Theorem 4.12, p. 26). Besides, for every $\alpha \in BV_0([0, 1], \mathbb{R})$, the Riemann-Stieltjes integral, $\int_0^1 \tilde{f} d\alpha$, exists (see [16]), since \tilde{f} is continuous. Given $\alpha \in BV_0([0, 1], \mathbb{R})$, we will show that

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0_+} \left\langle \frac{1}{\varepsilon} [\tilde{f}(t+\varepsilon) - \tilde{f}(t)], \alpha \right\rangle &= \lim_{\varepsilon \rightarrow 0_+} \int_0^1 \frac{1}{\varepsilon} [\tilde{f}(t+\varepsilon) - \tilde{f}(t)](s) d\alpha(s) \\ &= [\alpha(1) - \alpha(t+)], \end{aligned}$$

where $\alpha(t+)$ denotes the right lateral limit of α at $t \in [0, 1[$. We have

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0_+} \int_0^1 \frac{1}{\varepsilon} [\tilde{f}(t+\varepsilon) - \tilde{f}(t)](s) d\alpha(s) &= \lim_{\varepsilon \rightarrow 0_+} \int_0^1 \frac{1}{\varepsilon} [(t+\varepsilon) \wedge s - t \wedge s] d\alpha(s) \\ &= \lim_{\varepsilon \rightarrow 0_+} \int_t^{t+\varepsilon} \frac{1}{\varepsilon} (s-t) d\alpha(s) + \lim_{\varepsilon \rightarrow 0_+} \int_{t+\varepsilon}^1 \frac{1}{\varepsilon} [(t+\varepsilon) - t] d\alpha(s) \\ &= \lim_{\varepsilon \rightarrow 0_+} \int_t^{t+\varepsilon} \frac{1}{\varepsilon} (s-t) d\alpha(s) + \alpha(1) - \alpha(t+). \end{aligned}$$

But

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0_+} \int_t^{t+\varepsilon} \frac{1}{\varepsilon} (s-t) d\alpha(s) &= \lim_{\varepsilon \rightarrow 0_+} \frac{1}{\varepsilon} \left[\int_t^{t+\varepsilon} s d\alpha(s) - \int_t^{t+\varepsilon} t d\alpha(s) \right] \\ &= \lim_{\varepsilon \rightarrow 0_+} \frac{1}{\varepsilon} \left[s\alpha(s) \Big|_t^{t+\varepsilon} - \int_t^{t+\varepsilon} \alpha(s) ds - t\alpha(t+\varepsilon) + t\alpha(t) \right] \\ &= \alpha(t+) - \lim_{\varepsilon \rightarrow 0_+} \frac{1}{\varepsilon} \int_t^{t+\varepsilon} \alpha(s) ds = 0, \end{aligned}$$

where we applied the integration by parts formula to obtain the second equality. Hence,

$$\lim_{\varepsilon \rightarrow 0+} \int_0^1 \frac{1}{\varepsilon} [\tilde{f}(t+\varepsilon) - \tilde{f}(t)](s) d\alpha(s) = \alpha(1) - \alpha(t+).$$

In a similar way, it can be proved that

$$\left\langle \frac{1}{\varepsilon} [\tilde{f}(t) - \tilde{f}(t-\varepsilon)], \alpha \right\rangle \longrightarrow \alpha(t-) - \alpha(1),$$

as $\varepsilon \rightarrow 0+$, where $\alpha(t-)$ denotes the left lateral limit of α at $t \in]0, 1]$. Therefore, we showed that \tilde{f} is not weakly differentiable. \square

As we mentioned before, the inclusion $\mathcal{L}_1([a, b], X) \subset KMS([a, b], X)$ always holds. When $X = G^-(]0, 1], \mathbb{R})$, for instance, one can find a function $f \in KMS([a, b], X) \setminus \mathcal{L}_1([a, b], X)$ (see Example 3.1). In general, $KMS([a, b], X) \setminus \mathcal{L}_1([a, b], X) \neq \emptyset$ for X of infinite dimension as we show next.

Proposition 3.1 (Hönig). *If X is an infinite dimensional Banach space, then there exists $f \in KMS([a, b], X) \setminus \mathcal{L}_1([a, b], X)$.*

PROOF. Let $\dim X$ denote the dimension of X . If $\dim X = \infty$, then the Theorem of Dvoretzky-Rogers implies there exists a sequence $(x_n)_{n \in \mathbb{N}}$ in X which is summable but not absolutely summable. Thus, if we define a function $f : [1, \infty] \rightarrow X$ by $f(t) = x_n$, whenever $n \leq t < n+1$, then $(KMS) \int_a^b f = \sum_n x_n \in X$ if the integral exists (here, $(KMS) \int$ denotes the KMS integral). On the other hand, $f \notin \mathcal{L}_1([a, b], X)$, since $(L) \int_a^b \|f\| = \|x_1\| + \|x_2\| + \|x_3\| \dots = \infty$. \square

The next example exhibits a function which is integrable in the sense of Kurzweil but not in Henstock's sense. It also shows that the Monotone Convergence Theorem, which holds for monotone ordered normed space-valued Kurzweil integrals ([8]), may not be valid for Henstock integrals.

Example 3.2. Consider the space

$$Z = l_2(\mathbb{N} \times \mathbb{N}) = \left\{ z = (z_{ij})_{i,j \in \mathbb{N}}, z_{ij} \in \mathbb{R}; \sum_{i,j=1}^{\infty} |z_{ij}|^2 < \infty \right\}$$

equipped with the norm

$$z \mapsto \|z\|_2 = \left(\sum_{i,j=1}^{\infty} |z_{ij}|^2 \right)^{1/2}$$

and the function

$$f : [0, 1] \rightarrow Z$$

given by $f = \sum_{i=1}^{\infty} f_i$, where $f_i(t) = 2^i e_{ij}$ whenever $\frac{j}{2^i} \leq t < \frac{j}{2^i} + \frac{1}{2^{2i}}$, $j = 0, 1, 2, \dots, 2^i - 1$, and $f_i(t) = 0$ otherwise. By e_{ij} we mean the doubly infinite set of orthonormal vectors of Z . We have

$$f_1(t) = \begin{cases} 2e_{10}; & 0 \leq t < 1/4, \\ 2e_{11}; & 1/2 \leq t < 3/4, \\ 0; & 1/4 \leq t < 1/2 \text{ or } 3/4 \leq t \leq 1. \end{cases}$$

Hence,

$$\int_0^1 f_1 = \int_0^{1/4} 2e_{10} + \int_{1/2}^{3/4} 2e_{11} = \frac{1}{2} e_{10} + \frac{1}{2} e_{11}$$

and therefore,

$$\begin{aligned} \|f_1\|_A &= \sup_{0 \leq t \leq 1} \left\| \int_0^t f_1 \right\|_2 = \left\| \int_0^1 f_1 \right\|_2 = \left\| \frac{1}{2} e_{10} + \frac{1}{2} e_{11} \right\|_2 \\ &= \left[\left(\frac{1}{2} \right)^2 + \left(\frac{1}{2} \right)^2 \right]^{\frac{1}{2}} = \left(\frac{1}{2} \right)^{\frac{1}{2}}. \end{aligned}$$

Also,

$$f_2(t) = \begin{cases} 4e_{20}; & 0 \leq t < 1/16, \\ 4e_{21}; & 1/4 \leq t < 5/16, \\ 4e_{22}; & 1/2 \leq t < 9/16, \\ 4e_{23}; & 3/4 \leq t < 13/16, \\ 0; & \text{otherwise.} \end{cases}$$

Then,

$$\int_0^1 f_2 = \frac{1}{4} e_{20} + \frac{1}{4} e_{21} + \frac{1}{4} e_{22} + \frac{1}{4} e_{23}$$

and

$$\begin{aligned} \|f_1 + f_2\|_A &= \sup_{0 \leq t \leq 1} \left\| \int_0^t (f_1 + f_2) \right\|_2 = \left\| \int_0^1 f_1 + \int_0^1 f_2 \right\|_2 = \\ &= \left\| \frac{1}{2} e_{10} + \frac{1}{2} e_{11} + \frac{1}{4} e_{20} + \frac{1}{4} e_{21} + \frac{1}{4} e_{22} + \frac{1}{4} e_{23} \right\|_2 = \left[\frac{1}{2} + \frac{1}{4} \right]^{\frac{1}{2}}. \end{aligned}$$

By induction, it can be proved that

$$\|f_1 + f_2 + \dots + f_n\|_A = \left[\frac{1}{2} + \frac{1}{2^2} + \dots + \frac{1}{2^n} \right]^{\frac{1}{2}} < 1.$$

for every $n \in \mathbb{N}$. Thus, if we define $g_n = \sum_{i=1}^n f_i$, for every $n \in \mathbb{N}$, then the sequence $(\|g_n\|_A)_{n \in \mathbb{N}}$ is bounded. Besides, $g_n(t) \leq g_{n+1}(t) \leq f(t)$ for all $n \in \mathbb{N}$ and $t \in [0, 1]$. Hence the Monotone Convergence Theorem (see [8]) implies $f \in K([0, 1], Z)$ and $\int_0^1 g_n \rightarrow (K) \int_0^1 f$ as $n \rightarrow \infty$.

Since the Monotone Convergence Theorem also holds for the Kurzweil-McShane integral with obvious adaptations, it follows that $f \in KMS([0, 1], Z)$.

On the other hand, although $g_n \in H([0, 1], Z)$ for every $n \in \mathbb{N}$, Birkhoff asserted in [1] that the indefinite integral f of f is nowhere differentiable and, therefore, $f \notin H([0, 1], Z)$ by the Fundamental Theorem of Calculus for the Henstock integral (see [7]). \square

It is known that the space of all equivalence classes of real-valued Kurzweil (or Henstock) integrable functions, equipped with the Alexiewicz norm, is non-complete ([2]). More generally, $K([a, b], X)_A$ and $H([a, b], X)_A$ are non-complete spaces. However such spaces are ultrabornological ([9]) and, therefore, they have good functional analytic properties (see [18] for instance). The next example shows a Cauchy sequence, in the Alexiewicz norm, of Henstock integrable functions which is not convergent.

Example 3.3. Consider functions

$$f_n : [0, 1] \rightarrow l_2(\mathbb{N} \times \mathbb{N}), \quad n \in \mathbb{N}$$

defined by $f_n = \sum_{i=1}^n g_i$, where $g_i(t) = e_{ij}$ whenever $\frac{j-1}{2^i} \leq t < \frac{j}{2^i}$, $j = 1, 2, \dots, 2^i$, and $g_i(t) = 0$ otherwise. We have

$$g_1(t) = \begin{cases} e_{11}; & 0 \leq t < 1/2, \\ e_{12}; & 1/2 \leq t < 1, \\ 0; & t = 1. \end{cases}$$

Hence,

$$\begin{aligned} \|g_1\|_A &= \sup_{0 \leq t \leq 1} \left\| \int_0^t g_1 \right\|_2 = \left\| \int_0^1 g_1 \right\|_2 = \left\| \frac{1}{2} e_{11} + \frac{1}{2} e_{12} \right\|_2 \\ &= \left[\left(\frac{1}{2} \right)^2 + \left(\frac{1}{2} \right)^2 \right]^{\frac{1}{2}} = \left(\frac{1}{2} \right)^{\frac{1}{2}}. \end{aligned}$$

Also,

$$g_2(t) = \begin{cases} e_{21}; & 0 \leq t < 1/4, \\ e_{22}; & 1/4 \leq t < 1/2, \\ e_{23}; & 1/2 \leq t < 3/4, \\ e_{24}; & 3/4 \leq t < 1, \\ 0; & t = 1. \end{cases}$$

Then,

$$\int_0^1 g_2 = \int_0^{\frac{1}{4}} e_{21} + \int_{\frac{1}{4}}^{\frac{1}{2}} e_{22} + \int_{\frac{1}{2}}^{\frac{3}{4}} e_{23} + \int_{\frac{3}{4}}^1 e_{24} = \frac{1}{4} (e_{21} + e_{22} + e_{23} + e_{24}).$$

and therefore

$$\|g_2\|_A = \sup_{0 \leq t \leq 1} \left\| \int_0^t g_2 \right\|_2 = \left\| \int_0^1 g_2 \right\|_2 = \left(4 \frac{1}{4^2} \right)^{\frac{1}{2}} = \left(\frac{1}{4} \right)^{\frac{1}{2}}.$$

By induction, one can show that

$$\|g_i\|_A = \left\| \sum_{j=1}^{2^i} \int_{\frac{j-1}{2^i}}^{\frac{j}{2^i}} e_{ij} \right\|_2 = \left[2^i \left(\frac{1}{2^i} \right)^2 \right]^{\frac{1}{2}} = \frac{1}{2^{\frac{i}{2}}},$$

for every $i \in \mathbb{N}$. Then

$$\|f_n - f_m\|_A = \left\| \sum_{i=n+1}^m g_i \right\|_A \leq \sum_{i=n+1}^m \frac{1}{2^{\frac{i}{2}}}$$

which goes to zero for sufficiently large $n, m \in \mathbb{N}$, with $n > m$. Thus $(f_n)_{n \in \mathbb{N}}$ is a $\|\cdot\|_A$ -Cauchy sequence.

On the other hand,

$$\|f_n(t)\|_2 = \|g_1(t) + g_2(t) + \dots + g_n(t)\|_2 = \sqrt{n},$$

for every $t \in [0, 1]$. Hence there is no function $f(t) \in l_2(\mathbb{N} \times \mathbb{N})$, $t \in [0, 1]$, such that $\lim_{n \rightarrow \infty} \|f_n - f\|_A = 0$. \square

The next example presents a Banach space-valued function which is both Henstock and Kurzweil-McShane integrable but is not absolutely integrable.

Example 3.4. Let $f : [0, 1] \rightarrow l_2(\mathbb{N})$ be given by $f(t) = \frac{2^i}{i} e_i$, whenever $\frac{1}{2^i} \leq t < \frac{1}{2^{i-1}}$, $i = 1, 2, \dots$. Then

$$\int_{\frac{1}{2^i}}^{\frac{1}{2^{i-1}}} \frac{2^i}{i} e_i dt = \frac{1}{i} e_i$$

which is summable in $l_2(\mathbb{N})$. Since the Henstock integral contains its improper integrals (and the same applies to the Kurzweil integral), we have

$f \in H([0, 1], l_2(\mathbb{N}))$. However, $f \notin \mathcal{L}_1([0, 1], l_2(\mathbb{N}))$ because the sequence $(\frac{1}{i} e_i)_{i \in \mathbb{N}}$ is not summable in $\mathcal{L}_1([0, 1], l_2(\mathbb{N}))$. By the Monotone Convergence Theorem for the Kurzweil-McShane integral (which follows the ideas of [8] with obvious adaptations), $f \in KMS([0, 1], l_2(\mathbb{N}))$. But $f \notin RMS([0, 1], l_2(\mathbb{N}))$, since f is not bounded. \square

The example that follows shows a function of the unit square to $l_2(\mathbb{N} \times \mathbb{N})$ not satisfying the Fubini Theorem.

Example 3.5. Consider the function $f : [0, 1] \times [0, 1] \rightarrow l_2(\mathbb{N} \times \mathbb{N})$ given by $f(s, t) = 2^i g_i(t)$ on $2^{-i} \leq s < 2^{-i+1}$, $i = 1, 2, 3, \dots$, and $f(s, t) = 0$ where not otherwise defined, where $g_i(t) = e_{ij}$ whenever $\frac{j-1}{2^i} \leq t < \frac{j}{2^i}$, $j = 1, 2, \dots, 2^i$, and $g_i(t) = 0$ otherwise. Then, $f(s, t)$ is integrable over $[0, 1] \times [0, 1]$ with

$$\int \int_{[0,1] \times [0,1]} f(s, t) ds dt = \sum_{i=1}^{\infty} \sum_{j=1}^{2^i} \frac{1}{2^i} e_{ij}.$$

The integral with respect to s on a single line $t = \text{constant}$ exists, but the integral with respect to t on a single line $s = \text{constant}$ does not because

$$\int_0^1 f(s, t) dt = 2e_{1j_1} + 4e_{2j_2} + 8e_{3j_3} + \dots$$

for some j_1, j_2, j_3, \dots \square

The next example presents a function $f : [0, 1] \rightarrow l_2(\mathbb{N})$ such that $\|f(t)\|_2 = 1$ for every $t \in [0, 1]$, but $\|f\|_A < \varepsilon$ for a given $\varepsilon > 0$.

Example 3.6. Let $\varepsilon > 0$, $n \in \mathbb{N}$ and $f : [0, 1] \rightarrow l_2(\mathbb{N})$ be defined by $f(t) = e_n$, whenever $\frac{k-1}{n^2} \leq t < \frac{k}{n^2}$, $k = 1, 2, \dots, n^2$, and $f(t) = 0$ otherwise. Hence

$$\begin{aligned} \|f\|_A &= \left\| (K) \int_0^1 f(t) dt \right\|_2 = \left\| \sum_{k=1}^{n^2} \int_{\frac{k-1}{n^2}}^{\frac{k}{n^2}} e_n dt \right\|_2 = \left\| \sum_{k=1}^{n^2} \frac{1}{n^2} e_k \right\|_2 \\ &= \left(\frac{1}{n^4} \cdot n^2 \right)^{\frac{1}{2}} = \frac{1}{n}. \end{aligned}$$

Then taking $n > \frac{1}{\varepsilon}$, we have $\|f\|_A < \varepsilon$. \square

Example 3.7 in the sequel is a Birkhoff-type example due to Hönig. It gives a sequence of functions $f_n : [0, 1] \rightarrow l_2(\mathbb{N})$ such that $\sup_n \|f_n\|_A < \infty$ but $\|f_n(t)\|_2 \uparrow \infty$, for every $t \in [a, b]$.

Example 3.7. Let 1_D denote the characteristic function of a set $D \subset [0, 1]$. We define a sequence of functions $f_n : [0, 1] \rightarrow l_2(\mathbb{N})$, $n \in \mathbb{N}$, as follows: $f_n = \sum_{i=1}^n g_i$, where

$$g_i = \sum_{j=1}^{2^{i-1}} 1_{\left[\frac{j-1}{2^{i-1}}, \frac{j}{2^{i-1}}\right]} e_{2^{i-1}+j-1}, \quad i = 1, 2, \dots$$

Then $\sup_{n \rightarrow \infty} \|f_n\|_A < \infty$ and, for every $t \in [a, b]$ and every $n \in \mathbb{N}$, $\|f_n(t)\|_2 < \|f_{n+1}(t)\|_2$ and $\|f_n(t)\|_2 \rightarrow \infty$. \square

4 Appendix

The integrals introduced by J. Kurzweil ([19]) and independently by R. Henstock ([12]) in the late fifties give a Riemannian definition of the Denjoy-Perron integral which encompasses the Newton, Riemann and Lebesgue integrals. In 1969, McShane showed that a small change in this definition leads to the Lebesgue integral.

The Kurzweil and Henstock integrals can be immediately extended to Banach space-valued functions. The extension of the McShane integral made by Gordon, [10], gives a more general integral than that of Bochner-Lebesgue. But the variational Henstock-McShane definition for functions defined on a compact interval of the real line and taking values in a Banach space gives precisely the Bochner-Lebesgue integral. This fact was proved by Congxin and Xiabo ([4]) and independently by Hönig ([17]). Later, Di Piazza and Musal generalized this result ([5]).

Because reference [17] is unavailable to the majority of the mathematicians, we include its results in this Appendix. Unlike the proof of Congxin and Xiabo ([4]), which is based on the Frechet differentiability of the Bochner-Lebesgue integral, the idea of Hönig ([17]) to prove the equivalence of the Bochner-Lebesgue and the Henstock-McShane integrals uses the fact that the indefinite integral of Henstock-McShane and absolutely Henstock integrable functions are of bounded variation. In this manner, the proof in ([17]) seems to be more simple.

We say that a function $f : [a, b] \rightarrow X$ is *Bochner-Lebesgue integrable* (we write $f \in \mathcal{L}_1([a, b], X)$), if there exists a sequence $(f_n)_{n \in \mathbb{N}}$ of simple functions, $f_n : [a, b] \rightarrow X$, $n \in \mathbb{N}$, such that

- (i) $f_n \rightarrow f$ almost everywhere (i.e., $\lim_{n \rightarrow \infty} \|f_n(t) - f(t)\| = 0$ for almost every $t \in [a, b]$), and
- (ii) $\lim_{n, m \rightarrow \infty} (L) \int_a^b \|f_n(t) - f_m(t)\| dt = 0$.

We define $(L) \int_a^b f(t) dt = \lim_{n \rightarrow \infty} (L) \int_a^b f_n(t) dt$ and $\|f\|_1 = (L) \int_a^b \|f(t)\| dt$. The space of all equivalence classes of Bochner-Lebesgue integrable functions, equipped with the norm $\|f\|_1$, is complete.

We say that $f : [a, b] \rightarrow X$ is measurable, whenever there is a sequence of simple functions $f_n : [a, b] \rightarrow X$ such that $f_n \rightarrow f$ almost everywhere. When this is the case,

$$f \in \mathcal{L}_1([a, b], X) \quad \text{if and only if} \quad (L) \int_a^b \|f(t)\| dt < \infty \quad (3)$$

(see [29]).

Our next goal is to show that the integrals of Bochner-Lebesgue and Henstock-McShane coincide, that is, $\mathcal{L}_1([a, b], X) = HMS([a, b], X)$. In this manner, we will prove that the inclusions $\mathcal{L}_1([a, b], X) \subset HMS([a, b], X)$ and $HMS([a, b], X) \subset \mathcal{L}_1([a, b], X)$ hold and we will show that the integrals coincide when defined.

We let $(KMS) \int_a^b f$ denote the integral of a function $f \in KMS([a, b], X)$.

Lemma 4.1. *Given a sequence $(f_n)_{n \in \mathbb{N}}$ in $KMS([a, b], X)$ and a function $f : [a, b] \rightarrow X$, suppose there exists $\lim_{n \rightarrow \infty} (L) \int_a^b \|f_n(t) - f(t)\| dt = 0$. Then $f \in KMS([a, b], X)$ and*

$$\lim_{n \rightarrow \infty} (KMS) \int_a^b f_n(t) dt = (KMS) \int_a^b f(t) dt.$$

PROOF. Given $\varepsilon > 0$, take n_ε such that for $m, n \geq n_\varepsilon$,

$$(KMS) \int_a^b \|f_n(t) - f_m(t)\| dt < \varepsilon$$

and take a gauge δ of $[a, b]$ such that for every δ -fine $(\xi_i, t_i) \in STD_{[a, b]}$,

$$\sum_i \|f_{n_\varepsilon}(\xi_i) - f(\xi_i)\| (t_i - t_{i-1}) < \varepsilon. \quad (4)$$

The limit $I = \lim_{n \rightarrow \infty} (KMS) \int_a^b f_n(t) dt$ exists, since for $m, n \geq n_\varepsilon$,

$$\begin{aligned} & \left\| (KMS) \int_a^b f_n(t) dt - (KMS) \int_a^b f_m(t) dt \right\| \leq \\ & \leq (KMS) \int_a^b \|f_n(t) - f(t)\| dt + (KMS) \int_a^b \|f(t) - f_m(t)\| dt \leq 2\varepsilon. \end{aligned}$$

Hence, if $I_n = (KMS) \int_a^b f_n(t)dt$, then

$$\begin{aligned} & \left\| \sum_i f(\xi_i)(t_i - t_{i-1}) - I \right\| \leq \left\| \sum_i [f(\xi_i) - f_{n_\varepsilon}(\xi_i)](t_i - t_{i-1}) \right\| + \\ & \quad + \left\| \sum_i f_{n_\varepsilon}(\xi_i)(t_i - t_{i-1}) - I_{n_\varepsilon} \right\| + \|I_{n_\varepsilon} - I\| \leq \\ & \leq \sum_i \|f(\xi_i) - f_{n_\varepsilon}(\xi_i)\| (t_i - t_{i-1}) + \left\| \sum_i f_{n_\varepsilon}(\xi_i)(t_i - t_{i-1}) - I_{n_\varepsilon} \right\| + \|I_{n_\varepsilon} - I\|. \end{aligned} \quad (5)$$

Then the first summand in (5) is smaller than ε by (4), the third summand is smaller than ε by the definition of n_ε and, if we refine the gauge δ we may suppose, by the definition of I_{n_ε} , that the second summand is smaller than ε and the proof is complete. \square

We show next that Lemma 4.1 remains valid if we replace KMS by HMS.

Lemma 4.2. *Consider a sequence $(f_n)_{n \in \mathbb{N}}$ in $HMS([a, b], X)$ and let $f : [a, b] \rightarrow X$. If $\lim_n (L) \int_a^b \|f_n(t) - f(t)\| dt = 0$, then $f \in HMS([a, b], X)$ and*

$$\lim_n (KMS) \int_a^b f_n(t)dt = (KMS) \int_a^b f(t)dt.$$

PROOF. By Lemma 4.1, $f \in KMS([a, b], X)$ and we have the convergence of the integrals. It remains to prove that $f \in HMS([a, b], X)$, that is, for every $\varepsilon > 0$ there exists a gauge δ of $[a, b]$ such that for every δ -fine $(\xi_i, t_i) \in STD_{[a, b]}$,

$$\sum_i \left\| (KMS) \int_{t_{i-1}}^{t_i} f(t)dt - f(\xi_i)(t_i - t_{i-1}) \right\| \leq \varepsilon.$$

But,

$$\begin{aligned} & \sum_i \left\| (KMS) \int_{t_{i-1}}^{t_i} f(t)dt - f(\xi_i)(t_i - t_{i-1}) \right\| \leq \\ & \leq \sum_i \left\| (KMS) \int_{t_{i-1}}^{t_i} [f(t) - f_n(t)] dt \right\| + \\ & + \sum_i \left\| (KMS) \int_{t_{i-1}}^{t_i} f_n(t)dt - f_n(\xi_i)(t_i - t_{i-1}) \right\| + \end{aligned}$$

$$+ \sum_i \|f_n(\xi_i) - f(\xi_i)\| (t_i - t_{i-1}). \quad (6)$$

Because $\int_a^b \|f_n(t) - f(t)\| dt \rightarrow 0$, there exists $n_\varepsilon > 0$ such that the first summand in (6) is smaller than $\varepsilon/3$ for all $n \geq n_\varepsilon$. Choose an $n \geq n_\varepsilon$. Then we can take δ such that the third summand is smaller than $\varepsilon/3$, since it approaches $\int_a^b \|f_n(t) - f(t)\| dt$. Also, because $f_n \in HMS([a, b], X)$, we may refine δ so that the second summand becomes smaller than $\varepsilon/3$ and we finished the proof. \square

Lemma 4.3. $\mathcal{L}_1([a, b], X) \subset KMS([a, b], X)$.

For a proof of Lemma 4.3, see Theorem 16 in [10] for instance.

Now we are able to prove the inclusion

Theorem 4.1. $\mathcal{L}_1([a, b], X) \subset HMS([a, b], X)$.

PROOF. By Lemma 4.3, $\mathcal{L}_1([a, b], X) \subset KMS([a, b], X)$. Then, following the steps of the proof of Lemma 4.3 and using Lemma 4.2, we obtain the result. \square

Let $BV([a, b], X)$ denote the space of all functions $f : [a, b] \rightarrow X$ of bounded variation. We show next that the indefinite integral of any function of $HMS([a, b], X)$ belongs to $BV([a, b], X)$.

Lemma 4.4. If $f \in HMS([a, b], X)$, then $\tilde{f} \in BV([a, b], X)$.

PROOF. It is enough to show that every $\xi \in [a, b]$ has a neighborhood where \tilde{f} is of bounded variation. By hypothesis, given $\varepsilon > 0$, there exists a gauge δ of $[a, b]$ such that for every δ -fine semi-tagged division $d = (\xi_i, t_i)$ of $[a, b]$,

$$\sum_i \left\| \tilde{f}(t_i) - \tilde{f}(t_{i-1}) - f(\xi_i)(t_i - t_{i-1}) \right\| < \varepsilon. \quad (7)$$

Since $g = f$ almost everywhere implies $g \in HMS([a, b], X)$ and $\tilde{g} = \tilde{f}$ (this fact follows by straightforward adaptation of [11], Theorem 9.10 for Banach space-valued functions; see also [7]), we may change f on a set of measure zero and its indefinite integral does not change. We suppose, therefore, that $f(\xi) = 0$.

Let $s_0 < s_1 < \dots < s_m$ be any division of $[\xi - \delta(\xi), \xi + \delta(\xi)]$. If we take $\xi_j = \xi$ for $j = 1, 2, \dots, m$, then (ξ_j, s_j) is a δ -fine semi-tagged division of $[\xi - \delta(\xi), \xi + \delta(\xi)]$ and therefore from (7) and fact that $f(\xi_j) = f(\xi) = 0$ for all j , we have

$$\sum_{j=1}^m \left\| \tilde{f}(s_j) - \tilde{f}(s_{j-1}) \right\| \leq \varepsilon$$

and the proof is complete. \square

Lemma 4.5. *Suppose $f \in H([a, b], X)$. The following properties are equivalent:*

(i) *f is absolutely integrable;*

(ii) *$\tilde{f} \in BV([a, b], X)$.*

PROOF. (i) \Rightarrow (ii). Suppose f is absolutely integrable. Since the variation of \tilde{f} , $V(\tilde{f})$, is given by

$$V(\tilde{f}) = \sup \left\{ \sum_i \left\| \tilde{f}(t_i) - \tilde{f}(t_{i-1}) \right\| ; (t_i) \in D_{[a,b]} \right\}$$

we have

$$\begin{aligned} \sum_i \left\| \tilde{f}(t_i) - \tilde{f}(t_{i-1}) \right\| &= \sum_i \left\| (K) \int_{t_{i-1}}^{t_i} f(t) dt \right\| \leq \\ &\leq \sum_i (K) \int_{t_{i-1}}^{t_i} \|f(t)\| dt = (K) \int_a^b \|f(t)\| dt. \end{aligned}$$

(ii) \Rightarrow (i). Suppose $\tilde{f} \in BV([a, b], X)$. We will prove that the integral $(K) \int_a^b \|f(t)\| dt$ exists and $(K) \int_a^b \|f(t)\| dt = V(\tilde{f})$. Given $\varepsilon > 0$, we need to find a gauge δ of $[a, b]$ such that

$$\left| \sum_i \|f(\xi_i)\| (t_i - t_{i-1}) - V(\tilde{f}) \right| < \varepsilon,$$

whenever $(\xi_i, t_i) \in TD_{[a,b]}$ is δ -fine. But

$$\begin{aligned} &\left| \sum_i \|f(\xi_i)\| (t_i - t_{i-1}) - V(\tilde{f}) \right| \leq \\ &\leq \sum_i \left| \|f(\xi_i)\| (t_i - t_{i-1}) - \left\| (K) \int_{t_{i-1}}^{t_i} f(t) dt \right\| \right| + \\ &\quad + \left| \sum_i \left\| (K) \int_{t_{i-1}}^{t_i} f(t) dt \right\| - V(\tilde{f}) \right| \\ &\leq \sum_i \left| \|f(\xi_i)\| (t_i - t_{i-1}) - \left\| (K) \int_{t_{i-1}}^{t_i} f(t) dt \right\| \right| + \left| \sum_i \left\| \tilde{f}(t_i) - \tilde{f}(t_{i-1}) \right\| - V(\tilde{f}) \right|. \end{aligned} \tag{8}$$

By the definition of $V(\tilde{f})$, we may take $(t_i) \in D_{[a,b]}$ such that the last summand in (8) is smaller than $\varepsilon/2$. Because $f \in H([a, b], X)$, we may take a gauge δ such that for every δ -fine $(\xi_i, t_i) \in TD_{[a,b]}$, the first summand in (8) is also smaller than $\varepsilon/2$ (and we may suppose that the points chosen for the second summand are the points of the δ -fine tagged division (ξ_i, t_i)). \square

The next result follows from the fact that $HMS([a, b], X) \subset H([a, b], X)$ and Lemmas 4.4 and 4.5.

Corollary 4.1. *All functions of $HMS([a, b], X)$ are absolutely integrable.*

Lemma 4.6. *All functions of $H([a, b], X)$ are measurable.*

For a proof of Lemma 4.6, see Theorem 9 in [3] for instance.

Finally, we can prove the inclusion

Theorem 4.2. $HMS([a, b], X) \subset \mathcal{L}_1([a, b], X)$.

PROOF. The result follows from the facts that all functions of $H([a, b], X)$ and hence of $HMS([a, b], X)$ are measurable (Lemma 4.6) and all functions of $HMS([a, b], X)$ are absolutely integrable (Corollary 4.1) (see [29]). \square

Acknowledgement. The author would like to thank S. Schwabik, P. Muldowney and specially the referee for their helpful suggestions and information.

References

- [1] G. Birkhoff, *Integration of functions on Banach spaces*, Trans. Amer. Math. Soc., **38** (1935), 357–378.
- [2] B. Bongiorno, *Relatively weakly compact sets in the Denjoy Space*, J. Math. Study, **27** (1994), 37–43.
- [3] S. Cao, *The Henstock integral for Banach-valued functions*, SEA Bull. Math., **16**(1) (1992), 35–40.
- [4] W. Congxin and Y. Xiabo, *A Riemann-type definition of the Bochner integral*, J. Math. Studies, **27** (1994), 32–36.
- [5] L. Di Piazza and K. Musial, *A characterization of variationally McShane integrable Banach-space valued functions*, Illinois J. Math., **45**(1) (2001), 279–289.
- [6] M. Federson and R. Bianconi, *Linear integral equations of Volterra concerning the integral of Henstock*, Real Anal. Exchange, **25**(1) (2000), 389–417.

- [7] M. Federson, *The Fundamental Theorem of Calculus for multidimensional Banach space-valued Henstock vector integrals*, Real Analysis Exchange, **25**(1) (1999–2000), 469–480.
- [8] M. Federson, *The Monotone Convergence Theorem for multidimensional abstract Kurzweil vector integrals*, Czech. Math. J., **52**(127) 2 (2002), 429–439.
- [9] A. Gilioli, *Natural ultrabornological non-complete, normed function spaces*, Arch. Math., **61** (1993), 465–477.
- [10] R. A. Gordon, *The McShane integral of Banach-valued functions*, Illinois J. Math., **34** (1990), 557–567.
- [11] R. A. Gordon, *The Integrals of Lebesgue, Denjoy, Perron, and Henstock*, Graduate Studies in Math., AMS, Providence, Rhode Island, 1994.
- [12] R. Henstock, *Definitions of Riemann-type of the variational integrals*, Proc. London Math. Soc., **11**(3) (1961), 402–418.
- [13] R. Henstock, *A Riemann-type integral of Lebesgue power*, Canad. J. Math., **20** (1968), 79–87.
- [14] R. Henstock, *The General Theory of Integration*, Oxford Math. Monog., Clarendon Press, Oxford, 1991.
- [15] C. S. Hönig, *The Abstract Riemann-Stieltjes Integral and its Applications to Linear Differential Equations with generalized boundary conditions*, Notas do Instituto de Matemática e Estatística da Universidade de São Paulo, Série Matemática, 1, 1973.
- [16] C. S. Hönig, *Volterra-Stieltjes integral equations*, Math. Studies, 16, North-Holland Publ. Comp., Amsterdam, 1975.
- [17] C. S. Hönig, *A Riemannian characterization of the Bochner-Lebesgue integral*, Seminário Brasileiro de Análise, **35** (1992), 351–358.
- [18] G. Kothe, *Topological Vector Spaces*, vol. I and II, Springer, New York, 1969, 1979.
- [19] J. Kurzweil, *Generalized ordinary differential equations and continuous dependence on a parameter*, Czech. Math. J., **82**(7) (1957), 418–448.
- [20] J. Kurzweil, *Nichtabsolut Konvergente Integrale*, Leipzig, 1980.

- [21] P. Y. Lee, *Lanzhou Lectures on Henstock Integration*, World Sci., Singapore, 1989.
- [22] J. Mawhin, *Introduction à l'Analyse*, Cabay, Louvain-la-Neuve, 1983.
- [23] R. M. McLeod, *The Generalized Riemann Integral*, Carus Math. Monog. 20, The Math. Assoc. of America, 1980.
- [24] E. J. McShane, *A Riemann-type integral that includes Lebesgue-Stieltjes, Bochner and stochastic integrals*, AMS Memoirs, **88**, (1969).
- [25] E. J. McShane, *A unified theory of integration*, Am. Math. Monthly, **80** (1973), 349–359.
- [26] W. F. Pfeffer, *The Riemann Approach to Integration*, Cambridge, 1993.
- [27] H. L. Royden, *Real Analysis*, Second Edition, Macmillan Comp., 1968.
- [28] C. Swartz and J. DePree, *Introduction to Real Analysis*, John Wiley and Sons, Inc., New York, 1988.
- [29] K. Yosida, *Functional Analysis*, Springer, 1966.