

Krzysztof Chris Ciesielski, Department of Mathematics, West Virginia University, Morgantown, WV 26506-6310 and Department of Radiology, MIPG, University of Pennsylvania, Blockley Hall – 4th Floor, 423 Guardian Drive, Philadelphia, PA 19104-6021. email: KCies@math.wvu.edu

Jakub Jasinski, Department of Mathematics, University of Scranton, Scranton, PA 18510-4666. email: jakub.jasinski@scranton.edu

SMOOTH PEANO FUNCTIONS FOR PERFECT SUBSETS OF THE REAL LINE

Abstract

In this paper we investigate for which closed subsets P of the real line \mathbb{R} there exists a continuous map from P onto P^2 and, if such a function exists, how smooth can it be. We show that there exists an infinitely many times differentiable function $f: \mathbb{R} \rightarrow \mathbb{R}^2$ which maps an unbounded perfect set P onto P^2 . At the same time, no continuously differentiable function $f: \mathbb{R} \rightarrow \mathbb{R}^2$ can map a compact perfect set onto its square. Finally, we show that a disconnected compact perfect set P admits a continuous function from P onto P^2 if, and only if, P has uncountably many connected components.

1 Introduction and overview

Let P be a nonempty subset of the set \mathbb{R} of real numbers. If P has no isolated points and $n, m \in \{1, 2, 3, \dots\}$, then we consider the following classes of smooth functions from P to \mathbb{R}^m : \mathcal{D}^n of n -times differentiable functions and \mathcal{C}^n of continuously n -times differentiable functions. In addition, \mathcal{C}^0 will stand for the class of all continuous functions and \mathcal{C}^∞ for the class of functions differentiable infinitely many times. For every $n < \omega$ we have $\mathcal{C}^\infty \subset \mathcal{C}^{n+1} \subset \mathcal{D}^{n+1} \subset \mathcal{C}^n$.

A nonempty set $P \subseteq \mathbb{R}$ is called *perfect* if it is closed and has no isolated points. We say that a function $f: P \rightarrow \mathbb{R}^2$ is *Peano* if it is onto P^2 , that is,

Mathematical Reviews subject classification: Primary: 26A30; Secondary: 26B05
Key words: Peano curve, space filling curve, differentiability
Received by the editors March 4, 2013
Communicated by: Paul D. Humke

when $f[P] = P^2$. For example, the classic result of Peano [7] states that there exists a Peano function $f: [0, 1] \rightarrow [0, 1]^2$ of class \mathcal{C}^0 . More on this topic can be found in Sagan [9].

It is worth noting that some Peano functions $f: P \rightarrow \mathbb{R}^2$ of a given smoothness class can be extended to the entire functions $\widehat{f}: \mathbb{R} \rightarrow \mathbb{R}^2$ of the same class.

Proposition 1.1. *Let $P \subset \mathbb{R}$ be a perfect set.*

- (a) *Any \mathcal{C}^0 Peano function $f: P \rightarrow P^2$ may be extended to a \mathcal{C}^0 function $\widehat{f}: \mathbb{R} \rightarrow \mathbb{R}^2$.*
- (b) *Any \mathcal{D}^1 Peano function $f: P \rightarrow P^2$ may be extended to a \mathcal{D}^1 function $\widehat{f}: \mathbb{R} \rightarrow \mathbb{R}^2$.*

PROOF. (a) follows from the Generalized Tietze extension theorem, see e.g. [5, p. 151]. Part (b) follows from the following extension theorem due to V. Jarník [2]: “Every differentiable function f from a perfect set $P \subset \mathbb{R}$ into \mathbb{R} can be extended to a differentiable function $\widehat{f}: \mathbb{R} \rightarrow \mathbb{R}$.” More on Jarník’s theorem can be found in [4]. The theorem has also been independently proved in [8, theorem 4.5]. \square

Proposition 1.1 shows that for the functions from classes \mathcal{C}^0 and \mathcal{D}^1 , the existence of a Peano function for a perfect set $P \subset \mathbb{R}$ is equivalent to the existence of a function $f: \mathbb{R} \rightarrow \mathbb{R}^2$ of the same class with $f \upharpoonright P$ being Peano.




Figure 1: $f(0) = 0$ and $f(x) = (a_n)^2$ for $x \in [a_n, b_n]$.

Remark 1.2. For the functions of the higher classes of smoothness such simple equivalence is not achievable. Indeed, in general, a \mathcal{C}^1 function f from a perfect set $P \subset [0, 1]$ into \mathbb{R} need not be extendable to an entire \mathcal{C}^1 function $\widehat{f}: [0, 1] \rightarrow \mathbb{R}$, even if f is of the \mathcal{C}^∞ class.

Perhaps the simplest example supporting our Remark 1.2 is the function f defined on the set $P = \{0\} \cup \bigcup_{n=1}^{\infty} [a_n, b_n]$, where $a_n = 2^{-n}$ and $b_n \in (a_n, a_{n-1})$, as $f(0) = 0$ and $f(x) = (a_n)^2$ for $x \in [a_n, b_n]$. See Figure 1. Then, $f'(x) = 0$ for every $x \in P$, so f is \mathcal{C}^∞ . However, if we choose b_{n+1} 's such that the quotient $\frac{f(a_n) - f(b_{n+1})}{a_n - b_{n+1}} = \frac{(2^{-n})^2 - (2^{-n-1})^2}{2^{-n} - b_{n+1}}$ equals 1, $b_{n+1} = \frac{2^{n+2} - 3}{2^{2n+2}}$ works, then by the mean value theorem any differentiable extension $\widehat{f}: [0, 1] \rightarrow \mathbb{R}$ of f will have discontinuous derivative at 0.

Remark 1.2 shows that for the functions of at least \mathcal{C}^1 smoothness, it makes a difference, if we construct the Peano functions as the restrictions of the entire smooth functions or just on the set P . We pay attention to these details in what follows.

The following theorem summarizes all the results on the Peano functions for the subsets of \mathbb{R} presently known to us.

Theorem 1.3. *Let P be a closed subset of \mathbb{R} .*

- (a) *There exists a \mathcal{C}^0 Peano function f from P onto P^2 if, and only if, P is either connected or it has uncountably many components.*
- (b) *If P is perfect and has positive Lebesgue measure, then there is no \mathcal{D}^1 Peano function f from P onto P^2 .*
- (c) *If $f: \mathbb{R} \rightarrow \mathbb{R}^2$ is a \mathcal{C}^1 function and $P \subseteq \mathbb{R}$ is a compact perfect set, then $P^2 \not\subset f[P]$. Hence, $f \upharpoonright P$ is not Peano.*
- (d) *There exists a \mathcal{C}^∞ function $f: \mathbb{R} \rightarrow \mathbb{R}^2$ and a perfect unbounded subset P of \mathbb{R} such that $f[P] = P^2$, that is, $f \upharpoonright P$ is Peano.*

PROOF. (a) is proved in Theorem 4.1.

(b) Let $f = \langle f_1, f_2 \rangle: P \rightarrow P^2$ be differentiable. Morayne [6, theorem 3] showed (using the fact that \mathcal{D}^1 functions satisfy the Banach condition (T_2)) that $f[P]$ must have the planar Lebesgue measure zero. In particular, if P has positive measure, then $P^2 \not\subset f[P]$.

(c) is proved in Theorem 3.1.

(d) is proved in Theorem 2.2. □

2 A C^∞ function $f: \mathbb{R} \rightarrow \mathbb{R}^2$ with a Peano restriction $f \upharpoonright P$ for some perfect set $P \subset \mathbb{R}$

The idea is to construct a sequence $\langle P_k \subseteq [3k, 3k+2]: k < \omega \rangle$ of perfect sets such that for every $\ell, \ell' < k$ there exists a C^∞ function $f_{\ell, \ell'}^k$ from $[3k, 3k+2]$ into \mathbb{R}^2 which maps P_k onto $P_\ell \times P_{\ell'}$, see Figures 2 and 4. Then, the set $P = \bigcup_{k < \omega} P_k$ will be as required, since for any given sequence $\langle \langle \ell_k, \ell'_k \rangle: 0 < k < \omega \rangle$ of all pairs of natural numbers with $\ell_k, \ell'_k < k$, the function $\hat{f} = \bigcup_{0 < k < \omega} f_{\ell_k, \ell'_k}^k$ is C^∞ and it maps $\bigcup_{0 < k < \omega} P_k$ onto P^2 . Such an \hat{f} can easily be extended to the desired C^∞ function $f: \mathbb{R} \rightarrow \mathbb{R}^2$.




Figure 2: An f_{ℓ_k, ℓ'_k}^k fragment of the function f .

The construction of the sets P_k will naturally provide continuous mappings $\bar{f}_{\ell, \ell'}^k$ from P_k onto $P_\ell \times P_{\ell'}$. The difficulty will be to ensure that these functions are not only C^∞ , but that they can be also extended to the C^∞ functions $f_{\ell, \ell'}^k: [3k, 3k+2] \rightarrow \mathbb{R}^2$. The tool to insure the extendability is provided by the following Lemma 2.1. Notice, that the lemma can be considered as a version of Whitney extension theorem [10].¹

Note also, that no analytic function $f: \mathbb{R} \rightarrow \mathbb{R}^2$ can have a Peano restriction to any perfect set (since the coordinates, $f_1, f_2: \mathbb{R} \rightarrow \mathbb{R}$, of a Peano function need to be constant on some perfect subsets).

Lemma 2.1. *Every real-valued function g_0 from a compact nowhere dense set $K \subset \mathbb{R}$ having the property that for every $k < \omega$ there exists a $\delta_k \in (0, 1)$ such that*

¹*Added in proof.* Actually, Lemma 2.1 follows from [10, thm. 1 p. 65], since “ g_0 is of class C^∞ in K in terms of the functions $f_k \equiv 0$.” The authors like to thank Prof. Jan Kolar for pointing this out.

(P_k) $|g_0(x) - g_0(y)| < |x - y|^{k+1}$ for all $x, y \in K$ with $0 < |x - y| < \delta_k$

can be extended to a C^∞ function $g: \mathbb{R} \rightarrow \mathbb{R}$. Moreover, $g'(x) = 0$ for all $x \in K$.

PROOF. Let $\psi: \mathbb{R} \rightarrow \mathbb{R}$ be a monotone C^∞ map such that $\psi[(-\infty, 0)] = \{0\}$ and $\psi[(1, \infty)] = \{1\}$. For $k < \omega$ let

$$M_k = \sup \{ |\psi^{(i)}(x)| : x \in [0, 1] \ \& \ i \leq k \} \in [1, \infty).$$

Let \mathcal{K} be a family of all connected bounded components (a, b) of $\mathbb{R} \setminus K$. Let $g: \mathbb{R} \rightarrow \mathbb{R}$ be an extension of g_0 such that g is constant on the closure of each unbounded component of $\mathbb{R} \setminus K$ and on each $(a, b) \in \mathcal{K}$ function g is defined by a formula

$$g(x) = (g_0(b) - g_0(a))\psi\left(\frac{x-a}{b-a}\right) + g_0(a).$$

In other words, g on (a, b) is a function $\psi \upharpoonright (0, 1)$ shifted and linearly rescaled in such a way that $g \upharpoonright [a, b]$ is continuous. We will show that such defined g is our desired C^∞ function.

Clearly, the restriction $g|_{\mathbb{R} \setminus K}$ of g is infinitely many times differentiable at any $x \in \mathbb{R} \setminus K$. We need to show that the same is true for any $x \in K$. For this, we will show, by induction on $k \geq 1$, that

(I_k) for every $x \in K$, the k -th derivative $g^{(k)}(x)$ exists and is equal 0.

The inductive argument is based on the following estimate, where $k \geq 1$:

(S_k) $\left| \frac{g^{(k-1)}(y) - g^{(k-1)}(z)}{y-z} \right| < M_k(b-a)$ provided $(a, b) \in \mathcal{K}$, $b-a < \delta_k$, and $y, z \in [a, b]$ are distinct.

Let $k \geq 1$. To see (S_k), take y and z as in its assumption. Then,


$$\left| \frac{g^{(k-1)}(y) - g^{(k-1)}(z)}{y-z} \right| \leq \sup_{x \in (a,b)} |g^{(k)}(x)| \quad (1)$$

$$= \sup_{x \in (a,b)} \frac{|g(b) - g(a)|}{|b-a|^k} \left| \psi^{(k)}\left(\frac{x-a}{b-a}\right) \right| \quad (2)$$

$$\leq \frac{|g(b) - g(a)|}{|b-a|^k} M_k \quad (3)$$

$$< \frac{|b-a|^{k+1}}{|b-a|^k} M_k = M_k(b-a), \quad (4)$$

where (1) follows from the Mean Value Theorem, (2) from the fact that $g^{(k)}(x) = \frac{d^k}{dx^k} [(g(b) - g(a))\psi\left(\frac{x-a}{b-a}\right) + g(a)] = \frac{g(b)-g(a)}{(b-a)^k} \psi^{(k)}\left(\frac{x-a}{b-a}\right)$ for every $x \in$

Figure 3: $b - a < \varepsilon/M_1 < b_1 - a_1$.

(a, b) , (3) from the definition of M_k , while (4) is concluded from (P_k) used with $x = b$ and $y = a$.

To show (I_1) , fix an $x_0 \in K$ and an $\varepsilon > 0$. We will find a $\delta > 0$ for which

$$\left| \frac{g(y) - g(x_0)}{y - x_0} \right| < \varepsilon \text{ provided } x_0 < y < x_0 + \delta. \quad (5)$$

If x_0 is equal to the left endpoint of some component interval of $\mathbb{R} \setminus K$, then the existence δ follows from our definition of the function g on such intervals, specifically because $\psi'(0) = 0$. So, assume that this is not the case, that is, that $(x_0, x_0 + \eta) \cap K \neq \emptyset$ for every $\eta > 0$. Let $\delta \in (0, \min\{\varepsilon, \delta_1\})$ be such that $(x_0, x_0 + \delta)$ is disjoint with every $(a_1, b_1) \in \mathcal{K}$ for which $b_1 - a_1 \geq \varepsilon/M_1$. See Figure 3. We will show that such δ works.

So, fix a $y \in (x_0, x_0 + \delta)$ and let $z = \sup K \cap [x_0, y]$. Since $|z - x_0| < \delta < \delta_1$, by (P_1) we have $\left| \frac{g(z) - g(x_0)}{z - x_0} \right| < \frac{|z - x_0|^{1+1}}{|z - x_0|} = |z - x_0| < \delta < \varepsilon$. If $z = y$, this completes the proof of (5). So, assume that $z < y$. Then, there exists an $(a, b) \in \mathcal{K}$ for which $z = a$ and $y \in (a, b)$. Notice that, by the choice of δ , we have $b - a < \varepsilon/M_1$, see Figure 3. Hence, by (S_1) , we have $\left| \frac{g(y) - g(z)}{y - z} \right| < M_1(b - a) < \varepsilon$. Combining this with $\left| \frac{g(z) - g(x_0)}{z - x_0} \right| < \varepsilon$, we obtain $\left| \frac{g(y) - g(x_0)}{y - x_0} \right| \leq \max \left\{ \left| \frac{g(y) - g(z)}{y - z} \right|, \left| \frac{g(z) - g(x_0)}{z - x_0} \right| \right\} < \varepsilon$, finishing the proof of the property (5).

Similarly, we prove that there exists a $\delta > 0$ for which $\left| \frac{g(y) - g(x_0)}{y - x_0} \right| < \varepsilon$ provided $x_0 - \delta < y < x_0$. This completes the argument for (I_1) .

Next, assume that for some $k \geq 2$ the property (I_{k-1}) holds. We need to show (I_k) . So, fix an $x_0 \in K$ and an $\varepsilon > 0$. We will find a $\delta > 0$ for which

$$\left| \frac{g^{(k-1)}(y) - g^{(k-1)}(x_0)}{y - x_0} \right| < \varepsilon \text{ provided } x_0 < y < x_0 + \delta. \quad (6)$$

If x_0 is equal to the left endpoint of some component interval of $\mathbb{R} \setminus K$, then the existence of δ follow from our definition of function g on such intervals.

So, assume that this is not the case, that is, that $(x_0, x_0 + \eta) \cap K \neq \emptyset$ for every $\eta > 0$. Let $\delta \in (0, \min\{\varepsilon, \delta_k\})$ be such that $(x_0, x_0 + \delta)$ is disjoint with every $(a_1, b_1) \in \mathcal{K}$ for which $b_1 - a_1 \geq \varepsilon/M_k$. We will show that such δ works.

Fix a $y \in (x_0, x_0 + \delta)$. If $y \in K$, then $\left| \frac{g^{(k-1)}(y) - g^{(k-1)}(x_0)}{y - x_0} \right| = 0 < \varepsilon$ follows from (I_{k-1}) . So, we assume that $y \in (a, b)$ for some $(a, b) \in \mathcal{K}$. Then,

$$\left| \frac{g^{(k-1)}(y) - g^{(k-1)}(x_0)}{y - x_0} \right| = \left| \frac{g^{(k-1)}(y) - g^{(k-1)}(a)}{y - x_0} \right| \quad (7)$$

$$\leq \left| \frac{g^{(k-1)}(y) - g^{(k-1)}(a)}{y - a} \right| < M_k(b - a) < \varepsilon, \quad (8)$$

where (7) follows from $g^{(k-1)}(x_0) = 0 = g^{(k-1)}(a)$, which is implied by (I_{k-1}) , while (8) follows from (S_k) , since the choice of $\delta < \delta_k$ implies $b - a < \varepsilon/M_k$. This completes the proof of (6).

Similarly, we prove that there is a $\delta > 0$ for which $\left| \frac{g^{(k-1)}(y) - g^{(k-1)}(x_0)}{y - x_0} \right| < \varepsilon$ provided $x_0 - \delta < y < x_0$. This completes the argument for (I_k) and concludes the proof of the lemma. \square

Theorem 2.2. *There exist C^∞ functions $f_1, f_2: \mathbb{R} \rightarrow \mathbb{R}$ and a perfect set $P \subset \mathbb{R}$ such that $f = \langle f_1, f_2 \rangle$ maps P onto P^2 , that is, $f \upharpoonright P$ is a Peano function.*

PROOF. The construction will follow the outline indicated at the beginning of the section.

Perhaps the simplest continuous Peano-like function is the following map $h = \langle h^{\text{odd}}, h^{\text{even}} \rangle: 2^\omega \rightarrow (2^\omega)^2$, whose coordinate functions are the projections defined as $h^{\text{odd}}(s)(i) = s(2i + 1)$ and $h^{\text{even}}(s)(i) = s(2i)$. If we identify 2^ω with the Cantor ternary set $C = \left\{ \sum_{i < \omega} \frac{2s(i)}{3^{i+1}} : s \in 2^\omega \right\}$, then h becomes a continuous Peano function, from C onto C^2 . However, the compression of terms performed by h^{odd} and h^{even} gives us

$$\limsup_{s \rightarrow t} \left| \frac{h^{\text{odd}}(s) - h^{\text{odd}}(t)}{s - t} \right| = \infty.$$

Hence, h is not differentiable. In Section 3 we observe that this is a common problem for all compact sets.

To compensate for this compression, we define the sets P_k inductively, creating each P_k by “thickening” P_{k-1} in such a way, that the “condensed” coordinate projections of P_k , via analogs of the maps h^{odd} and h^{even} , may still be

mapped onto P_l in a differentiable way as long as $l < k$. Notice that while the “thickening” must be essential enough to obtain the above-mentioned requirement, it cannot be too radical, since the produced sets P_k must be of measure zero. This balancing act will be facilitated by the following functions p_k .

For every $k < \omega$ choose an increasing function $p_k: \omega \rightarrow [1, \infty)$ such that

$$\lim_{i \rightarrow \infty} \frac{p_\ell(i)}{p_k(2i)} = \lim_{i \rightarrow \infty} \frac{p_\ell(i)}{p_k(2i+1)} = \infty \text{ for every } \ell < k < \omega. \quad (9)$$

For example, the formula $p_k(i) = (i+1)^{2^{-k}}$ insures (9), as for every $i > 0$ we have

$$\frac{p_\ell(i)}{p_k(2i)} \geq \frac{p_\ell(i)}{p_k(2i+1)} = \frac{(i+1)^{2^{-\ell}}}{(2i+1)^{2^{-k}}} \geq \frac{i^{2^{-\ell}}}{(3i)^{2^{-k}}} = \frac{1}{3^{2^{-k}}} \frac{i^{2^{-\ell}}}{i^{2^{-k}}} = \frac{1}{3^{2^{-k}}} i^{2^{-\ell} - 2^{-k}},$$

and $\lim_{i \rightarrow \infty} \frac{1}{3^{2^{-k}}} i^{2^{-\ell} - 2^{-k}} = \infty$ since $2^{-\ell} - 2^{-k} > 0$.

For $k < \omega$ define $h_k: 2^\omega \rightarrow [3k, 3k+2]$ as $h_k(s) = 3k + \sum_{n=0}^{\infty} s(n)3^{-np_k(n)}$. Notice, that h_k is a continuous embedding. Moreover, for every $i < \omega$ we have $\sum_{n=i}^{\infty} 3^{-np_k(n)} \leq \sum_{n=i}^{\infty} 3^{-np_k(i)} \leq 3^{-ip_k(i)} \sum_{n=0}^{\infty} 3^{-n} = \frac{3}{2} 3^{-ip_k(i)}$. In particular, for every distinct $s, t \in 2^\omega$, if $i = \min\{n < \omega: s(n) \neq t(n)\}$, then

$$\frac{1}{2} 3^{-ip_k(i)} \leq |h_k(s) - h_k(t)| \leq \sum_{n=i}^{\infty} 3^{-np_k(n)} \leq \frac{3}{2} 3^{-ip_k(i)}, \quad (10)$$

where the first of the inequalities is justified by the following estimation,

$$\begin{aligned} |h_k(s) - h_k(t)| &= \left| \sum_{n=i}^{\infty} (s(n) - t(n))3^{-np_k(n)} \right| \geq 3^{-ip_k(i)} - \sum_{n=i+1}^{\infty} 3^{-np_k(n)} \\ &\geq 3^{-ip_k(i)} - \frac{3}{2} 3^{-(i+1)p_k(i+1)} \geq 3^{-ip_k(i)} - \frac{3}{2} 3^{-(i+1)p_k(i)} \\ &\geq 3^{-ip_k(i)} - \frac{1}{2} 3^{-ip_k(i)}. \end{aligned}$$

Let $P_k = h_k[2^\omega]$ and put $P = \bigcup_{k < \omega} P_k$. Clearly P is a perfect subset of \mathbb{R} . We will show that it satisfies the theorem.

For every $\ell < k < \omega$ let $h_{k,\ell}^{\text{odd}} = h_\ell \circ h^{\text{odd}} \circ h_k^{-1}$. It is easy to see that $h_{k,\ell}^{\text{odd}}$ is a continuous function from P_k onto P_ℓ . The key fact is that $h_{k,\ell}^{\text{odd}}$ satisfies the assumptions of Lemma 2.1, that is, for every $m < \omega$ there exists a $\delta_m \in (0, 1)$ such that

$$|h_{k,\ell}^{\text{odd}}(x) - h_{k,\ell}^{\text{odd}}(y)| < |x - y|^{m+1} \text{ for all } x, y \in P_k \text{ with } 0 < |x - y| < \delta_m. \quad (11)$$

Clearly, for any $\delta_m \in (0, 1)$, the condition (11) holds for any distinct $x, y \in P_k$ with $h_{k,\ell}^{\text{odd}}(x) = h_{k,\ell}^{\text{odd}}(y)$. Therefore, we are interested only in the case when $h_{k,\ell}^{\text{odd}}(x) \neq h_{k,\ell}^{\text{odd}}(y)$. Now, since $P_k = h_k[2^\omega]$, there exist $s, t \in 2^\omega$ with $x = h_k(s)$ and $y = h_k(t)$ and then $h_\ell(h^{\text{odd}}(s)) = h_{k,\ell}^{\text{odd}}(x) \neq h_{k,\ell}^{\text{odd}}(y) = h_\ell(h^{\text{odd}}(t))$. Since h_ℓ is injective, this implies that $h^{\text{odd}}(s) \neq h^{\text{odd}}(t)$. In short, we need to study $s, t \in 2^\omega$ for which $h^{\text{odd}}(s) \neq h^{\text{odd}}(t)$.

So, fix $s, t \in 2^\omega$ for which $h^{\text{odd}}(s) \neq h^{\text{odd}}(t)$ and define

$$x = h_k(s) \text{ and } y = h_k(t). \quad (12)$$

Let $i = \min\{n < \omega : h^{\text{odd}}(s)(n) \neq h^{\text{odd}}(t)(n)\}$. By the formula (10) we have the inequality $|h_\ell(h^{\text{odd}}(s)) - h_\ell(h^{\text{odd}}(t))| \leq \frac{3}{2} 3^{-ip_\ell(i)}$. Moreover, we have $s(2i+1) = h^{\text{odd}}(s)(i) \neq h^{\text{odd}}(t)(i) = t(2i+1)$. It follows that the number $i_1 = \min\{n < \omega : s(n) \neq t(n)\}$ is $\leq 2i+1$ and, again by the formula (10), we have $|x - y| = |h_k(s) - h_k(t)| \geq \frac{1}{2} 3^{-i_1 p_k(i_1)} \geq 3^{-(2i+1)p_k(2i+1)-1}$. In particular

$$\begin{aligned} |h_{k,\ell}^{\text{odd}}(x) - h_{k,\ell}^{\text{odd}}(y)| &\leq \frac{3}{2} 3^{-ip_\ell(i)} \\ &= \frac{3}{2} \left(3^{-(2i+1)p_k(2i+1)-1} \right)^{\frac{ip_\ell(i)}{(2i+1)p_k(2i+1)+1}} \\ &\leq \frac{3}{2} |x - y|^{\frac{ip_\ell(i)}{(2i+1)p_k(2i+1)+1}}. \end{aligned}$$

But, by (9), for every $m < \omega$ there is an $i_m < \omega$ with $\frac{ip_\ell(i)}{(2i+1)p_k(2i+1)+1} \geq m+2$ for all $i \geq i_m$. Moreover, since function h_k^{-1} is uniformly continuous, there is a $\delta_m \in (0, 1/2)$ such that $|h_k(s) - h_k(t)| < \delta_m$ implies that $s(j) = t(j)$ for all $j \leq 2i_m + 1$. Notice that this δ_m insures (11).

Indeed, if $|h_{k,\ell}^{\text{odd}}(x) - h_{k,\ell}^{\text{odd}}(y)| = 0$, then the condition certainly holds. Otherwise, with $s = h_k^{-1}(x)$ and $t = h_k^{-1}(y)$, we have $h^{\text{odd}}(s) \neq h^{\text{odd}}(t)$ and the choice of δ_m insures that $i = \min\{n < \omega : h^{\text{odd}}(s)(n) \neq h^{\text{odd}}(t)(n)\}$ is greater than i_m . So,

$$|h_{k,\ell}^{\text{odd}}(x) - h_{k,\ell}^{\text{odd}}(y)| \leq \frac{3}{2} |x - y|^{\frac{ip_\ell(i)}{(2i+1)p_k(2i+1)+1}} \leq \frac{3}{2} |x - y|^{m+2} < |x - y|^{m+1}$$

completing the proof of (11). In a similar manner, whenever $l < k < \omega$ we define $h_{k,\ell}^{\text{even}} = h_\ell \circ h^{\text{even}} \circ h_k^{-1}$, and obtain that

$$h_{k,\ell}^{\text{even}} \text{ satisfies the assumptions of Lemma 2.1.} \quad (13)$$

Let $\langle \langle \ell_k, \ell'_k \rangle : k = 1, 2, 3, \dots \rangle$ be a list of pairs from $\omega \times \omega$ such that for all $k \geq 1$, $\ell_k < k$ and $\ell'_k < k$. For each $k \geq 1$ define \bar{f}_1 on P_k as $h_{k,\ell_k}^{\text{odd}}$ and \bar{f}_2




Figure 4: We will define f so that $f \upharpoonright P_k = \langle h_{l_k}, h_{l'_k} \rangle \circ \langle h^{\text{odd}}, h^{\text{even}} \rangle \circ h_k^{-1}$.

on P_k as $h_{k, \ell'_k}^{\text{even}}$. In addition, we define \bar{f}_1 and \bar{f}_2 on P_0 as constant equal 0. Since sets P_k are separated, (11) and (13) ensure that \bar{f}_1 and \bar{f}_2 satisfy the assumptions of Lemma 2.1. Let $f_1: \mathbb{R} \rightarrow \mathbb{R}$ and $f_2: \mathbb{R} \rightarrow \mathbb{R}$ be C^∞ extensions of \bar{f}_1 and \bar{f}_2 , respectively. The proof will be complete as soon as we show that $f = \langle f_1, f_2 \rangle$ maps P onto P^2 . We have $f \upharpoonright P_k = \langle h_{l_k}, h_{l'_k} \rangle \circ \langle h^{\text{odd}}, h^{\text{even}} \rangle \circ h_k^{-1}$, see Figure 4. Since h_k^{-1} maps P_k onto 2^ω , $\langle h^{\text{odd}}, h^{\text{even}} \rangle$ maps 2^ω onto $2^\omega \times 2^\omega$, $h_{l_k}[2^\omega] = P_{l_k}$, and $h_{l'_k}[2^\omega] = P_{l'_k}$, we have $f[P_k] = P_{l_k} \times P_{l'_k}$. Therefore, $f[P] = \bigcup_{k < \omega} f[P_k] = \{0\} \cup \bigcup_{k=1}^\infty P_{l_k} \times P_{l'_k} = P^2$, completing the proof. \square

3 There is no C^1 function $f: \mathbb{R} \rightarrow \mathbb{R}^2$ with Peano restriction to a compact perfect set

Theorem 3.1. *For any compact perfect $P \subset \mathbb{R}$ and any C^1 function $f: \mathbb{R} \rightarrow \mathbb{R}^2$ we have $P^2 \notin f[P]$.*

The proof is based on the following two lemmas.

Lemma 3.2. *Let P be a perfect subset of \mathbb{R} and $f = \langle f_1, f_2 \rangle$ be a continuous function from P into \mathbb{R}^2 such that the coordinate function f_1 is differentiable. If $E = \{x \in P: f'_1(x) \neq 0\}$, then $f[E] \cap P^2$ is meager in P^2 .*

PROOF. Since the derivative of a coordinate function $f_1: P \rightarrow \mathbb{R}$ is Baire class one (see e.g. [8]), the set E is σ -compact and so is $f[E]$. Also, for every compact $K \subset E$, every level set $(f_1 \upharpoonright K)^{-1}(y) = \{x \in K: f_1(x) = y\}$ of $f_1 \upharpoonright K$ is finite. In particular, each vertical section of $f[K] = \{\langle f_1(x), f_2(x) \rangle: x \in K\}$ is finite, so $f[K] \cap P^2$ is nowhere dense in P^2 . \square

Lemma 3.3. *Let $g: \mathbb{R} \rightarrow \mathbb{R}$ be a C^1 function. If P is a compact perfect subset of \mathbb{R} such that $P \subset g[P]$, then there exists an $x \in P$ such that $|g'(x)| \geq 1$.*

PROOF. By way of contradiction, assume that $|g'(x)| < 1$ for every $x \in P$. Since P is compact and g' continuous, there exists an $M < 1$ such that $|g'(x)| < M$ for all $x \in P$. Notice that there exists a $\delta > 0$ such that

$$\left| \frac{g(x)-g(y)}{x-y} \right| < M \text{ for every } x, y \in P \text{ with } 0 < |x-y| \leq \delta. \quad (14)$$

Indeed, otherwise for every $n < \omega$ there exist $x_n, y_n \in P$ for which we have $0 < y_n - x_n \leq 2^{-n}$ and $\left| \frac{g(x_n)-g(y_n)}{x_n-y_n} \right| \geq M$. By the mean value theorem, there exist points $\xi_n \in (x_n, y_n)$ for which $|g'(\xi_n)| \geq M$. Choosing a subsequence, if necessary, we can assume that $\langle x_n \rangle_n$ converges to an $x \in P$. Then also $\langle \xi_n \rangle_n$ converges to x , which contradicts continuity of g' , since $\langle |g'(\xi_n)| \rangle_n$ does not converge to $|g'(x)| < M$.

For every $k < \omega$ let \mathcal{U}_k be a collection of the families $\{I_j: j < k\}$ of intervals such that each interval I_j has length $|I_j| \leq \delta$ and $P \subset \bigcup_{j < k} I_j$. Fix a $k < \omega$ for which the \mathcal{U}_k is not empty and let $L = \inf \{ \sum_{j < k} |I_j| : \{I_j: j < k\} \in \mathcal{U}_k \}$. Notice, that $L > 0$, even if P has measure 0. In fact, if P_0 is any subset of P containing $k+1$ points, then L is greater than or equal to the minimal distance between distinct points in P_0 .

Choose $\{I_j: j < k\} \in \mathcal{U}_k$ with $\sum_{j < k} |I_j| < L/M$. For every $j < k$ let J_j be the shortest interval containing $g[P \cap I_j]$. Then, by (14), $|J_j| \leq M|I_j|$. In particular, $\sum_{j < k} |J_j| \leq \sum_{j < k} M|I_j| < L$, so $\bigcup_{j < k} J_j \supset \bigcup_{j < k} g[P \cap I_j] = g[P]$ does not cover P . \square

PROOF OF THEOREM 3.1. Let $P \subseteq \mathbb{R}$ be compact and $f = \langle f_1, f_2 \rangle: \mathbb{R} \rightarrow \mathbb{R}^2$ be of class C^1 . By way of contradiction assume that $P^2 \subset f[P]$, and let $P_0 = \{x \in P: f'_1(x) = 0\}$. Then P_0 is closed, since f'_1 is continuous. Let $E = P \setminus P_0$. Then, by Lemma 3.2, $f[E]$ is meager in P^2 , so $f[P_0] \supset P^2 \setminus f[E]$ is dense in P^2 . Therefore, $P^2 \subset f[P_0]$, as $f[P_0]$ is compact.

Next, let E_0 be the set of all isolated points of P_0 and let $P_1 = P_0 \setminus E_0$. Then, P_1 is compact perfect and E_0 is countable. Therefore, as above, we conclude that $P^2 \subset f[P_1] \subset f_1[P_1] \times f_2[P_1]$. Hence, $P_1 \subset P \subset f_1[P_1]$.

Applying Lemma 3.3 to $g = f_1$ and P_1 , we conclude that there is an $x \in P_1$ such that $f'_1(x) \geq 1$. But this contradicts the definition of $P_0 \supset P_1$. \square

4 Compact sets $P \subset \mathbb{R}$ with C^0 Peano functions $f: P \rightarrow P^2$

The goal of this section is to give a full characterization of compact subsets P of \mathbb{R} for which there exists a C^0 Peano function $f: P \rightarrow P^2$. This is provided by the following theorem.

Theorem 4.1. *Let $P \subset \mathbb{R}$ be compact and let κ be the number of connected components in P . Then there exists a \mathcal{C}^0 Peano function $f: P \rightarrow P^2$ if, and only if, either $\kappa = 1$ or $\kappa = \mathfrak{c}$.*

Actually, since the classical Peano curve covers the case when P is connected ($\kappa = 1$) only disconnected sets P are of true interest in this result. For such sets the theorem can be reformulated as follows.

Corollary 4.2. *A disconnected compact set $P \subset \mathbb{R}$ admits a \mathcal{C}^0 Peano function $f: P \rightarrow P^2$ if, and only if, P has uncountably many components.*

The proof of the theorem will be based on the following two lemmas. To formulate them, we need to recall the following classical definitions. See Kechris [3, pp. 33-34].

For an $X \subseteq \mathbb{R}$ let $(X)'$ be the set of all accumulation points of X . For the ordinal numbers $\alpha, \lambda < \omega_1$, where λ is a limit ordinal, we define

$$X^{(0)} = X, X^{(\alpha+1)} = (X^{(\alpha)})', \text{ and } X^{(\lambda)} = \bigcap_{\alpha < \lambda} X^{(\alpha)}. \quad (15)$$

For a closed countable set $X \subset \mathbb{R}$, we define its *Cantor-Bendixon rank*, denoted $|X|_{CB}$, to be the least ordinal number $\alpha < \omega_1$ such that $X^{(\alpha)} = \emptyset$.

Lemma 4.3. *If $P \subset \mathbb{R}$ is a countable compact set and a function $f: P \rightarrow \mathbb{R}$ is countable, then $|f[P]|_{CB} \leq |P|_{CB}$.*

PROOF. We will show, by induction on β , that the condition

$$(I_\beta) \quad f[P]^{(\beta)} \subseteq f[P^{(\beta)}]$$

holds for every $\beta < \omega_1$. This clearly implies the result.

So, assume that, for some $\beta < \omega_1$, the inclusion $f[P]^{(\alpha)} \subseteq f[P^{(\alpha)}]$ holds for all $\alpha < \beta$. We need to show (I_β) . We will consider three cases.

$\beta = 0$: Then $f[P]^{(\beta)} = f[P] = f[P^{(\beta)}]$, so (I_β) holds.

$\beta > 0$ is a limit ordinal number: First notice that

$$(\bullet) \quad \bigcap_{\alpha < \beta} f[P^{(\alpha)}] \subseteq f[\bigcap_{\alpha < \beta} P^{(\alpha)}].$$

To see this, fix a point $y \in \bigcap_{\alpha < \beta} f[P^{(\alpha)}]$ and choose an increasing sequence $\langle \alpha_n < \beta : n < \omega \rangle$ cofinal with β , that is, such that $\lim_n \alpha_n = \beta$. Then, for every $n < \omega$, there exists an $x_n \in P^{(\alpha_n)} \subseteq P$ such that $y = f(x_n)$. By compactness of P , choosing a subsequence if necessary, we can assume that $\langle x_n \rangle_n$ converges to some $x \in P$. Since the sequence $\langle P^{(\alpha_n)} \rangle_n$ is decreasing, we have

$x \in \bigcap_{n < \omega} P^{(\alpha_n)} = \bigcap_{\alpha < \beta} P^{(\alpha)}$. Therefore, $y = f(x) \in f[\bigcap_{\alpha < \beta} P^{(\alpha)}]$, as required for proving (\bullet) .

Now, by (\bullet) ,

$$f[P]^{(\beta)} = \bigcap_{\alpha < \beta} f[P]^{(\alpha)} \subseteq \bigcap_{\alpha < \beta} f[P^{(\alpha)}] \subseteq f[\bigcap_{\alpha < \beta} P^{(\alpha)}] = f[P^{(\beta)}],$$

where the first inclusion is justified by (I_α) . So, once again, (I_β) holds.

β is a successor ordinal: Suppose $\beta = \alpha + 1$ and fix a $y \in f[P]^{(\beta)} = (f[P]^{(\alpha)})'$. Then, there exists a one-to-one sequence $\langle y_n \in f[P]^{(\alpha)} : n < \omega \rangle$ converging to y . By the inductive assumption $y_n \in f[P]^{(\alpha)} \subseteq f[P^{(\alpha)}]$, so, for every $n < \omega$, there exists an $x_n \in P^{(\alpha)}$ with $y_n = f(x_n)$. Since the sequence $\langle y_n : n < \omega \rangle$ is one-to-one, so is $\langle x_n \in P^{(\alpha)} : n < \omega \rangle$. By compactness of $P^{(\alpha)}$, choosing a subsequence if necessary, we can assume that $\langle x_n \rangle_n$ converges to some $x \in P^{(\alpha)}$. Since $\langle x_n \rangle_n$ is one-to-one, $x \in (P^{(\alpha)})' = P^{(\beta)}$. Finally, $f(x) = f(\lim_n x_n) = \lim_n f(x_n) = \lim_n y_n = y$, so $y = f(x) \in f[P^{(\beta)}]$, as needed for the proof of (I_β) . \square

Lemma 4.4. *Let P be a countable compact subset of \mathbb{R} . If P is infinite, then $|P|_{CB} < |P \times P|_{CB}$.*

PROOF. Let $|P|_{CB} = \beta$. The compactness of P implies that β is a successor ordinal, say $\beta = \alpha + 1$. We need to show that $((P \times P)^{(\alpha)})' = (P \times P)^{(\alpha+1)} \neq \emptyset$.

Notice, that $X' \times Y \subseteq (X \times Y)'$ for every $X, Y \subset \mathbb{R}$. From this, an obvious inductive argument shows that $X^{(\alpha)} \times Y \subseteq (X \times Y)^{(\alpha)}$. In particular, we have $P^{(\alpha)} \times P \subseteq (P \times P)^{(\alpha)}$. Thus, it is enough to show that $(P^{(\alpha)} \times P)' \neq \emptyset$. But this is obvious, since $P^{(\alpha)} \neq \emptyset$ and P is infinite. \square

PROOF OF THEOREM 4.1. The argument naturally leads to the following four cases.

$\kappa = 1$: In this case the classical Peano curve works.

$\kappa > 1$ is finite: Let $f: P \rightarrow \mathbb{R}^2$ be continuous. Then $f[P]$ can have at most κ -many components. Since P^2 has κ^2 components and $\kappa^2 > \kappa$, $f[P]$ cannot be equal P^2 .

κ is countable infinite: This means that $\kappa = \omega$. We need to show that there is no \mathcal{C}^0 Peano function $f: P \rightarrow P^2$.

First we note that this is true when P is totally disconnected (i.e., it has only one-point components):

- (*) if an infinite compact totally disconnected set P has countably many components, then there is no continuous function from P onto $P^2 = P \times P$.

Indeed, if $f: P \rightarrow \mathbb{R}^2$ is continuous then, by Lemma 4.3, $|f[P]|_{CB} \leq |P|_{CB}$. So, $f[P]$ cannot be equal P^2 since, by Lemma 4.4, $|P|_{CB} < |P^2|_{CB}$. The general case will be reduced to (*).

By way of contradiction, suppose that there exists a continuous function $f = \langle f_1, f_2 \rangle$ from P onto P^2 . Let \sim be an equivalence relation defined as: $x \sim y$ if, and only if, x and y belong to the same component of P . The equivalence class of $x \in P$ with respect to \sim will be denoted $[x]$. Let $P/\sim = \{[x]: x \in P\}$ be the quotient space, that is, $U \subseteq P/\sim$ is declared open if, and only if, the set $\hat{U} = \bigcup\{[x]: [x] \in U\}$ is open in P . Notice that P/\sim is homeomorphic to a subset of \mathbb{R} , since

P/\sim is compact, Hausdorff, totally disconnected.

Indeed, if $\{U_j: j \in J\}$ is an open cover of P/\sim , then $\{\hat{U}_j: j \in J\}$ is an open cover of P . So, there is a finite $J_0 \subseteq J$ such that $\{\hat{U}_j: j \in J_0\}$ covers P . Therefore, $\{U_j: j \in J_0\}$ is a cover of P/\sim , implying compactness of P/\sim . To see the other two properties, take $x, y \in P$ with $[x] \neq [y]$. We can assume that $x < y$. Then, there exists an $r \in \mathbb{R} \setminus P$ such that $[x] \subset (-\infty, r)$ and $[y] \subset (r, \infty)$. In particular, if $U = P \cap (r, \infty)$, then \hat{U} is a clopen subset of P/\sim containing $[x]$ but not $[y]$. It is worth noting that our space P/\sim falls into a broader class of quotient spaces which are metrizable, see e.g. [1, theorem 4.2.13].

Let $i \in \{1, 2\}$. Since f_i is a continuous function from P into itself, we have $f_i([x]) = [f_i(x)]$ for every $x \in P$. In particular, the function $g_i: (P/\sim) \rightarrow (P/\sim)$ given by $g_i([x]) = [f_i(x)]$ is well defined and it is continuous, since for every U open in P/\sim , the set $W = g_i^{-1}(U)$ is open in P/\sim , as $\hat{W} = f_i^{-1}(\hat{U})$.

The above shows that function $g = \langle g_1, g_2 \rangle: (P/\sim) \rightarrow (P/\sim)^2$ is well defined and continuous. Moreover, it is onto $(P/\sim)^2$, since $f[P] = P^2$. The space P/\sim is countable so this contradicts (*), completing the proof of this case.

κ is uncountable: In this case $\kappa = \mathfrak{c}$. Recall, that 2^ω can be mapped onto any compact metric space, see e.g. [3, theorem 4.18]. In particular, there exists a continuous function 2^ω onto P^2 .

Also, there exists a continuous function g from P onto 2^ω . Indeed, we can define a Cantor-like tree $\{P_s: s \in 2^{<\omega}\}$ of compact subsets of P such that $P_\emptyset = P$ and every P_s is split into two clopen subsets, P_{s0} and P_{s1} , each containing uncountably many components of P . For $t \in 2^\omega$ put $g(x) = t$ if, any only if, $x \in \bigcap_{n < \omega} P_{t \upharpoonright n}$. Then g is as required.

Finally notice that $f = h \circ g$ is continuous and maps P onto P^2 . □

5 Final remarks and open problems

Although we proved that for a compact perfect $P \subset \mathbb{R}$ there is no Peano function f from P onto P^2 which can be extended to a \mathcal{C}^1 function $\hat{f}: \mathbb{R} \rightarrow \mathbb{R}^2$, the argument used in the proof of Theorem 3.1 does not work without the extendability assumption of f . Of course, by Proposition 1.1(b), the extendability would play no role if we could prove a version of Theorem 3.1 with the class \mathcal{C}^1 replaced by \mathcal{D}^1 . But, once again, our argument does not seem to generalize to this case.

In light of this discussion, the following question seems to be of interest.

Problem 1. Does there exist a compact perfect set $P \subset \mathbb{R}$ and a \mathcal{D}^1 function f from P onto P^2 ? If so, can such an f be \mathcal{C}^1 ? (See Remark 1.2.)

Also, Theorem 4.1 gives a full characterization of compact sets P admitting \mathcal{C}^0 Peano functions. It would be interesting to find analogous characterization that includes also the unbounded closed sets. However, if there exists such a characterization (in terms of a structure of connected components), it seems it would be quite complicated in nature.

Finally, in the example given in Theorem 2.2, the \mathcal{C}^∞ Peano function f from P onto P^2 is extendable to a \mathcal{C}^∞ function $\hat{f}: \mathbb{R} \rightarrow \mathbb{R}^2$. Is this always the case? More precisely it seems to us that the following question should have a negative answer.

Problem 2. Let $P \subset \mathbb{R}$ be a perfect subset of \mathbb{R} for which there is a \mathcal{C}^∞ function from P onto P^2 . Does this imply that there exists a \mathcal{C}^∞ function $f: \mathbb{R} \rightarrow \mathbb{R}^2$ such that $f[P] = P^2$?

References

- [1] R. Engelking, *General Topology*, Heldermann-Verlag, New York, 1989.
- [2] V. Jarník, *Sur l'extension du domaine de définition des fonctions d'une variable, qui laisse intacte la dérivabilité de la fonction*, Bull. Internat. Acad. Sci. Boheme, (1923) 1–5.
- [3] A. Kechris, *Classical Descriptive Set Theory*, Springer-Verlag, New York, 1995.
- [4] M. Koc and L. Zajíček, *A joint generalization of Whitney's C^1 extension theorem and Aversa–Laczkovich–Preiss' extension theorem*, J. Math. Anal. Appl., **388** (2012), 1027–1039.

- [5] K. Kuratowski, *Topology I*, Academic Press, New York, 1966.
- [6] M. Morayne, *On differentiability of Peano type functions*, Colloq. Math., **53** (1987), 129–131.
- [7] G. Peano, *Sur une courbe, qui remplit toute une aire plane*, Math. Ann., **36** (1890), 157–160.
- [8] G. Petruska and M. Laczkovich, *Baire 1 functions, approximately continuous functions and derivatives*, Acta Math. Acad. Sci. Hung., **25** (1974), 189–212.
- [9] H. Sagan, *Space-Filling Curves*, Springer-Verlag, New York, 1994.
- [10] H. Whitney, *Analytic extensions of differentiable functions defined in closed sets*, Trans. Amer. Math. Soc., **36** (1934), 63–89.