

*Research Article*

**On Minimal Norms on  $M_n$**

Madjid Mirzavaziri and Mohammad Sal Moslehian

Received 18 June 2007; Accepted 19 August 2007

Recommended by Allan C. Peterson

We show that for each minimal norm  $N(\cdot)$  on the algebra  $\mathcal{M}_n$  of all  $n \times n$  complex matrices, there exist norms  $\|\cdot\|_1$  and  $\|\cdot\|_2$  on  $\mathbb{C}^n$  such that  $N(A) = \max \{\|Ax\|_2 : \|x\|_1 = 1, x \in \mathbb{C}^n\}$  for all  $A \in \mathcal{M}_n$ . This may be regarded as an extension of a known result on characterization of minimal algebra norms.

Copyright © 2007 M. Mirzavaziri and M. S. Moslehian. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**1. Introduction**

Let  $\mathcal{M}_n$  denote the algebra of all  $n \times n$  complex matrices  $A$  with entries in  $\mathbb{C}$ , together with the usual matrix operations. By an algebra norm (or a matrix norm) we mean a norm  $\|\cdot\|$  on  $\mathcal{M}_n$  such that  $\|AB\| \leq \|A\|\|B\|$  for all  $A, B \in \mathcal{M}_n$ . It is easy to see that the norm  $\|A\|_\sigma = \sum_{i,j=1}^n |\alpha_{ij}|$  is an algebra norm, but the norm  $\|A\|_m = \max \{|\alpha_{i,j}| : 1 \leq i, j \leq n\}$  is not an algebra norm, (see [1]).

Let  $\|\cdot\|_1$  and  $\|\cdot\|_2$  be two norms on  $\mathbb{C}^n$ . Then the norm  $\|\cdot\|_{1,2}$  on  $\mathcal{M}_n$  defined by  $\|A\|_{1,2} := \max \{\|Ax\|_2 : \|x\|_1 = 1\}$  is called the generalized induced (or g-ind) norm constructed via  $\|\cdot\|_1$  and  $\|\cdot\|_2$ . If  $\|\cdot\|_1 = \|\cdot\|_2$ , then  $\|\cdot\|_{1,1}$  is called an induced norm.

It is known that  $\|A\|_C = \max \{\sum_{i=1}^n |\alpha_{i,j}| : j \leq n\}$ ,  $\|A\|_R = \max \{\sum_{j=1}^n |\alpha_{i,j}| : 1 \leq i \leq n\}$  and the spectral norm  $\|A\|_S = \max \{\sqrt{\lambda} : \lambda \text{ is an eigenvalue of } A^*A\}$  are induced by  $\ell_1, \ell_\infty$ , and  $\ell_2$ , respectively, (cf. [2]). Recall that the  $\ell_p$ -norm ( $1 \leq p \leq \infty$ ) on  $\mathbb{C}^n$  is defined by

$$\ell_p(x) = \ell_p\left(\sum_{i=1}^n x_i e_i\right) = \begin{cases} \left(\sum_{i=1}^n |x_i|^p\right)^{1/p}, & 1 \leq p < \infty, \\ \max \{|x_1|, \dots, |x_n|\}, & p = \infty. \end{cases} \tag{1.1}$$

## 2 Abstract and Applied Analysis

It is known that the algebra norm  $\|A\| = \max\{\|A\|_C, \|A\|_R\}$  is not induced, and it is not hard to show that it is not  $g$ -ind too (cf. Corollary 3.2.6 of [3]).

A norm  $N(\cdot)$  on  $\mathcal{M}_n$  is called minimal if for any norm  $\|\cdot\|$  on  $\mathcal{M}_n$  satisfying  $\|\cdot\| \leq N(\cdot)$ , we have  $\|\cdot\| = N(\cdot)$ . It is known [3, Theorem 3.2.3] that an algebra norm is an induced norm if and only if it is a minimal element in the set of all algebra norms. Note that a generalized induced norm may not be minimal. For instance, put  $\|\cdot\|_\alpha = \ell_\infty(\cdot)$ ,  $\|\cdot\|_\beta = 2\ell_2(\cdot)$ , and  $\|\cdot\|_\gamma = \ell_2(\cdot)$ . Then  $\|\cdot\|_\gamma \leq \|\cdot\|_{\alpha,\beta}$  but  $\|\cdot\|_{\gamma,\beta} \neq \|\cdot\|_{\alpha,\beta}$ .

In [1], the authors investigate generalized induced norms. In particular, they examine the problem that “for any norm  $\|\cdot\|$  on  $\mathcal{M}_n$ , are there two norms  $\|\cdot\|_1$  and  $\|\cdot\|_2$  on  $\mathbb{C}^n$  such that  $\|A\| = \max\{\|Ax\|_2 : \|x\|_1 = 1\}$  for all  $A \in \mathcal{M}_n$ ?” In this short note, we utilize some ideas of [1] to study the minimal norms on  $\mathcal{M}_n$ . More precisely, we show that for each minimal norm  $N(\cdot)$  on the algebra  $\mathcal{M}_n$  of all  $n \times n$  complex matrices, there exist norms  $\|\cdot\|_1$  and  $\|\cdot\|_2$  on  $\mathbb{C}^n$  such that  $N(A) = \max\{\|Ax\|_2 : \|x\|_1 = 1, x \in \mathbb{C}^n\}$  for all  $A \in \mathcal{M}_n$ . In particular, if  $N(\cdot)$  is an algebra norm, then  $\|\cdot\|_1 = \|\cdot\|_2$ . This may be regarded as an extension of the above known result on characterization of minimal algebra norms.

### 2. Main result

For  $x \in \mathbb{C}^n$  and  $1 \leq j \leq n$ , let  $C_{x,j} \in \mathcal{M}_n$  be defined by the operator  $C_{x,j}(y) = y_j x$ . Hence  $C_{x,j}$  is the  $n \times n$  matrix with  $x$  in the  $j$  column and 0 elsewhere. Define  $C_x \in \mathcal{M}_n$  by  $C_x = \sum_{j=1}^n C_{x,j}$ . Hence  $C_x$  is the  $n \times n$  matrix whose all columns are  $x$ .

If  $\|\cdot\|_{1,2}$  is a generalized induced norm on  $\mathcal{M}_n$  obtained via  $\|\cdot\|_1$  and  $\|\cdot\|_2$  then  $\|C_x\|_{1,2} = \alpha \|x\|_2$ , where  $\alpha = \max\{|\sum_{j=1}^n y_j| : \|(y_1, \dots, y_j, \dots, y_n)\|_1 = 1\}$ .

To achieve our goal, we need the following lemmas.

**LEMMA 2.1** [1, Theorem 2.7]. *Let  $\|\cdot\|_1$  and  $\|\cdot\|_2$  be two norms on  $\mathbb{C}^n$ . Then  $\|\cdot\|_{1,2}$  is an algebra norm on  $\mathcal{M}_n$  if and only if  $\|\cdot\|_1 \leq \|\cdot\|_2$ .*

**LEMMA 2.2** [1, Corollary 2.5].  *$\|\cdot\|_{1,2} = \|\cdot\|_{3,4}$  if and only if there exists  $\gamma > 0$  such that  $\|\cdot\|_1 = \gamma \|\cdot\|_3$  and  $\|\cdot\|_2 = \gamma \|\cdot\|_4$ .*

**THEOREM 2.3.** *Let  $N(\cdot)$  be a minimal norm on  $\mathcal{M}_n$ , then  $N(\cdot) = \|\cdot\|_{1,2}$  for some  $\|\cdot\|_1$  and  $\|\cdot\|_2$  on  $\mathbb{C}^n$ . Moreover, if  $N(\cdot)$  is an algebra norm, then  $\|\cdot\|_1 = \|\cdot\|_2$ .*

*Proof.* For  $x \in \mathbb{C}^n$ , set

$$\begin{aligned} \|x\|_1 &= \max\{N(C_{Ax}) : N(A) = 1, A \in \mathcal{M}_n\}, \\ \|x\|_2 &= N(C_x). \end{aligned} \tag{2.1}$$

We will show that  $\|\cdot\|_1$  and  $\|\cdot\|_2$  are norms on  $\mathbb{C}^n$ .

To see that  $\|\cdot\|_1$  is a norm, let  $x \in \mathbb{C}^n$ . Then  $\|x\|_1 = 0$  if and only if  $N(C_{Ax}) = 0$  for all matrix  $A$  with  $N(A) = 1$ , and this holds if and only if  $Ax = 0$  for all  $A$ , or equivalently  $x = 0$ .

For  $\alpha \in \mathbb{C}^n$  and  $x, y \in \mathbb{C}^n$ , we have

$$\begin{aligned}
\|\alpha x\|_1 &= \max \{N(C_{A(\alpha x)}) : N(A) = 1, A \in \mathcal{M}_n\} \\
&= \max \{N(\alpha C_{Ax}) : N(A) = 1, A \in \mathcal{M}_n\} \\
&= \max \{|\alpha|N(C_{Ax}) : N(A) = 1, A \in \mathcal{M}_n\} \\
&= |\alpha| \max \{N(C_{Ax}) : N(A) = 1, A \in \mathcal{M}_n\} \\
&= |\alpha| \|x\|_1, \\
\|x + y\|_1 &= \max \{N(C_{A(x+y)}) : N(A) = 1, A \in \mathcal{M}_n\} \\
&= \max \{N(C_{Ax} + C_{Ay}) : N(A) = 1, A \in \mathcal{M}_n\} \\
&\leq \max \{N(C_{Ax}) : N(A) = 1, A \in \mathcal{M}_n\} \\
&\quad + \max \{N(C_{Ay}) : N(A) = 1, A \in \mathcal{M}_n\} \\
&= \|x\|_1 + \|y\|_1.
\end{aligned} \tag{2.2}$$

To see that  $\|\cdot\|_2$  is a norm, let  $x \in \mathbb{C}^n$ . Then  $\|x\|_2 = 0$  if and only if  $C_x = 0$  and this holds if and only if  $x = 0$ .

For  $\alpha \in \mathbb{C}^n$  and  $x, y \in \mathbb{C}^n$ , we have

$$\begin{aligned}
\|\alpha x\|_2 &= N(C_{\alpha x}) = N(\alpha C_x) = |\alpha|N(C_x) = |\alpha| \|x\|_2, \\
\|x + y\|_2 &= N(C_{x+y}) = N(C_x + C_y) \leq N(C_x) + N(C_y) = \|x\|_2 + \|y\|_2.
\end{aligned} \tag{2.3}$$

Now let  $A \in \mathcal{M}_n \setminus \{0\}$ . Then  $N(A/N(A)) = 1$  so that

$$\left\| \frac{A}{N(A)}(x) \right\|_2 = N(C_{(A/N(A))(x)}) \leq \|x\|_1, \tag{2.4}$$

whence

$$\|Ax\|_2 \leq N(A) \|x\|_1. \tag{2.5}$$

Therefore  $\|A\|_{1,2} \leq N(A)$ . Since  $N(\cdot)$  is a minimal norm, we conclude that  $\|A\|_{1,2} = N(A)$ .

If  $N(A)$  is an algebra norm, then Lemma 2.1 implies that  $\|\cdot\|_1 \leq \|\cdot\|_2$ .

Next, let  $A \in \mathcal{M}_n$ . It follows from  $\|Ax\|_1 \leq \|A\|_{1,1} \|x\|_1 \leq \|A\|_{1,1} \|x\|_2$ , ( $x \in \mathbb{C}^n$ ) that  $\|A\|_{2,1} \leq \|A\|_{1,1}$ . In a similar fashion, one can get

$$\|\cdot\|_{2,1} \leq \|\cdot\|_{k,k} \leq \|\cdot\|_{1,2} \quad (k = 1, 2). \tag{2.6}$$

By the minimality of  $\|\cdot\|_{1,2}$ , we deduce that  $\|\cdot\|_{1,2} = \|\cdot\|_{1,1}$ . It then follows from Lemma 2.2 that  $\|\cdot\|_1 = \|\cdot\|_2$ .  $\square$

## References

- [1] S. Hejazian, M. Mirzavaziri, and M. S. Moslehian, "Generalized induced norms," *Czechoslovak Mathematical Journal*, vol. 57, no. 1, pp. 127–133, 2007.
- [2] R. A. Horn and C. R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, UK, 1994.
- [3] G. R. Belitskiĭ and Yu. I. Lyubich, *Matrix Norms and Their Applications*, vol. 36 of *Operator Theory: Advances and Applications*, Birkhäuser, Basel, Switzerland, 1988.

Madjid Mirzavaziri: Department of Mathematics, Ferdowsi University, P.O. Box 1159, Mashhad 91775, Iran; Banach Mathematical Research Group (BMRG), Mashhad, Iran  
*Email address:* madjid@mirzavaziri.com

Mohammad Sal Moslehian: Department of Mathematics, Ferdowsi University, P.O. Box 1159, Mashhad 91775, Iran; Centre of Excellence in Analysis on Algebraic Structures (CEAAS), Ferdowsi University, Iran  
*Email address:* moslehian@ferdowsi.um.ac.ir