

Research Article

Some New Bounds for Mathieu's Series

Abdolhossein Hoorfar and Feng Qi

Received 9 February 2007; Revised 3 June 2007; Accepted 6 August 2007

Recommended by Lance L. Littlejohn

Two upper and lower bounds for Mathieu's series are established, which refine to a certain extent a sharp double inequality obtained by Alzer-Brenner-Ruehr in 1998. Moreover, the very closer lower and upper bounds for $\zeta(3)$ are deduced.

Copyright © 2007 A. Hoorfar and F. Qi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In 1890, Mathieu in [1] defined $S(r)$ for $r > 0$ by

$$S(r) = \sum_{n=1}^{\infty} \frac{2n}{(n^2 + r^2)^2} \quad (1.1)$$

and conjectured that $S(r) < 1/r^2$. We call formula (1.1) Mathieu's series.

There has been a lot of literature about the estimations of $S(r)$ for more than 100 years till 1998, for example, [2–14] and the references therein. In [9], Makai proved that

$$\frac{1}{r^2 + 1/2} < S(r) < \frac{1}{r^2}. \quad (1.2)$$

In 1998, Alzer et al. presented in [2] that

$$\frac{1}{r^2 + 1/2\zeta(3)} < S(r) < \frac{1}{r^2 + 1/6}, \quad (1.3)$$

where ζ denotes the zeta function and the constants $1/2\zeta(3)$ and $1/6$ in (1.3) are the best possible.

2 Abstract and Applied Analysis

After 2000, among other things, several open problems on the estimations and integral representations of generalized Mathieu's series were posed in [15–17] by Guo and Qi. Stimulated by or originated from these open problems, a lot of articles such as [18–37] have been published in variant reputable journals by many mathematicians all over the world.

In this article, by utilizing the well-known telescope technique ever used in [9, 38], we would like to improve or refine the sharp double inequality (1.3) and to establish a very closer double inequality for $\zeta(3)$.

Our main results are the following four theorems.

THEOREM 1.1. For $r > 0$,

$$S(r) > \frac{1}{r^2 + 1/6 + (r^2 + 6)/3(9r^2 + 8)} = \frac{1}{r^2 + 1/2 - 2(4r^2 + 1)/3(9r^2 + 8)}. \quad (1.4)$$

Remark 1.2. By standard argument, it is showed readily that inequality (1.4) is better than the left-hand side inequality in (1.3) when $r > 2\sqrt{(5\zeta(3) - 6)/(27 - 11\zeta(3))} = 0.05\dots$

THEOREM 1.3. For $r > 0$,

$$\begin{aligned} \frac{1}{r^2 + 1/6 + 5/6(2r^2 + 3)} &= \frac{1}{r^2 + 1/2 - (4r^2 + 1)/6(2r^2 + 3)} < S(r) \\ &< \frac{1}{r^2 + 1/2 - (4r^2 + 1)/2(2r^2 - 3 + 4\sqrt{r^4 + 2r^2 + 5})}. \end{aligned} \quad (1.5)$$

Remark 1.4. It is not difficult to verify that the left-hand side inequality in (1.5) is better than the left-hand side inequality in (1.3) when $r > \sqrt{(8\zeta(3) - 9)/2[3 - \zeta(3)]} = 0.41\dots$ and that the right-hand side inequality in (1.5) is better than the right-hand side inequality in (1.3) when $r < \sqrt{239/16} = 3.86\dots$

It is important to remark that inequality (1.4) and the left-hand side inequality in (1.5) do not include each other, which can be proved straightforwardly.

THEOREM 1.5. For $r > 0$,

$$S(r) < \frac{1}{\sqrt{r^4 + 2r^2 + 2} - 1}. \quad (1.6)$$

Remark 1.6. It is easy to deduce that inequality (1.6) is better than the right-hand side inequality in (1.3) when $0 < r < \sqrt{23/12} = 1.38\dots$

THEOREM 1.7. For $m \in \mathbb{N}$, let $S_3(m) = \sum_{n=1}^m (1/n^3)$. Then

$$\frac{1}{2m^2 + 2m + 1 - 1/6(m^2 + m + 3/2)} < \zeta(3) - S_3(m) < \frac{1}{2m^2 + 2m + 1 - 1/6(m^2 + m + 1)}. \quad (1.7)$$

Remark 1.8. Calculation by Mathematica 5.2 shows that

$$\zeta(3) = 1.202056903159594285399\dots \quad (1.8)$$

If taking m from 1 to 9, the sums of the right side term in (1.7) and $S_3(m)$ are

$$\begin{array}{lll}
 1.202247191011235955, & 1.202064220183486239, & 1.202057560382342322, \\
 1.202057003155139651, & 1.202056924652726768, & 1.202056909039779896, \\
 1.202056905080018071, & 1.202056903877571143, & 1.202056903458154800.
 \end{array} \tag{1.9}$$

If taking m from 1 to 9, the sums of the left side term in (1.7) and $S_3(m)$ are

$$\begin{array}{lll}
 1.201923076923076923, & 1.202054794520547945, & 1.202056799882886839, \\
 1.202056893315403149, & 1.202056901714344462, & 1.202056902872941459, \\
 1.202056903088695828, & 1.202056903138840387, & 1.202056903152657143.
 \end{array} \tag{1.10}$$

These numerical computations by mathematic 5.2 reveals that inequalities in (1.7) give much accurate approximations from left and right.

COROLLARY 1.9. *If $1 \leq \delta < 3/2$ and $m \geq \sqrt{(3\delta^2 - \delta + 1/12)/(6 - 4\delta)} - 1$, then*

$$\zeta(3) < S_3(m) + \frac{1}{2m^2 + 2m + 1 - 1/6(m^2 + m + \delta)}. \tag{1.11}$$

Remark 1.10. In [39, 40], the number $\zeta(3)$ was estimated by using Jordan’s inequality and its refinements. In [41, 42], some more general conclusions were obtained.

Remark 1.11. Finally, an open problem is posed: find the best possible constants a and b such that

$$\frac{1}{r^2 + 1/2 - (4r^2 + 1)/12(r^2 + a)} < S(r) < \frac{1}{r^2 + 1/2 - (4r^2 + 1)/12(r^2 + b)} \tag{1.12}$$

holds true for all $r > 0$.

It is clear that $a \leq 3/2$ and $b \geq 1/4$.

2. Proofs of theorems and corollary

Now we are in a position to prove our theorems and corollary.

Proof of Theorem 1.1. For $n \in \mathbb{N}$, let

$$w_n(r) = n(n - 1) + r^2 + \frac{1}{2} - \frac{\theta}{n^2 + \gamma}, \tag{2.1}$$

where $\theta = (1/3)(r^2 + 1/4)$ and γ is a possible and undetermined positive function of r such that

$$\frac{1}{w_n(r)} - \frac{1}{w_{n+1}(r)} \leq \frac{2n}{(n^2 + r^2)^2}. \tag{2.2}$$

4 Abstract and Applied Analysis

Straightforward computation yields that

$$\frac{1}{w_n(r)} - \frac{1}{w_{n+1}(r)} = \frac{2n\{1 + \theta(1 + 1/2n)/(n^2 + \gamma)[(n+1)^2 + \gamma]\}}{(n^2 + r^2)^2 + \theta Q(n, r, \gamma)/(n^2 + \gamma)[(n+1)^2 + \gamma]}, \quad (2.3)$$

where

$$Q(n, r, \gamma) = n^4 + 4n^3 + (4\gamma - 2r^2 - 1)n^2 + (6\gamma - 2r^2 - 2)n + 3\gamma^2 + 2(1 - r^2)\gamma - \frac{2r^2}{3} - \frac{5}{12}. \quad (2.4)$$

It is easy to see that if

$$\frac{1 + 1/2n}{Q(n, r, \gamma)} \leq \frac{1}{(n^2 + r^2)^2}, \quad (2.5)$$

then inequality (2.2) holds. Further, inequality (2.5) is equivalent to

$$n^4 + 4n^3 + (4\gamma - 2r^2 - 1)n^2 + (6\gamma - 2r^2 - 2)n + 3\gamma^2 + 2(1 - r^2)\gamma - \frac{2r^2}{3} - \frac{5}{12} \geq \left(1 + \frac{1}{2n}\right)(n^2 + r^2)^2, \quad (2.6)$$

which can be rewritten as

$$7n^3 + (8\gamma - 8r^2 - 2)n^2 + (12\gamma - 6r^2 - 4)n + 6\gamma^2 + 4(1 - r^2)\gamma - 2r^4 - \frac{4r^2}{3} - \frac{5}{6} - \frac{r^4}{n} \geq 0, \quad (2.7)$$

which can be further rearranged as

$$f(n, \gamma) \triangleq (n-1) \left[7n^2 + (8\gamma - 8r^2 + 5)n + 20\gamma - 14r^2 + 1 + \frac{r^4}{n} \right] + 6\gamma^2 + 4(6 - r^2)\gamma - 3r^4 - \frac{46}{3}r^2 + \frac{1}{6} \geq 0. \quad (2.8)$$

Direct computation reveals that

$$f\left(n, \frac{9r^2}{8}\right) = (n-1) \left[7n^2 + (r^2 + 5)n + \frac{17}{2}r^2 + 1 + \frac{r^4}{n} \right] + \frac{3}{32}r^4 + \frac{35}{3}r^2 + \frac{1}{6} > 0, \quad (2.9)$$

but

$$f(n, r^2) = (n-1) \left(7n^2 + 5n + 6r^2 + \frac{r^4}{n} \right) - r^4 + \frac{26}{3}r^2 + \frac{1}{6} \quad (2.10)$$

is negative if r is large enough. Consequently, if taking $\gamma = 9r^2/8$, then inequality (2.2) is valid. Summing up on both sides of (2.2), with respect to $n = 1, 2, \dots$, leads to (1.4). The proof of Theorem 1.1 is finished. \square

Proof of Theorem 1.3. Now let us consider the sequence

$$v_n(r) = n(n-1) + r^2 + \frac{1}{2} - \frac{\theta}{n(n-1) + \beta} \tag{2.11}$$

for $n \in \mathbb{N}$, where θ and β are two undetermined functions of r , in order that

$$\frac{1}{v_n(r)} - \frac{1}{v_{n+1}(r)} < \frac{2n}{(n^2 + r^2)^2}. \tag{2.12}$$

Direct calculation yields that

$$\frac{1}{v_n(r)} - \frac{1}{v_{n+1}(r)} = \frac{2n + 2\theta n / (n^2 - n + \beta)(n^2 + n + \beta)}{(n^2 + r^2)^2 + P(n, r, \theta, \beta) / (n^2 - n + \beta)(n^2 + n + \beta)}, \tag{2.13}$$

where

$$P(n, r, \theta, \beta) = \left(r^2 + \frac{1}{4} - 2\theta\right)n^4 + \left(r^2 + \frac{1}{4}\right)\beta^2 - \theta\beta(2r^2 + 1) + \theta^2 + \left[\left(r^2 + \frac{1}{4}\right)(2\beta - 1) - \theta(2\beta + 2r^2 + 3)\right]n^2. \tag{2.14}$$

Letting $r^2 + 1/4 - 2\theta = \theta$ and

$$\left(r^2 + \frac{1}{4}\right)(2\beta - 1) - \theta(2\beta + 2r^2 + 3) = 2\theta r^2 \tag{2.15}$$

give

$$\theta = \frac{1}{3}\left(r^2 + \frac{1}{4}\right), \quad \beta = r^2 + \frac{3}{2}. \tag{2.16}$$

Consequently,

$$\begin{aligned} P(n, r, \theta, \beta) &= \theta n^4 + 2\theta r^2 n^2 + 3\theta\beta^2 - \theta\beta(2r^2 + 1) + \theta^2 \\ &= \theta(n^2 + r^2)^2 + \theta[3\beta^2 - \beta(2r^2 + 1) + \theta - r^4] \\ &= \theta(n^2 + r^2)^2 + \frac{16}{3}\theta(r^2 + 1). \end{aligned} \tag{2.17}$$

As a result,

$$\begin{aligned} \frac{1}{v_2(r)} - \frac{1}{v_{n+1}(r)} &= \frac{2n + 2\theta n / (n^2 - n + \beta)(n^2 + n + \beta)}{(n^2 + r^2)^2 + (\theta(n^2 + r^2)^2 + 16\theta(r^2 + 1)/3) / (n^2 - n + \beta)(n^2 + n + \beta)} \\ &< \frac{2n + 2\theta n / (n^2 - n + \beta)(n^2 + n + \beta)}{(n^2 + r^2)^2 + \theta(n^2 + r^2)^2 / (n^2 - n + \beta)(n^2 + n + \beta)} = \frac{2n}{(n^2 + r^2)^2}. \end{aligned} \tag{2.18}$$

Summing up on both sides of the above inequality with respect to $n \in \mathbb{N}$ leads to

$$S(r) > \frac{1}{\nu_1} = \frac{1}{r^2 + 1/2 - \theta/\beta} = \frac{1}{r^2 + 1/2 - (4r^2 + 1)/(12r^2 + 18)}. \quad (2.19)$$

As mentioned above, taking $\theta = (1/3)(r^2 + 1/4)$ and simplifying yield that

$$\begin{aligned} P(n, r, \theta, \beta) &= \theta(n^2 + r^2)^2 - \theta[(4r^2 + 6 - 4\beta)n^2 - 3\beta^2 + (2r^2 + 1)\beta + r^4 - \theta] \\ &= \theta(n^2 + r^2)^2 - \theta(4r^2 + 6 - 4\beta)(n^2 - 1) + \theta R, \end{aligned} \quad (2.20)$$

where

$$R = 3\beta^2 - (2r^2 - 3)\beta - r^4 - \frac{11}{3}r^2 - \frac{71}{12}. \quad (2.21)$$

Now choosing $\beta > 0$ such that $R = 0$ gives

$$\beta = \frac{2r^2 - 3 + 4\sqrt{r^4 + 2r^2 + 5}}{6}. \quad (2.22)$$

It is observed that

$$4r^2 + 6 - 4\beta = \frac{8}{3}(r^2 + 3 - \sqrt{r^4 + 2r^2 + 5}) > 0 \quad (2.23)$$

and, for $n \in \mathbb{N}$,

$$P(n, r, \theta, \beta) = \theta(n^2 + r^2)^2 - \frac{8\theta}{3}(r^2 + 3 - \sqrt{r^4 + 2r^2 + 5})(n^2 - 1) \geq \theta(n^2 + r^2)^2. \quad (2.24)$$

Therefore,

$$\frac{1}{\nu_n(r)} - \frac{1}{\nu_{n+1}(r)} > \frac{2n}{(n^2 + r^2)^2}. \quad (2.25)$$

Summing up on both sides from $n = 1$ to ∞ gives

$$\frac{1}{\nu_1(r)} = \frac{1}{r^2 + 1/2 - \theta/\beta} = \frac{1}{r^2 + 1/2 - (4r^2 + 1)/2(2r^2 - 3 + 4\sqrt{r^4 + 2r^2 + 5})} > S(r). \quad (2.26)$$

The proof of Theorem 1.3 is complete. □

Proof of Theorem 1.5. Let $u_n(r) = n(n - 1) + r^2 + \mu(r)$ for $n \in \mathbb{N}$, where

$$\mu(r) = \sqrt{(r^2 + 1)^2 + 1} - (r^2 + 1) > 0. \quad (2.27)$$

Then,

$$\frac{1}{u_n(r)} - \frac{1}{u_{n+1}(r)} = \frac{2n}{(n^2 + r^2)^2 - [1 - 2\mu(r)]n^2 + \mu^2(r) + 2r^2\mu(r)}. \quad (2.28)$$

From (2.27), it is deduced that $\mu^2(r) + 2r^2\mu(r) = 1 - 2\mu(r) > 0$. Hence,

$$\frac{1}{u_n(r)} - \frac{1}{u_{n+1}(r)} = \frac{2n}{(n^2 + r^2)^2 - [1 - 2\mu(r)](n^2 - 1)} \geq \frac{2n}{(n^2 + r^2)^2}, \quad (2.29)$$

and then

$$\sum_{n=1}^{\infty} \frac{2n}{(n^2 + r^2)^2} < \frac{1}{u_1} = \frac{1}{r^2 + \mu(r)} = \frac{1}{\sqrt{r^4 + 2r^2 + 2} - 1}. \quad (2.30)$$

The proof of Theorem 1.5 is complete. \square

Proof of Theorem 1.7. Let

$$t_n = 2n^2 - 2n + 1 - \frac{1}{6(n^2 - n + \delta)}, \quad (2.31)$$

where δ is a fixed positive number and $n \in \mathbb{N}$. Direct computation gives

$$\frac{1}{t_n} - \frac{1}{t_{n+1}} = \frac{4n + 2n/6(n^2 - n + \delta)(n^2 + n + \delta)}{4n^4 + (2n^4 + (8\delta - 12)n^2 + 6\delta^2 - 2\delta + 1/6)/6(n^2 - n + \delta)(n^2 + n + \delta)}. \quad (2.32)$$

If $\delta = 3/2$, then $8\delta - 12 = 0$ and

$$\frac{1}{t_n} - \frac{1}{t_{n+1}} = \frac{4n + 2n/6(n^2 - n + 3/2)(n^2 + n + 3/2)}{4n^4 + (2n^4 + 32/3)/6(n^2 - n + 3/2)(n^2 + n + 3/2)} < \frac{1}{n^3}. \quad (2.33)$$

Summing up on both sides of the above inequality for n from $m + 1$ to infinity produces

$$\frac{1}{t_{m+1}} = \frac{1}{2m^2 + 1 - 1/6(m^2 + m + 3/2)} < \sum_{n=m+1}^{\infty} \frac{1}{n^3}. \quad (2.34)$$

Adding $S_3(m)$ on both sides of the above inequality leads to the left-hand side inequality in (1.7).

If $\delta = 1$ and $n > 1$, then

$$\frac{1}{t_n} - \frac{1}{t_{n+1}} = \frac{4n + 2n/6(n^2 - n + 1)(n^2 + n + 1)}{4n^4 + (2n^4 - [4(n^2 - 1) - 1/6])/6(n^2 - n + 1)(n^2 + n + 1)} > \frac{1}{n^3}. \quad (2.35)$$

Summing up on both sides of the above inequality for n from $m + 1$ to infinity yields that

$$\frac{1}{2m^2 + 2m + 1 - 1/2(m^2 + m + 1)} > \sum_{n=m+1}^{\infty} \frac{1}{n^3}. \quad (2.36)$$

This is equivalent to the right-hand side inequality in (1.7). Theorem 1.7 is proved. \square

Proof of Corollary 1.9. It is easy to see that

$$2n^4 + (8\delta - 12)n^2 + 6\delta^2 - 2\delta + \frac{1}{6} = 2n^4 - (12 - 8\delta) \left(n^2 - \frac{3\delta^2 - \delta + 1/12}{6 - 4\delta} \right). \quad (2.37)$$

If $1 \leq \delta < 3/2$ and $n \geq \sqrt{(3\delta^2 - \delta + 1/12)/(6 - 4\delta)}$, from (2.32), it is deduced that

$$\frac{1}{t_n} - \frac{1}{t_{n+1}} \geq \frac{1}{n^3}. \quad (2.38)$$

By the same argument as mentioned above, when $m \geq \sqrt{(3\delta^2 - \delta + 1/12)/(6 - 4\delta)} - 1$, inequality

$$\frac{1}{t_{m+1}} = \frac{1}{2m^2 + 2m + 1 - 1/6(m^2 + m + \delta)} > \sum_{n=m+1}^{\infty} \frac{1}{n^3} \quad (2.39)$$

is obtained, which is equivalent to (1.11). The proof of Corollary 1.9 is complete. \square

Acknowledgment

The authors appreciate heartily two anonymous referees for many kind and valuable comments on this paper.

References

- [1] E. Mathieu, *Traité de physique mathématique, VI-VII: Théorie de l'élasticité des corps solides*, Gauthier-Villars, Paris, France, 1890.
- [2] H. Alzer, J. L. Brenner, and O. G. Ruehr, "On Mathieu's inequality," *Journal of Mathematical Analysis and Applications*, vol. 218, no. 2, pp. 607–610, 1998.
- [3] L. Berg, "Über eine Abschätzung von Mathieu," *Mathematische Nachrichten*, vol. 7, no. 5, pp. 257–259, 1952.
- [4] P. H. Diananda, "On some inequalities related to Mathieu's," *Publikacije Elektrotehničkog Fakulteta Univerzitet u Beogradu*, vol. 716–734, pp. 22–24, 1981.
- [5] P. H. Diananda, "Some inequalities related to an inequality of Mathieu," *Mathematische Annalen*, vol. 250, no. 2, pp. 95–98, 1980.
- [6] Á. Elbert, "Asymptotic expansion and continued fraction for Mathieu's series," *Periodica Mathematica Hungarica*, vol. 13, no. 1, pp. 1–8, 1982.
- [7] O. Emersleben, "Über die Reihe $\sum_{k=1}^{\infty} k/(k^2 + c^2)^2$," *Mathematische Annalen*, vol. 125, no. 1, pp. 165–171, 1952.
- [8] A. Jakimovski and D. C. Russell, "Mercerian theorems involving Cesàro means of positive order," *Monatshefte für Mathematik*, vol. 96, no. 2, pp. 119–131, 1983.
- [9] E. Makai, "On the inequality of Mathieu," *Publicationes Mathematicae Debrecen*, vol. 5, pp. 204–205, 1957.
- [10] D. C. Russell, "A note on Mathieu's inequality," *Aequationes Mathematicae*, vol. 36, no. 2-3, pp. 294–302, 1988.
- [11] K. Schröder, "Das Problem der eingespannten rechteckigen elastischen Platte. I. Die biharmonische Randwertaufgabe für das Rechteck," *Mathematische Annalen*, vol. 121, pp. 247–326, 1949.
- [12] J. G. van der Corput and L. O. Heflinger, "On the inequality of Mathieu," *Indagationes Mathematicae*, vol. 18, pp. 15–20, 1956.
- [13] C. L. Wang and X. H. Wang, "Refinements of Mathieu's inequality," *Chinese Science Bulletin*, vol. 26, no. 5, p. 315, 1981 (Chinese).
- [14] C. L. Wang and X. H. Wang, "Refinements of the Mathieu inequality," *Journal of Mathematical Research & Exposition*, vol. 1, no. 1, pp. 107–112, 1981 (Chinese).
- [15] B.-N. Guo, "Note on Mathieu's inequality," *RGMIA Research Report Collection*, vol. 3, no. 3, article 5, pp. 389–392, 2000.

- [16] F. Qi, "Inequalities for Mathieu's series," *RGMI*A Research Report Collection, vol. 4, no. 2, article 3, pp. 187–193, 2001.
- [17] F. Qi, "Integral expression and inequalities of Mathieu type series," *RGMI*A Research Report Collection, vol. 6, no. 2, article 10, 2003.
- [18] P. Cerone, "Bounding Mathieu type series," *RGMI*A Research Report Collection, vol. 6, no. 3, article 7, 2003.
- [19] P. Cerone and C. T. Lenard, "On integral forms of generalised Mathieu series," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 5, article 100, pp. 1–11, 2003.
- [20] P. Cerone and C. T. Lenard, "On integral forms of generalised Mathieu series," *RGMI*A Research Report Collection, vol. 6, no. 2, article 19, 2003.
- [21] Ch.-P. Chen and F. Qi, "On an evaluation of upper bound of Mathieu series," *Studies in College Mathematics*, vol. 6, no. 1, pp. 48–49, 2003 (Chinese).
- [22] B. Drašćić and T. K. Pogány, "On integral representation of Bessel function of the first kind," *Journal of Mathematical Analysis and Applications*, vol. 308, no. 2, pp. 775–780, 2005.
- [23] B. Drašćić and T. K. Pogány, "On integral representation of first kind Bessel function," *RGMI*A Research Report Collection, vol. 7, no. 3, article 18, 2004.
- [24] B. Drašćić and T. K. Pogány, "Testing Alzer's inequality for Mathieu series $S(r)$," *Mathematica Macedonica*, vol. 2, pp. 1–4, 2004.
- [25] I. Gavrea, "Some remarks on Mathieu's series," in *Mathematical Analysis and Approximation Theory*, pp. 113–117, Burg, Sibiu, Romania, 2002.
- [26] A.-Q. Liu, T.-F. Hu, and W. Li, "Notes on Mathieu's series," *Journal of Jiaozuo Institute of Technology*, vol. 20, no. 4, pp. 302–304, 2001 (Chinese).
- [27] T. K. Pogány, "Integral representation of a series which includes the Mathieu \mathbf{a} -series," *Journal of Mathematical Analysis and Applications*, vol. 296, no. 1, pp. 309–313, 2004.
- [28] T. K. Pogány, "Integral representation of Mathieu (\mathbf{a}, λ) -series," *Integral Transforms and Special Functions*, vol. 16, no. 8, pp. 685–689, 2005.
- [29] T. K. Pogány, "Integral representation of Mathieu (\mathbf{a}, λ) -series," *RGMI*A Research Report Collection, vol. 7, no. 1, article 9, 2004.
- [30] T. K. Pogány, H. M. Srivastava, and Ž. Tomovski, "Some families of Mathieu \mathbf{a} -series and alternating Mathieu \mathbf{a} -series," *Applied Mathematics and Computation*, vol. 173, no. 1, pp. 69–108, 2006.
- [31] T. K. Pogány and Ž. Tomovski, "On multiple generalized Mathieu series," *Integral Transforms and Special Functions*, vol. 17, no. 4, pp. 285–293, 2006.
- [32] F. Qi, "An integral expression and some inequalities of Mathieu type series," *Rostocker Mathematisches Kolloquium*, no. 58, pp. 37–46, 2004.
- [33] F. Qi, Ch.-P. Chen, and B.-N. Guo, "Notes on double inequalities of Mathieu's series," *International Journal of Mathematics and Mathematical Sciences*, vol. 2005, no. 16, pp. 2547–2554, 2005.
- [34] H. M. Srivastava and Ž. Tomovski, "Some problems and solutions involving Mathieu's series and its generalizations," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 5, no. 2, article 45, pp. 1–13, 2004.
- [35] Ž. Tomovski, "New double inequalities for Mathieu type series," *Publikacije Elektrotehničkog Fakulteta Univerzitet u Beogradu*, vol. 15, pp. 80–84, 2004.
- [36] Ž. Tomovski, "New double inequalities for Mathieu type series," *RGMI*A Research Report Collection, vol. 6, no. 2, article 17, 2003.
- [37] Ž. Tomovski and K. Trenčevski, "On an open problem of Bai-Ni Guo and Feng Qi," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 2, article 29, pp. 1–7, 2003.

- [38] A. Hoorfar and F. Qi, "Some new bounds for Mathieu's series," *RGMI*A Research Report Collection, vol. 10, no. 1, article 12, 2007.
- [39] Q.-M. Luo, Z.-L. Wei, and F. Qi, "Lower and upper bounds of $\zeta(3)$," *Advanced Studies in Contemporary Mathematics*, vol. 6, no. 1, pp. 47–51, 2003.
- [40] Q.-M. Luo, Z.-L. Wei, and F. Qi, "Lower and upper bounds of $\zeta(3)$," *RGMI*A Research Report Collection, vol. 4, no. 4, article 7, pp. 565–569, 2003.
- [41] Q.-M. Luo, B.-N. Guo, and F. Qi, "On evaluation of Riemann zeta function $\zeta(s)$," *Advanced Studies in Contemporary Mathematics*, vol. 7, no. 2, pp. 135–144, 2003.
- [42] Q.-M. Luo, B.-N. Guo, and F. Qi, "On evaluation of Riemann zeta function $\zeta(s)$," *RGMI*A Research Report Collection, vol. 6, no. 1, article 8, 2003.

Abdolhossein Hoorfar: Department of Irrigation Engineering, College of Agriculture and Natural Resources, University of Tehran, Karaj 31587-77871, Iran
Email address: hoorfar@ut.ac.ir

Feng Qi: Research Institute of Mathematical Inequality Theory, Henan Polytechnic University, Jiaozuo City, Henan Province 454010, China
Email addresses: qifeng618@gmail.com; qifeng618@hotmail.com; qifeng618@qq.com