

## Research Article

# Weighted Composition Operators from $F(p, q, s)$ Spaces to $H_\mu^\infty$ Spaces

**Xiangling Zhu**

*Department of Mathematics, JiaYing University, Meizhou, GuangDong 514015, China*

Correspondence should be addressed to Xiangling Zhu, jyuzxl@163.com

Received 25 January 2009; Accepted 5 February 2009

Recommended by Stevo Stevic

Let  $H(B)$  denote the space of all holomorphic functions on the unit ball  $B$ . Let  $u \in H(B)$  and  $\varphi$  be a holomorphic self-map of  $B$ . In this paper, we investigate the boundedness and compactness of the weighted composition operator  $uC_\varphi$  from the general function space  $F(p, q, s)$  to the weighted-type space  $H_\mu^\infty$  in the unit ball.

Copyright © 2009 Xiangling Zhu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

Let  $B$  be the unit ball of  $\mathbb{C}^n$ . Let  $z = (z_1, \dots, z_n)$  and let  $w = (w_1, \dots, w_n)$  be points in  $\mathbb{C}^n$ , we write

$$|z| = \sqrt{|z_1|^2 + \dots + |z_n|^2}, \quad \langle z, w \rangle = z_1 \bar{w}_1 + \dots + z_n \bar{w}_n. \quad (1.1)$$

Thus  $B = \{z \in \mathbb{C}^n : |z| < 1\}$ . Let  $dv$  be the normalized Lebesgue measure of  $B$ , that is,  $v(B) = 1$ . Let  $H(B)$  be the space of all holomorphic functions on  $B$ . For  $f \in H(B)$ , let

$$\Re f(z) = \sum_{j=1}^n z_j \frac{\partial f}{\partial z_j}(z) \quad (1.2)$$

represent the radial derivative of  $f \in H(B)$ . For  $a, z \in B$ ,  $a \neq 0$ , let  $\varphi_a$  denote the Möbius transformation of  $B$  taking 0 to  $a$ , which is defined by

$$\varphi_a(z) = \frac{a - P_a(z) - \sqrt{1 - |z|^2} Q_a(z)}{1 - \langle z, a \rangle}, \quad (1.3)$$

where  $P_a(z)$  is the orthogonal projection of  $z$  onto the one dimensional subspace of  $\mathbb{C}^n$  spanned by  $a$ , and  $Q_a(z) = z - P_a(z)$ .

A positive continuous function  $\mu$  on  $[0, 1)$  is called normal, if there exist positive numbers  $s$  and  $t$ ,  $0 < s < t$ , and  $\delta \in [0, 1)$  such that

$$\begin{aligned} \frac{\mu(r)}{(1-r)^s} \text{ is decreasing on } [\delta, 1), \quad \lim_{r \rightarrow 1} \frac{\mu(r)}{(1-r)^s} = 0; \\ \frac{\mu(r)}{(1-r)^t} \text{ is increasing on } [\delta, 1), \quad \lim_{r \rightarrow 1} \frac{\mu(r)}{(1-r)^t} = \infty \end{aligned} \quad (1.4)$$

(see [1]).

Let  $\mu$  be a normal function on  $[0, 1)$ . An  $f \in H(B)$  is said to belong to the weighted-type space  $H_\mu^\infty = H_\mu^\infty(B)$ , if

$$\|f\|_{H_\mu^\infty} = \sup_{z \in B} \mu(|z|)|f(z)| < \infty, \quad (1.5)$$

where  $\mu$  is normal on  $[0, 1)$  (see, e.g., [2–4]).  $H_\mu^\infty$  is a Banach space with the norm  $\|\cdot\|_{H_\mu^\infty}$ . We denote by  $H_{\mu,0}^\infty$  the subspace of  $H_\mu^\infty$  consisting of those  $f \in H_\mu^\infty$  such that

$$\lim_{|z| \rightarrow 1} \mu(|z|)|f(z)| = 0. \quad (1.6)$$

When  $\mu(r) = (1-r^2)^\alpha$ , the induced spaces  $H_\mu^\infty$  and  $H_{\mu,0}^\infty$  become the (classical) weighted space  $H_\alpha^\infty$  and  $H_{\alpha,0}^\infty$ , respectively.

For  $\alpha > 0$ , recall that the  $\alpha$ -Bloch space  $\mathcal{B}^\alpha = \mathcal{B}^\alpha(B)$  is the space of all  $f \in H(B)$  for which (see [5])

$$b_\alpha(f) = \sup_{z \in B} (1-|z|^2)^\alpha |\Re f(z)| < \infty. \quad (1.7)$$

Under the norm  $\|f\|_{\mathcal{B}^\alpha} = |f(0)| + b_\alpha(f)$ ,  $\mathcal{B}^\alpha$  is a Banach space. When  $\alpha = 1$ , we get the classical Bloch space  $\mathcal{B}$ . For more information of the Bloch space and the  $\alpha$ -Bloch space (see, e.g., [5–8] and the references therein).

For  $p \in (0, \infty)$ , the weighted Bergman space  $A^p(B)$  is the space of all holomorphic functions  $f$  on  $B$  for which

$$\|f\|_{A^p}^p = \int_B |f(z)|^p d\nu(z) < \infty. \quad (1.8)$$

The Hardy space  $H^p(B)$  ( $0 < p < \infty$ ) on the unit ball is defined by

$$H^p(B) = \left\{ f \mid f \in H(B), \|f\|_{H^p(B)} = \sup_{0 \leq r < 1} M_p(f, r) < \infty \right\}, \quad (1.9)$$

where

$$M_p(f, r) = \left( \int_{\partial B} |f(r\zeta)|^p d\sigma(\zeta) \right)^{1/p} \tag{1.10}$$

and  $d\sigma$  is the normalized surface measure on  $\partial B$ .

For  $0 < p < \infty$ , the  $Q_p$  space is defined by (see [9])

$$Q_p(B) = \left\{ f \in H(B) : \|f\|_{Q_p}^2 = \sup_{a \in B} \int_B |\tilde{\nabla} f(z)|^2 G^p(z, a) \frac{dv(z)}{(1 - |z|^2)^{n+1}} < \infty \right\}. \tag{1.11}$$

Here  $\tilde{\nabla} f(z) = \nabla(f \circ \varphi_z)(0)$  denotes the invariant gradient of  $f$ , and  $G(z, a)$  is the invariant Green function defined by  $G(z, a) = g(\varphi_a(z))$ , where

$$g(z) = \frac{n+1}{2n} \int_{|z|}^1 (1-t^2)^{n-1} t^{-2n+1} dt. \tag{1.12}$$

Let  $0 < p, s < \infty, -n-1 < q < \infty$ . A function  $f \in H(B)$  is said to belong to  $F(p, q, s) = F(p, q, s)(B)$  (see [10–12]) if

$$\|f\|_{F(p,q,s)}^p = |f(0)|^p + \sup_{a \in B} \int_B |\nabla f(z)|^p (1 - |z|^2)^q G^s(z, a) dv(z) < \infty. \tag{1.13}$$

$F(p, q, s)$  is called the general function space since we can get many function spaces, such as Hardy space, Bergman space, Bloch space,  $Q_p$  space, if we take special parameters of  $p, q, s$ . For example,  $F(2, 1, 0) = H^2$ ,  $F(p, p, 0) = A^p$ , and  $F(2, 0, s) = Q_s$ . If  $q + s \leq -1$ , then  $F(p, q, s)$  is the space of constant functions. For the setting of the unit disk, see [13].

Let  $u \in H(B)$  and  $\varphi$  be a holomorphic self-map of  $B$ . For  $f \in H(B)$ , the weighted composition operator  $uC_\varphi$  is defined by

$$(uC_\varphi f)(z) = u(z)f(\varphi(z)), \quad z \in B. \tag{1.14}$$

The weighted composition operator is the generalization of a multiplication operator and a composition operator, which is defined by  $(C_\varphi f)(z) = f(\varphi(z))$ . The main subject in the study of composition operators is to describe operator theoretic properties of  $C_\varphi$  in terms of function theoretic properties of  $\varphi$ . The book [14] is a good reference for the theory of composition operators. Recall that a linear operator is said to be bounded if the image of a bounded set is a bounded set, while a linear operator is compact if it takes bounded sets to sets with compact closure.

In the setting of the unit ball, we studied the boundedness and compactness of the weighted composition operator between Bergman-type spaces and  $H^\infty$  in [15]. More general results can be found in [16]. Some necessary and sufficient conditions for the weighted composition operator to be bounded or compact between the Bloch space and  $H^\infty$  are given in [17]. In the setting of the unit polydisk  $D^n$ , some necessary and sufficient conditions for a weighted composition operator to be bounded or compact between the Bloch space and

$H^\infty(D^n)$  are given in [18, 19] (see, also [20] for the case of composition operators). Some related results can be found, for example, in [2, 3, 6, 21–31].

In the present paper, we are mainly concerned about the boundedness and compactness of the weighted composition operator from  $F(p, q, s)$  to the space  $H_\mu^\infty$ . Some necessary and sufficient conditions for the weighted composition operator  $uC_\varphi$  to be bounded and compact are given.

Constants are denoted by  $C$  in this paper, they are positive and may differ from one occurrence to the other.  $a \leq b$  means that there is a positive constant  $C$  such that  $a \leq Cb$ . Moreover, if both  $a \leq b$  and  $b \leq a$  hold, then one says that  $a \asymp b$ .

## 2. Main Results and Proofs

In order to prove our results, we need some auxiliary results which are incorporated in the following lemmas.

**Lemma 2.1** (see [12]). *For  $0 < p, s < \infty, -n - 1 < q < \infty, q + s > -1$ , if  $f \in F(p, q, s)$ , then  $f \in \mathcal{B}^{(n+1+q)/p}$  and*

$$\|f\|_{\mathcal{B}^{(n+1+q)/p}} \leq C\|f\|_{F(p,q,s)}. \quad (2.1)$$

The following lemma can be found, for example, in [32].

**Lemma 2.2.** *If  $f \in \mathcal{B}^\alpha$ , then*

$$|f(z)| \leq C \begin{cases} \|f\|_{\mathcal{B}^\alpha}, & 0 < \alpha < 1, \\ \|f\|_{\mathcal{B}^\alpha} \ln \frac{e}{1-|z|^2}, & \alpha = 1, \\ \frac{\|f\|_{\mathcal{B}^\alpha}}{(1-|z|^2)^{\alpha-1}}, & \alpha > 1, \end{cases} \quad (2.2)$$

for some  $C$  independent of  $f$ .

**Lemma 2.3.** *Assume that  $\mu$  is normal. A closed set  $K$  in  $H_{\mu,0}^\infty$  is compact if and only if it is bounded and satisfies*

$$\lim_{|z| \rightarrow 1} \sup_{f \in K} \mu(|z|)|f(z)| = 0. \quad (2.3)$$

The proof of Lemma 2.3 is similar to the proof of Lemma 1 of [33]. We omit the details.

**Lemma 2.4.** *Assume that  $u \in H(B)$ ,  $\varphi$  is a holomorphic self-map of  $B$ ,  $\mu$  is a normal function on  $[0, 1)$ ,  $0 < p, s < \infty, -n - 1 < q < \infty, q + s > -1$ . Then  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is compact if and only if  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is bounded and for any bounded sequence  $(f_k)_{k \in \mathbb{N}}$  in  $F(p, q, s)$  which converges to zero uniformly on compact subsets of  $B$  as  $k \rightarrow \infty$ , one has  $\|uC_\varphi f_k\|_{H_\mu^\infty} \rightarrow 0$  as  $k \rightarrow \infty$ .*

Lemma 2.4 follows by standard arguments similar to those outlined in [14, Proposition 3.11] (see, also, the corresponding lemmas in [20, 34, 35]). We omit the details.

Note that when  $p > n + 1 + q$ , the function in  $F(p, q, s)$  is indeed Lipschitz continuous by Lemmas 2.1 and 2.2. By Arzela-Ascoli theorem, similarly to the proof of Lemma 3.6 of [26], we have the following result.

**Lemma 2.5.** *Let  $0 < p, s < \infty, -n - 1 < q < \infty, q + s > -1$ , and  $p > n + 1 + q$ . Let  $(f_k)$  be a bounded sequence in  $F(p, q, s)$  which converges to 0 uniformly on compact subsets of  $B$ , then*

$$\lim_{k \rightarrow \infty} \sup_{z \in B} |f_k(z)| = 0. \quad (2.4)$$

### 2.1. Case $p < n + 1 + q$

In this subsection, we consider the case  $p < n + 1 + q$ . Our first result is the following theorem.

**Theorem 2.6.** *Assume that  $u \in H(B)$ ,  $\varphi$  is a holomorphic self-map of  $B$ ,  $\mu$  is a normal function on  $[0, 1)$ ,  $0 < p, s < \infty, -n - 1 < q < \infty, q + s > -1, p < n + 1 + q$ . Then  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is bounded if and only if*

$$\sup_{z \in B} \frac{\mu(|z|)|u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}} < \infty. \quad (2.5)$$

Moreover, when  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is bounded, the following relationship

$$\|uC_\varphi\|_{F(p,q,s) \rightarrow H_\mu^\infty} \asymp \sup_{z \in B} \frac{\mu(|z|)|u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}} \quad (2.6)$$

holds.

*Proof.* Assume that (2.5) holds. For any  $f \in F(p, q, s)$ , by Lemmas 2.1 and 2.2,

$$\begin{aligned} \|uC_\varphi f\|_{H_\mu^\infty} &= \sup_{z \in B} \mu(|z|) |(uC_\varphi f)(z)| \\ &= \sup_{z \in B} \mu(|z|) |f(\varphi(z))| |u(z)| \\ &\leq C \|f\|_{\mathcal{B}^{(n+1+q)/p}} \sup_{z \in B} \frac{\mu(|z|)|u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}} \\ &\leq C \|f\|_{F(p,q,s)} \sup_{z \in B} \frac{\mu(|z|)|u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}}. \end{aligned} \quad (2.7)$$

Therefore, (2.5) implies that  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is bounded.

Conversely, suppose that  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is bounded. For  $b \in B$ , let

$$f_b(z) = \frac{1 - |b|^2}{(1 - \langle z, b \rangle)^{(n+1+q)/p}}. \quad (2.8)$$

It is easy to see that

$$f_{\varphi(w)}(\varphi(w)) = \frac{1}{(1 - |\varphi(w)|^2)^{(n+1+q)/p-1}}, \quad |\Re f_{\varphi(w)}(\varphi(w))| = \frac{(n+1+q)|\varphi(w)|^2}{p(1 - |\varphi(w)|^2)^{(n+1+q)/p}}. \quad (2.9)$$

If  $\varphi(w) = 0$ , then  $f_{\varphi(w)} \equiv 1$  obviously belongs to  $F(p, q, s)$ . From [12], we know that  $f_b \in F(p, q, s)$ ; moreover, there is a positive constant  $K$  such that  $\sup_{b \in B} \|f_b\|_{F(p, q, s)} \leq K$ . Therefore,

$$\begin{aligned} \sup_{z \in B} \mu(|z|) |f_{\varphi(w)}(\varphi(z))u(z)| &= \sup_{z \in B} \mu(|z|) |(uC_\varphi f_{\varphi(w)})(z)| \\ &= \|uC_\varphi f_{\varphi(w)}\|_{H_\mu^\infty} \leq K \|uC_\varphi\|_{F(p, q, s) \rightarrow H_\mu^\infty}, \end{aligned} \quad (2.10)$$

for every  $w \in B$ , from which we get

$$\sup_{w \in B} \frac{\mu(|w|)|u(w)|}{(1 - |\varphi(w)|^2)^{(n+1+q)/p-1}} \leq K \|uC_\varphi\|_{F(p, q, s) \rightarrow H_\mu^\infty} < \infty, \quad (2.11)$$

that is, (2.5) follows. From (2.7) and (2.11), we see that (2.6) holds. The proof of this theorem is finished.  $\square$

**Theorem 2.7.** *Assume that  $u \in H(B)$ ,  $\varphi$  is a holomorphic self-map of  $B$ ,  $\mu$  is a normal function on  $[0, 1)$ ,  $0 < p, s < \infty$ ,  $-n - 1 < q < \infty$ ,  $q + s > -1$ ,  $p < n + 1 + q$ . Then  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is compact if and only if  $u \in H_\mu^\infty$  and*

$$\lim_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}} = 0. \quad (2.12)$$

*Proof.* First assume that  $u \in H_\mu^\infty$  and the condition in (2.12) hold. In order to prove that  $uC_\varphi$  is compact, according to Lemma 2.4 it suffices to show that if  $(f_k)_{k \in \mathbb{N}}$  is bounded in  $F(p, q, s)$  and converges to 0 uniformly on compact subsets of  $B$  as  $k \rightarrow \infty$ , then  $\|uC_\varphi f_k\|_{H_\mu^\infty} \rightarrow 0$  as  $k \rightarrow \infty$ .

Now assume that  $(f_k)_{k \in \mathbb{N}}$  is a sequence in  $F(p, q, s)$  such that  $\sup_{k \in \mathbb{N}} \|f_k\|_{F(p, q, s)} \leq L$  and  $f_k \rightarrow 0$  uniformly on compact subsets of  $B$  as  $k \rightarrow \infty$ . From (2.12), we have that for every  $\varepsilon > 0$ , there is a constant  $\delta \in (0, 1)$  such that

$$\frac{\mu(|z|)|u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}} < \varepsilon, \quad (2.13)$$

when  $\delta < |\varphi(z)| < 1$ . By Lemmas 2.1 and 2.2, we have

$$\begin{aligned}
\|uC_\varphi f_k\|_{H_\mu^\infty} &= \sup_{z \in B} \mu(|z|) |(uC_\varphi f_k)(z)| \\
&= \sup_{z \in B} \mu(|z|) |u(z)| |f_k(\varphi(z))| \\
&\leq \sup_{\varphi(z) \in \overline{B(0, \delta)}} \mu(|z|) |u(z)| |f_k(\varphi(z))| \\
&\quad + C \sup_{\varphi(z) \in B \setminus \overline{B(0, \delta)}} \frac{\mu(|z|) |u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}} \|f_k\|_{F(p, q, s)} \\
&\leq M_1 \sup_{\varphi(z) \in \overline{B(0, \delta)}} |f_k(\varphi(z))| + CL\varepsilon,
\end{aligned} \tag{2.14}$$

where  $M_1 := \sup_{z \in B} \mu(|z|) |u(z)| < \infty$ . Using the fact that  $f_k \rightarrow 0$  uniformly on compact subsets of  $B$  as  $k \rightarrow \infty$ , we obtain

$$M_1 \limsup_{k \rightarrow \infty} \sup_{\varphi(z) \in \overline{B(0, \delta)}} |f_k(\varphi(z))| = 0. \tag{2.15}$$

Therefore,

$$\limsup_{k \rightarrow \infty} \|uC_\varphi f_k\|_{H_\mu^\infty} \leq CL\varepsilon. \tag{2.16}$$

Since  $\varepsilon$  is an arbitrary positive number, we have that  $\lim_{k \rightarrow \infty} \|uC_\varphi f_k\|_{H_\mu^\infty} = 0$ , and, therefore,  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is compact by Lemma 2.4.

Conversely, suppose  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is compact. Let  $(z_k)_{k \in \mathbb{N}}$  be a sequence in  $B$  such that  $|\varphi(z_k)| \rightarrow 1$  as  $k \rightarrow \infty$  (if such a sequence does not exist that condition (2.12) is vacuously satisfied). Set

$$f_k(z) = \frac{1 - |\varphi(z_k)|^2}{(1 - \langle z, \varphi(z_k) \rangle)^{(q+n+1)/p}}, \quad k \in \mathbb{N}. \tag{2.17}$$

From the proof of Theorem 2.6, we see that  $f_k \in F(p, q, s)$  for every  $k \in \mathbb{N}$ ; moreover,  $\sup_{k \in \mathbb{N}} \|f_k\|_{F(p, q, s)} \leq C$ . Beside this,  $f_k$  converges to 0 uniformly on compact subsets of  $B$  as  $k \rightarrow \infty$ . Since  $uC_\varphi$  is compact, by Lemma 2.4 we have that  $\|uC_\varphi f_k\|_{H_\mu^\infty} \rightarrow 0$  as  $k \rightarrow \infty$ . Thus

$$\begin{aligned}
(1 - |\varphi(z_k)|^2)^{1-(q+n+1)/p} \mu(|z_k|) |u(z_k)| &= \mu(|z_k|) |u(z_k)| |f_k(\varphi(z_k))| \\
&\leq \sup_{z \in B} \mu(|z|) |f_k(\varphi(z))| |u(z)| \\
&= \sup_{z \in B} \mu(|z|) |(uC_\varphi f_k)(z)| \\
&= \|uC_\varphi f_k\|_{H_\mu^\infty} \rightarrow 0,
\end{aligned} \tag{2.18}$$

as  $k \rightarrow \infty$ , from which we obtain (2.12), finishing the proof of the theorem.  $\square$

**Theorem 2.8.** Assume that  $u \in H(B)$ ,  $\varphi$  is a holomorphic self-map of  $B$ ,  $\mu$  is a normal function on  $[0, 1)$ ,  $0 < p$ ,  $s < \infty$ ,  $-n - 1 < q < \infty$ ,  $q + s > -1$ ,  $p < n + 1 + q$ . Then  $uC_\varphi : F(p, q, s) \rightarrow H_{\mu,0}^\infty$  is compact if and only if

$$\lim_{|z| \rightarrow 1} \frac{\mu(|z|)|u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}} = 0. \quad (2.19)$$

*Proof.* Suppose that (2.19) holds. From Lemma 2.3, we see that  $uC_\varphi : F(p, q, s) \rightarrow H_{\mu,0}^\infty$  is compact if and only if

$$\lim_{|z| \rightarrow 1} \sup_{\|f\|_{F(p,q,s)} \leq 1} \mu(|z|)|(uC_\varphi f)(z)| = 0. \quad (2.20)$$

On the other hand, by Lemmas 2.1 and 2.2, we have that

$$\mu(|z|)|(uC_\varphi f)(z)| \leq C\|f\|_{F(p,q,s)} \frac{\mu(|z|)|u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}}. \quad (2.21)$$

Taking the supremum in (2.21) over the the unit ball in the space  $F(p, q, s)$ , then letting  $|z| \rightarrow 1$  and applying (2.19) the result follows.

Conversely, suppose that  $uC_\varphi : F(p, q, s) \rightarrow H_{\mu,0}^\infty$  is compact. Then  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is compact and hence by Theorem 2.7,

$$\lim_{|\varphi(z)| \rightarrow 1} \frac{\mu(|z|)|u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}} = 0. \quad (2.22)$$

In addition,  $uC_\varphi : F(p, q, s) \rightarrow H_{\mu,0}^\infty$  is bounded. Taking  $f(z) = 1$ , then employing the boundedness of  $uC_\varphi : F(p, q, s) \rightarrow H_{\mu,0}^\infty$ , we get

$$\lim_{|z| \rightarrow 1} \mu(|z|)|u(z)| = 0. \quad (2.23)$$

By (2.22), for every  $\varepsilon > 0$ , there exists a  $\delta \in (0, 1)$ ,

$$\frac{\mu(|z|)|u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}} < \varepsilon, \quad (2.24)$$

when  $\delta < |\varphi(z)| < 1$ . By (2.23), for above chosen  $\varepsilon$ , there exists  $r \in (0, 1)$ ,

$$\mu(|z|)|u(z)| \leq \varepsilon(1 - \delta^2)^{(n+1+q)/p-1}, \quad (2.25)$$

when  $r < |z| < 1$ .

Therefore, when  $r < |z| < 1$  and  $\delta < |\varphi(z)| < 1$ , we have that

$$\frac{\mu(|z|)|u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}} < \varepsilon. \tag{2.26}$$

If  $|\varphi(z)| \leq \delta$  and  $r < |z| < 1$ , we obtain

$$\frac{\mu(|z|)|u(z)|}{(1 - |\varphi(z)|^2)^{(n+1+q)/p-1}} \leq \frac{1}{(1 - \delta^2)^{(n+1+q)/p-1}} \mu(|z|)|u(z)| < \varepsilon. \tag{2.27}$$

Combing (2.26) with (2.27), we get (2.19), as desired. □

### 2.2. Case $p = n + 1 + q$

**Theorem 2.9.** *Assume that  $u \in H(B)$ ,  $\varphi$  is a holomorphic self-map of  $B$ ,  $\mu$  is a normal function on  $[0, 1)$ ,  $0 < p, s < \infty, -n - 1 < q < \infty, q + s > -1, p = n + 1 + q$ . Then  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is bounded if and only if*

$$\sup_{z \in B} \mu(|z|)|u(z)| \ln \frac{e}{1 - |\varphi(z)|^2} < \infty. \tag{2.28}$$

Moreover, the following relationship

$$\|uC_\varphi\|_{F(p,q,s) \rightarrow H_\mu^\infty} \asymp \sup_{z \in B} \mu(|z|)|u(z)| \ln \frac{e}{1 - |\varphi(z)|^2} \tag{2.29}$$

holds.

*Proof.* Suppose that (2.28) holds. For any  $f \in F(p, q, s) \subseteq \mathcal{B}$ , by Lemmas 2.1 and 2.2, we have

$$\begin{aligned} \|uC_\varphi f\|_{H_\mu^\infty} &= \sup_{z \in B} \mu(|z|)|(uC_\varphi f)(z)| \\ &= \sup_{z \in B} \mu(|z|)|f(\varphi(z))||u(z)| \\ &\leq C \|f\|_{F(p,q,s)} \sup_{z \in B} \mu(|z|)|u(z)| \ln \frac{e}{1 - |\varphi(z)|^2}. \end{aligned} \tag{2.30}$$

Therefore, (2.28) implies that  $uC_\varphi$  is a bounded operator from  $F(p, q, s)$  to  $H_\mu^\infty$ .

Conversely, suppose that  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is bounded. For  $b \in B$ , let

$$f_b(z) = \ln \frac{e}{1 - \langle z, b \rangle}, \quad z \in B. \tag{2.31}$$

Then by [12] we see that  $f_b \in F(p, q, s)$ ; moreover, there is a positive constant  $K$  such that  $\sup_{b \in B} \|f_b\|_{F(p, q, s)} \leq K$ . Hence

$$\mu(|w|)|u(w)| \ln \frac{e}{1 - |\varphi(w)|^2} = \mu(|w|)|f_{\varphi(w)}(\varphi(w))||u(w)| \leq \|uC_\varphi f_{\varphi(w)}\|_{H_\mu^\infty}, \quad (2.32)$$

for every  $w \in B$ , that is, we get

$$\begin{aligned} \sup_{w \in B} \mu(|w|)|u(w)| \ln \frac{e}{1 - |\varphi(w)|^2} &\leq C \|uC_\varphi f_{\varphi(w)}\|_{H_\mu^\infty} \\ &\leq CK \|uC_\varphi\|_{F(p, q, s) \rightarrow H_\mu^\infty} < \infty. \end{aligned} \quad (2.33)$$

From (2.30) and (2.33), (2.29) follows. The proof is finished.  $\square$

**Theorem 2.10.** *Assume that  $u \in H(B)$ ,  $\varphi$  is a holomorphic self-map of  $B$ ,  $\mu$  is a normal function on  $[0, 1)$ ,  $0 < p$ ,  $s < \infty$ ,  $-n - 1 < q < \infty$ ,  $q + s > -1$ ,  $p = n + 1 + q$ . Then  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is compact if and only if  $u \in H_\mu^\infty$  and*

$$\lim_{|\varphi(z)| \rightarrow 1} \mu(|z|)|u(z)| \ln \frac{e}{1 - |\varphi(z)|^2} = 0. \quad (2.34)$$

*Proof.* Assume that  $u \in H_\mu^\infty$  and (2.34) hold, and that  $(f_k)_{k \in \mathbb{N}}$  is bounded in  $F(p, q, s)$  and converges to 0 uniformly on compact subsets of  $B$  as  $k \rightarrow \infty$ . We have that, for every  $\varepsilon > 0$ , there is a  $\delta \in (0, 1)$  such that

$$\mu(|z|)|u(z)| \ln \frac{e}{1 - |\varphi(z)|^2} < \varepsilon, \quad (2.35)$$

when  $\delta < |\varphi(z)| < 1$ .

In addition

$$\begin{aligned} \|uC_\varphi f_k\|_{H_\mu^\infty} &= \sup_{z \in B} \mu(|z|)|(uC_\varphi f_k)(z)| \\ &= \sup_{z \in B} \mu(|z|)|u(z)||f_k(\varphi(z))| \\ &\leq \sup_{\varphi(z) \in \overline{B(0, \delta)}} \mu(|z|)|u(z)||f_k(\varphi(z))| \\ &\quad + C \|f_k\|_{F(p, q, s)} \sup_{\varphi(z) \in B \setminus \overline{B(0, \delta)}} \mu(|z|)|u(z)| \ln \frac{e}{1 - |\varphi(z)|^2} \\ &\leq C \sup_{\varphi(z) \in \overline{B(0, \delta)}} |f_k(\varphi(z))| + C\varepsilon. \end{aligned} \quad (2.36)$$

Similar to the proof of Theorem 2.7, we obtain  $\|uC_\varphi f_k\|_{H_\mu^\infty} \rightarrow 0$  as  $k \rightarrow \infty$ . Therefore,  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is compact by Lemma 2.4.

Conversely, suppose that  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is compact. Assume that  $(z_k)_{k \in \mathbb{N}}$  is a sequence in  $B$  such that  $|\varphi(z_k)| \rightarrow 1$  as  $k \rightarrow \infty$  (if such a sequence does not exist that condition (2.34) is vacuously satisfied). Set

$$f_k(z) = \left( \ln \frac{e}{1 - |\varphi(z_k)|^2} \right)^{-1} \left( \ln \frac{e}{1 - \langle z, \varphi(z_k) \rangle} \right)^2, \quad k \in \mathbb{N}. \quad (2.37)$$

After some calculations or from [12], we see that  $\sup_{k \in \mathbb{N}} \|f_k\|_{F(p,q,s)} \leq C$  for some positive  $C$  independent of  $k$ , and  $f_k$  converges to 0 uniformly on compact subsets of  $B$  as  $k \rightarrow \infty$ . Since  $uC_\varphi$  is compact, by Lemma 2.4, we have  $\|uC_\varphi f_k\|_{H_\mu^\infty} \rightarrow 0$  as  $k \rightarrow \infty$ . Thus

$$\begin{aligned} \mu(|z_k|)|u(z_k)| \ln \frac{e}{1 - |\varphi(z_k)|^2} &\leq \sup_{z \in B} \mu(|z|)|f_k(\varphi(z_k))||u(z_k)| \\ &= \sup_{z \in B} \mu(|z|)|(uC_\varphi f_k)(z)| \\ &= \|uC_\varphi f_k\|_{H_\mu^\infty} \rightarrow 0, \end{aligned} \quad (2.38)$$

as  $k \rightarrow \infty$ , which is equivalent to (2.34). The proof of this theorem is finished.  $\square$

Similarly to the proof of Theorem 2.8, we can obtain the following results. We omit the details of the proof.

**Theorem 2.11.** *Assume that  $u \in H(B)$ ,  $\varphi$  is a holomorphic self-map of  $B$ ,  $\mu$  is a normal function on  $[0, 1)$ ,  $0 < p$ ,  $s < \infty$ ,  $-n - 1 < q < \infty$ ,  $q + s > -1$ ,  $p = n + 1 + q$ . Then  $uC_\varphi : F(p, q, s) \rightarrow H_{\mu,0}^\infty$  is compact if and only if*

$$\lim_{|z| \rightarrow 1} \mu(|z|)|u(z)| \ln \frac{e}{1 - |\varphi(z)|^2} < \infty. \quad (2.39)$$

### 2.3. Case $p > n + 1 + q$

**Theorem 2.12.** *Assume that  $u \in H(B)$ ,  $\varphi$  is a holomorphic self-map of  $B$ ,  $\mu$  is a normal function on  $[0, 1)$ ,  $0 < p$ ,  $s < \infty$ ,  $-n - 1 < q < \infty$ ,  $q + s > -1$ ,  $p > n + 1 + q$ . Then the following statements are equivalent:*

- (i)  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is bounded;
- (ii)  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is compact;
- (iii)  $u \in H_\mu^\infty$ .

*Proof.* (ii) $\Rightarrow$ (i) This implication is obvious.

(i) $\Rightarrow$ (iii) Taking  $f(z) = 1$ , then using the boundedness of  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  the implication follows.

(iii) $\Rightarrow$ (ii) Suppose that  $u \in H_\mu^\infty$ . For an  $f \in F(p, q, s)$ , by Lemma 2.1, we see that  $f$  is continuous on the closed unit ball and so is bounded in  $B$ . Therefore,

$$\mu(|z|)|(uC_\varphi f)(z)| = \mu(|z|)|f(\varphi(z))||u(z)| \leq C\|f\|_{F(p,q,s)}\mu(|z|)|u(z)|. \quad (2.40)$$

From the above inequality, we see that  $uC_\varphi : F(p, q, s) \rightarrow H_\mu^\infty$  is bounded. Let  $(f_k)_{k \in \mathbb{N}}$  be any bounded sequence in  $F(p, q, s)$  and  $f_k \rightarrow 0$  uniformly on compact subsets of  $B$  as  $k \rightarrow \infty$ . Employing Lemma 2.5, we have

$$\|uC_\varphi f_k\|_{H_\mu^\infty} = \sup_{z \in B} \mu(|z|) |f_k(\varphi(z))u(z)| \leq \|u\|_{H_\mu^\infty} \sup_{z \in B} |f_k(\varphi(z))| \rightarrow 0, \quad (2.41)$$

as  $k \rightarrow \infty$ . Then the result follows from Lemma 2.3.  $\square$

**Theorem 2.13.** *Assume that  $u \in H(B)$ ,  $\varphi$  is a holomorphic self-map of  $B$ ,  $\mu$  is a normal function on  $[0, 1)$ ,  $0 < p$ ,  $s < \infty$ ,  $-n - 1 < q < \infty$ ,  $q + s > -1$ ,  $p > n + 1 + q$ . Then the following statements are equivalent:*

- (i)  $uC_\varphi : F(p, q, s) \rightarrow H_{\mu,0}^\infty$  is bounded;
- (ii)  $uC_\varphi : F(p, q, s) \rightarrow H_{\mu,0}^\infty$  is compact;
- (iii)  $u \in H_{\mu,0}^\infty$ .

*Proof.* (ii) $\Rightarrow$ (i) It is obvious.

(i) $\Rightarrow$ (iii) Taking  $f(z) = 1$ , then using the boundedness of  $uC_\varphi : F(p, q, s) \rightarrow H_{\mu,0}^\infty$ , we get the desired result.

(iii) $\Rightarrow$ (ii) Suppose that  $u \in H_{\mu,0}^\infty$ . For any  $f \in F(p, q, s)$  with  $\|f\|_{F(p,q,s)} \leq 1$ , we have

$$\mu(|z|) |(uC_\varphi f)(z)| \leq C \|f\|_{F(p,q,s)} \mu(|z|) |u(z)| \leq C \mu(|z|) |u(z)|, \quad (2.42)$$

from which we obtain

$$\lim_{|z| \rightarrow 1} \sup_{\|f\|_{F(p,q,s)} \leq 1} \mu(|z|) |(uC_\varphi f)(z)| \leq C \lim_{|z| \rightarrow 1} \mu(|z|) |u(z)| = 0. \quad (2.43)$$

Using Lemma 2.3, we see that  $uC_\varphi : F(p, q, s) \rightarrow H_{\mu,0}^\infty$  is compact, as desired.  $\square$

## Acknowledgment

The author of this paper is supported by Educational Commission of Guangdong Province, China.

## References

- [1] A. L. Shields and D. L. Williams, "Bonded projections, duality, and multipliers in spaces of analytic functions," *Transactions of the American Mathematical Society*, vol. 162, pp. 287–302, 1971.
- [2] X. Fu and X. Zhu, "Weighted composition operators on some weighted spaces in the unit ball," *Abstract and Applied Analysis*, vol. 2008, Article ID 605807, 8 pages, 2008.
- [3] S. Stević, "Norm of weighted composition operators from Bloch space to  $H_\mu^\infty$  on the unit ball," *Ars Combinatoria*, vol. 88, pp. 125–127, 2008.
- [4] S. Stević, "On a new integral-type operator from the weighted Bergman space to the Bloch-type space on the unit ball," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 154263, 14 pages, 2008.

- [5] K. Zhu, *Spaces of Holomorphic Functions in the Unit Ball*, vol. 226 of *Graduate Texts in Mathematics*, Springer, New York, NY, USA, 2005.
- [6] D. D. Clahane and S. Stević, "Norm equivalence and composition operators between Bloch/Lipschitz spaces of the ball," *Journal of Inequalities and Applications*, vol. 2006, Article ID 61018, 11 pages, 2006.
- [7] S. Li and S. Stević, "Some characterizations of the Besov space and the  $\alpha$ -Bloch space," *Journal of Mathematical Analysis and Applications*, vol. 346, no. 1, pp. 262–273, 2008.
- [8] S. Li and H. Wulan, "Characterizations of  $\alpha$ -Bloch spaces on the unit ball," *Journal of Mathematical Analysis and Applications*, vol. 343, no. 1, pp. 58–63, 2008.
- [9] C. Ouyang, W. Yang, and R. Zhao, "Möbius invariant  $Q_p$  spaces associated with the Green's function on the unit ball of  $\mathbb{C}^n$ ," *Pacific Journal of Mathematics*, vol. 182, no. 1, pp. 69–99, 1998.
- [10] S. Li, "Riemann-Stieltjes operators from  $F(p, q, s)$  spaces to  $\alpha$ -Bloch spaces on the unit ball," *Journal of Inequalities and Applications*, vol. 2006, Article ID 27874, 14 pages, 2006.
- [11] S. Li and S. Stević, "Compactness of Riemann-Stieltjes operators between  $F(p, q, s)$  spaces and  $\alpha$ -Bloch spaces," *Publicationes Mathematicae Debrecen*, vol. 72, no. 1-2, pp. 111–128, 2008.
- [12] X. J. Zhang, "Multipliers on some holomorphic function spaces," *Chinese Annals of Mathematics. Series A*, vol. 26, no. 4, pp. 477–486, 2005.
- [13] R. Zhao, "On a general family of function spaces," *Annales Academiæ Scientiarum Fennicæ. Series A I. Mathematica. Dissertationes*, no. 105, p. 56, 1996.
- [14] C. C. Cowen and B. D. MacCluer, *Composition Operators on Spaces of Analytic Functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, Fla, USA, 1995.
- [15] X. Zhu, "Weighted composition operators between  $H^\infty$  and Bergman type spaces," *Communications of the Korean Mathematical Society*, vol. 21, no. 4, pp. 719–727, 2006.
- [16] S. Stević, "Weighted composition operators between mixed norm spaces and  $H_\alpha^\infty$  spaces in the unit ball," *Journal of Inequalities and Applications*, vol. 2007, Article ID 28629, 9 pages, 2007.
- [17] S. Li and S. Stević, "Weighted composition operators between  $H^\infty$  and  $\alpha$ -Bloch spaces in the unit ball," *Taiwanese Journal of Mathematics*, vol. 12, no. 7, pp. 1625–1639, 2008.
- [18] S. Li and S. Stević, "Weighted composition operators from  $\alpha$ -Bloch space to  $H^\infty$  on the polydisc," *Numerical Functional Analysis and Optimization*, vol. 28, no. 7-8, pp. 911–925, 2007.
- [19] S. Li and S. Stević, "Weighted composition operators from  $H^\infty$  to the Bloch space on the polydisc," *Abstract and Applied Analysis*, vol. 2007, Article ID 48478, 13 pages, 2007.
- [20] S. Stević, "Composition operators between  $H^\infty$  and  $\alpha$ -Bloch spaces on the polydisc," *Zeitschrift für Analysis und ihre Anwendungen*, vol. 25, no. 4, pp. 457–466, 2006.
- [21] L. Jiang and C. Ouyang, "Cyclic behavior of linear fractional composition operators in the unit ball of  $\mathbb{C}^n$ ," *Journal of Mathematical Analysis and Applications*, vol. 341, no. 1, pp. 601–612, 2008.
- [22] S. Li and S. Stević, "Composition followed by differentiation from mixed norm spaces to  $\alpha$ -Bloch spaces," *Matematicheskij Sbornik*, vol. 199, no. 12, pp. 117–128, 2008.
- [23] S. Li and S. Stević, "Products of Volterra type operator and composition operator from  $H^\infty$  and Bloch spaces to Zygmund spaces," *Journal of Mathematical Analysis and Applications*, vol. 345, no. 1, pp. 40–52, 2008.
- [24] S. Li and S. Stević, "Weighted composition operators from Zygmund spaces into Bloch spaces," *Applied Mathematics and Computation*, vol. 206, no. 2, pp. 825–831, 2008.
- [25] L. Luo and S.-I. Ueki, "Weighted composition operators between weighted Bergman spaces and Hardy spaces on the unit ball of  $\mathbb{C}^n$ ," *Journal of Mathematical Analysis and Applications*, vol. 326, no. 1, pp. 88–100, 2007.
- [26] S. Ohno, K. Stroethoff, and R. Zhao, "Weighted composition operators between Bloch-type spaces," *The Rocky Mountain Journal of Mathematics*, vol. 33, no. 1, pp. 191–215, 2003.
- [27] S. Stević, "Essential norms of weighted composition operators from the  $\alpha$ -Bloch space to a weighted-type space on the unit ball," *Abstract and Applied Analysis*, vol. 2008, Article ID 279691, 11 pages, 2008.
- [28] S. Stević, "Norms of some operators from Bergman spaces to weighted and Bloch-type spaces," *Utilitas Mathematica*, vol. 76, pp. 59–64, 2008.
- [29] S.-I. Ueki and L. Luo, "Compact weighted composition operators and multiplication operators between Hardy spaces," *Abstract and Applied Analysis*, vol. 2008, Article ID 196498, 12 pages, 2008.
- [30] S. Ye, "Weighted composition operator between the little  $\alpha$ -Bloch spaces and the logarithmic Bloch," *Journal of Computational Analysis and Applications*, vol. 10, no. 2, pp. 243–252, 2008.
- [31] X. Zhu, "Generalized weighted composition operators from Bloch type spaces to weighted Bergman spaces," *Indian Journal of Mathematics*, vol. 49, no. 2, pp. 139–150, 2007.
- [32] S. Stević, "On an integral operator on the unit ball in  $\mathbb{C}^n$ ," *Journal of Inequalities and Applications*, no. 1, pp. 81–88, 2005.

- [33] K. Madigan and A. Matheson, "Compact composition operators on the Bloch space," *Transactions of the American Mathematical Society*, vol. 347, no. 7, pp. 2679–2687, 1995.
- [34] S. Li and S. Stević, "Riemann-Stieltjes-type integral operators on the unit ball in  $\mathbb{C}^n$ ," *Complex Variables and Elliptic Equations*, vol. 52, no. 6, pp. 495–517, 2007.
- [35] S. Stević, "Boundedness and compactness of an integral operator in a mixed norm space on the polydisk," *Sibirskii Matematicheskii Zhurnal*, vol. 48, no. 3, pp. 694–706, 2007.