

Research Article

On Stability of Linear Delay Differential Equations under Perron's Condition

J. Diblík^{1,2} and A. Zafer³

¹ Department of Mathematics and Descriptive Geometry, Faculty of Civil Engineering, Brno University of Technology, Brno 602 00, Czech Republic

² Department of Mathematics, Faculty of Electrical Engineering and Communication, Brno University of Technology, Brno 616 00, Czech Republic

³ Department of Mathematics, Middle East Technical University, 06531 Ankara, Turkey

Correspondence should be addressed to J. Diblík, diblik@feec.vutbr.cz

Received 18 January 2011; Accepted 22 February 2011

Academic Editor: Miroslava Růžičková

Copyright © 2011 J. Diblík and A. Zafer. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The stability of the zero solution of a system of first-order linear functional differential equations with nonconstant delay is considered. Sufficient conditions for stability, uniform stability, asymptotic stability, and uniform asymptotic stability are established.

1. Introduction

We begin with a classical result for the linear system

$$x' = A(t)x, \quad (\text{L1})$$

where A is an $n \times n$ matrix function defined and continuous on $[0, \infty)$. By $C_B[0, \infty)$, we will denote the set of bounded functions defined and continuous on $[0, \infty)$ and by $|\cdot|$ the Euclidean norm.

In 1930, Perron first formulated the following definition being named after him.

Definition 1.1 (see [1]). System (L1) is said to satisfy Perron's condition (P1) if, for any given vector function $f \in C_B[0, \infty)$, the solution $x(t)$ of

$$x' = A(t)x + f(t), \quad x(0) = 0 \quad (\text{N1})$$

is bounded.

The following theorem by Bellman [2] is well known.

Theorem 1.2 (see [2]). *If (P1) holds and $|A(t)| \leq M_1$ for some positive number M_1 , then the zero solution of (L1) is uniformly asymptotically stable.*

The proof is accomplished by making use of the basic properties of a fundamental matrix, the Banach-Steinhaus theorem, and the adjoint system

$$x' = -A^T(t)x, \quad (1.1)$$

where A^T denotes the transpose of A .

It is shown by an example in [3] that Theorem 1.2 may not be valid if the function f appearing in (N1) is replaced by a constant vector. However, such a theorem is later obtained in [4] under a *Perron-like* condition.

Theorem 1.2 is extended by Halanay [5] to linear delay systems of the form

$$x'(t) = A(t)x(t) + B(t)x(t - \tau), \quad (L2)$$

where A, B are $n \times n$ matrix functions defined and continuous on $[0, \infty)$ and τ is a positive real number.

Definition 1.3. System (L2) is said to satisfy Perron's condition (P2) if for any given vector function $f \in C_B[0, \infty)$, the solution $x(t)$ of

$$x'(t) = A(t)x(t) + B(t)x(t - \tau) + f(t) \quad (N2)$$

satisfying $x(t) = 0, t \leq 0$, is bounded.

Theorem 1.4 (see [5]). *If (P2) holds, $|A(t)| \leq M_1$, and $|B(t)| \leq M_2$ for some positive numbers M_1 and M_2 , then the zero solution of (L2) is uniformly asymptotically stable.*

The method used to prove Theorem 1.4 is similar to Bellman's except that the adjoint system

$$y'(t) = -A^T(t)y(t) - B^T(t + \tau)y(t + \tau) \quad (1.2)$$

is not constructed with respect to an inner product but the functional

$$F(x, y)(t) = \int_t^{t+\tau} y^T(s)B(s)x(s - \tau)ds + x^T(t)y(t). \quad (1.3)$$

For some extensions to impulsive differential equations, we refer the reader in particular to [6, 7].

In this paper, we consider the more general linear delay system

$$x'(t) = A(t)x(t) + B(t)x(g(t)), \quad (1.4)$$

where A and B are $n \times n$ matrix functions defined and continuous on $[0, \infty)$ and g is a continuously differentiable increasing function defined on $[0, \infty)$ satisfying $g(t) < t$ and $g'(t) \leq 1$. We set $h := g^{-1}$. Obviously, $h \in C^1[0, \infty)$ and increases on $[0, \infty)$ and $h(t) > t$.

Perron's condition takes the following form.

Definition 1.5. System (1.4) is said to satisfy Perron's condition (P) if, for any given vector function $f \in C_B[0, \infty)$, the solution $x(t)$ of

$$x'(t) = A(t)x(t) + B(t)x(g(t)) + f(t) \quad (1.5)$$

satisfying $x(t) = 0, t \leq 0$ is bounded.

A natural question is whether the zero solution of (1.4) is uniformly asymptotically stable under Perron's condition (P). It turns out that the answer depends on the delay function g .

The paper is organized as follows. In Section 2, we only state our results; the proofs are included in Section 5. We define an adjoint system and give a variation of parameters formula in Section 3 to be needed in proving the main results. Section 4 contains also some lemmas concerning Perron's condition and a relation useful for changing the order of integration.

2. Stability Theorems

The conclusion obtained by Bellman and Halanay for systems (L1) and (L2), respectively, is quite strong. We are only able to prove the stability of the zero solution for more general equation (1.4) under Perron's condition. To get uniform stability or asymptotic stability or uniform asymptotic stability, we impose restrictions on the delay function.

For our purpose, we denote

$$\begin{aligned} h_*(t) &:= h(t) - t, \quad t \geq 0, \\ g_*(t, t_0) &:= \sup_{r \in [h(t_0), t]} \{r - g(r)\}, \quad t, t_0 \geq 0. \end{aligned} \quad (2.1)$$

Theorem 2.1. *Let (P) hold. If there are positive numbers M_1 and M_2 such that*

$$|A(t)| \leq M_1, \quad |B(t)| \leq M_2 \quad \forall t \geq 0, \quad (2.2)$$

then the zero solution of (1.4) is stable.

Theorem 2.2. *Let (P) hold. If (2.2) is satisfied and if there exists a positive real number M_3 such that*

$$h_*(t) \leq M_3 \quad \forall t \geq 0, \quad (2.3)$$

then the zero solution of (1.4) is uniformly stable.

Theorem 2.3. *Let (P) hold. If (2.2) and*

$$\limsup_{t \rightarrow \infty} \frac{g^*(t, t_0)}{t - t_0} = 0 \quad \text{for each } t_0 \geq 0 \quad (2.4)$$

are satisfied, then the zero solution of (1.4) is asymptotically stable.

Theorem 2.4. *Let (P) hold. If (2.2), (2.3), and*

$$\limsup_{t \rightarrow \infty} \frac{g^*(t, t_0)}{t - t_0} = 0 \quad \text{uniformly for } t_0 \geq 0 \quad (2.5)$$

are satisfied, then the zero solution of (1.4) is uniformly asymptotically stable.

Remark 2.5. Note that if $g(t) = t - \tau$, then $h(t) = t + \tau$ and hence the conditions (2.3), (2.4), and (2.5) are automatically satisfied. In this case, all theorems become equivalent, that is, the zero solution is uniformly asymptotically stable. Thus, the results obtained by Bellman and Halanay are recovered.

3. Variation of Parameters Formula

To establish a variation of parameters formula to represent the solutions of (1.5), one needs an adjoint system. The following lemma helps to define the adjoint of (1.4).

Lemma 3.1. *Let $x(t)$ be a solution of (1.4). If $y(t)$ is a solution of*

$$y'(t) = -A^T(t)y(t) - B^T(h(t))y(h(t))h'(t), \quad (3.1)$$

then

$$\frac{d}{dt}F(x(t), y(t)) = 0, \quad (3.2)$$

where

$$F(x, y)(t) = \int_t^{h(t)} y^T(s)B(s)x(g(s))ds + x^T(t)y(t). \quad (3.3)$$

Proof. Verify directly. □

Definition 3.2. The system (3.1) is said to be adjoint to system (1.4).

It is easy to see that the adjoint of system (3.1) is system (1.4); thus the systems are mutually adjoint to each other.

Lemma 3.3. Let $Y(t, s)$ be a matrix solution of (3.1) for $t < s$ satisfying $Y(s, s) = I$ and $Y(t, s) = 0$ for $t > s$. Then $x(t)$ is a solution of (1.5) if and only if

$$x(t) = Y^T(s, t)x(s) + \int_{g(s)}^s Y^T(h(\beta), t)B(h(\beta))x(\beta)h'(\beta)d\beta + \int_s^t Y^T(\beta, t)f(\beta)d\beta. \quad (3.4)$$

Proof. Replacing t by β in (1.5) and then integrating the resulting equation multiplied by $Y^T(\beta, t)$ over $\beta \in [s, t]$, we have

$$\begin{aligned} & \int_s^t Y^T(\beta, t)A(\beta)x(\beta)d\beta + \int_s^t Y^T(\beta, t)B(\beta)x(g(\beta))d\beta + \int_s^t Y^T(\beta, t)f(\beta)d\beta \\ &= \int_s^t Y^T(\beta, t)x'(\beta)d\beta \\ &= x(t) - Y^T(s, t)x(s) - \int_s^t \left[\frac{\partial}{\partial \beta} Y^T(\beta, t) \right] x(\beta)d\beta \\ &= x(t) - Y^T(s, t)x(s) + \int_s^t \left[Y^T(\beta, t)A(\beta) + Y^T(h(\beta), t)B(h(\beta))h'(\beta) \right] x(\beta)d\beta \\ &= x(t) - Y^T(s, t)x(s) + \int_s^t Y^T(\beta, t)A(\beta)x(\beta)d\beta + \int_{h(s)}^{h(t)} Y^T(\beta, t)B(\beta)x(g(\beta))d\beta. \end{aligned} \quad (3.5)$$

Comparing both sides and using

$$\int_t^{h(t)} Y^T(\beta, t)B(\beta)x(g(\beta))d\beta = 0, \quad (3.6)$$

which is true in view of $Y(\beta, t) = 0$ for $\beta > t$, we get

$$x(t) = Y^T(s, t)x(s) - \int_{h(s)}^s Y^T(\beta, t)B(\beta)x(g(\beta))d\beta + \int_s^t Y^T(\beta, t)f(\beta)d\beta \quad (3.7)$$

and hence

$$x(t) = Y^T(s, t)x(s) + \int_{g(s)}^s Y^T(h(\beta), t)B(h(\beta))x(\beta)h'(\beta)d\beta + \int_s^t Y^T(\beta, t)f(\beta)d\beta. \quad (3.8)$$

□

It is not difficult to see from (3.4) that if $X(t, s)$ is a matrix solution of (1.4) for $t > s$ satisfying $X(s, s) = I$ and $X(t, s) = 0$ for $t < s$, then

$$X(t, s) = Y^T(s, t). \quad (3.9)$$

Using this relation in Lemma 3.3 leads to the following variation of parameters formula.

Lemma 3.4. Let $X(t, s)$ be a matrix solution of (1.4) for $t > s$ satisfying $X(s, s) = I$ and $X(t, s) = 0$ for $t < s$. Then $x(t)$ is a solution of (1.5) if and only if

$$x(t) = X(t, s)x(s) + \int_{g(s)}^s X(t, h(\beta))B(h(\beta))x(\beta)h'(\beta)d\beta + \int_s^t X(t, \beta)f(\beta)d\beta. \quad (3.10)$$

4. Auxiliary Results

Lemma 4.1. If (P) holds, then there is a positive number K_1 such that

$$\int_0^t |X(t, s)|ds \leq K_1 \quad \forall t > 0. \quad (4.1)$$

Proof. The proof follows as in [5]. We provide only the steps for the reader's convenience. Define

$$\begin{aligned} (Sf)(t) &= \int_0^t X(t, \beta)f(\beta)d\beta, \quad f \in C_B[0, \infty), \\ S_k(f) &= \int_0^{t_k} X(t_k, \beta)f(\beta)d\beta, \quad f \in C_B[0, \infty), \end{aligned} \quad (4.2)$$

for each rational number $t_k, k \in \mathbb{N}$.

In view of (P), the family of continuous linear operators $\{S_k\}$ from $C_B[0, \infty)$ to $C_B[0, \infty)$ is pointwise-bounded. For the space of bounded continuous functions $C_B[0, \infty)$, the usual sup norm $\|\cdot\|$ is used.

By the Banach-Steinhaus theorem, the family is uniformly bounded. Thus, there is a positive number M such that $\|S_k(f)\| \leq M\|f\|$ for every $f \in C_B[0, \infty)$.

As the rational numbers are dense in the real numbers, for each t there is t_k such that $t_k \rightarrow t$ as $k \rightarrow \infty$ and so

$$\left| \int_0^t X(t, \beta)f(\beta)d\beta \right| \leq M\|f\| \quad \forall f \in C_B[0, \infty). \quad (4.3)$$

The final step involves choosing a sequence of functions and using a limiting process. \square

Lemma 4.2. If (2.2) and (4.1) are true, then there is a positive number K_2 such that

$$|Y(s, t)| \leq K_2 \quad \forall 0 \leq s < t. \quad (4.4)$$

Proof. From (3.1), we have

$$Y(s, t) = I + \int_s^t A^T(\beta)Y(\beta, t)d\beta + \int_s^t B^T(h(\beta))Y(h(\beta), t)h'(\beta)d\beta. \quad (4.5)$$

Hence, by using (4.1), we see that for all $0 \leq s < t$,

$$|Y(s, t)| \leq 1 + M_1 K_1 + M_2 K_1 =: K_2. \tag{4.6}$$

□

Lemma 4.3. *Let $G(r, t)$ be a continuous function satisfying $G(r, t) = 0$ for $r > t$. Then*

$$\int_{t_0}^t \left[\int_s^{h(s)} G(r, t) dr \right] ds = \int_{h(t_0)}^t (r - g(r)) G(r, t) dr + \int_{t_0}^{h(t_0)} (r - t_0) G(r, t) dr. \tag{4.7}$$

5. Proofs of Theorems

Let $t_0 \geq 0$ be given. For a given continuous vector function ϕ defined on $[g(t_0), t_0]$, let $x(t) = x(t, t_0, \phi)$ denote the solution of (1.4) satisfying

$$x(t) = \phi(t), \quad t \leq t_0. \tag{5.1}$$

As usual,

$$\|\phi\|_g = \sup_{t \in [g(t_0), t_0]} |\phi(t)|. \tag{5.2}$$

Proof of Theorem 2.1. From Lemma 3.3, we may write

$$x(t) = Y^T(t_0, t)\phi(t_0) + \int_{g(t_0)}^{t_0} Y^T(h(\beta), t) B(h(\beta)) \phi(\beta) h'(\beta) d\beta. \tag{5.3}$$

In view of Lemma 4.2, it follows that

$$|x(t)| \leq (K_2 + (h(t_0) - t_0) K_2 M_2) \|\phi\|_g. \tag{5.4}$$

Hence, the zero solution is stable. □

Proof of Theorem 2.2. Using (2.3) in (5.4), we get

$$|x(t)| \leq K_3 \|\phi\|_g, \quad K_3 = K_2 + K_2 M_2 M_3, \tag{5.5}$$

from which the uniform stability follows. □

Proof of Theorem 2.3. By Theorem 2.1, the zero solution is stable. We need to show the attractivity property.

From Lemma 3.3, for $s \geq t_0$, we can write

$$x(t, t_0, \phi) = Y^T(s, t)x(s, t_0, \phi) + \int_{g(s)}^s G(h(\beta), t)x(\beta, t_0, \phi)h'(\beta) d\beta, \tag{5.6}$$

where

$$G(s, t) = Y^T(s, t)B(s). \quad (5.7)$$

Integrating with respect to s from t_0 to t , we have

$$(t - t_0)x(t, t_0, \phi) = \int_{t_0}^t \left[Y^T(s, t)x(s, t_0, \phi) + \int_s^{h(s)} G(r, t)x(g(r), t_0, \phi) dr \right] ds. \quad (5.8)$$

We change the order of integration by employing Lemma 4.3. After some rearrangements, we obtain

$$\begin{aligned} (t - t_0)x(t, t_0, \phi) &= \int_{t_0}^t Y^T(s, t)x(s, t_0, \phi) ds + \int_{h(t_0)}^t (s - g(s))G(s, t)x(g(s), t_0, \phi) ds \\ &\quad + \int_{t_0}^{h(t_0)} (s - t_0)G(s, t)x(g(s), t_0, \phi) ds. \end{aligned} \quad (5.9)$$

It follows that

$$(t - t_0)|x(t, t_0, \phi)| \leq K_1 K_3 \|\phi\|_g + g_*(t, t_0) M_2 K_1 \|\phi\|_g + h_*(t_0) M_2 K_1 \|\phi\|_g. \quad (5.10)$$

In view of condition (2.4), we see from (5.10) that

$$\lim_{t \rightarrow \infty} |x(t, t_0, \phi)| = 0. \quad (5.11)$$

□

Proof of Theorem 2.4. By Theorem 2.2, the zero solution is uniformly stable. From (5.10) and (2.3), we have

$$(t - t_0)|x(t, t_0, \phi)| \leq K_1 K_3 \|\phi\|_g + g_*(t, t_0) M_2 K_1 \|\phi\|_g + M_3 M_2 K_1 \|\phi\|_g. \quad (5.12)$$

Using condition (2.4) in the above inequality, we see that the zero solution is uniformly asymptotically stable as $t \rightarrow \infty$. □

Acknowledgments

This research was supported by Grant P201/11/0768 of the Czech Grant Agency (Prague), by the Council of Czech Government MSM 0021630503 and MSM 00216 30519, and by Grant FEKT-S-11-2-921 of Faculty of Electrical Engineering and Communication, Brno University of Technology.

References

- [1] O. Perron, "Die stabilitätsfrage bei differentialgleichungen," *Mathematische Zeitschrift*, vol. 32, no. 1, pp. 703–728, 1930.
- [2] R. Bellman, "On an application of a Banach-Steinhaus theorem to the study of the boundedness of solutions of non-linear differential and difference equations," *Annals of Mathematics*, vol. 49, pp. 515–522, 1948.
- [3] J. Kloch, "An illustrative example for the Perron condition," *Annales Polonici Mathematici*, vol. 35, no. 1, pp. 11–14, 1978.
- [4] R. Reissig, "A Perron-like stability criterion for linear systems," *Archiv der Mathematik*, vol. 34, no. 1, pp. 53–59, 1980.
- [5] A. Halanay, *Differential Equations: Stability, Oscillations, Time Lags*, Academic Press, New York, NY, USA, 1966.
- [6] M. U. Akhmet, J. Alzabut, and A. Zafer, "Perron's theorem for linear impulsive differential equations with distributed delay," *Journal of Computational and Applied Mathematics*, vol. 193, no. 1, pp. 204–218, 2006.
- [7] A. Anokhin, L. Berezansky, and E. Braverman, "Exponential stability of linear delay impulsive differential equations," *Journal of Mathematical Analysis and Applications*, vol. 193, no. 3, pp. 923–941, 1995.