

Research Article

Sharp Bounds for Seiffert Mean in Terms of Contraharmonic Mean

Yu-Ming Chu¹ and Shou-Wei Hou²

¹ Department of Mathematics, Huzhou Teachers College, Huzhou 313000, China

² Department of Mathematics, Hangzhou Normal University, Hangzhou 310012, China

Correspondence should be addressed to Yu-Ming Chu, chuyuming2005@yahoo.com.cn

Received 17 September 2011; Accepted 5 November 2011

Academic Editor: Muhammad Aslam Noor

Copyright © 2012 Y.-M. Chu and S.-W. Hou. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We find the greatest value α and the least value β in $(1/2, 1)$ such that the double inequality $C(\alpha a + (1 - \alpha)b, \alpha b + (1 - \alpha)a) < T(a, b) < C(\beta a + (1 - \beta)b, \beta b + (1 - \beta)a)$ holds for all $a, b > 0$ with $a \neq b$. Here, $T(a, b) = (a - b) / [2 \arctan((a - b) / (a + b))]$ and $C(a, b) = (a^2 + b^2) / (a + b)$ are the Seiffert and contraharmonic means of a and b , respectively.

1. Introduction

For $a, b > 0$ with $a \neq b$, the Seiffert mean $T(a, b)$ and contraharmonic mean $C(a, b)$ are defined by

$$T(a, b) = \frac{a - b}{2 \arctan((a - b) / (a + b))}, \quad (1.1)$$

$$C(a, b) = \frac{a^2 + b^2}{a + b}, \quad (1.2)$$

respectively. Recently, both mean values have been the subject of intensive research. In particular, many remarkable inequalities and properties for these means can be found in the literature [1–12].

Let $A(a, b) = (a + b) / 2$, $G(a, b) = \sqrt{ab}$, $S(a, b) = \sqrt{(a^2 + b^2) / 2}$, and let $M_p(a, b) = ((a^p + b^p) / 2)^{1/p}$ ($p \neq 0$) and $M_0(a, b) = \sqrt{ab}$ be the arithmetic, geometric, square root, and p th power means of two positive numbers a and b , respectively. Then it is well known that

$M_p(a, b)$ is continuous and strictly increasing with respect to $p \in \mathbb{R}$ for fixed $a, b > 0$ with $a \neq b$, and the inequalities

$$G(a, b) = M_0(a, b) < A(a, b) = M_1(a, b) < S(a, b) = M_2(a, b) < C(a, b) \quad (1.3)$$

hold for all $a, b > 0$ with $a \neq b$.

Seiffert [12] proved that the double inequality

$$A(a, b) = M_1(a, b) < T(a, b) < M_2(a, b) = S(a, b) \quad (1.4)$$

holds for all $a, b > 0$ with $a \neq b$.

Hästö [13] proved that the function $T(1, x)/M_p(1, x)$ is increasing in $(0, \infty)$ if $p \leq 1$.

In [14], the authors found the greatest value p and the least value q such that the double inequality $H_p(a, b) < T(a, b) < H_q(a, b)$ holds for all $a, b > 0$ with $a \neq b$. Here, $H_k(a, b) = ((a^k + (ab)^{k/2} + b^k)/3)^{1/k}$ ($k \neq 0$), and $H_0(a, b) = \sqrt{ab}$ is the k th power-type Heron mean of a and b .

Wang et al. [15] answered the question: what are the best possible parameters λ and μ such that the double inequality $L_\lambda(a, b) < T(a, b) < L_\mu(a, b)$ holds for all $a, b > 0$ with $a \neq b$, where $L_r(a, b) = (a^{r+1} + b^{r+1})/(a^r + b^r)$ is the r th Lehmer mean of a and b .

In [16, 17], the authors proved that the inequalities

$$\begin{aligned} \alpha_1 T(a, b) + (1 - \alpha_1)G(a, b) &< A(a, b) < \beta_1 T(a, b) + (1 - \beta_1)G(a, b), \\ \alpha_2 S(a, b) + (1 - \alpha_2)A(a, b) &< T(a, b) < \beta_2 S(a, b) + (1 - \beta_2)A(a, b), \\ S^{\alpha_3}(a, b)A^{1-\alpha_3}(a, b) &< T(a, b) < S^{\beta_3}(a, b)A^{1-\beta_3}(a, b) \end{aligned} \quad (1.5)$$

hold for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_1 \leq 3/5$, $\beta_1 \geq \pi/4$, $\alpha_2 \leq (4 - \pi)/[(\sqrt{2} - 1)\pi]$, $\beta_2 \geq 2/3$, $\alpha_3 \leq 2/3$ and $\beta_3 \geq 4 - 2 \log \pi / \log 2$.

For fixed $a, b > 0$ with $a \neq b$, let $x \in [1/2, 1]$ and

$$J(x) = C(xa + (1 - x)b, xb + (1 - x)a). \quad (1.6)$$

Then it is not difficult to verify that $J(x)$ is continuous and strictly increasing in $[1/2, 1]$. Note that $J(1/2) = A(a, b) < T(a, b)$ and $J(1) = C(a, b) > T(a, b)$. Therefore, it is natural to ask what are the greatest value α and the least value β in $(1/2, 1)$ such that the double inequality

$$C(\alpha a + (1 - \alpha)b, \alpha b + (1 - \alpha)a) < T(a, b) < C(\beta a + (1 - \beta)b, \beta b + (1 - \beta)a) \quad (1.7)$$

holds for all $a, b > 0$ with $a \neq b$. The main purpose of this paper is to answer this question. Our main result is the following Theorem 1.1.

Theorem 1.1. *If $\alpha, \beta \in (1/2, 1)$, then the double inequality*

$$C(\alpha a + (1 - \alpha)b, \alpha b + (1 - \alpha)a) < T(a, b) < C(\beta a + (1 - \beta)b, \beta b + (1 - \beta)a) \quad (1.8)$$

holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha \leq (1 + \sqrt{4/\pi - 1})/2$ and $\beta \geq (3 + \sqrt{3})/6$.

2. Proof of Theorem 1.1

Proof of Theorem 1.1. Let $\lambda = (1 + \sqrt{4/\pi - 1})/2$ and $\mu = (3 + \sqrt{3})/6$. We first proof that the inequalities

$$T(a, b) > C(\lambda a + (1 - \lambda)b, \lambda b + (1 - \lambda)a), \tag{2.1}$$

$$T(a, b) < C(\mu a + (1 - \mu)b, \mu b + (1 - \mu)a) \tag{2.2}$$

hold for all $a, b > 0$ with $a \neq b$.

From (1.1) and (1.2) we clearly see that both $T(a, b)$ and $C(a, b)$ are symmetric and homogenous of degree 1. Without loss of generality, we assume that $a > b$. Let $t = a/b > 1$ and $p \in (1/2, 1)$, then from (1.1) and (1.2) one has

$$\begin{aligned} & C(pa + (1 - p)b, pb + (1 - p)a) - T(a, b) \\ &= b \frac{[pt + (1 - p)]^2 + [(1 - p)t + p]^2}{2(t + 1) \arctan((t - 1)/(t + 1))} \\ & \times \left\{ 2 \arctan\left(\frac{t - 1}{t + 1}\right) - \frac{t^2 - 1}{[pt + (1 - p)]^2 + [(1 - p)t + p]^2} \right\}. \end{aligned} \tag{2.3}$$

Let

$$f(t) = 2 \arctan\left(\frac{t - 1}{t + 1}\right) - \frac{t^2 - 1}{[pt + (1 - p)]^2 + [(1 - p)t + p]^2}. \tag{2.4}$$

Then simple computations lead to

$$f(1) = 0, \tag{2.5}$$

$$\lim_{t \rightarrow +\infty} f(t) = \frac{\pi}{2} - \frac{1}{p^2 + (1 - p)^2}, \tag{2.6}$$

$$f'(t) = \frac{2f_1(t)}{\left\{ [pt + (1 - p)]^2 + [(1 - p)t + p]^2 \right\}^2 (t + 1)}, \tag{2.7}$$

where

$$\begin{aligned} f_1(t) &= (4p^4 - 8p^3 + 10p^2 - 6p + 1)t^4 - 2(2p - 1)^2(2p^2 - 2p + 1)t^3 \\ &+ 2(12p^4 - 24p^3 + 18p^2 - 6p + 1)t^2 \end{aligned} \tag{2.8}$$

$$\begin{aligned} & - 2(2p - 1)^2(2p^2 - 2p + 1)t + 4p^4 - 8p^3 + 10p^2 - 6p + 1, \\ & f_1(1) = 0. \end{aligned} \tag{2.9}$$

Let $f_2(t) = f'_1(t)/2$, $f_3(t) = f'_2(t)/2$, $f_4(t) = f'_3(t)/3$. Then from (2.8) we get

$$f_2(t) = 2(4p^4 - 8p^3 + 10p^2 - 6p + 1)t^3 - 3(2p - 1)^2(2p^2 - 2p + 1)t^2 + 2(12p^4 - 24p^3 + 18p^2 - 6p + 1)t - (2p - 1)^2(2p^2 - 2p + 1), \quad (2.10)$$

$$f_2(1) = 0, \quad (2.11)$$

$$f_3(t) = 3(4p^4 - 8p^3 + 10p^2 - 6p + 1)t^2 - 3(2p - 1)^2(2p^2 - 2p + 1)t + 12p^4 - 24p^3 + 18p^2 - 6p + 1, \quad (2.12)$$

$$f_3(1) = 6p^2 - 6p + 1, \quad (2.13)$$

$$f_4(t) = 2(4p^4 - 8p^3 + 10p^2 - 6p + 1)t - (2p - 1)^2(2p^2 - 2p + 1), \quad (2.14)$$

$$f_4(1) = 6p^2 - 6p + 1. \quad (2.15)$$

We divide the proof into two cases.

Case 1 ($p = \lambda = (1 + \sqrt{4/\pi - 1})/2$). Then (2.6), (2.13), and (2.15) lead to

$$\lim_{t \rightarrow +\infty} f(t) = 0, \quad (2.16)$$

$$f_3(1) = -\frac{2(\pi - 3)}{\pi} < 0, \quad (2.17)$$

$$f_4(1) = -\frac{2(\pi - 3)}{\pi} < 0. \quad (2.18)$$

Note that

$$4p^4 - 8p^3 + 10p^2 - 6p + 1 = \frac{4 + 2\pi - \pi^2}{\pi^2} > 0. \quad (2.19)$$

It follows from (2.8), (2.10), (2.12), (2.14), and (2.19) that

$$\lim_{t \rightarrow +\infty} f_1(t) = +\infty, \quad (2.20)$$

$$\lim_{t \rightarrow +\infty} f_2(t) = +\infty, \quad (2.21)$$

$$\lim_{t \rightarrow +\infty} f_3(t) = +\infty, \quad (2.22)$$

$$\lim_{t \rightarrow +\infty} f_4(t) = +\infty. \quad (2.23)$$

From (2.14) and inequality (2.19), we clearly see that $f_4(t)$ is strictly increasing in $[1, +\infty)$. Then (2.18) and (2.23) lead to the conclusion that there exists $t_0 > 1$ such that $f_4(t) < 0$ for $t \in [1, t_0)$ and $f_4(t) > 0$ for $t \in (t_0, +\infty)$. Hence, $f_3(t)$ is strictly decreasing in $[1, t_0]$ and strictly increasing in $[t_0, +\infty)$.

It follows from (2.17) and (2.22) together with the piecewise monotonicity of $f_3(t)$ that there exists $t_1 > t_0 > 1$ such that $f_2(t)$ is strictly decreasing in $[1, t_1]$ and strictly increasing in $[t_1, +\infty)$.

From (2.11) and (2.21) together with the piecewise monotonicity of $f_2(t)$, we conclude that there exists $t_2 > t_1 > 1$ such that $f_1(t)$ is strictly decreasing in $[1, t_2]$ and strictly increasing in $[t_2, +\infty)$.

Equations (2.7), (2.9), and (2.20) together with the piecewise monotonicity of $f_1(t)$ imply that there exists $t_3 > t_2 > 1$ such that $f(t)$ is strictly decreasing in $[1, t_3]$ and strictly increasing in $[t_3, +\infty)$.

Therefore, inequality (2.1) follows from (2.3)–(2.5) and (2.16) together with the piecewise monotonicity of $f(t)$.

Case 2 ($p = \mu = (3 + \sqrt{3})/6$). Then (2.8) leads to

$$f_1(t) = \frac{(t-1)^4}{9} > 0 \tag{2.24}$$

for $t > 1$.

Inequality (2.24) and (2.7) imply that $f(t)$ is strictly increasing in $[1, +\infty)$. Therefore, inequality (2.2) follows from (2.3)–(2.5) together with the monotonicity of $f(t)$.

From inequalities (2.1) and (2.2) together with the monotonicity of $J(x) = C(xa + (1-x)b, xb + (1-x)a)$ in $[1/2, 1]$, we know that inequality (1.8) holds for all $\alpha \leq (1 + \sqrt{4/\pi - 1})/2$, $\beta \geq (3 + \sqrt{3})/6$, and all $a, b > 0$ with $a \neq b$.

Next, we prove that $\lambda = (1 + \sqrt{4/\pi - 1})/2$ is the best possible parameter in $[1/2, 1]$ such that inequality (2.1) holds for all $a, b > 0$ with $a \neq b$.

For any $1 > p > \lambda = (1 + \sqrt{4/\pi - 1})/2$, from (2.6) one has

$$\lim_{t \rightarrow +\infty} f(t) = \frac{\pi}{2} - \frac{1}{p^2 + (1-p)^2} > 0. \tag{2.25}$$

Equations (2.3) and (2.4) together with inequality (2.25) imply that for any $1 > p > \lambda = (1 + \sqrt{4/\pi - 1})/2$ there exists $T_0 = T_0(p) > 1$ such that

$$C(pa + (1-p)b, pb + (1-p)a) > T(a, b) \tag{2.26}$$

for $a/b \in (T_0, +\infty)$.

Finally, we prove that $\mu = (3 + \sqrt{3})/6$ is the best possible parameter such that inequality (2.2) holds for all $a, b > 0$ with $a \neq b$.

For any $1/2 < p < \mu = (3 + \sqrt{3})/6$, from (2.13) one has

$$f_3(1) = 6p^2 - 6p + 1 < 0. \tag{2.27}$$

From inequality (2.27) and the continuity of $f_3(t)$, we know that there exists $\delta = \delta(p) > 0$ such that

$$f_3(t) < 0 \tag{2.28}$$

for $t \in (1, 1 + \delta)$.

Equations (2.3)–(2.5), (2.7), (2.9), and (2.11) together with inequality (2.28) imply that for any $1/2 < p < \mu = (3 + \sqrt{3})/6$ there exists $\delta = \delta(p) > 0$ such that

$$T(a, b) > C(pa + (1 - p)b, pb + (1 - p)a) \quad (2.29)$$

for $a/b \in (1, 1 + \delta)$. □

Acknowledgments

This research was supported by the National Natural Science Foundation of China under Grants 11071069 and 11171307, and the Innovation Team Foundation of the Department of Education of Zhejiang Province under Grant T200924.

References

- [1] Y.-M. Chu, M.-K. Wang, S.-L. Qiu, and Y.-F. Qiu, "Sharp generalized Seiffert mean bounds for Toader mean," *Abstract and Applied Analysis*, vol. 2011, Article ID 605259, 8 pages, 2011.
- [2] S.-W. Hou and Y.-M. Chu, "Optimal convex combination bounds of root-square and harmonic root-square means for Seiffert mean," *International Journal of Mathematical Analysis*, vol. 5, no. 39, pp. 1897–1904, 2011.
- [3] H. Liu and X.-J. Meng, "The optimal convex combination bounds for Seiffert's mean," *Journal of Inequalities and Applications*, vol. 2011, Article ID 686834, 9 pages, 2011.
- [4] J. Pahikkala, "On contraharmonic mean and Pythagorean triples," *Elementary Math*, vol. 65, no. 2, pp. 62–67, 2010.
- [5] S. Toader and Gh. Toader, "Complementaries of Greek means with respect to Gini means," *International Journal of Applied Mathematics & Statistics*, vol. 11, no. N07, pp. 187–192, 2007.
- [6] E. Neuman and J. Sándor, "On the Schwab-Borchardt mean II," *Mathematica Pannonica*, vol. 17, no. 1, pp. 49–59, 2006.
- [7] Y. Lim, "The inverse mean problem of geometric mean and contraharmonic means," *Linear Algebra and its Applications*, vol. 408, no. 1–3, pp. 221–229, 2005.
- [8] S. Toader and Gh. Toader, "Generalized complementaries of Greek means," *Pure Mathematics and Applications*, vol. 15, no. 2–3, pp. 335–342, 2004.
- [9] E. Neuman and J. Sándor, "On the Schwab-Borchardt mean," *Mathematica Pannonica*, vol. 14, no. 2, pp. 253–266, 2003.
- [10] J. R. Siler, "Mean crowds and Pythagorean triples," *Fibonacci Quarterly*, vol. 36, no. 4, pp. 323–326, 1998.
- [11] W. N. Anderson Jr., M. E. Mays, T. D. Morley, and G. E. Trapp, "The contraharmonic mean of HSD matrices," *SIAM Journal on Algebraic and Discrete Methods*, vol. 8, no. 4, pp. 674–682, 1987.
- [12] H. J. Seiffert, "Aufgabe β 16," *Die Wurzel*, vol. 29, pp. 221–222, 1995.
- [13] P. A. Hästö, "A monotonicity property of ratios of symmetric homogeneous means," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 3, no. 5, article 71, 23 pages, 2002.
- [14] Y.-M. Chu, M.-K. Wang, and Y.-F. Qiu, "An optimal double inequality between power-type Heron and Seiffert means," *Journal of Inequalities and Applications*, vol. 2010, Article ID 146945, 11 pages, 2010.
- [15] M.-K. Wang, Y.-F. Qiu, and Y.-M. Chu, "Sharp bounds for Seiffert means in terms of Lehmer means," *Journal of Mathematical Inequalities*, vol. 4, no. 4, pp. 581–586, 2010.
- [16] Y.-M. Chu, C. Zong, and G.-D. Wang, "Optimal convex combination bounds of Seiffert and geometric means for the arithmetic mean," *Journal of Mathematical Inequalities*, vol. 5, no. 3, pp. 429–434, 2011.
- [17] Y.-M. Chu, M.-K. Wang, and W.-M. Gong, "Two sharp double inequalities for Seiffert mean," *Journal of Inequalities and Applications*, vol. 2011, article 44, 2011.