

## Research Article

# Some Spaces of Double Sequences Obtained through Invariant Mean and Related Concepts

S. A. Mohiuddine and Abdullah Alotaibi

Department of Mathematics, Faculty of Science, King Abdulaziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia

Correspondence should be addressed to S. A. Mohiuddine; mohiuddine@gmail.com

Received 29 November 2012; Accepted 11 March 2013

Academic Editor: Elena Braverman

Copyright © 2013 S. A. Mohiuddine and A. Alotaibi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce some double sequences spaces involving the notions of invariant mean (or  $\sigma$ -mean) and  $\sigma$ -convergence for double sequences while the idea of  $\sigma$ -convergence for double sequences was introduced by Çakan et al. 2006, by using the notion of invariant mean. We determine here some inclusion relations and topological results for these new double sequence spaces.

## 1. Preliminaries, Background, and Notation

In 1900, Pringsheim [1] presented the following notion of convergence for double sequences. A double sequence  $x = (x_{jk})$  is said to converge to the limit  $L$  in Pringsheim's sense (shortly,  $p$ -convergent to  $L$ ) if for every  $\varepsilon > 0$  there exists an integer  $N$  such that  $|x_{jk} - L| < \varepsilon$  whenever  $j, k > N$ . In this case,  $L$  is called the  $p$ -limit of  $x$ .

A double sequence  $x = (x_{jk})$  of real or complex numbers is said to be bounded if  $\|x\|_\infty = \sup_{j,k} |x_{jk}| < \infty$ . The space of all bounded double sequences is denoted by  $\mathcal{M}_u$ .

If  $x \in \mathcal{M}_u$  and is  $p$ -convergent to  $L$ , then  $x$  is said to be boundedly  $p$ -convergent to  $L$  (shortly,  $bp$ -convergent to  $L$ ). In this case,  $L$  is called the  $bp$ -limit of  $x$ . The assumption of  $p$ -convergent was made because a double sequence on which  $p$ -convergent is not necessarily bounded.

In general, for any notion of convergence  $\nu$ , the space of all  $\nu$ -convergent double sequences will be denoted by  $\mathcal{C}_\nu$ , the space of all  $\nu$ -convergent to 0 double sequences by  $\mathcal{C}_{\nu 0}$  and the limit of a  $\nu$ -convergent double sequence  $x$  by  $\nu\text{-lim}_{j,k} x_{jk}$ , where  $\nu \in \{p, bp\}$ .

Let  $\Omega$  denote the vector space of all double sequences with the vector space operations defined coordinatewise. Vector subspaces of  $\Omega$  are called double sequence spaces. In addition

to the above-mentioned double sequence spaces, we consider the double sequence space as

$$\mathcal{L}_u := \left\{ x \in \Omega \mid \|x\|_1 := \sum_{j,k} |x_{jk}| < \infty \right\} \quad (1)$$

of all absolutely summable double sequences.

All considered double sequence spaces are supposed to contain

$$\Phi := \text{span} \{ e^{jk} \mid j, k \in \mathbb{N} \}, \quad (2)$$

where

$$e_{il}^{jk} = \begin{cases} 1; & \text{if } (j, k) = (i, \ell), \\ 0; & \text{otherwise.} \end{cases} \quad (3)$$

We denote the pointwise sums  $\sum_{j,k} e^{jk}$ ,  $\sum_j e^{jk}$  ( $k \in \mathbb{N}$ ), and  $\sum_k e^{jk}$  ( $j \in \mathbb{N}$ ) by  $e$ ,  $e^k$ , and  $e_j$ , respectively.

Let  $\sigma$  be a one-to-one mapping from the set  $\mathbb{N}$  of natural numbers into itself. A continuous linear functional  $\varphi$  on the space  $\ell_\infty$  of bounded single sequences is said to be an invariant mean or a  $\sigma$ -mean if and only if (i)  $\varphi(x) \geq 0$  when

the sequence  $x = (x_k)$  has  $x_k \geq 0$  for all  $k$ , (ii)  $\varphi(e) = 1$ , where  $e = (1, 1, 1, \dots)$ , and (iii)  $\varphi(x) = \varphi((x_{\sigma(k)})$  for all  $x \in \ell_\infty$ .

Throughout this paper we consider the mapping  $\sigma$  which has no finite orbits, that is,  $\sigma^p(k) \neq k$  for all integer  $k \geq 0$  and  $p \geq 1$ , where  $\sigma^p(k)$  denotes the  $p$ th iterate of  $\sigma$  at  $k$ . Note that a  $\sigma$ -mean extends the limit functional on the space  $c$  of convergent single sequences in the sense that  $\varphi(x) = \lim x$  for all  $x \in c$ , (see [2]). Consequently,  $c \subset V_\sigma$  the set of bounded sequences all of whose  $\sigma$ -means are equal. We say that a sequence  $x = (x_k)$  is  $\sigma$ -convergent if and only if  $x \in V_\sigma$ . Using this concept, Schaefer [3] defined and characterized  $\sigma$ -conservative,  $\sigma$ -regular, and  $\sigma$ -coercive matrices for single sequences. If  $\sigma$  is translation then  $V_\sigma$  is reduced to the set  $f$  of almost convergent sequences [4]. Recently, Mohiuddine [5] has obtained an application of almost convergence for single sequences in approximation theorems and proved some related results.

In 2006, Çakan et al. [6] presented the following definition of  $\sigma$ -convergence for double sequences and further studied by Mursaleen and Mohiuddine [7–9]. A double sequence  $x = (x_{jk})$  of real numbers is said to be  $\sigma$ -convergent to a number  $L$  if and only if  $x \in \mathcal{V}_\sigma$ , where

$$\mathcal{V}_\sigma = \left\{ x \in \mathcal{M}_u : \lim_{p,q \rightarrow \infty} \zeta_{pqst}(x) = L \text{ uniformly in } s, t; \right. \\ \left. L = \sigma\text{-}\lim x \right\}, \\ \zeta_{pqst}(x) = \frac{1}{(p+1)(q+1)} \sum_{j=0}^p \sum_{k=0}^q x_{\sigma^j(s), \sigma^k(t)}, \tag{4}$$

while here the limit means  $bp$ -limit. Let us denote by  $\mathcal{V}_\sigma$  the space of  $\sigma$ -convergent double sequences  $x = (x_{jk})$ . For  $\sigma(n) = n + 1$ , the set  $\mathcal{V}_\sigma$  is reduced to the set  $\mathcal{F}$  of almost convergent double sequences [10]. Note that  $\mathcal{E}_{bp} \subset \mathcal{V}_\sigma \subset \mathcal{M}_u$ .

Maddox [11] has defined the concepts of strong almost convergence and  $M$ -convergence for single sequences and established inclusion relation between strong almost convergence,  $M$ -convergence, and almost convergence for single sequence. Başarir [12] extended the notion of strong almost convergence from single sequences to double sequences and proved some interesting results involving this idea and the notion of almost convergence for double sequences. In the recent past, Mursaleen and Mohiuddine [13] presented the notions of absolute and strong  $\sigma$ -convergence for double sequences. A bounded double sequence  $x = (x_{jk})$  is said to be *strongly  $\sigma$ -convergent* if there exists a number  $\ell$  such that

$$\frac{1}{(p+1)(q+1)} \sum_{j=0}^p \sum_{k=0}^q |x_{\sigma^j(s), \sigma^k(t)} - \ell| \rightarrow 0 \\ \text{as } p, q \rightarrow \infty \text{ uniformly in } s, t, \tag{5}$$

while here the limit means  $bp$ -limit. In this case, we write  $[\mathcal{V}_\sigma]\text{-}\lim x = \ell$ . Let us denote by  $[\mathcal{V}_\sigma]$  the set of all strongly  $\sigma$ -convergent sequences  $x = (x_{jk})$ . If  $\sigma$  is translation then  $[\mathcal{V}_\sigma]$  is reduced to the set  $[\mathcal{F}]$  of strong almost convergence double sequences due to Başarir [12].

For more details of spaces for single and double sequences and related concepts, we refer to [14–31] and references therein.

In this paper, we define and study some new spaces involving the idea of invariant mean and  $\sigma$ -convergence for double sequences and establish a relation between these spaces. Further, we extend above spaces to more general spaces by considering the double sequences  $\alpha = (\alpha_{jk})$  such that  $\alpha_{jk} > 0$  for all  $j, k$  and  $\sup_{j,k} \alpha_{jk} = H < \infty$  and prove some topological results.

## 2. The Double Sequence Spaces

We construct the following spaces involving the idea of invariant mean and  $\sigma$ -convergence for double sequences:

$$\mathcal{W}_\sigma = \left\{ x = (x_{jk}) : \frac{1}{(m+1)(n+1)} \right. \\ \left. \times \sum_{p=0}^m \sum_{q=0}^n \zeta_{pqst}(x - \ell E) \rightarrow 0 \text{ as } m, n \rightarrow \infty, \right. \\ \left. \text{uniformly in } s, t, \text{ for some } \ell \right\}, \\ [\mathcal{W}_\sigma] = \left\{ x = (x_{jk}) : \frac{1}{(m+1)(n+1)} \right. \\ \left. \times \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst}(x - \ell E)| \rightarrow 0 \text{ as } m, n \rightarrow \infty, \right. \\ \left. \text{uniformly in } s, t, \text{ for some } \ell \right\}, \\ \mathcal{W}'_\sigma = \left\{ x = (x_{jk}) : \right. \\ \left. \sum_{p=0}^\infty \sum_{q=0}^\infty |\xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t}| \right. \\ \left. \text{converges uniformly in } s, t \right\}, \\ \mathcal{W}''_\sigma = \left\{ x = (x_{jk}) : \right. \\ \left. \sup_{s,t} \sum_{p=0}^\infty \sum_{q=0}^\infty |\xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} \right. \\ \left. + \xi_{p-1,q-1,s,t}| < \infty \right\}, \tag{6}$$

where  $E = (e_{jk})$  with  $e_{jk} = 1$  for all  $j, k$ ;

$$\xi_{mnst} = \xi_{mnst}(x) = \frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n \zeta_{pqst}(x),$$

$$\begin{aligned}
 \xi_{0,0,s,t}(x) &= \zeta_{0,0,s,t} = x_{s,t}, \\
 \xi_{-1,0,s,t}(x) &= \zeta_{-1,0,s,t}(x) = x_{s-1,t}, \\
 \xi_{0,-1,s,t}(x) &= \zeta_{0,-1,s,t}(x) = x_{s,t-1}, \\
 \xi_{-1,-1,s,t}(x) &= \zeta_{-1,-1,s,t}(x) = x_{s-1,t-1}.
 \end{aligned}
 \tag{7}$$

Remark 1. If  $[\mathcal{W}_\sigma]$ - $\lim x = \ell$ , that is,

$$\frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst} - \ell| \rightarrow 0
 \tag{8}$$

as  $m, n \rightarrow \infty$ , uniformly in  $s, t$ ; then

$$\begin{aligned}
 \frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n \left| \frac{1}{p+1} \sum_{j=0}^p \zeta_{jqst} - \ell \right| &\rightarrow 0, \\
 \frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n \left| \frac{1}{q+1} \sum_{k=0}^q \zeta_{pkst} - \ell \right| &\rightarrow 0.
 \end{aligned}
 \tag{9}$$

We remark that by using Abel's transformation for single series

$$\sum_{i=1}^m v_i (u_i \mp u_{i+1}) = \sum_{i=1}^m u_i (v_i \mp v_{i-1}) \mp u_{m+1} v_m.
 \tag{10}$$

We get Abel's transformation for double series

$$\begin{aligned}
 &\sum_{j=1}^p \sum_{k=1}^q v_{jk} (u_{jk} - u_{j+1,k} - u_{j,k+1} + u_{j+1,k+1}) \\
 &= \sum_{j=1}^p \sum_{k=1}^q u_{jk} (\Delta_{11} v_{jk}) - \sum_{j=1}^p u_{j,q+1} (\Delta_{10} v_{jq}) \\
 &\quad - \sum_{k=1}^q u_{p+1,k} (\Delta_{01} v_{pk}) + u_{p+1,q+1} v_{pq},
 \end{aligned}
 \tag{11}$$

where

$$\begin{aligned}
 \Delta_{10} v_{jq} &= v_{jq} - v_{j-1,q}, & \Delta_{01} v_{pk} &= v_{pk} - v_{p,k-1} \\
 \Delta_{11} v_{jk} &= v_{jk} - v_{j-1,k} - v_{j,k-1} + v_{j-1,k-1}.
 \end{aligned}
 \tag{12}$$

In the recent past, Altay and Başar [32] also presented another form of Abel's transformation for double series.

### 3. Inclusion Relations

In the following theorem, we establish a relationship between spaces defined in Section 2. Before proceeding further, first we prove the following lemmas which will be used to prove our inclusion relations.

Lemma 2. Consider that  $[\mathcal{W}_\sigma]$ - $\lim x = \ell$  if and only if

$$\text{(L1) } \mathcal{W}_\sigma\text{-}\lim x = \ell;$$

$$\text{(L2) } (1/uv) \sum_{m=1}^u \sum_{n=1}^v |\Omega_{mnst}^\dagger - \ell| \rightarrow 0 \text{ as } u, v \rightarrow \infty \text{ (uniformly in } s, t);$$

$$\text{(L3) } (1/uv) \sum_{m=1}^u \sum_{n=1}^v |\Omega_{mnst}^\ddagger - \ell| \rightarrow 0 \text{ as } u, v \rightarrow \infty \text{ (uniformly in } s, t);$$

$$\text{(L4) } (1/uv) \sum_{m=1}^u \sum_{n=1}^v |\zeta_{mnst} + \xi_{mnst} - \Omega_{mnst}^\dagger - \Omega_{mnst}^\ddagger| \rightarrow 0 \text{ as } u, v \rightarrow \infty \text{ (uniformly in } s, t),$$

where

$$\Omega_{mnst}^\dagger = \frac{1}{(m+1)} \sum_{p=0}^m \zeta_{pnst},
 \tag{13}$$

$$\Omega_{mnst}^\ddagger = \frac{1}{(n+1)} \sum_{q=0}^n \zeta_{mqst}.$$

Proof. Suppose that  $[\mathcal{W}_\sigma]$ - $\lim x = \ell$ . Thus, we have  $\mathcal{W}_\sigma$ - $\lim x = \ell$ , that is, (L1) holds. We see that conditions (L2) and (L3) follows from the Remark 1. Write

$$\begin{aligned}
 &\frac{1}{uv} \sum_{m=1}^u \sum_{n=1}^v |\zeta_{mnst} + \xi_{mnst} - \Omega_{mnst}^\dagger - \Omega_{mnst}^\ddagger| \\
 &= \frac{1}{uv} \sum_{m=1}^u \sum_{n=1}^v |\zeta_{mnst} - \ell + \xi_{mnst} - \ell - \Omega_{mnst}^\dagger + \ell - \Omega_{mnst}^\ddagger + \ell| \\
 &\leq \frac{1}{uv} \sum_{m=1}^u \sum_{n=1}^v |\zeta_{mnst} - \ell| + \frac{1}{uv} \sum_{m=1}^u \sum_{n=1}^v |\xi_{mnst} - \ell| \\
 &\quad + \frac{1}{uv} \sum_{m=1}^u \sum_{n=1}^v |\Omega_{mnst}^\dagger - \ell| + \frac{1}{uv} \sum_{m=1}^u \sum_{n=1}^v |\Omega_{mnst}^\ddagger - \ell| \\
 &= \Sigma_1 + \Sigma_2 + \Sigma_3 + \Sigma_4 \quad (\text{say}).
 \end{aligned}
 \tag{14}$$

By our assumption, that is,  $[\mathcal{W}_\sigma]$ - $\lim x = \ell$ ,  $\Sigma_1 \rightarrow 0$  as  $u, v \rightarrow \infty$  uniformly in  $s, t$ . The condition (L1) implies that  $\xi_{mnst}$  tends to zero as  $m, n$  tending to  $\infty$  uniformly in  $s, t$ ; therefore  $\Sigma_2 \rightarrow 0$  as  $u, v \rightarrow \infty$  uniformly in  $s, t$  and  $\Sigma_3, \Sigma_4 \rightarrow 0$  as  $u, v \rightarrow \infty$  uniformly in  $s, t$  by the conclusion (L2) and (L3), respectively. Thus, (14) tends to zero as  $u, v \rightarrow \infty$  uniformly in  $s, t$ , that is, (L4) holds.

Conversely, let (L1)–(L4) hold. Then,

$$\begin{aligned}
 &\frac{1}{uv} \sum_{m=1}^u \sum_{n=1}^v |\zeta_{mnst} - \ell| \\
 &\leq \frac{1}{uv} \sum_{m=1}^u \sum_{n=1}^v |\zeta_{mnst} + \xi_{mnst} - \Omega_{mnst}^\dagger - \Omega_{mnst}^\ddagger| \\
 &\quad + \frac{1}{uv} \sum_{m=1}^u \sum_{n=1}^v |\xi_{mnst} - \ell| \\
 &\quad + \frac{1}{uv} \sum_{m=1}^u \sum_{n=1}^v |\Omega_{mnst}^\dagger - \ell| \\
 &\quad + \frac{1}{uv} \sum_{m=1}^u \sum_{n=1}^v |\Omega_{mnst}^\ddagger - \ell|
 \end{aligned}
 \tag{15}$$

$$\rightarrow 0 (u, v \rightarrow \infty) \text{ uniformly in } s, t.$$

□

**Lemma 3.** One has

$$\begin{aligned} & \zeta_{mnst} + \xi_{mnst} - \Omega_{mnst}^\dagger - \Omega_{mnst}^\ddagger \\ &= mn [\xi_{mnst} - \xi_{m-1,n,s,t} - \xi_{m,n-1,s,t} + \xi_{m-1,n-1,s,t}]. \end{aligned} \quad (16)$$

*Proof.* Since

$$\begin{aligned} & \xi_{mnst} - \xi_{m-1,n,s,t} - \xi_{m,n-1,s,t} + \xi_{m-1,n-1,s,t} \\ &= \left[ \frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n \zeta_{pqst} - \frac{1}{m(n+1)} \sum_{p=0}^{m-1} \sum_{q=0}^n \zeta_{pqst} \right] \\ & \quad - \left[ \frac{1}{(m+1)n} \sum_{p=0}^m \sum_{q=0}^{n-1} \zeta_{pqst} - \frac{1}{mn} \sum_{p=0}^{m-1} \sum_{q=0}^{n-1} \zeta_{pqst} \right]. \end{aligned} \quad (17)$$

First, we solve the expression in the first bracket

$$\begin{aligned} & \left[ \frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n \zeta_{pqst} - \frac{1}{m(n+1)} \sum_{p=0}^{m-1} \sum_{q=0}^n \zeta_{pqst} \right] \\ &= \frac{1}{m(m+1)(n+1)} \\ & \quad \times \left[ \sum_{q=0}^n \left( m \sum_{p=0}^m \zeta_{pqst} - (m+1) \sum_{p=0}^{m-1} \zeta_{pqst} \right) \right] \\ &= \frac{1}{m(m+1)(n+1)} \sum_{q=0}^n \left[ m \zeta_{mqst} - \sum_{p=0}^{m-1} \zeta_{pqst} \right] \\ &= \frac{1}{m(m+1)(n+1)} \sum_{q=0}^n \left[ (m+1) \zeta_{mqst} - \sum_{p=0}^m \zeta_{pqst} \right] \\ &= \frac{1}{m(n+1)} \sum_{q=0}^n \left[ \zeta_{mqst} - \frac{1}{(m+1)} \sum_{p=0}^m \zeta_{pqst} \right] \\ &= \frac{1}{m(n+1)} \sum_{q=0}^n \zeta_{mqst} - \frac{1}{m(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n \zeta_{pqst} \\ &= \frac{1}{m(n+1)} \sum_{q=0}^n \zeta_{mqst} - \frac{1}{m} \xi_{mnst}. \end{aligned} \quad (18)$$

Now, the expression in the second bracket

$$\begin{aligned} & \frac{1}{(m+1)n} \sum_{p=0}^m \sum_{q=0}^{n-1} \zeta_{pqst} - \frac{1}{mn} \sum_{p=0}^{m-1} \sum_{q=0}^{n-1} \zeta_{pqst} \\ &= \frac{1}{n} \sum_{q=0}^{n-1} \left[ \frac{1}{m(m+1)} \left\{ m \sum_{p=0}^m \zeta_{pqst} - (m+1) \sum_{p=0}^{m-1} \zeta_{pqst} \right\} \right] \\ &= \frac{1}{n} \sum_{q=0}^{n-1} \left[ \frac{1}{m(m+1)} \left\{ m \zeta_{mqst} - \sum_{p=0}^{m-1} \zeta_{pqst} \right\} \right] \end{aligned}$$

$$\begin{aligned} &= \frac{1}{n} \sum_{q=0}^{n-1} \left[ \frac{1}{m(m+1)} \left\{ (m+1) \zeta_{mqst} - \sum_{p=0}^m \zeta_{pqst} \right\} \right] \\ &= \frac{1}{mn} \sum_{q=0}^{n-1} \zeta_{mqst} - \frac{1}{m(m+1)n} \sum_{p=0}^m \sum_{q=0}^{n-1} \zeta_{pqst} \\ &= \frac{1}{mn} \sum_{q=0}^{n-1} \zeta_{mqst} - \frac{1}{m} \xi_{m,n-1,s,t}. \end{aligned} \quad (19)$$

Substituting (18) and (19) in (17), we get

$$\begin{aligned} & \xi_{mnst} - \xi_{m-1,n,s,t} - \xi_{m,n-1,s,t} + \xi_{m-1,n-1,s,t} \\ &= \frac{1}{m(n+1)} \sum_{q=0}^n \zeta_{mqst} - \frac{1}{mn} \sum_{q=0}^{n-1} \zeta_{mqst} - \frac{1}{m} \xi_{mnst} \\ & \quad + \frac{1}{m} \xi_{m,n-1,s,t} \\ &= \frac{1}{mn(n+1)} \left[ n \sum_{q=0}^n \zeta_{mqst} - (n+1) \sum_{q=0}^{n-1} \zeta_{mqst} \right] \\ & \quad - \frac{1}{m} (\xi_{mnst} - \xi_{m,n-1,s,t}) \\ &= \frac{1}{mn(n+1)} \left[ n \zeta_{mnst} - \sum_{q=0}^{n-1} \zeta_{mqst} \right] \\ & \quad - \frac{1}{m} (\xi_{mnst} - \xi_{m,n-1,s,t}) \\ &= \frac{1}{mn(n+1)} \left[ (n+1) \zeta_{mnst} - \sum_{q=0}^n \zeta_{mqst} \right] \\ & \quad - \frac{1}{m} (\xi_{mnst} - \xi_{m,n-1,s,t}) \\ &= \frac{1}{mn} \left[ \zeta_{mnst} - \frac{1}{(n+1)} \sum_{q=0}^n \zeta_{mqst} \right] \\ & \quad - \frac{1}{m} (\xi_{mnst} - \xi_{m,n-1,s,t}). \end{aligned} \quad (20)$$

We know that

$$\begin{aligned} \xi_{mnst} &= \frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n \zeta_{pqst} \\ &= \frac{1}{(m+1)(n+1)} \left[ \sum_{p=0}^{m-1} \sum_{q=0}^n \zeta_{pqst} + \sum_{q=0}^n \zeta_{mqst} \right], \quad (21) \\ \xi_{m-1,n,s,t} &= \frac{1}{m(n+1)} \sum_{p=0}^{m-1} \sum_{q=0}^n \zeta_{pqst}. \end{aligned}$$

From (21), we have

$$(m + 1) \xi_{mnst} - m \xi_{m-1,n,s,t} = \frac{1}{(n + 1)} \sum_{q=0}^n \zeta_{mqst}. \quad (22)$$

Thus, (20) becomes

$$\begin{aligned} & \xi_{mnst} - \xi_{m-1,n,s,t} - \xi_{m,n-1,s,t} + \xi_{m-1,n-1,s,t} \\ &= \frac{1}{mn} [\zeta_{mnst} - (m + 1) \xi_{mnst} + m \xi_{m-1,n,s,t}] \\ & \quad - \frac{1}{m} (\xi_{mnst} - \xi_{m,n-1,s,t}) \\ &= \frac{1}{mn} [\zeta_{mnst} - \xi_{mnst} - m (\xi_{mnst} - \xi_{m-1,n,s,t}) \\ & \quad - n (\xi_{mnst} - \xi_{m,n-1,s,t})]. \end{aligned} \quad (23)$$

Also (22) can be written as

$$m (\xi_{mnst} - \xi_{m-1,n,s,t}) = \frac{1}{(n + 1)} \sum_{q=0}^n \zeta_{mqst} - \xi_{mnst}. \quad (24)$$

Similarly, we can write

$$n (\xi_{mnst} - \xi_{m,n-1,s,t}) = \frac{1}{(m + 1)} \sum_{q=0}^m \zeta_{pnst} - \xi_{mnst}. \quad (25)$$

Using (24) and (25) in (23), we get

$$\begin{aligned} & \xi_{mnst} - \xi_{m-1,n,s,t} - \xi_{m,n-1,s,t} + \xi_{m-1,n-1,s,t} \\ &= \frac{1}{mn} \left[ \zeta_{mnst} + \xi_{mnst} - \frac{1}{(m + 1)} \sum_{p=0}^m \zeta_{pnst} - \frac{1}{(n + 1)} \sum_{q=0}^n \zeta_{mqst} \right]. \end{aligned} \quad (26)$$

This implies that

$$\begin{aligned} & \zeta_{mnst} + \xi_{mnst} - \Omega_{mnst}^\dagger - \Omega_{mnst}^\ddagger \\ &= mn [\xi_{mnst} - \xi_{m-1,n,s,t} - \xi_{m,n-1,s,t} + \xi_{m-1,n-1,s,t}]. \end{aligned} \quad (27)$$

**Theorem 4.** One has the following inclusions and the limit is preserved in each case:

- (i)  $[\mathcal{V}'_\sigma] \subset [\mathcal{W}'_\sigma] \subset \mathcal{W}'_\sigma$ ,
- (ii)  $\mathcal{W}'_\sigma \subset [\mathcal{W}'_\sigma]$  if the conditions (L2) and (L3) of Lemma 2 hold,
- (iii)  $\mathcal{W}'_\sigma \subset \mathcal{W}''_\sigma$ .

*Proof.* (i) Let  $x \in [\mathcal{V}'_\sigma]$  with  $[\mathcal{V}'_\sigma]$ -lim  $x = \ell$ , say. Then,

$$\zeta_{pqst} (|x - \ell|) \rightarrow 0 \quad \text{as } p, q \rightarrow \infty, \text{ uniformly in } s, t. \quad (28)$$

This implies that

$$\begin{aligned} & \frac{1}{(m + 1)(n + 1)} \sum_{p=0}^m \sum_{q=0}^n \zeta_{pqst} (|x - \ell|) \rightarrow 0 \\ & \text{as } p, q \rightarrow \infty, \text{ uniformly in } s, t. \end{aligned} \quad (29)$$

Also, we have

$$\begin{aligned} & \frac{1}{(m + 1)(n + 1)} \left| \sum_{p=0}^m \sum_{q=0}^n \zeta_{pqst} (x - \ell) \right| \\ & \leq \frac{1}{(m + 1)(n + 1)} \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst} (x - \ell)| \\ & \leq \frac{1}{(m + 1)(n + 1)} \sum_{p=0}^m \sum_{q=0}^n \zeta_{pqst} (|x - \ell|). \end{aligned} \quad (30)$$

Hence,  $[\mathcal{V}'_\sigma] \subset [\mathcal{W}'_\sigma] \subset \mathcal{W}'_\sigma$  and

$$[\mathcal{W}'_\sigma]\text{-lim } x = \mathcal{W}'_\sigma\text{-lim } x = \ell. \quad (31)$$

(ii) We have to show that  $\mathcal{W}'_\sigma \subset [\mathcal{W}'_\sigma]$ . If  $x \in \mathcal{W}'_\sigma$ , then we have

$$\begin{aligned} \mathfrak{S}_{mnst} &= \sum_{p=m}^\infty \sum_{q=n}^\infty |\xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t}| \\ & \rightarrow 0 \end{aligned} \quad (32)$$

as  $m, n \rightarrow \infty$ , uniformly in  $s, t$ ; and

$$\xi_{pqst} \rightarrow \ell \quad (\text{say}) \quad \text{as } p, q \rightarrow \infty, \text{ uniformly in } s, t, \quad (33)$$

that is,  $\mathcal{W}'_\sigma$ -lim  $x = \ell$ .

In order to prove that  $x \in [\mathcal{W}'_\sigma]$ , it is enough to show that condition (L4) of Lemma 2 holds. Now,

$$\begin{aligned} & \mathfrak{S}_{mnst} - \mathfrak{S}_{m,n+1,s,t} - \mathfrak{S}_{m+1,n,s,t} + \mathfrak{S}_{m+1,n+1,s,t} \\ &= \sum_{p=m}^\infty \sum_{q=n}^\infty |\xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t}| \\ & \quad - \sum_{p=m}^\infty \sum_{q=n+1}^\infty |\xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t}| \\ & \quad - \sum_{p=m+1}^\infty \sum_{q=n}^\infty |\xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t}| \\ & \quad + \sum_{p=m+1}^\infty \sum_{q=n+1}^\infty |\xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t}| \\ &= \sum_{p=m}^\infty \left[ \sum_{q=n}^\infty - \sum_{q=n+1}^\infty \right] \\ & \quad \times |\xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t}| \\ & \quad - \sum_{p=m+1}^\infty \left[ \sum_{q=n}^\infty - \sum_{q=n+1}^\infty \right] \\ & \quad \times |\xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t}| \end{aligned}$$

$$\begin{aligned}
&= \sum_{p=m}^{\infty} \left| \xi_{pnst} - \xi_{p-1,n,s,t} - \xi_{p,n-1,s,t} + \xi_{p-1,n-1,s,t} \right| \\
&\quad - \sum_{p=m+1}^{\infty} \left| \xi_{pnst} - \xi_{p-1,n,s,t} - \xi_{p,n-1,s,t} + \xi_{p-1,n-1,s,t} \right| \\
&= \left[ \sum_{p=m}^{\infty} - \sum_{p=m+1}^{\infty} \right] \left| \xi_{pnst} - \xi_{p-1,n,s,t} - \xi_{p,n-1,s,t} + \xi_{p-1,n-1,s,t} \right| \\
&= \left| \xi_{mnst} - \xi_{m-1,n,s,t} - \xi_{m,n-1,s,t} + \xi_{m-1,n-1,s,t} \right|. \tag{34}
\end{aligned}$$

Replacing  $m$  and  $n$  by  $p$  and  $q$ , respectively, we have

$$\begin{aligned}
&\mathfrak{F}_{pqst} - \mathfrak{F}_{p,q+1,s,t} - \mathfrak{F}_{p+1,q,s,t} + \mathfrak{F}_{p+1,q+1,s,t} \\
&= \left| \xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t} \right|. \tag{35}
\end{aligned}$$

By Lemma 3, we have

$$\begin{aligned}
&\zeta_{pqst} + \xi_{pqst} - \Omega_{pqst}^{\dagger} - \Omega_{pqst}^{\ddagger} \\
&= pq \left[ \xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t} \right]. \tag{36}
\end{aligned}$$

So that we have

$$\begin{aligned}
&\frac{1}{mn} \sum_{p=1}^m \sum_{q=1}^n \left| \zeta_{pqst} + \xi_{pqst} - \Omega_{pqst}^{\dagger} - \Omega_{pqst}^{\ddagger} \right| \\
&= \frac{1}{mn} \sum_{p=1}^m \sum_{q=1}^n pq \left[ \mathfrak{F}_{pqst} - \mathfrak{F}_{p,q+1,s,t} \right. \\
&\quad \left. - \mathfrak{F}_{p+1,q,s,t} + \mathfrak{F}_{p+1,q+1,s,t} \right]. \tag{37}
\end{aligned}$$

By using Abel's transformation for double series in the right hand side of above equation, we have

$$\begin{aligned}
&\frac{1}{mn} \sum_{p=1}^m \sum_{q=1}^n \left| \zeta_{pqst} + \xi_{pqst} - \Omega_{pqst}^{\dagger} - \Omega_{pqst}^{\ddagger} \right| \\
&= \frac{1}{mn} \left[ \sum_{p=1}^m \sum_{q=1}^n \mathfrak{F}_{pqst} - m \sum_{q=1}^n \mathfrak{F}_{m+1,q,s,t} \right. \\
&\quad \left. - n \sum_{p=1}^m \mathfrak{F}_{p,n+1,s,t} + mn \mathfrak{F}_{m+1,n+1,s,t} \right] \tag{38}
\end{aligned}$$

$\rightarrow 0$  as  $m, n \rightarrow \infty$ , uniformly in  $s, t$  (by (32)). Hence, by Lemma 2,  $x \in [\mathcal{W}'_{\sigma}]$ .

(iii) Let  $x \in \mathcal{W}'_{\sigma}$ , and we have to show that

$$\sup_{s,t} \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} \left| \xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t} \right| \leq K, \tag{39}$$

where  $K$  is an absolute constant. Since  $x \in \mathcal{W}'_{\sigma}$ , there exist integers  $p_0, q_0$  such that

$$\sum_{p>p_0} \sum_{q>q_0} \left| \xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t} \right| < 1, \quad \forall s, t. \tag{40}$$

Hence, it is left to show that for fixed  $p, q$

$$\left| \xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t} \right| \leq K, \quad \forall s, t. \tag{41}$$

From (40), we have

$$\left| \xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t} \right| < 1, \tag{42}$$

for every fixed  $p > p_0, q > q_0$  and for all  $s, t$ . Since

$$\begin{aligned}
&m(m+1)n(n+1) \left( \xi_{mnst} - \xi_{m-1,n,s,t} - \xi_{m,n-1,s,t} + \xi_{m-1,n-1,s,t} \right) \\
&= \sum_{p=1}^m \sum_{q=1}^n pq \left( \zeta_{pqst} - \zeta_{p-1,q,s,t} - \zeta_{p,q-1,s,t} + \zeta_{p-1,q-1,s,t} \right). \tag{43}
\end{aligned}$$

Accordingly,

$$\begin{aligned}
&mn \left[ (m+1)(n+1) \right. \\
&\quad \times \left( \xi_{mnst} - \xi_{m-1,n,s,t} - \xi_{m,n-1,s,t} + \xi_{m-1,n-1,s,t} \right) \\
&\quad - (m-1)(n+1) \\
&\quad \times \left( \xi_{m-1,n,s,t} - \xi_{m-2,n,s,t} - \xi_{m-1,n-1,s,t} + \xi_{m-2,n-1,s,t} \right) \\
&\quad - (m+1)(n-1) \\
&\quad \times \left( \xi_{m,n-1,s,t} - \xi_{m-1,n-1,s,t} - \xi_{m,n-2,s,t} + \xi_{m-1,n-2,s,t} \right) \\
&\quad \left. + (m-1)(n-1) \right. \\
&\quad \left. \times \left( \xi_{m-1,n-1,s,t} - \xi_{m-2,n-1,s,t} - \xi_{m-1,n-2,s,t} + \xi_{m-2,n-2,s,t} \right) \right] \\
&= \sum_{q=1}^n q \left[ \sum_{p=1}^m p \left( \zeta_{pqst} - \zeta_{p-1,q,s,t} - \zeta_{p,q-1,s,t} + \zeta_{p-1,q-1,s,t} \right) \right. \\
&\quad \left. - \sum_{p=1}^{m-1} p \left( \zeta_{pqst} - \zeta_{p-1,q,s,t} - \zeta_{p,q-1,s,t} + \zeta_{p-1,q-1,s,t} \right) \right] \\
&\quad - \sum_{q=1}^{n-1} q \left[ \sum_{p=1}^m p \left( \zeta_{pqst} - \zeta_{p-1,q,s,t} - \zeta_{p,q-1,s,t} + \zeta_{p-1,q-1,s,t} \right) \right. \\
&\quad \left. - \sum_{p=1}^{m-1} p \left( \zeta_{pqst} - \zeta_{p-1,q,s,t} \right. \right. \\
&\quad \left. \left. - \zeta_{p,q-1,s,t} + \zeta_{p-1,q-1,s,t} \right) \right] \\
&= \sum_{q=1}^n q \left[ m \left( \zeta_{mqst} - \zeta_{m-1,q,s,t} - \zeta_{m,q-1,s,t} + \zeta_{m-1,q-1,s,t} \right) \right. \\
&\quad \left. - \sum_{q=1}^{n-1} q \left[ m \left( \zeta_{mqst} - \zeta_{m-1,q,s,t} - \zeta_{m,q-1,s,t} + \zeta_{m-1,q-1,s,t} \right) \right] \right] \tag{44}
\end{aligned}$$

This implies that

$$\begin{aligned}
 & (m+1)(n+1)(\xi_{mnst} - \xi_{m-1,n,s,t} - \xi_{m,n-1,s,t} + \xi_{m-1,n-1,s,t}) \\
 & - (m-1)(n+1) \\
 & \times (\xi_{m-1,n,s,t} - \xi_{m-2,n,s,t} - \xi_{m-1,n-1,s,t} + \xi_{m-2,n-1,s,t}) \\
 & - (m+1)(n-1) \\
 & \times (\xi_{m,n-1,s,t} - \xi_{m-1,n-1,s,t} - \xi_{m,n-2,s,t} + \xi_{m-1,n-2,s,t}) \\
 & + (m-1)(n-1) \\
 & \times (\xi_{m-1,n-1,s,t} - \xi_{m-2,n-1,s,t} - \xi_{m-1,n-2,s,t} + \xi_{m-2,n-2,s,t}) \\
 & = [\zeta_{mnst} - \zeta_{m-1,n,s,t} - \zeta_{m,n-1,s,t} + \zeta_{m-1,n-1,s,t}]. \tag{45}
 \end{aligned}$$

Using (42) and (45), we have

$$|\zeta_{mnst} - \zeta_{m-1,n,s,t} - \zeta_{m,n-1,s,t} + \zeta_{m-1,n-1,s,t}| \leq K(m, n) \tag{46}$$

for every fixed  $m > p_0$ ,  $n > q_0$  and for all  $s, t$ , where  $K(m, n)$  is a constant depending upon  $m, n$ .

Now, for any given infinite double series  $\sum_s \sum_t a_{st}$  denoted as “ $a$ ”, let us write

$$x_{jk} = \sum_{s=0}^j \sum_{t=0}^k a_{st}, \quad j, k = 1, 2, \dots \tag{47}$$

and  $\sigma$  be monotonically increasing. For simplicity in notation, we denote

$$\begin{aligned}
 \phi_{mnst} &= \zeta_{mnst}(a) - \zeta_{m-1,n,s,t}(a) - \zeta_{m,n-1,s,t}(a) \\
 & + \zeta_{m-1,n-1,s,t}(a). \tag{48}
 \end{aligned}$$

Again from the definition of  $\zeta_{mnst}$ , it is easy to obtain

$$\phi_{mnst} = \frac{1}{m(m+1)n(n+1)} \sum_{j=1}^m \sum_{k=1}^n jk \left( \sum_{u=d_j}^{d'_j} \sum_{v=h_k}^{h'_k} a_{uv} \right) \tag{49}$$

for all  $m, n \geq 1$  and  $\phi_{0,0,s,t}(a) = a_{st}$  with  $d_j = \sigma^{j-1}(s) + 1$ ,  $d'_j = \sigma^j(s)$ ,  $h_k = \sigma^{k-1}(t) + 1$ ,  $h'_k = \sigma^k(t)$ . Further, we calculate

$$\begin{aligned}
 & m(m+1)n(n+1)\phi_{mnst} - m(m-1)n(n+1)\phi_{m-1,n,s,t} \\
 & - m(m+1)n(n-1)\phi_{m,n-1,s,t} \\
 & + m(m-1)n(n-1)\phi_{m-1,n-1,s,t}
 \end{aligned}$$

$$\begin{aligned}
 & = \left\{ \sum_{j=1}^m \sum_{k=1}^n jk - \sum_{j=1}^{m-1} \sum_{k=1}^n jk - \sum_{j=1}^m \sum_{k=1}^{n-1} jk + \sum_{j=1}^{m-1} \sum_{k=1}^{n-1} jk \right\} \\
 & \times \left( \sum_{u=d_j}^{d'_j} \sum_{v=h_k}^{h'_k} a_{uv} \right) \\
 & = \left\{ \sum_{k=1}^n k \left[ \sum_{j=1}^m j - \sum_{j=1}^{m-1} j \right] - \sum_{k=1}^{n-1} k \left[ \sum_{j=1}^m j + \sum_{j=1}^{m-1} j \right] \right\} \\
 & \times \left( \sum_{u=d_j}^{d'_j} \sum_{v=h_k}^{h'_k} a_{uv} \right) \\
 & = \left\{ m \sum_{k=1}^n k - m \sum_{k=1}^{n-1} k \right\} \left( \sum_{u=d_m}^{d'_m} \sum_{v=h_n}^{h'_n} a_{uv} \right) \\
 & = mn \left( \sum_{u=d_m}^{d'_m} \sum_{v=h_n}^{h'_n} a_{uv} \right). \tag{50}
 \end{aligned}$$

Thus, we have

$$\begin{aligned}
 \sum_{u=d_m}^{d'_m} \sum_{v=h_n}^{h'_n} a_{uv} &= (m+1)(n+1)\phi_{m,n,s,t} \\
 & - (m-1)(n+1)\phi_{m-1,n,s,t} \\
 & - (m+1)(n-1)\phi_{m,n-1,s,t} \\
 & + (m-1)(n-1)\phi_{m-1,n-1,s,t}. \tag{51}
 \end{aligned}$$

Hence, it follows from (46) that for each fixed  $m > p_0$ ,  $n > q_0$ ,

$$\left| \sum_{u=d_m}^{d'_m} \sum_{v=h_n}^{h'_n} a_{uv} \right| \leq K(m, n), \quad \forall s, t. \tag{52}$$

Hence, it follows from (52) that

$$|a_{uv}| \leq K \quad \forall u, v, \tag{53}$$

where  $K$  is independent of  $u, v$ . By (49), we have

$$\begin{aligned}
 & |\zeta_{mnst} - \zeta_{m-1,n,s,t} - \zeta_{m,n-1,s,t} + \zeta_{m-1,n-1,s,t}| \\
 & \leq K, \quad \forall m, n, s, t. \tag{54}
 \end{aligned}$$

Also from (43) and (54), we have

$$|\xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t}| \leq K, \quad \forall p, q, s, t. \tag{55}$$

□

### 4. Topological Results

Here, we extend the spaces  $[\mathcal{W}_\sigma]$ ,  $\mathcal{W}'_\sigma$ ,  $\mathcal{W}''_\sigma$  to more general spaces, respectively, denoted by  $[\mathcal{W}_\sigma(\alpha)]$ ,  $\mathcal{W}'_\sigma(\alpha)$ ,  $\mathcal{W}''_\sigma(\alpha)$ ,

where  $\alpha = (\alpha_{jk})$  is a double sequence of positive real numbers for all  $j, k$  and  $\sup_{j,k} \alpha_{jk} = H < \infty$ . First, we recall the notion of paranorm as follows.

A *paranorm* is a function  $g : X \rightarrow \mathbb{R}$  defined on a linear space  $X$  such that for all  $x, y, z \in X$

(P1)  $g(x) = 0$  if  $x = \theta$

(P2)  $g(-x) = g(x)$

(P3)  $g(x + y) \leq g(x) + g(y)$

(P4) If  $(\alpha_n)$  is a sequence of scalars with  $\alpha_n \rightarrow \alpha_0$  ( $n \rightarrow \infty$ ) and  $x_n, a \in X$  with  $x_n \rightarrow a$  ( $n \rightarrow \infty$ ) in the sense that  $g(x_n - a) \rightarrow 0$  ( $n \rightarrow \infty$ ), then  $\alpha_n x_n \rightarrow \alpha_0 a$  ( $n \rightarrow \infty$ ), in the sense that  $g(\alpha_n x_n - \alpha_0 a) \rightarrow 0$  ( $n \rightarrow \infty$ ).

A paranorm  $g$  for which  $g(x) = 0$  implies  $x = \theta$  is called a *total paranorm* on  $X$ , and the pair  $(X, g)$  is called a *total paranormed space*. Note that each seminorm (norm)  $p$  on  $X$  is a paranorm (total) but converse needs not be true.

A paranormed space  $(X, g)$  is a topological linear space with the topology given by the paranorm  $g$ .

Now, we define the following spaces:

$$[\mathcal{W}_\sigma(\alpha)] = \left\{ x = (x_{jk}) : \frac{1}{(m+1)(n+1)} \times \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst}(x - \ell E)|^{\alpha_{pq}} \rightarrow 0, \right. \\ \left. \text{as } m, n \rightarrow \infty, \text{ uniformly in } s, t \right\},$$

$$\mathcal{W}'_\sigma(\alpha) = \left\{ x = (x_{jk}) : \sum_{p=0}^\infty \sum_{q=0}^\infty |\xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t}|^{\alpha_{pq}} \right. \\ \left. \text{converges uniformly in } s, t \right\},$$

$$\mathcal{W}''_\sigma(\alpha) = \left\{ x = (x_{jk}) : \sup_{s,t} \sum_{p=0}^\infty \sum_{q=0}^\infty |\xi_{pqst} - \xi_{p-1,q,s,t} - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t}|^{\alpha_{pq}} < \infty \right\}.$$

(56)

We remark that if  $\alpha = (\alpha_{jk})$  is a constant sequence, then we write  $[\mathcal{W}_{\sigma,\alpha}]$  in place of  $[\mathcal{W}_\sigma(\alpha)]$ .

**Theorem 5.** Let  $\alpha = (\alpha_{pq})$  be a bounded double sequence of strictly positive real numbers. Then,  $[\mathcal{W}_\sigma(\alpha)]$  is a complete linear topological space paranormed by

$$g(x) = \sup_{m,n,s,t} \left( \frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst}(x)|^{\alpha_{pq}} \right)^{1/M},$$

(57)

where  $M = \max(1, \sup_{p,q} \alpha_{pq})$ . If  $\alpha \geq 1$  then  $[\mathcal{W}_{\sigma,\alpha}]$  is a Banach space and  $[\mathcal{W}_{\sigma,\alpha}]$  is a  $\alpha$ -normed space if  $0 < \alpha < 1$ .

*Proof.* Let  $(x_{pq})$  and  $(y_{pq})$  be two double sequences. Then,

$$|x_{pq} + y_{pq}|^{\alpha_{pq}} \leq K (|x_{pq}|^{\alpha_{pq}} + |y_{pq}|^{\alpha_{pq}}),$$

(58)

where  $K = \max(1, 2^{H-1})$  and  $H = \sup_{p,q} \alpha_{pq}$ . Since

$$|\lambda|^{\alpha_{pq}} \leq \max(1, |\lambda|^H),$$

(59)

therefore

$$\frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst}(\lambda x + \mu y)|^{\alpha_{pq}} \\ \leq KK_1 \frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst}(x)|^{\alpha_{pq}} \\ + KK_2 \frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst}(y)|^{\alpha_{pq}},$$

(60)

where  $K_1 = \max(1, |\lambda|^H)$  and  $K_2 = \max(1, |\mu|^H)$ . From (60), we have that if  $x, y \in [\mathcal{W}_\sigma(\alpha)]$ , then  $\lambda x + \mu y \in [\mathcal{W}_\sigma(\alpha)]$ . Thus,  $[\mathcal{W}_\sigma(\alpha)]$  is a linear space. Without loss of generality, we can take

$$[\mathcal{W}_\sigma(\alpha)]\text{-}\lim x = 0.$$

(61)

Clearly,  $g(0) = 0, g(-x) = g(x)$ . From (58) and Minkowski's inequality, we have

$$\left( \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst}(x+y)|^{\alpha_{pq}} \right)^{1/M} \\ \leq \left[ \sum_{p=0}^m \sum_{q=0}^n (|\zeta_{pqst}(x)|^{\alpha_{pq}/M} + |\zeta_{pqst}(y)|^{\alpha_{pq}/M}) \right]^{1/M} \\ \leq \left( \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst}(x)|^{\alpha_{pq}} \right)^{1/M} \\ + \left( \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst}(y)|^{\alpha_{pq}} \right)^{1/M}.$$

(62)

Hence,

$$g(x+y) \leq g(x) + g(y).$$

(63)



Since  $\alpha = (\alpha_{mn})$  is bounded away from zero, there exists a constant  $\delta > 0$  such that  $\alpha_{mn} \geq \delta$  for all  $m, n$ . Now for  $|\lambda| \leq 1$ ,  $|\lambda|^{\alpha_{st}} \leq |\lambda|^\delta$  and so

$$g(\lambda x) \leq |\lambda|^{\delta/M} g(x), \tag{64}$$

that is, the scalar multiplication is continuous. Hence,  $g$  is a paranorm on  $[\mathcal{W}'_\sigma(\alpha)]$ .

Let  $(x^b)$  be a Cauchy sequence in  $[\mathcal{W}'_\sigma(\alpha)]$ , that is,

$$g(x^d - x^b) \rightarrow 0 \quad \text{as } b, d \rightarrow \infty. \tag{65}$$

Since

$$\frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst}(x^d - x^b)|^{\alpha_{pq}} \leq [g(x^d - x^b)]^M, \tag{66}$$

it follows that

$$|\zeta_{pqst}(x^d - x^b)|^{\alpha_{pq}} \rightarrow 0 \quad \text{as } b, d \rightarrow \infty \quad \forall p, q, s, t. \tag{67}$$

In particular,

$$\begin{aligned} |\zeta_{0,0,s,t}(x^d - x^b)| &= |x_{st}^d - x_{st}^b| \rightarrow 0 \\ &\text{as } b, d \rightarrow \infty \text{ for fixed } s, t. \end{aligned} \tag{68}$$

Hence,  $(x^d)$  is a Cauchy sequence in  $\mathbb{C}$ . Since  $\mathbb{C}$  is complete, there exists  $x = (x_{jk}) \in \mathbb{C}$  such that  $x^d \rightarrow x$  coordinatewise as  $d \rightarrow \infty$ . It follows from (66) that given  $\epsilon > 0$ , there exists  $d_0$  such that

$$\left( \frac{1}{(m+1)(n+1)} \sum_{p=0}^m \sum_{q=0}^n |\zeta_{pqst}(x^d - x^b)|^{\alpha_{pq}} \right)^{1/M} < \epsilon, \tag{69}$$

for  $b, d > d_0$ . Now taking  $b \rightarrow \infty$  and  $\sup_{s,t,m,n}$  in (69), we have  $g(x^d - x) \leq \epsilon$  for  $d > d_0$ . This proves that  $x^d \rightarrow x$  and  $x = (x_{jk}) \in [\mathcal{W}'_\sigma(\alpha)]$ . Hence  $[\mathcal{W}'_\sigma(\alpha)]$  is complete. If  $\alpha$  is a constant then it is easy to prove the rest of the theorem.  $\square$

**Theorem 6.** *One has the following:*

$\mathcal{W}'_\sigma(\alpha)$  is a complete paranormed space, paranormed by

$$\begin{aligned} h(x) = \sup_{s,t} \left( \sum_{p=0}^\infty \sum_{q=0}^\infty |\xi_{pqst} - \xi_{p-1,q,s,t} \right. \\ \left. - \xi_{p,q-1,s,t} + \xi_{p-1,q-1,s,t}|^{\alpha_{pq}} \right)^{1/M} \end{aligned} \tag{70}$$

which is defined on  $\mathcal{W}'_\sigma(\alpha)$ . If  $\alpha \geq 1$  then  $\mathcal{W}'_{\sigma,\alpha}$  is a Banach space and if  $0 < \alpha < 1$ ,  $\mathcal{W}'_{\sigma,\alpha}$  is  $\alpha$ -normed space.

*Proof.* In order for the paranorm in (70) be defined, we require that

$$\mathcal{W}'_\sigma(\alpha) \subset \mathcal{W}''_\sigma(\alpha) \tag{71}$$

which is proved in the next theorem (i.e., Theorem 7). Using the standard technique as in the previous theorem, we can prove that  $h$  is subadditive.

Now, we have to prove the continuity of scalar multiplication. Suppose that  $x = (x_{jk}) \in \mathcal{W}'_\sigma(\alpha)$ . Then, for  $\epsilon > 0$  there exist integers  $M, N > 0$  such that

$$\begin{aligned} \sum_{p=M}^\infty \sum_{q=N}^\infty |\xi_{pqst}(x) - \xi_{p-1,q,s,t}(x) \\ - \xi_{p,q-1,s,t}(x) + \xi_{p-1,q-1,s,t}(x)|^{\alpha_{pq}} < \epsilon. \end{aligned} \tag{72}$$

If  $|\lambda| \leq 1$ , then by (72) we have

$$\begin{aligned} \sum_{p=M}^\infty \sum_{q=N}^\infty |\xi_{pqst}(\lambda x) - \xi_{p-1,q,s,t}(\lambda x) \\ - \xi_{p,q-1,s,t}(\lambda x) + \xi_{p-1,q-1,s,t}(\lambda x)|^{\alpha_{pq}} \\ \leq \sum_{p=M}^\infty \sum_{q=N}^\infty |\xi_{pqst}(x) - \xi_{p-1,q,s,t}(x) \\ - \xi_{p,q-1,s,t}(x) + \xi_{p-1,q-1,s,t}(x)|^{\alpha_{pq}} < \epsilon. \end{aligned} \tag{73}$$

Since for fixed  $M, N$

$$\begin{aligned} \sum_{p=0}^{M-1} \sum_{q=0}^{N-1} |\xi_{pqst}(\lambda x) - \xi_{p-1,q,s,t}(\lambda x) - \xi_{p,q-1,s,t}(\lambda x) \\ + \xi_{p-1,q-1,s,t}(\lambda x)|^{\alpha_{pq}} \rightarrow 0 \end{aligned} \tag{74}$$

as  $\lambda \rightarrow 0$ , it follows from (73) and (74) that for fixed  $x = (x_{jk})$ ,  $h(\lambda x) \rightarrow 0$  as  $\lambda \rightarrow 0$ . Also, since  $|\lambda|^{\alpha_{pq}} \leq \max(1, |\lambda|^H)$  implies that

$$h(\lambda x) \leq (\sup |\lambda|^{\alpha_{pq}})^{1/M} h(x). \tag{75}$$

It follows that for fixed  $\lambda$ ,  $h(\lambda x) \rightarrow 0$  as  $x \rightarrow 0$ . This proves the continuity of scalar multiplication. Hence,  $h$  is a paranorm. The proof of the completeness of  $\mathcal{W}'_\sigma(\alpha)$  can be achieved by using the same technique as in Theorem 5.  $\square$

**Theorem 7.** *Suppose that  $\alpha = (\alpha_{pq})$  is bounded double sequence of strictly positive real numbers. Then,*

- (i)  $\mathcal{W}''_\sigma(\alpha)$  is a complete linear space paranormed by the function  $h$  defined in (70). In particular, if  $\alpha$  is a constant,  $\mathcal{W}''_{\sigma,\alpha}$  is a Banach space for  $\alpha \geq 1$  and  $\alpha$ -normed space for  $0 < \alpha < 1$ ,
- (ii)  $\mathcal{W}'_\sigma(\alpha)$  is a closed subspace of  $\mathcal{W}''_\sigma(\alpha)$ .

*Proof.* (i) Proceeding along the same lines as in Theorem 6, except for the proof of continuity of scalar multiplication. If  $x = (x_{jk}) \in \mathcal{W}''_\sigma(\alpha)$ , we cannot assert (72) as in the case when  $x = (x_{jk}) \in \mathcal{W}'_\sigma(\alpha)$ . Since  $\alpha$  is bounded away from zero, there

exists  $\delta > 0$  such that  $\alpha_{mn} \geq \delta$  for all  $m, n$ . Hence,  $|\lambda| \leq 1$  implies  $|\lambda|^{\alpha_{pq}} \leq |\lambda|^\delta$ . Since

$$h(\lambda x) \leq |\lambda|^{\alpha_{pq}} h(x), \quad (76)$$

continuity of scalar multiplication follows.

(ii) For this, first we have to show that

$$\mathcal{W}'_\sigma(\alpha) \subset \mathcal{W}''_\sigma(\alpha). \quad (77)$$

Let  $x = (x_{jk}) \in \mathcal{W}'_\sigma(\alpha)$ . Then, there exist integers  $M, N$  such that

$$\sum_{p=M}^{\infty} \sum_{q=N}^{\infty} |\xi_{pqst}(x) - \xi_{p-1,q,s,t}(x) - \xi_{p,q-1,s,t}(x) + \xi_{p-1,q-1,s,t}(x)|^{\alpha_{pq}} < 1. \quad (78)$$

By an argument similar to Theorem 4(iii), we obtain (77). Since  $\mathcal{W}'_\sigma(\alpha)$  and  $\mathcal{W}''_\sigma(\alpha)$  are complete with the same metric, we have (ii).  $\square$

## Acknowledgement

This work was funded by the Deanship of Scientific Research (DSR), King Abdulaziz University, Jeddah, under Grant no. (130-068-D1433). The authors, therefore, acknowledge with thanks DSR technical and financial support.

## References

- [1] A. Pringsheim, "Zur Theorie der zweifach unendlichen Zahlenfolgen," *Mathematische Annalen*, vol. 53, no. 3, pp. 289–321, 1900.
- [2] M. Mursaleen, "On some new invariant matrix methods of summability," *The Quarterly Journal of Mathematics*, vol. 34, no. 133, pp. 77–86, 1983.
- [3] P. Schaefer, "Infinite matrices and invariant means," *Proceedings of the American Mathematical Society*, vol. 36, pp. 104–110, 1972.
- [4] G. G. Lorentz, "A contribution to the theory of divergent sequences," *Acta Mathematica*, vol. 80, pp. 167–190, 1948.
- [5] S. A. Mohiuddine, "An application of almost convergence in approximation theorems," *Applied Mathematics Letters*, vol. 24, no. 11, pp. 1856–1860, 2011.
- [6] C. Çakan, B. Altay, and M. Mursaleen, "The  $\sigma$ -convergence and  $\sigma$ -core of double sequences," *Applied Mathematics Letters*, vol. 19, no. 10, pp. 1122–1128, 2006.
- [7] M. Mursaleen and S. A. Mohiuddine, "Double  $\sigma$ -multiplicative matrices," *Journal of Mathematical Analysis and Applications*, vol. 327, no. 2, pp. 991–996, 2007.
- [8] M. Mursaleen and S. A. Mohiuddine, "Regularly  $\sigma$ -conservative and  $\sigma$ -coercive four dimensional matrices," *Computers & Mathematics with Applications*, vol. 56, no. 6, pp. 1580–1586, 2008.
- [9] M. Mursaleen and S. A. Mohiuddine, "On  $\sigma$ -conservative and boundedly  $\sigma$ -conservative four-dimensional matrices," *Computers & Mathematics with Applications*, vol. 59, no. 2, pp. 880–885, 2010.
- [10] F. Móricz and B. E. Rhoades, "Almost convergence of double sequences and strong regularity of summability matrices," *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 104, no. 2, pp. 283–294, 1988.
- [11] I. J. Maddox, "A new type of convergence," *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 83, no. 1, pp. 61–64, 1978.
- [12] M. Başarır, "On the strong almost convergence of double sequences," *Periodica Mathematica Hungarica*, vol. 30, no. 3, pp. 177–181, 1995.
- [13] M. Mursaleen and S. A. Mohiuddine, "Some new double sequence spaces of invariant means," *Glasnik Matematički*, vol. 45, no. 65, pp. 139–153, 2010.
- [14] A. Alotaibi, M. Mursaleen, and M. A. Alghamdi, "Invariant and absolute invariant means of double sequences," *Journal of Function Spaces and Applications*, vol. 2012, Article ID 465364, 9 pages, 2012.
- [15] C. Aydin and F. Başar, "Some new paranormed sequence spaces," *Information Sciences*, vol. 160, no. 1–4, pp. 27–40, 2004.
- [16] C. Aydın and F. Başar, "Some new difference sequence spaces," *Applied Mathematics and Computation*, vol. 157, no. 3, pp. 677–693, 2004.
- [17] F. Başar and M. Kirişçi, "Almost convergence and generalized difference matrix," *Computers & Mathematics with Applications*, vol. 61, no. 3, pp. 602–611, 2011.
- [18] C. Çakan, B. Altay, and H. Coşkun, " $\sigma$ -regular matrices and a  $\sigma$ -core theorem for double sequences," *Hacettepe Journal of Mathematics and Statistics*, vol. 38, no. 1, pp. 51–58, 2009.
- [19] G. Das and S. K. Sahoo, "On some sequence spaces," *Journal of Mathematical Analysis and Applications*, vol. 164, no. 2, pp. 381–398, 1992.
- [20] K. Kayaduman and C. Çakan, "The cesàro core of double sequences," *Abstract and Applied Analysis*, vol. 2011, Article ID 950364, 9 pages, 2011.
- [21] M. Mursaleen, "Almost strongly regular matrices and a core theorem for double sequences," *Journal of Mathematical Analysis and Applications*, vol. 293, no. 2, pp. 523–531, 2004.
- [22] M. Mursaleen, "Some matrix transformations on sequence spaces of invariant means," *Hacettepe Journal of Mathematics and Statistics*, vol. 38, no. 3, pp. 259–264, 2009.
- [23] M. Mursaleen and O. H. H. Edely, "Almost convergence and a core theorem for double sequences," *Journal of Mathematical Analysis and Applications*, vol. 293, no. 2, pp. 532–540, 2004.
- [24] M. Mursaleen, A. M. Jarrah, and S. A. Mohiuddine, "Almost convergence through the generalized de la Vallée-Poussin mean," *Iranian Journal of Science and Technology. Transaction A*, vol. 33, no. 2, pp. 169–177, 2009.
- [25] M. Mursaleen and S. A. Mohiuddine, "Almost bounded variation of double sequences and some four dimensional summability matrices," *Publicationes Mathematicae Debrecen*, vol. 75, no. 3-4, pp. 495–508, 2009.
- [26] M. Mursaleen and S. A. Mohiuddine, "Some inequalities on sublinear functionals related to the invariant mean for double sequences," *Mathematical Inequalities & Applications*, vol. 13, no. 1, pp. 157–163, 2010.
- [27] M. Mursaleen and S. A. Mohiuddine, "Invariant mean and some core theorems for double sequences," *Taiwanese Journal of Mathematics*, vol. 14, no. 1, pp. 21–33, 2010.
- [28] M. Mursaleen and S. A. Mohiuddine, "Some matrix transformations of convex and paranormed sequence spaces into the spaces of invariant means," *Journal of Function Spaces and Applications*, vol. 2012, Article ID 612671, 10 pages, 2012.
- [29] M. Mursaleen and S. A. Mohiuddine, "Banach limit and some new spaces of double sequences," *Turkish Journal of Mathematics*, vol. 36, no. 1, pp. 121–130, 2012.

- [30] M. Zeltser, M. Mursaleen, and S. A. Mohiuddine, "On almost conservative matrix methods for double sequence spaces," *Publicationes Mathematicae Debrecen*, vol. 75, no. 3-4, pp. 387–399, 2009.
- [31] G. A. Anastassiou, M. Mursaleen, and S. A. Mohiuddine, "Some approximation theorems for functions of two variables through almost convergence of double sequences," *Journal of Computational Analysis and Applications*, vol. 13, no. 1, pp. 37–46, 2011.
- [32] B. Altay and F. Başar, "Some new spaces of double sequences," *Journal of Mathematical Analysis and Applications*, vol. 309, no. 1, pp. 70–90, 2005.