

Research Article

Local C^r Stability for Iterative Roots of Orientation-Preserving Self-Mappings on the Interval

Yingying Zeng

Department of Mathematics, Sichuan University, Chengdu, Sichuan 610064, China

Correspondence should be addressed to Yingying Zeng; mathyyz@163.com

Received 9 April 2014; Accepted 1 May 2014; Published 21 May 2014

Academic Editor: Yongkun Li

Copyright © 2014 Yingying Zeng. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Stability of iterative roots is important in their numerical computation. It is known that under some conditions iterative roots of orientation-preserving self-mappings are both globally C^0 stable and locally C^1 stable but globally C^1 unstable. Although the global C^1 instability implies the general global C^r ($r \geq 2$) instability, the local C^1 stability does not guarantee the local C^r ($r \geq 2$) stability. In this paper we generally prove the local C^r ($r \geq 2$) stability for iterative roots. For this purpose we need a uniform estimate for the approximation to the conjugation in C^r linearization, which is given by improving the method used for the C^1 case.

1. Introduction

Let $I := [0, 1]$ and let $C^r(I, I)$, $r \geq 0$, be the set of all C^r self-mappings defined on I . An iterative root of order k of a given self-mapping $F : I \rightarrow I$ is a self-mapping $f : I \rightarrow I$ such that



$$f^k(x) = F(x), \quad \forall x \in I, \quad (1)$$

where f^k denotes the k th iterate of f , defined by $f^k(x) := f(f^{k-1}(x))$ and $f^0(x) = x$ for all $x \in I$ inductively. The study of iterative roots was started long ago, at least about two hundred years ago when Babbage published his paper [1]. In recent decades, regarded as a weak version of the embedding flow problem for dynamical systems [2, 3], the problem of iterative roots attracted great attention in the field of dynamical systems [3, 4] and functional equations [5–8]. Based on the work for monotonic mappings [6, 7], advances have been made to nonmonotonic cases [8–11], self-mappings on circles [12, 13], set-valued functions [14, 15], and high-dimensional mappings [16, 17].

Because of the potential in extensive applications (e.g., to information processing [18, 19] and graph theory [20]), numerical computation [21, 22] of iterative roots became an important task, which demands approximation to iterative roots and considers stability of iterative roots. In 2007 Xu

and Zhang [23] proved C^0 stability for iterative roots on a closed interval with exact one fixed point at an endpoint. This result is substantially a local C^0 stability because the stability is totally decided by the behaviors of the iterative root in a sufficiently small neighborhood of the fixed point. In [24] results of global C^0 stability are given, where the global sense means the stability on a closed interval bounded by two fixed points. Recently, it was proved in [25] that iterative roots of every orientation-preserving self-mapping on the interval are locally C^1 stable but globally C^1 unstable.

In this paper we generally consider C^r ($r \geq 1$) stability of iterative roots. It is clear that the global C^1 instability given in [25] implies the general global C^r ($r \geq 2$) instability because C^1 approximation is the most fundamental requirement for C^r approximation. However, the above result of local C^1 stability does not guarantee the local C^r ($r \geq 2$) stability. In this paper we concentrate on the local C^r ($r \geq 2$) stability for iterative roots of orientation-preserving self-mapping on I . Clearly, the given mapping is a strictly increasing function. The local C^r ($r \geq 2$) stability is proved by approximation to the conjugation in C^r linearization. In order to give an estimate to the approximation uniformly with respect to the order of iteration, we improve the method used in [25] to obtain lower growth rate for given functions under iteration.

FIGURE 1: $h \in \mathcal{H}_-^r(\lambda)$.FIGURE 2: $h \in \mathcal{H}_+^r(\lambda)$.

2. Preliminaries

In order to state our results clearly, we first pay attention to the existence of C^r ($r \geq 2$) iterative roots of increasing C^r self-mappings on a compact interval including exactly one fixed point which is hyperbolic. In some sense, this is a local case. For each $\lambda \in (0, 1)$ and integer $r \geq 0$, let

$$\mathcal{H}_-^r(\lambda) := \left\{ h \in C^r(I, I) : h(0) = 0, h'(0) = \lambda, h'(x) > 0, \right. \\ \left. h(x) < x, \forall x \in (0, 1] \right\},$$

$$\mathcal{H}_+^r(\lambda) := \left\{ h \in C^r(I, I) : h(1) = 1, h'(1) = \lambda, h'(x) > 0, \right. \\ \left. h(x) > x, \forall x \in [0, 1) \right\}$$

(2)

(cf. Figures 1 and 2) together with the norm

$$\|h\|_r := \sup_{x \in I} |h(x)| + \cdots + \sup_{x \in I} |h^{(r)}(x)|. \quad (3)$$

In what follows we only discuss the first class because $\mathcal{H}_+^r(\lambda)$ can be converted to $\mathcal{H}_-^r(\lambda)$ by considering $G(x) = 1 - F(1 - x)$.

Given integers $k, r \geq 2$, a function F belonging to the class $\bigcup_{\lambda \in (0, 1)} \mathcal{H}_-^r(\lambda)$ has a unique k th order C^r iterative root f defined on I , which is strictly increasing and is given by the formula

$$f(x) := \varphi^{-1}(\lambda^{1/k} \varphi(x)), \quad (4)$$

where $\varphi : I \rightarrow \mathbb{R}$ is the principal solution of Schröder's equation

$$\varphi(F(x)) = \lambda \varphi(x). \quad (5)$$

The principle solution is given by $\varphi(x) = \lim_{n \rightarrow +\infty} \lambda^{-n} F^n(x)$, satisfying C^r differentiable in I with $\varphi'(0) = 1$ and $\varphi(0) = 0$ and strictly increasing by Theorem 6.1 in [6]. Moreover, the proofs of Theorem 3.5.1 in [7] and Theorem 4.5 in [6] show that

$$\varphi^{(r)}(x) = \lim_{n \rightarrow +\infty} \lambda^{-n} (F^n)^{(r)}(x). \quad (6)$$

Note that the aim of this paper is to consider the local C^r ($r \geq 2$) stability for iterative roots. Then we recall the formula for higher order derivatives of composition ([26, page 3]). Namely, for integer $m \geq 1$,

$$(G \circ H)^{(m)}(x) = \sum_{\omega \in \Omega(m)} c_\omega G^{(p)}(H(x)) \prod_{q=1}^p H^{(j_q)}(x), \quad (7)$$

where $\Omega(m) := \{(p; j_1, \dots, j_p) : 1 \leq p \leq m, j_1 + \cdots + j_p = m \text{ and } j_q \geq 1 \text{ for all } q = 1, \dots, p\}$. Here c_ω is a positive universal constant, which is independent of G and H . Then we have the following lemma.

Lemma 1. Let $F \in \mathcal{H}_-^r(\lambda)$ with some $r \geq 2$ and $\lambda \in (0, 1)$ and let (F_i) be a sequence of functions in $\mathcal{H}_-^r(\lambda)$ satisfying condition

$$\lim_{i \rightarrow +\infty} \|F_i - F\|_r = 0. \quad (8)$$

Then, for a given number $\mu \in (\lambda, 1)$, there exist an $L > 1$, an $\varepsilon > 0$, and an $N_0 \in \mathbb{N}$ such that

$$\left| (F^n)^{(m)}(x) \right| \leq L\lambda^n, \quad \left| (F_i^n)^{(m)}(x) \right| \leq L\lambda^n, \quad (9)$$

$$\left| F_i^n(x) - F^n(x) \right| \leq \left\{ \left(1 + \frac{L\lambda}{\mu - \lambda} \right) \mu^{n-1} - \frac{L}{\mu - \lambda} \lambda^n \right\} \|F_i - F\|_1 \quad (10)$$

for all $i \geq N_0$, $n \in \mathbb{N}$, and $0 \leq m \leq r$ and for all $x \in I_\varepsilon := [0, \varepsilon]$.

Proof. Let $K := 2\|F\|_r$, and choose a sufficiently small $\varepsilon > 0$ such that $\lambda + K\varepsilon < 1$ and $\sup_{x \in I_\varepsilon} |F'(x)| < \mu$. Then by the mean value theorem,

$$\begin{aligned} |F(x)| &= |F(x) - F(0)| = |F'(\xi_x)| |x| \\ &\leq |F'(0)| |x| + |F'(\xi_x) - F'(0)| |x| \\ &\leq \lambda |x| + K|x|^2 \end{aligned} \quad (11)$$

for all $x \in I$. In particular, $|F^{n-1}(x)| \leq (\lambda + K\varepsilon)^{n-1}$ for all $n \geq 1$ and $x \in I_\varepsilon$. It follows that

$$\begin{aligned} |F^n(x)| &\leq \lambda |F^{n-1}(x)| + K|F^{n-1}(x)|^2 \\ &\leq \{\lambda + K(\lambda + K\varepsilon)^{n-1}\} |F^{n-1}(x)| \\ &\leq \prod_{s=0}^{n-1} \{\lambda + K(\lambda + K\varepsilon)^s\} |x| \\ &= \lambda^n \prod_{s=0}^{n-1} \{1 + K\lambda^{-1}(\lambda + K\varepsilon)^s\} |x| \\ &\leq L_0 \lambda^n, \quad \forall x \in I_\varepsilon, \end{aligned} \quad (12)$$

where $L_0 := \prod_{s=0}^{+\infty} \{1 + K\lambda^{-1}(\lambda + \eta\varepsilon)^s\} \in (1, +\infty)$.

Then we give the proof of the first inequality given in (9) by induction on m greater than 1. From (7), write

$$\begin{aligned} (F^n)^{(m)}(x) &= F'(F^{n-1}(x)) (F^{n-1})^{(m)}(x) \\ &\quad + \sum_{\substack{p \geq 2 \\ \omega \in \Omega(m)}} c_\omega F^{(p)}(F^{n-1}(x)) \prod_{q=1}^p (F^{n-1})^{(j_q)}(x). \end{aligned} \quad (13)$$

When $m = 1$, the second term is absent. Applying (12) and by induction, we have

$$\begin{aligned} |(F^n)'(x)| &= |F'(F^{n-1}(x)) (F^{n-1})'(x)| \\ &\leq \{|F'(0)| + |F'(F^{n-1}(x)) - F'(0)|\} |(F^{n-1})'(x)| \\ &\leq \{\lambda + KL_0 \lambda^{n-1}\} |(F^{n-1})'(x)| \\ &\leq \prod_{s=0}^{n-1} \{\lambda + KL_0 \lambda^s\} |x| \\ &= \lambda^n \prod_{s=0}^{n-1} \{1 + KL_0 \lambda^{s-1}\} |x| \leq L_1 \lambda^n, \quad \forall x \in I_\varepsilon, \end{aligned} \quad (14)$$

where $L_1 := \prod_{s=0}^{+\infty} \{1 + KL_0 \lambda^{s-1}\} \in (1, +\infty)$. Further, assume that the first inequality in (9) holds for all $m \leq \ell < r$. Let $\Theta := \sum_{p \geq 2, \omega \in \Omega(\ell+1)} c_\omega K \prod_{q=1}^\ell L_q$. Noting that $\lambda \in (0, 1)$ and

$$\begin{aligned} |F'(F^{n-1}(x))| &\leq |F'(0)| + |F'(F^{n-1}(x)) - F'(0)| \\ &\leq \lambda + KL_0 \lambda^{n-1}, \quad \forall x \in I_\varepsilon, \end{aligned} \quad (15)$$

we get

$$\begin{aligned} &|(F^n)^{(\ell+1)}(x)| \\ &= \left| F'(F^{n-1}(x)) (F^{n-1})^{(\ell+1)}(x) \right. \\ &\quad \left. + \sum_{\substack{p \geq 2 \\ \omega \in \Omega(\ell+1)}} c_\omega F^{(p)}(F^{n-1}(x)) \prod_{q=1}^p (F^{n-1})^{(j_q)}(x) \right| \\ &\leq (\lambda + KL_0 \lambda^{n-1}) |(F^{n-1})^{(\ell+1)}(x)| + \Theta \lambda^{2(n-1)} \\ &\leq \lambda^n \prod_{s=-1}^{n-2} (1 + KL_0 \lambda^s) + \Theta \sum_{t=0}^{n-1} \lambda^{2t} \prod_{s=t+1}^{n-1} (\lambda + KL_0 \lambda^s) \\ &= \lambda^n \prod_{s=-1}^{n-2} (1 + KL_0 \lambda^s) + \Theta \lambda^n \sum_{t=0}^{n-1} \lambda^{t-1} \prod_{s=t}^{n-2} (1 + KL_0 \lambda^s) \\ &\leq L_{\ell+1} \lambda^n \end{aligned} \quad (16)$$

inductively, where

$$\begin{aligned} L_{\ell+1} &:= \left\{ \prod_{s=-1}^{+\infty} (1 + L_0 K \lambda^s) \right. \\ &\quad \left. + \Theta \sum_{t=0}^{+\infty} \lambda^{t-1} \prod_{s=0}^{+\infty} (1 + KL_0 \lambda^s) \right\} \in (1, +\infty). \end{aligned} \quad (17)$$

Put $L := \max\{L_0, L_1, \dots, L_r\}$. Then the first inequality given in (9) is proved.

It follows from (8) that there is an $N_0 \in \mathbb{N}$ such that if $i \geq N_0$, then $\|F_i\|_r \leq 2\|F\|_r = K$ and $\sup_{x \in I_\varepsilon} |F_i'(x)| < \mu$. Thus, by the same procedure as before, we can prove the second inequality given in (9).

In the following, we are going to prove inequality (10). It is clear that (10) holds when $n = 1$. Further, assume that (10) holds for $n = \ell \in \mathbb{N}$. It follows that

$$\begin{aligned} &|F_i^{\ell+1}(x) - F^{\ell+1}(x)| \\ &= |F_i(F_i^\ell(x)) - F(F^\ell(x))| \\ &\leq |F_i(F_i^\ell(x)) - F_i(F^\ell(x))| \\ &\quad + |F_i(F^\ell(x)) - F(F^\ell(x))| \\ &\leq \sup_{\xi \in I_\varepsilon} |F_i'(\xi)| \cdot |F_i^\ell(x) - F^\ell(x)| \\ &\quad + \sup_{\xi \in I_\varepsilon} |(F_i - F)'(\xi)| \cdot |F^\ell(x)| \\ &\leq \mu |F_i^\ell(x) - F^\ell(x)| + L \lambda^\ell \|F_i - F\|_1 \end{aligned}$$

$$\begin{aligned} &\leq \mu \left\{ \left(1 + \frac{L\lambda}{\mu - \lambda} \right) \mu^{\ell-1} - \frac{L}{\mu - \lambda} \lambda^\ell \right\} \\ &\quad \times \|F_i - F\|_1 + L\lambda^\ell \|F_i - F\|_1 \\ &\leq \left\{ \left(1 + \frac{L\lambda}{\mu - \lambda} \right) \mu^\ell - \frac{L}{\mu - \lambda} \lambda^{\ell+1} \right\} \|F_i - F\|_1 \end{aligned} \tag{18}$$

for all $i \geq N_0$ and for all $x \in I_\varepsilon$. Thus, we can obtain (10) by induction. This completes the proof. \square

In Lemma 1 we gave a better estimate for F^n and F_i^n and their derivatives than that given in [25, Lemma 2.1]. In Lemma 1 the growth rate on n is much lower in the sense that the constant $L\lambda^n$ given in (9) tends to 0 as $n \rightarrow +\infty$ faster than the constant μ_0^n given in (2.4) of [25].

3. The Main Result

Our aim of this section is to prove the following stability result.

Theorem 2. *Given integers $k, r \geq 2$, let $F \in \mathcal{H}_-^{r+1}(\lambda)$ with some $\lambda \in (0, 1)$ and let (F_i) be a sequence of functions in $\mathcal{H}_-^{r+1}(\lambda)$. If*

$$\begin{aligned} &\lim_{i \rightarrow +\infty} \|F_i - F\|_{r+1} = 0, \\ &F^{(m)}(0) = F_i^{(m)}(0), \quad \forall m = 2, \dots, r, \end{aligned} \tag{19}$$

then

$$\lim_{i \rightarrow +\infty} \|f_i - f\|_r = 0, \tag{20}$$

where f and f_i are unique k th order C^r iterative roots of F and F_i , respectively, defined on I .

In order to prove Theorem 2 we need the following lemma on C^r stability of Schröder's equation.

Lemma 3. *Given an integer $r \geq 2$, let $F \in \mathcal{H}_-^{r+1}(\lambda)$ with some $\lambda \in (0, 1)$ and let (F_i) be a sequence of functions in $\mathcal{H}_-^{r+1}(\lambda)$ satisfying condition (19). Then*

$$\lim_{i \rightarrow +\infty} \|\varphi_i - \varphi\|_r = 0, \tag{21}$$

where $\varphi : I \rightarrow \mathbb{R}$ and $\varphi_i : I \rightarrow \mathbb{R}$ are the principal solutions of Schröder's equations $\varphi(F(x)) = \lambda\varphi(x)$ and $\varphi_i(F_i(x)) = \lambda\varphi_i(x)$, respectively.

Proof. From (6) we can see that

$$\begin{aligned} \varphi^{(m)}(x) &= \lim_{n \rightarrow +\infty} \lambda^{-n} (F^n)^{(m)}(x), \\ \varphi_i^{(m)}(x) &= \lim_{n \rightarrow +\infty} \lambda^{-n} (F_i^n)^{(m)}(x) \end{aligned} \tag{22}$$

for all $m \in [0, r] \cap \mathbb{Z}$ and $x \in I$. In what follows we intend to discuss our results in a sufficiently small interval I_ε first

and extend them to the whole interval I , where ε is given in Lemma 1.

In order to prove the convergence of the sequence (φ_i) in I_ε , we claim that there exists a constant M_m , which is independent of n , such that

$$\lambda^{-n} \left| (F_i^n)^{(m)}(x) - (F^n)^{(m)}(x) \right| \leq M_m \|F_i - F\|_{r+1}, \tag{23}$$

$$\forall x \in I_\varepsilon,$$

where $m = 0, 1, \dots, r$. If (23) is true, then we get

$$\begin{aligned} \left| \varphi_i^{(m)}(x) - \varphi^{(m)}(x) \right| &= \lim_{n \rightarrow +\infty} \lambda^{-n} \left| (F_i^n)^{(m)}(x) - (F^n)^{(m)}(x) \right| \\ &\leq M_m \|F_i - F\|_{r+1}, \quad \forall x \in I_\varepsilon, \end{aligned} \tag{24}$$

implying the stability in I_ε . Next, we extend the result (24) from I_ε to the whole interval I . As indicated in [25], we have

$$\lim_{i \rightarrow +\infty} \|F_i^n - F^n\|_{r+1} = 0, \quad \forall n \in \mathbb{N}, \tag{25}$$

by (8) because the composition operator is continuous by Example 4.4.5 in [27]. Moreover, since 0 is the unique stable fixed point of F in I and by (8), there is an integer $N \in \mathbb{N}$ such that $F^N(x), F_i^N(x) \in I_\varepsilon$ for all $i \in \mathbb{N}$ and $x \in I$. Then, according to Schröder's equation, we can obtain the formulae

$$\begin{aligned} \varphi(x) &= \lambda^{-N} \tilde{\varphi}(F^N(x)), & \varphi_i(x) &= \lambda^{-N} \tilde{\varphi}_i(F_i^N(x)), \\ & & & \forall x \in I, \end{aligned} \tag{26}$$

where $\tilde{\varphi} := \varphi|_{I_\varepsilon}$ and $\tilde{\varphi}_i := \varphi_i|_{I_\varepsilon}$. Then by Lemma 1 and from (24), (26), and the uniform continuity of $\varphi^{(m)}$, we get $\lim_{i \rightarrow +\infty} \|\varphi_i - \varphi\|_0 = 0$ and

$$\begin{aligned} &\left| \varphi_i^{(m)}(x) - \varphi^{(m)}(x) \right| \\ &= \lambda^{-N} \left| \sum_{\omega \in \Omega(m)} c_\omega \tilde{\varphi}_i^{(p)}(F_i^N(x)) \prod_{q=1}^p (F_i^N)^{(j_q)}(x) \right. \\ &\quad \left. - \sum_{\omega \in \Omega(m)} c_\omega \tilde{\varphi}^{(p)}(F^N(x)) \prod_{q=1}^p (F^N)^{(j_q)}(x) \right| \\ &\leq \lambda^{-N} \sum_{\omega \in \Omega(m)} c_\omega \left| \tilde{\varphi}_i^{(p)}(F_i^N(x)) \prod_{q=1}^p (F_i^N)^{(j_q)}(x) \right. \\ &\quad \left. - \tilde{\varphi}^{(p)}(F^N(x)) \prod_{q=1}^p (F^N)^{(j_q)}(x) \right| \\ &\leq \lambda^{-N} \sum_{\omega \in \Omega(m)} c_\omega \left\{ \left| \tilde{\varphi}_i^{(p)}(F_i^N(x)) - \tilde{\varphi}^{(p)}(F^N(x)) \right| \right. \\ &\quad \left. + \prod_{q=1}^p \left| (F_i^N)^{(j_q)}(x) - (F^N)^{(j_q)}(x) \right| \right\} \end{aligned}$$

$$\begin{aligned}
 & \times \left| \prod_{q=1}^p (F_i^N)^{(j_q)}(x) - \prod_{q=1}^p (F^N)^{(j_q)}(x) \right\} \\
 \leq & \lambda^{-N} \sum_{\omega \in \Omega(m)} c_\omega \{ L^p \lambda^{pN} \\
 & \times (|\bar{\varphi}_i^{(p)}(F_i^N(x)) - \bar{\varphi}^{(p)}(F^N(x))| \\
 & + |\bar{\varphi}^{(p)}(F_i^N(x)) - \bar{\varphi}^{(p)}(F^N(x))|) \\
 & + L^{p-1} \lambda^{(p-1)N} |\bar{\varphi}^{(p)}(F^N(x))| \\
 & \times \|F_i^N - F^N\|_{r+1} \} \\
 \leq & \lambda^{-N} \sum_{\omega \in \Omega(m)} c_\omega \{ L^p \lambda^{pN} \\
 & \times (M_p \|F_i - F\|_{r+1} \\
 & + |\bar{\varphi}^{(p)}(F_i^N(x)) - \bar{\varphi}^{(p)}(F^N(x))|) \\
 & + L^{p-1} \lambda^{(p-1)N} |\bar{\varphi}^{(p)}(F^N(x))| \\
 & \times \|F_i^N - F^N\|_{r+1} \} \rightarrow 0
 \end{aligned} \tag{27}$$

as $i \rightarrow +\infty$ for all $m = 1, \dots, r$. Hence, (21) is proved and the proof is completed.

In the following, we will prove the claimed (23) by induction on m . Clearly, (2.11) in [25] and what is indicated above the proof of Theorem 2.1 in [25] show that (23) holds for $m = 1$ and $m = 0$, respectively. Then we suppose that (23) is also satisfied for all $m \in [1, \ell]$, where $2 \leq \ell < r$, and we will prove (23) for $m = \ell + 1$. Our strategy is to prove that, for given $s \in \mathbb{N}$, there exists $M_{\ell+1}(s)$ such that

$$\lambda^{-s} \left| (F_i^s)^{(\ell+1)}(x) - (F^s)^{(\ell+1)}(x) \right| \leq M_{\ell+1}(s) \|F_i - F\|_{r+1} \tag{28}$$

and $(M_{\ell+1}(s))$ is a bounded sequence whose upper bound satisfies (23). Clearly, for $s = 0$ we can find constant $M_{\ell+1}(0)$ satisfying (28). Then, assume that there exists $M_{\ell+1}(s)$ such that (28) holds for any $s = 0, \dots, n - 1$. Applying Lemma 1 and the inductive hypothesis, we have

$$\begin{aligned}
 & \lambda^{-n} \left| (F_i^n)^{(\ell+1)}(x) - (F^n)^{(\ell+1)}(x) \right| \\
 = & \lambda^{-n} \left| F_i'(F_i^{n-1}(x)) (F_i^{n-1})^{(\ell+1)}(x) \right. \\
 & + \sum_{\substack{p \geq 2 \\ \omega \in \Omega(\ell+1)}} c_\omega F_i^{(p)}(F_i^{n-1}(x)) \prod_{q=1}^p (F_i^{n-1})^{(j_q)}(x) \\
 & \left. - F'(F^{n-1}(x)) (F^{n-1})^{(\ell+1)}(x) \right|
 \end{aligned}$$

$$\begin{aligned}
 & - \sum_{\substack{p \geq 2 \\ \omega \in \Omega(\ell+1)}} c_\omega F^{(p)}(F^{n-1}(x)) \prod_{q=1}^p (F^{n-1})^{(j_q)}(x) \Big| \\
 \leq & \lambda^{-n} \left\{ \left| F_i'(F_i^{n-1}(x)) (F_i^{n-1})^{(\ell+1)}(x) \right. \right. \\
 & \left. \left. - F'(F^{n-1}(x)) (F^{n-1})^{(\ell+1)}(x) \right| \right. \\
 & + \sum_{\substack{p \geq 2 \\ \omega \in \Omega(\ell+1)}} c_\omega \left| F_i^{(p)}(F_i^{n-1}(x)) \prod_{q=1}^p (F_i^{n-1})^{(j_q)}(x) \right. \\
 & \left. \left. - F^{(p)}(F^{n-1}(x)) \prod_{q=1}^p (F^{n-1})^{(j_q)}(x) \right| \right\} \\
 \leq & \lambda^{-1} M_{\ell+1}(n-1) |F_i'(F_i^{n-1}(x))| \|F_i - F\|_{r+1} \\
 & + L \lambda^{-1} A_{1,n-1}(x) \\
 & + \sum_{\substack{p \geq 2 \\ \omega \in \Omega(\ell+1)}} c_\omega \{ L^p \lambda^{(p-1)n-p} A_{p,n-1}(x) \\
 & + \sum_{q=1}^p M_{j_q} L^{p-1} \lambda^{(p-1)n-p} \\
 & \times |F^{(p)}(F^{n-1}(x))| \|F_i - F\|_{r+1} \},
 \end{aligned} \tag{29}$$

where $A_{p,n}(x) := |F_i^{(p)}(F_i^n(x)) - F^{(p)}(F^n(x))|$. By Lemma 1 and condition (19), we obtain that $|F_i'(F_i^{n-1}(x))| \leq |F_i'(0)| + |F_i'(F_i^{n-1}(x)) - F_i'(0)| \leq \lambda + L\eta\lambda^{n-1}$,

$$\begin{aligned}
 |F^{(p)}(F^{n-1}(x))| & \leq |F^{(p)}(0)| + |F^{(p)}(F^{n-1}(x)) - F^{(p)}(0)| \\
 & \leq \eta + L\eta\lambda^{n-1},
 \end{aligned}$$

$$\begin{aligned}
 A_{p,n}(x) & \leq |F_i^{(p)}(F_i^n(x)) - F^{(p)}(F_i^n(x)) - F_i^{(p)}(0) + F^{(p)}(0)| \\
 & + |F^{(p)}(F_i^n(x)) - F^{(p)}(F^n(x))| \\
 & \leq L\lambda^n \|F_i - F\|_{r+1} + \eta |F_i^n(x) - F^n(x)| \\
 & \leq L\lambda^n \|F_i - F\|_{r+1} + \eta \left\{ \left(1 + \frac{L\lambda}{\mu - \lambda} \right) \mu^{n-1} - \frac{L}{\mu - \lambda} \lambda^n \right\} \\
 & \quad \times \|F_i - F\|_1 \\
 = & \left\{ \left(1 + \frac{L\lambda}{\mu - \lambda} \right) \eta \mu^{n-1} + \left(1 - \frac{\eta}{\mu - \lambda} \right) L\lambda^n \right\} \|F_i - F\|_{r+1},
 \end{aligned} \tag{30}$$

where $\eta := 2\|F\|_{r+1}$ and μ is given in Lemma 1, for all $x \in I_\varepsilon$, $0 < p \leq r, i \geq N_0$, and $n \in \mathbb{N}$. It follows that

$$\begin{aligned} & \lambda^{-n} \left| (F_i^n)^{(\ell+1)}(x) - (F^n)^{(\ell+1)}(x) \right| \\ & \leq \left\{ (1 + L\eta\lambda^{n-2})M_{\ell+1}(n-1) + \left(1 + \frac{L\lambda}{\mu-\lambda}\right)L\eta\mu^{n-2}\lambda^{-1} \right. \\ & \quad \left. + \left(1 - \frac{\eta}{\mu-\lambda}\right)L^2\lambda^{n-2} \right\} \|F_i - F\|_{r+1} \\ & \quad + \sum_{\substack{p \geq 2 \\ \omega \in \Omega(\ell+1)}} c_\omega \left\{ \left(1 + \frac{L\lambda}{\mu-\lambda}\right)L^p\eta\mu^{n-2}\lambda^{(p-1)n-p} \right. \\ & \quad \left. + \left(1 - \frac{\eta}{\mu-\lambda}\right)L^{p+1}\lambda^{pn-p-1} \right. \\ & \quad \left. + (1 + L\lambda^{n-1})L^{p-1}\eta\lambda^{(p-1)n-p} \sum_{q=1}^p M_{j_q} \right\} \\ & \quad \times \|F_i - F\|_{r+1} \\ & \leq \left\{ (1 + L\eta\lambda^{n-2})M_{\ell+1}(n-1) + \left(1 + \frac{L\lambda}{\mu-\lambda}\right)L\eta\mu^{n-2}\lambda^{-1} \right. \\ & \quad \left. + \left(1 - \frac{\eta}{\mu-\lambda}\right)L^2\mu^{n-2} \right. \\ & \quad \left. + \sum_{\substack{p \geq 2 \\ \omega \in \Omega(\ell+1)}} c_\omega \left\{ \left(1 + \frac{L\lambda}{\mu-\lambda}\right)L^p\eta\mu^{2n-4} \right. \right. \\ & \quad \left. \left. + \left(1 - \frac{\eta}{\mu-\lambda}\right)L^{p+1}\mu^{2n-3} \right. \right. \\ & \quad \left. \left. + (1 + L)L^{p-1}\eta\mu^{n-2} \sum_{q=1}^p M_{j_q} \right\} \right\} \\ & \quad \times \|F_i - F\|_{r+1} \\ & \leq \left\{ (1 + L\eta\mu^{n-2})M_{\ell+1}(n-1) + \Gamma\mu^n \right\} \|F_i - F\|_{r+1} \end{aligned} \tag{31}$$

because $\lambda < \mu \in (0, 1)$, where

$$\begin{aligned} \Gamma := & \mu^4 \left\{ \left(1 + \frac{L\lambda}{\mu-\lambda}\right)L\eta\lambda^{-1} + \left(1 - \frac{\eta}{\mu-\lambda}\right)L^2 \right. \\ & \left. + \sum_{\substack{p \geq 2 \\ \omega \in \Omega(\ell+1)}} c_\omega \left\{ \left(1 + \frac{L\lambda}{\mu-\lambda}\right)L^p\eta \right. \right. \\ & \quad \left. \left. + \left(1 - \frac{\eta}{\mu-\lambda}\right)L^{p+1} \right. \right. \\ & \quad \left. \left. + (1 + L)L^{p-1}\eta \sum_{q=1}^p M_{j_q} \right\} \right\}. \end{aligned} \tag{32}$$

Let

$$\begin{aligned} M_{\ell+1}(n) & := (1 + L\eta\mu^{n-2})M_{\ell+1}(n-1) + \Gamma\mu^n \\ & = \prod_{j=1}^{n-2} (1 + L\eta\mu^j)M_{\ell+1}(2) + \Gamma \sum_{j=3}^n \mu^j \prod_{t=j-1}^{n-2} (1 + L\eta\mu^t). \end{aligned} \tag{33}$$

It is easy to check that

$$M_{\ell+1}(n) \leq \prod_{j=1}^{+\infty} (1 + L\eta\mu^j)M_{\ell+1}(2) + \Gamma \sum_{j=3}^{+\infty} \mu^j \prod_{t=0}^{+\infty} (1 + L\eta\mu^t) \tag{34}$$

for all $n \in \mathbb{N}$. By putting $M_{\ell+1} := \prod_{j=1}^{+\infty} (1 + L\eta\mu^j)M_{\ell+1}(2) + \Gamma \sum_{j=3}^{+\infty} \mu^j \prod_{t=0}^{+\infty} (1 + L\eta\mu^t)$, we can prove (23) for $m = \ell + 1$ and (23) is proved by induction. \square

Having this preparation, we can give a proof to the main result of this paper.

Proof of Theorem 2. By (4) the C^r iterative roots f and f_i for each $i \in \mathbb{N}$ can be presented by

$$\begin{aligned} f(x) & = \varphi^{-1}(\lambda^{1/k}\varphi(x)), & f_i(x) & = \varphi_i^{-1}(\lambda^{1/k}\varphi_i(x)), \\ & & & \forall x \in I, \end{aligned} \tag{35}$$

respectively. In order to prove the convergence of (f_i) in I , note that, for sufficiently large $i \in \mathbb{N}$ such that $\lambda^{1/k}\varphi_i(x) \in \varphi(I)$ for all $x \in I$, we have

$$\begin{aligned} & \left| (\varphi_i^{-1})^{(p)}(\lambda^{1/k}\varphi_i(x)) \prod_{q=1}^p \varphi_i^{(j_q)}(x) \right. \\ & \quad \left. - (\varphi^{-1})^{(p)}(\lambda^{1/k}\varphi(x)) \prod_{q=1}^p \varphi^{(j_q)}(x) \right| \\ & \leq \left| (\varphi_i^{-1})^{(p)}(\lambda^{1/k}\varphi_i(x)) - (\varphi^{-1})^{(p)}(\lambda^{1/k}\varphi(x)) \right| \\ & \quad \times \prod_{q=1}^p \left| \varphi_i^{(j_q)}(x) \right| + \left| (\varphi^{-1})^{(p)}(\lambda^{1/k}\varphi(x)) \right| \\ & \quad \times \left| \prod_{q=1}^p \varphi_i^{(j_q)}(x) - \prod_{q=1}^p \varphi^{(j_q)}(x) \right| \\ & \leq 2^p (\|\varphi\|_r)^p \left\{ \|\varphi_i^{-1} - \varphi^{-1}\|_r + \left| (\varphi^{-1})^{(p)}(\lambda^{1/k}\varphi_i(x)) \right. \right. \\ & \quad \left. \left. - (\varphi^{-1})^{(p)}(\lambda^{1/k}\varphi(x)) \right| \right\} \\ & \quad + p2^{p-1} (\|\varphi\|_r)^{p-1} \|\varphi^{-1}\|_r \|\varphi_i - \varphi\|_r \rightarrow 0 \end{aligned} \tag{36}$$

as $i \rightarrow +\infty$ for all integers $p \in [1, r]$ by (21) and the fact that the inverse operator is continuous and $(\varphi^{-1})^{(p)}$ is uniformly continuous. Hence, $\lim_{i \rightarrow +\infty} \|f_i - f\|_0 = 0$ and

$$\begin{aligned} & \left| (f_i)^{(m)}(x) - f^{(m)}(x) \right| \\ &= \left| \sum_{\omega \in \Omega(m)} c_\omega \lambda^{p/k} (\varphi_i^{-1})^{(p)} (\lambda^{1/k} \varphi_i(x)) \prod_{q=1}^p \varphi_i^{(j_q)}(x) \right. \\ & \quad \left. - \sum_{\omega \in \Omega(m)} c_\omega \lambda^{p/k} (\varphi^{-1})^{(p)} (\lambda^{1/k} \varphi(x)) \prod_{q=1}^p \varphi^{(j_q)}(x) \right| \\ &\leq \sum_{\omega \in \Omega(m)} c_\omega \lambda^{p/k} \left| (\varphi_i^{-1})^{(p)} (\lambda^{1/k} \varphi_i(x)) \prod_{q=1}^p \varphi_i^{(j_q)}(x) \right. \\ & \quad \left. - (\varphi^{-1})^{(p)} (\lambda^{1/k} \varphi(x)) \prod_{q=1}^p \varphi^{(j_q)}(x) \right| \rightarrow 0 \end{aligned} \tag{37}$$

as $i \rightarrow +\infty$ for all $m = 1, \dots, r$, which implies that $\lim_{i \rightarrow +\infty} \|f_i - f\|_r = 0$ and completes the proof. \square

Theorem 2 is also valid for $r = 1$, which is the same as Theorem 2.1 in [25] for $r = 1$. However, it is hard to use the estimates, for example, (2.4) and (2.5) in [25], to generalize the result to the general r parallel. In fact, we cannot use those estimates to give a uniform constant Γ with respect to n , the order of iteration, in (33). Using those estimates, corresponding to Γ given in (32), we obtain the quantity

$$\begin{aligned} \Gamma^*(n) := & \mu_0^4 \left\{ L^2 + (n-1)L\eta\lambda^{-1} \right. \\ & + \sum_{\substack{p \geq 2 \\ \omega \in \Omega(\ell+1)}} c_\omega \left\{ L^{p+1} + (n-1)L^p\eta \right. \\ & \left. \left. + (1+L)L^{p-1}\eta \sum_{q=1}^p M_{j_q} \right\} \right\}, \end{aligned} \tag{38}$$

which tends to $+\infty$ as $n \rightarrow +\infty$. For this reason it is hard to prove the boundedness of $M_{\ell+1}(n)$. As remarked after the proof of Lemma 1, our estimation in (9) and (10) enables us to give the boundedness of $M_{\ell+1}(n)$ and complete the proof of (23).

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of this paper.

References

[1] C. Babbage, "An essay towards the calculus of functions," *Philosophical Transactions of the Royal Society of London*, vol. 105, pp. 389–423, 1815.

[2] M. K. Fort, Jr., "The embedding of homeomorphisms in flows," *Proceedings of the American Mathematical Society*, vol. 6, pp. 960–967, 1955.

[3] J. Palis, Jr. and W. de Melo, *Geometric Theory of Dynamical Systems: An Introduction*, Springer, New York, NY, USA, 1982.

[4] G. Targoński, *Topics in Iteration Theory*, vol. 6, Vandenhoeck & Ruprecht, Göttingen, Germany, 1981.

[5] K. Baron and W. Jarczyk, "Recent results on functional equations in a single variable, perspectives and open problems," *Aequationes Mathematicae*, vol. 61, no. 1-2, pp. 1–48, 2001.

[6] M. Kuczma, *Functional Equations in a Single Variable*, vol. 46 of *Monograph in Mathematics*, PWN, Warsaw, Poland, 1968.

[7] M. Kuczma, B. Choczewski, and R. Ger, *Iterative Functional Equations*, Encyclopedia of Mathematics and Its Applications, Cambridge University Press, Cambridge, UK, 1990.

[8] J. Zhang, L. Yang, and W. Zhang, "Some advances on functional equations," *Advances in Mathematics*, vol. 24, no. 5, pp. 385–405, 1995.

[9] A. Blokh, E. Coven, M. Misiurewicz, and Z. Nitecki, "Roots of continuous piecewise monotone maps of an interval," *Acta Mathematica Universitatis Comenianae*, vol. 60, no. 1, pp. 3–10, 1991.

[10] L. Liu, W. Jarczyk, L. Li, and W. Zhang, "Iterative roots of piecewise monotonic functions of nonmonotonicity height not less than 2," *Nonlinear Analysis. Theory, Methods & Applications A*, vol. 75, no. 1, pp. 286–303, 2012.

[11] L. Liu and W. Zhang, "Non-monotonic iterative roots extended from characteristic intervals," *Journal of Mathematical Analysis and Applications*, vol. 378, no. 1, pp. 359–373, 2011.

[12] K. Ciepliński, "On the embeddability of a homeomorphism of the unit circle in disjoint iteration groups," *Publicationes Mathematicae Debrecen*, vol. 55, no. 3-4, pp. 363–383, 1999.

[13] M. C. Zdun, "On iterative roots of homeomorphisms of the circle," *Bulletin of the Polish Academy of Sciences. Mathematics*, vol. 48, no. 2, pp. 203–213, 2000.

[14] L. Li, J. Jarczyk, W. Jarczyk, and W. Zhang, "Iterative roots of mappings with a unique set-value point," *Publicationes Mathematicae Debrecen*, vol. 75, no. 1-2, pp. 203–220, 2009.

[15] T. Powierza, "Higher order set-valued iterative roots of bijections," *Publicationes Mathematicae Debrecen*, vol. 61, no. 3-4, pp. 315–324, 2002.

[16] Z. Leśniak, "On fractional iterates of a homeomorphism of the plane," *Annales Polonici Mathematici*, vol. 79, no. 2, pp. 129–137, 2002.

[17] T. Narayaninsamy, "Fractional iterates for n -dimensional maps," *Applied Mathematics and Computation*, vol. 98, no. 2-3, pp. 261–278, 1999.

[18] N. Iannella and L. Kindermann, "Finding iterative roots with a spiking neural network," *Information Processing Letters*, vol. 95, no. 6, pp. 545–551, 2005.

[19] L. Kindermann, "Computing iterative roots with neural networks," in *Proceedings of the 5th Conference Neural Information Processing*, vol. 2, pp. 713–715, 1998.

[20] T. Narayaninsamy, "A connection between fractional iteration and graph theory," *Applied Mathematics and Computation*, vol. 107, no. 2-3, pp. 181–202, 2000.

[21] J. Kobza, "Iterative functional equation $x(x(t)) = f(t)$ with $f(t)$ piecewise linear," *Journal of Computational and Applied Mathematics*, vol. 115, no. 1-2, pp. 331–347, 2000.

- [22] W. Zhang and W. Zhang, "Computing iterative roots of polygonal functions," *Journal of Computational and Applied Mathematics*, vol. 205, no. 1, pp. 497–508, 2007.
- [23] B. Xu and W. Zhang, "Construction of continuous solutions and stability for the polynomial-like iterative equation," *Journal of Mathematical Analysis and Applications*, vol. 325, no. 2, pp. 1160–1170, 2007.
- [24] W. Zhang and W. Zhang, "Continuity of iteration and approximation of iterative roots," *Journal of Computational and Applied Mathematics*, vol. 235, no. 5, pp. 1232–1244, 2011.
- [25] W. Zhang, Y. Zeng, W. Jarczyk, and W. Zhang, "Local C^1 stability versus global C^1 unstability for iterative roots," *Journal of Mathematical Analysis and Applications*, vol. 386, no. 1, pp. 75–82, 2012.
- [26] R. Abraham and J. Robbin, *Transversal Mappings and Flows*, Benjamin, New York, NY, USA, 1967.
- [27] R. S. Hamilton, "The inverse function theorem of Nash and Moser," *Bulletin of the American Mathematical Society. New Series*, vol. 7, no. 1, pp. 65–222, 1982.