

FROM LOGARITHMIC TO SUBDIFFUSIVE POLYNOMIAL FLUCTUATIONS FOR INTERNAL DLA AND RELATED GROWTH MODELS¹

BY AMINE ASSELAH AND ALEXANDRE GAUILLIÈRE

Université Paris-Est and Université de Provence

Dedicated to Joel Lebowitz, for his 80th birthday

We consider a cluster growth model on \mathbb{Z}^d , called internal diffusion limited aggregation (internal DLA). In this model, random walks start at the origin, one at a time, and stop moving when reaching a site not occupied by previous walks. It is known that the asymptotic shape of the cluster is spherical. When dimension is 2 or more, we prove that fluctuations with respect to a sphere are at most a power of the logarithm of its radius in dimension $d \geq 2$. In so doing, we introduce a closely related cluster growth model, that we call *the flashing process*, whose fluctuations are controlled easily and accurately. This process is coupled to internal DLA to yield the desired bound. Part of our proof adapts the approach of Lawler, Bramson and Griffeath, on another space scale, and uses a sharp estimate (written by Blachère in our Appendix) on the expected time spent by a random walk inside an annulus.

1. Introduction. The internal DLA cluster of volume N , say $A(N)$, is obtained inductively as follows. Initially, we assume that the explored region is empty, that is, $A(0) = \emptyset$. Then, consider N independent discrete-time random walks S_1, \dots, S_N starting from 0. For $k \leq N$, assume $A(k-1)$ is obtained, and define

$$\tau_k = \inf\{t \geq 0: S_k(t) \notin A(k-1)\} \quad \text{and} \quad A(k) = A(k-1) \cup \{S_k(\tau_k)\}.$$

In such a particle system, we call explorers the particles. We say that the k th explorer is *settled* on $S_k(\tau_k)$ after time τ_k , and is *unsettled* before time τ_k . The cluster $A(N)$ consists of the positions of the N settled explorers.

The mathematical model of internal DLA was introduced first in the chemical physics literature by Meakin and Deutch [13]. There are many industrial processes that look like internal DLA; see the nice review paper [7]. The most important

Received May 2010; revised April 2012.

¹Supported by GDRE 224 GREFI-MEFI, the French Ministry of Education through the ANR BLAN07-2184264 grant, by the European Research Council through the “Advanced Grant” PTRELSS 228032.

MSC2010 subject classifications. 60K35, 82B24, 60J45.

Key words and phrases. Internal diffusion limited aggregation, cluster growth, random walk, shape theorem, logarithmic fluctuations, subdiffusive fluctuations.

seems to be electropolishing, defined as *the improvement of surface finish of a metal effected by making it anodic in an appropriate solution*. There are actually two distinct industrial processes (i) *anodic leveling or smoothing* which corresponds to the elimination of surface roughness of height larger than 1 micron, and (ii) *anodic brightening* which refers to elimination of surface defects which are protruding by less than 1 micron. The latter phenomenon requires an understanding of atom removal from a crystal lattice. It was noted in [13] that, at a qualitative level, the model produces smooth clusters, and the authors wrote, “it is also of some fundamental significance to know just how smooth a surface formed by diffusion limited processes may be.”

Diaconis and Fulton [2] introduced internal DLA in mathematics. They allowed explorers to start on distinct sites, and showed that the law of the cluster was invariant under permutation of the order in which explorers were launched. This invariance, named *the abelian property*, was central in their motivation. They treat, among other things, the special one-dimensional case.

In dimension two or more, Lawler, Bramson and Griffeath [10] prove that in order to cover, without holes, a sphere of radius n , we need about the number of sites of \mathbb{Z}^d contained in this sphere. In other words, the asymptotic shape of the cluster is a sphere. Then, Lawler in [9] shows subdiffusive fluctuations. The latter result is formulated in terms of inner and outer errors, which we now introduce with some notation. We denote with $\|\cdot\|$ the Euclidean norm on \mathbb{R}^d . For any x in \mathbb{R}^d and r in \mathbb{R} , set

$$B(x, r) = \{y \in \mathbb{R}^d : \|y - x\| < r\} \quad \text{and} \quad \mathbb{B}(x, r) = B(x, r) \cap \mathbb{Z}^d.$$

For $\Lambda \subset \mathbb{Z}^d$, $|\Lambda|$ denotes the number of sites in Λ . The inner error $\delta_I(n)$ is such that

$$n - \delta_I(n) = \sup\{r \geq 0 : \mathbb{B}(0, r) \subset A(|\mathbb{B}(0, n)|)\}.$$

Also, the outer error $\delta_O(n)$ is such that

$$n + \delta_O(n) = \inf\{r \geq 0 : A(|\mathbb{B}(0, n)|) \subset \mathbb{B}(0, r)\}.$$

The main result of [9] reads as follows.

THEOREM 1.1 (Lawler). *Assume $d \geq 2$. Then*

$$(1.1) \quad P(\exists n(\omega) : \forall n \geq n(\omega) \delta_I(n) \leq n^{1/3} \log(n)^2) = 1$$

and

$$(1.2) \quad P(\exists n(\omega) : \forall n \geq n(\omega) \delta_O(n) \leq n^{1/3} \log(n)^4) = 1.$$

Since Lawler’s paper, published 15 years ago, no improvement of these estimates was achieved, but it is believed that fluctuations are on a much smaller scale than $n^{1/3}$. Moreover, (1.1) and (1.2) are almost sure upper bounds on errors, and

no lower bound on the inner or outer error has been established. Computer simulations [3, 14] suggest indeed that fluctuations are logarithmic. In addition, Levine and Peres studied a deterministic analogue of internal DLA, the rotor-router model, introduced by Propp [6]. They bound, in [12], the inner error $\delta_I(n)$ by $\log(n)$, and the outer error $\delta_O(n)$ by $n^{1-1/d}$.

Our main result is the following improvement of Theorem 1.1.

THEOREM 1.2. *Assume $d \geq 2$. There is a positive constant A_d such that*

$$(1.3) \quad P(\exists n(\omega) : \forall n \geq n(\omega) \delta_I(n) \leq A_d \log(n)) = 1$$

and

$$(1.4) \quad P(\exists n(\omega) : \forall n \geq n(\omega) \delta_O(n) \leq A_d \log^2(n)) = 1.$$

Note added in proof. At about the same time, and with an independent approach, Jerison, Levine and Sheffield [5] obtained similar results with an improved bound on the outer error in $d = 2$. Then, by refining our approach, we obtained in [1] a bound of order $\sqrt{\log(n)}$ for both internal and external errors in dimension three or more. Jerison, Levine and Sheffield [4] did the same by following their approach.

Our approach builds on the work of Lawler, Bramson and Griffeath [10], which we review later. It also deals with more general models of diffusion limited aggregation which we now describe. Indeed, we introduce a family of cluster growth models for which a control of the fluctuations of the cluster shape is easily obtained. These growth models are built so that the asymptotic shape is spherical, but still they exhibit a large diversity of fluctuations parametrized by a certain width ranging from a large constant to a power 1/3 of the radius of the asymptotic sphere. Moreover, all these clusters are coupled to internal DLA, and, as a consequence, we obtain logarithmic bounds on the fluctuations for internal DLA. We generalize internal DLA by allowing explorers to settle only at some special times. Thus, each explorer i is associated with a collection of times $\{\sigma_{i,k}, k \in \mathbb{N}\}$ and

$$\tau_i^* = \inf\{\sigma_{i,k} : S_i(\sigma_{i,k}) \notin A^*(i - 1)\} \quad \text{and} \quad A^*(i) = A^*(i - 1) \cup \{S_i(\tau_i^*)\}.$$

The internal DLA is recovered as we choose $\sigma_{i,k} = k$ for all $i = 1, \dots, N$ and $k \in \mathbb{N}$. We call $\{\sigma_{i,k}, k \in \mathbb{N}\}$ the *flashing times* associated to the i th explorer, and $\{S_i(\sigma_{i,k}), k \in \mathbb{N}\}$ its *flashing positions*.

In this paper, we consider stopping times of a special form, linked with the spherical nature of the internal DLA cluster. An illustration with one *flashing* explorer's trajectory is made in Figure 1.

The precise definition of the *flashing times* requires additional notation, which we postpone to Section 3. We describe here key features of flashing processes. We first choose a sequence of widths, say $\mathcal{H} = \{h_n, n \in \mathbb{N}\}$, and then partition \mathbb{Z}^d into concentric shells $\{S_n, n \in \mathbb{N}\}$, whose respective widths are $\{2h_n, n \in \mathbb{N}\}$. Each shell is in turn partitioned into cells, which are brick-like domain, of side length equal




FIG. 1. Cell decomposition, and flashing positions as stars.

to the width of the shell. The flashing times are chosen such that (i) an explorer flashes at most once in each shell, (ii) the flashing position, in a shell, is essentially uniform over the cell an explorer first hits upon entering the shell and (iii) when an explorer leaves a shell, it cannot afterward flash in it.

For a given sequence \mathcal{H} , we call the process just described the \mathcal{H} -flashing process. Note that feature (ii) is the seed of a deep difference with internal DLA. *The mechanism of covering a cell, for the flashing process, is very much the same as completing an album in the classical coupon-collector process.* Thus, we need of the order of $V \log(V)$ explorers to cover a cell of volume V . For internal DLA, with explorers started at the origin, we only need of order V explorers to cover a sphere of volume V as shown in [10], and we believe that we need a number of explorers of order $|\mathcal{C}|$ to cover a cell \mathcal{C} , even if they start on the boundary of the cell. In addition, feature (ii) allows us to *localize* the covering mechanism, in the sense that a particle entering a shell cannot flash outside the cell through which it entered that shell. Finally, feature (iii) is essential for having a useful coupling between flashing and internal DLA processes.

LEMMA 1.3. *Assume that N is an integer, and \mathcal{H} is a sequence of positive integers. There is a coupling between the two processes, using the same trajectories S_1, \dots, S_N such that*

$$(1.5) \quad A(N) = \bigcup_{i=1}^N \{S_i(T(i))\} \quad \text{and} \quad A^*(N) = \bigcup_{i=1}^N \{S_i(T^*(i))\}$$

and $T^*(i) \geq T(i)$ for all $i = 1, \dots, N$.

As a corollary of Lemma 1.3, we have the following useful result.

COROLLARY 1.4. *Under the hypotheses of the previous lemma, for $k \geq 1$:*

- *if $A^*(N) \subset \bigcup_{j < k} \mathcal{S}_j$, then $A(N) \subset \bigcup_{j < k} \mathcal{S}_j$;*
- *if $\bigcup_{j < k} \mathcal{S}_j \subset A^*(N)$, then $\bigcup_{j < k} \mathcal{S}_j \subset A(N)$.*

An \mathcal{H} -flashing process, with $h_j \geq h_0$ for $j \geq 0$, and h_0 a large constant, produces a cluster $A^*(N)$, for which we bound easily the inner error, $\delta_I^*(n)$. Then, to bound the outer error, $\delta_O^*(n)$, we follow the approach of [9], though with a slightly simpler proof.

PROPOSITION 1.5. *Assume that for $j \geq 1$, $h_j \leq h_{j+1} \leq (1 + \frac{1}{2j})h_j$, with a large h_0 . For a positive constant A_d^* , we have*

$$(1.6) \quad P(\exists n(\omega) : \forall n \geq n(\omega) \delta_I^*(n) \leq A_d^* h(n) \log(n)) = 1$$

and

$$(1.7) \quad P(\exists n(\omega) : \forall n \geq n(\omega) \delta_O^*(n) \leq A_d^* h(n) \log^2(n)) = 1,$$

where $h(n) = \max\{h_k \in \mathbb{R} : r_k \leq n\}$.

Finally, we establish lower bound on the inner and outer error.

PROPOSITION 1.6. *Assume that h_0 is large enough. Then, there is a constant a_d^* such that*

$$(1.8) \quad P(\exists n(\omega) : \forall n \geq n(\omega) \delta_I^*(n) \geq a_d^* h(n) \log(h(n))) = 1$$

and

$$(1.9) \quad P(\exists n(\omega) : \forall n \geq n(\omega) \delta_O^*(n) \geq a_d^* h(n) \log(n)) = 1.$$

Corollary 1.4 and Proposition 1.5, with the choice $h_j = h_0$ for all $j > 0$, imply Theorem 1.2 which deals with internal DLA.

Let us now review previous work on internal DLA.

On previous bounds for internal DLA. We describe the approach of [10], for establishing the upper bound for the inner error. It is convenient to consider explorers starting outside the origin with initial configuration denoted η . We denote also by $A(\Lambda, \eta)$ the cluster obtained from explorers initially on η , with an explored region $\Lambda \subset \mathbb{Z}^d$.

Now, for a site $z \in \mathbb{Z}^d$, we call $W(\eta, z)$ [resp., $M(\eta, z)$] the number of explorers (resp., of random walks) which visit z before settling. For an integer n , and η consisting of $|\mathbb{B}(0, n)|$ explorers at the origin, the authors of [10] first write

$$\{\mathbb{B}(0, r) \not\subset A(\emptyset, \eta)\} \subset \bigcup_{z \in \mathbb{B}(0, r)} \{W(\eta, z) = 0\}.$$

Then, they look for the largest value of r_n (in terms of n) which guarantees that $|\mathbb{B}(0, r_n)| \times \sup_{z \in \mathbb{B}(0, r_n)} P(W(\eta, z) = 0)$ be the term of a convergent series.

The approach of [10] is based on the following observations. (i) If explorers would not settle, they would just be independent random walks; (ii) exactly one explorer occupies each site of the cluster. Thus, the following equality holds in law:

$$W(\eta, z) + M(A(\emptyset, \eta), z) \geq M(\eta, z).$$

Now, an observation of Diaconis and Fulton [2] is that we can realize the cluster by sending many *exploration waves*. Let us illustrate this observation with two waves. We first stop the explorers on the external boundary of a ball of radius R , say $\partial\mathbb{B}(0, R)$. The cluster consisting of the positions of settled explorers is denoted $A_R(\emptyset, \eta)$, so that $A_R(\emptyset, \eta) \subset \mathbb{B}(0, R)$. The configuration with stopped explorers on $\partial\mathbb{B}(0, R)$ is denoted $\zeta_R(\eta)$. Then, the second wave consists in launching the explorers of $\zeta_R(\eta)$, with explored region $A_R(\emptyset, \eta)$. In other words, we have an equality in law

$$A(\emptyset, \eta) = A_R(\emptyset, \eta) \cup A(A_R(\emptyset, \eta), \zeta_R(\eta)).$$

Moreover, if the index R refers only to explorers (or walks) of the first wave, then for $z \in \mathbb{B}(0, R)$,

$$(1.10) \quad W_R(\eta, z) + M_R(A_R(\emptyset, \eta), z) \geq M_R(\eta, z).$$

The authors of [10] consider $R = n$ and $z \in \mathbb{B}(0, n)$. Since $W(\eta, z) \geq W_n(\eta, z)$, we have using (1.10), for any $\alpha > 0$,

$$(1.11) \quad P(W(\eta, z) = 0) \leq P(M_n(\eta, z) < \alpha) + P(M_n(\mathbb{B}(0, n), z) > \alpha).$$

We then look for sites z such that $E[M_n(\eta, z)] > \alpha > E[M_n(\mathbb{B}(0, n), z)]$ (and $\eta = |\mathbb{B}(0, n)|\delta_0$). Note that $M_n(\eta, z)$ and $M_n(\mathbb{B}(0, n), z)$ are sums of independent Bernoulli variables with well-known large deviation estimates. If we set

$$2\alpha = E[M_n(\eta, z)] + E[M_n(\mathbb{B}(0, n), z)]$$

and

$$\tilde{\mu}_n(z) = E[M_n(\eta, z)] - E[M_n(\mathbb{B}(0, n), z)],$$

then

$$(1.12) \quad \begin{aligned} P(M_n(\eta, z) < \alpha) &\leq \exp\left(-\frac{(E[M_n(\eta, z)] - \alpha)^2}{2E[M_n(\eta, z)]}\right) \\ &\leq \exp\left(-\frac{\tilde{\mu}_n^2(z)}{8E[M_n(\eta, z)]}\right). \end{aligned}$$

Lawler in [9] establishes that for $z \in \mathbb{B}(0, n)$,

$$E[M_n(\eta, z)] \sim n(n - \|z\|) \quad \text{and} \quad \tilde{\mu}_n(z) \sim (n - \|z\|)^2.$$

Replacing these values in (1.12), the bound $n - \|z\| \geq n^{1/3} \log(n)$ is such that $P(W_n(\eta, z) = 0)$ is the term of a convergent series.

We now sketch our main ideas leading to logarithmic fluctuations for internal DLA.

On logarithmic fluctuations. Our approach is inspired by Lawler, Bramson and Griffeath’s work [10]. We develop three original ideas: (i) we propose a cluster growth model, the *flashing process*, whose covering mechanism is simpler than internal DLA; (ii) we look at an intermediary scale, *the scale of cells*, since the deviations of the number of visits decrease with the cell-length; (iii) we build a coupling between *flashing process* and internal DLA which allows us to transport bounds from one model to the other.

Let us describe how the idea of an intermediary scale is used in the context of flashing processes. Recall that we first partition \mathbb{Z}^d into a sequence of concentric shells. Each shell is partitioned into cells whose side length equals the width of the shell. Now, we observe that a site has good chances to lie inside the cluster if some cell, say \mathcal{C} , about this site, is crossed by *many* explorers. The notation $W(\eta, \mathcal{C})$ refers to the number of explorers visiting \mathcal{C} , when their initial configuration is η . We drop the index n appearing in $W_n(\eta, z)$ since there are no more constraints on not escaping the ball $\mathbb{B}(0, n)$. Now, the coupon-collector nature of the covering mechanism suggests that for some positive constant α_d ,

$$(1.13) \quad \begin{aligned} W(\eta, \mathcal{C}) &\geq \alpha_d |\mathcal{C}| \times \log(|\mathcal{C}|) \\ &\implies \mathcal{C} \subset A(\emptyset, \eta) \text{ with a large probability.} \end{aligned}$$

We neglect in these heuristics the $\log(|\mathcal{C}|)$ term in (1.13).

Note that in [10], all the explorers start from the origin, whereas here, we only know that they *cross* \mathcal{C} . For internal DLA, estimating the probability that \mathcal{C} is not covered, when \mathcal{C} is large and $W(\eta, \mathcal{C}) \geq \alpha_d |\mathcal{C}|$ raises a difficulty which is absent when considering flashing processes.

We now make our argument more precise. For a scale h and an integer $K > 1$, to be determined, assume that $\mathbb{B}(0, n - Kh)$ is covered by settled explorers. Partition the shell $\mathcal{S} = \mathbb{B}(0, n - (K - 1)h) \setminus \mathbb{B}(0, n - Kh)$ into about $(n/h)^{d-1}$ cells, each of volume h^d . It is also convenient to stop the explorers as they reach the boundary of $\mathbb{B}(0, n - Kh)$. Thus, with such a stopped process, explorers are either settled inside $\mathbb{B}(0, n - Kh)$ or unsettled but stopped on its boundary, denoted $\partial\mathbb{B}(0, n - Kh)$. What we have called earlier *the number of explorers crossing \mathcal{C}* is taken here to be the unsettled explorers stopped on $\mathcal{C} \cap \partial\mathbb{B}(0, n - Kh)$.

Assuming (1.13) holds, it remains to show that the probability of the event $\{\exists \mathcal{C} \in \mathcal{S} : W(\eta, \mathcal{C}) < \alpha_d |\mathcal{C}|\}$ is small. We improve (1.11) by first using the independence between $W(\eta, \mathcal{C})$ and $M(\mathbb{B}(0, n - Kh), \mathcal{C})$, and then by replacing $A_R(\emptyset, \eta)$ by $\mathbb{B}(0, n - Kh)$ in (1.10) with $R = n - Kh$ and $\eta = |\mathbb{B}(0, n)|\mathbb{1}_0$,

$$(1.14) \quad W(\eta, \mathcal{C}) + M(\mathbb{B}(0, n - Kh), \mathcal{C}) \geq M(\eta, \mathcal{C}).$$

Also, we define

$$\mu(\mathcal{C}) = E[M(\eta, \mathcal{C})] - E[M(\mathbb{B}(0, n - Kh), \mathcal{C})].$$

Now, using that $M(\eta, z)$ and $M(\mathbb{B}(0, n - Kh), z)$ are sums of independent Bernoulli variables, we show that (1.14) implies a Gaussian-type lower tail

$$(1.15) \quad P(W(\eta, \mathcal{C}) < \alpha_d |\mathcal{C}|) \leq \exp\left(-\frac{(\mu(\mathcal{C}) - \alpha_d |\mathcal{C}|)^2}{c\nu(\mathcal{C})}\right)$$

for a positive constant c , and where $\nu(\mathcal{C})$

$$\nu(\mathcal{C}) = \text{var}(M(\eta, \mathcal{C})) - \text{var}(M(\mathbb{B}(0, n - Kh), \mathcal{C})).$$

We then show that both $\mu(\mathcal{C})$ and $\nu(\mathcal{C})$ are of order $K|\mathcal{C}|$. Then, $P(W(\eta, \mathcal{C}) < \alpha_d |\mathcal{C}|)$ is summable as soon as $K|\mathcal{C}| \geq A \log(n)$.

Outline of the paper. The rest of the paper is organized as follows. Section 2 introduces the main notation, and recalls known useful facts. In Section 3, we build the flashing process, give an alternative construction through exploration waves and sketch the proof of Lemma 1.3. In Section 4, we prove Propositions 1.5 and 1.6 using the construction in terms of exploration waves. In Section 5, we obtain a sharp estimate on the expected number of explorers crossing a given cell, and prove feature (ii) of the flashing times. Both proofs are based on classical potential theory estimates. Finally, in the Appendix, we give a proof of Lemma 1.3, and recall a result of Sébastien Blachère.

2. Notation and useful tools.

2.1. *Notation.* We say that $z, z' \in \mathbb{Z}^d$ are nearest neighbors when $\|z - z'\| = 1$, and we write $z \sim z'$. For any subset $\Lambda \subset \mathbb{Z}^d$, we define

$$\partial\Lambda = \{z \in \mathbb{Z}^d \setminus \Lambda : \exists z' \in \Lambda, z' \sim z\}.$$

For any $r \leq R$, we define the annulus

$$(2.1) \quad A(r, R) = B(0, R) \setminus B(0, r) \quad \text{and} \quad \mathbb{A}(r, R) = A(r, R) \cap \mathbb{Z}^d.$$

A trajectory S is a discrete nearest-neighbor path on \mathbb{Z}^d . That is, $S : \mathbb{N} \rightarrow \mathbb{Z}^d$ with $S(t) \sim S(t + 1)$ for all integer t . For a subset Λ in \mathbb{Z}^d , and a trajectory S , we define the hitting time of Λ as

$$H(\Lambda; S) = \min\{t \geq 0 : S(t) \in \Lambda\}.$$

We often omit S in the notation when no confusion is possible. We use the short-hand notation

$$B_n = B(0, n), \quad \mathbb{B}_n = \mathbb{B}(0, n), \quad H_R = H(B_R^c) \quad \text{and} \quad H_z = H(\{z\}).$$

For any a, b in \mathbb{R} we write $a \wedge b = \min\{a, b\}$, and $a \vee b = \max\{a, b\}$. Let Γ be a finite collection of trajectories on \mathbb{Z}^d . For $R > 0, z$ in \mathbb{Z}^d and Λ a subset of \mathbb{Z}^d , we call $M(\Gamma, R, z)$ [resp., $M(\Gamma, R, \Lambda)$] the number of trajectories which exit $\mathbb{B}(0, R)$ on z (resp., in Λ).

$$M(\Gamma, R, z) = \sum_{S \in \Gamma} \mathbf{1}_{\{S(H_R)=z\}} \quad \text{and} \quad M(\Gamma, R, \Lambda) = \sum_{z \in \Lambda} M(\Gamma, R, z).$$

When we deal with a collection of independent random trajectories, we rather specify its initial configuration $\eta \in \mathbb{N}^{\mathbb{Z}^d}$, so that $M(\eta, R, z)$ is the number of random walks starting from η and hitting $\mathbb{B}(0, R)^c$ on z . Two types of initial configurations are important here: (i) the configuration $n\mathbf{1}_{z^*}$ formed by n walkers starting on a given site z^* and (ii) for $\Lambda \subset \mathbb{Z}^d$, the configuration $\mathbf{1}_\Lambda$ that we simply identify with Λ . For any configuration $\eta \in \mathbb{N}^{\mathbb{Z}^d}$ we write

$$|\eta| = \sum_{z \in \mathbb{Z}^d} \eta(z).$$

For any $\Lambda \subset \mathbb{Z}^d$, we define Green’s function restricted to Λ, G_Λ , as follows. For $x, y \in \Lambda$, the expectation with respect to the law of the simple random walk started at x , is denoted with \mathbb{E}_x (the law is denoted \mathbb{P}_x) and

$$G_\Lambda(x, y) = \mathbb{E}_x \left[\sum_{0 \leq n < H(\Lambda^c)} \mathbf{1}_{\{S(n)=y\}} \right].$$

In dimension 3 or more, Green’s function on the whole space is well defined and denoted G . That is, for any $x, y \in \mathbb{Z}^d$,

$$G(x, y) = \mathbb{E}_x \left[\sum_{n \geq 0} \mathbf{1}_{\{S(n)=y\}} \right].$$

In dimension 2, the potential kernel plays the role of Green’s function

$$a(x, y) = \lim_{n \rightarrow \infty} \mathbb{E}_x \left[\sum_{l=0}^n (\mathbf{1}_{\{S(l) = x\}} - \mathbf{1}_{\{S(l) = y\}}) \right].$$

2.2. *Some useful tools.* We recall here some well-known facts. Some of them are proved for the reader’s convenience. This section can be skipped at a first reading.

In [10], the authors emphasized the fact that the spherical limiting shape of internal DLA was intimately linked to strong isotropy properties of Green’s function. This isotropy is expressed by the following asymptotics (Theorem 4.3.1 of [11]). In $d \geq 3$, there is a constant K_g , such that for any $z \neq 0$,

$$(2.2) \quad \left| G(0, z) - \frac{C_d}{\|z\|^{d-2}} \right| \leq \frac{K_g}{\|z\|^d} \quad \text{with } C_d = \frac{2}{v_d(d-2)},$$

where v_d stands for the volume of the Euclidean unit ball in \mathbb{R}^d . The first order expansion (2.2) is proved in [11] for general symmetric walks with finite $d + 3$ moments and vanishing third moment. All the estimates we use are eventually based on (2.2), and we emphasize the fact that the estimate is uniform in $\|z\|$. There is a similar expansion for the potential kernel. Theorem 4.4.4 of [11] establishes that for $z \neq 0$ (with γ the Euler constant),

$$(2.3) \quad \left| a(0, z) - \frac{2}{\pi} \log(\|z\|) - \frac{2\gamma + \log(8)}{\pi} \right| \leq \frac{K_g}{\|z\|^2}.$$

We recall a rough but useful result about the exit site distribution from a sphere. This is Lemma 1.7.4 of [8].

LEMMA 2.1. *There are two positive constants c_1, c_2 such that for any $z \in \partial B(0, n)$, and $n > 0$*

$$(2.4) \quad \frac{c_1}{n^{d-1}} \leq \mathbb{P}_0(S(H_n) = z) \leq \frac{c_2}{n^{d-1}}.$$

We now state an elementary lemma.

LEMMA 2.2. *Each z^* in $\mathbb{Z}^d \setminus \{0\}$ has a nearest-neighbor z (i.e., $z^* \sim z$) such that*

$$(2.5) \quad \|z\| \leq \|z^*\| - \frac{1}{2\sqrt{d}}.$$

PROOF. Without loss of generality we can assume that all the coordinates of z^* are nonnegative. Let us denote by b the maximum of these coordinates, and note that

$$(2.6) \quad \|z^*\|^2 \leq db^2 \quad \text{and} \quad b \geq 1.$$

Denote by z the nearest-neighbor obtained from z^* by decreasing by one unit a maximum coordinate. Using (2.6),

$$(2.7) \quad \|z^*\|^2 - \|z\|^2 = b^2 - (b - 1)^2 = 2b - 1 \geq b \geq \frac{\|z^*\|}{\sqrt{d}}.$$

Note that (2.5) follows from $2\|z^*\|(\|z^*\| - \|z\|) \geq \|z^*\|^2 - \|z\|^2$, and (2.7). \square

We state now a handy estimate dealing with sums of independent Bernoulli variables.

LEMMA 2.3. *Let $\{X_n, Y_n, n \in \mathbb{N}\}$ be independent 0–1 Bernoulli variables. For integers n, m let $S = X_1 + \dots + X_n$ and $S' = Y_1 + \dots + Y_m$. Define for $t \in \mathbb{R}$*

$$f(t) = e^t - 1 - t \quad \text{and} \quad g(t) = (e^t - 1)^2.$$

If $0 \leq t \leq \log(2)$, then

$$(2.8) \quad \frac{E[\exp(t(S - E[S]))]}{E[\exp(t(S' - E[S']))]} \leq \exp\left(f(t)E[S - S'] + g(t) \sum_{i=1}^m E[Y_i]^2\right).$$

Assume now that for $\kappa > 1$, $\sup_n E[Y_n] \leq \frac{\kappa-1}{\kappa}$. If $t \leq 0$, then

$$(2.9) \quad \frac{E[\exp(t(S - E[S]))]}{E[\exp(t(S' - E[S']))]} \leq \exp\left(f(t)E[S - S'] + \frac{\kappa}{2}g(t) \sum_{i=1}^m E[Y_i]^2\right).$$

PROOF. Let X be a Bernoulli variable, and $p = E[X]$. Using the inequality $e^x \geq 1 + x$ for $x \in \mathbb{R}$, we have

$$(2.10) \quad \begin{aligned} E[\exp(t(X - E[X]))] &= pe^{t(1-p)} + (1-p)e^{-tp} \\ &= e^{-pt}(1 + p(e^t - 1)) \\ &\leq \exp(f(t)E[X]). \end{aligned}$$

For a lower bound, we distinguish two cases.

First, assume $t \geq 0$. We claim that $\exp(x - x^2) \leq 1 + x$ for $0 \leq x \leq 1$. Indeed, we use three obvious inequalities: $e^x \geq 1 + x$ for $x \in \mathbb{R}$, (i) for $x \leq 1$, $1 + x + x^2 \geq e^x$, and (ii) $(1 + x^2)(1 + x) \geq 1 + x + x^2$. Thus

$$e^{x^2}(1 + x) \geq (1 + x^2)(1 + x) \geq 1 + x + x^2 \geq e^x.$$

This yields the claim. Now, set $x = p(e^t - 1)$, so that $x \leq 1$ when $e^t \leq 2$. The last inequality in (2.10) yields

$$(2.11) \quad \begin{aligned} E[\exp(t(X - E[X]))] &\geq \exp(-tp + p(e^t - 1) - p^2(e^t - 1)^2) \\ &= e^{f(t)p - g(t)p^2}. \end{aligned}$$

Assume now that $t \leq 0$, and for $\kappa > 1$, $p < \frac{\kappa-1}{\kappa}$. We claim that for $0 \leq x \leq \frac{\kappa-1}{\kappa}$,

$$(2.12) \quad \exp\left(-x - \frac{\kappa}{2}x^2\right) \leq 1 - x.$$

Indeed, we have an additional inequality (iii) $1 - x + \frac{x^2}{2} \geq \exp(-x)$ when $x \geq 0$. Note also that

$$\left(1 + \frac{\kappa}{2}x^2\right)(1 - x) \geq 1 - x + \frac{x^2}{2} \iff x \leq \frac{\kappa - 1}{\kappa}.$$

Thus

$$e^{\kappa x^2/2}(1 - x) \geq \left(1 + \frac{\kappa}{2}x^2\right)(1 - x) \geq 1 - x + \frac{x^2}{2} \geq e^{-x}.$$

Now, set $x = -p(e^t - 1) \geq 0$, so that $x \leq \frac{\kappa-1}{\kappa}$. We obtain

$$\begin{aligned}
 E[\exp(t(X - E[X]))] &\geq \exp\left(-tp + p(e^t - 1) - \frac{\kappa}{2}p^2(e^t - 1)^2\right) \\
 (2.13) \qquad \qquad \qquad &= e^{f(t)p - \kappa g(t)p^2/2}.
 \end{aligned}$$

Inequalities (2.8) and (2.9) follow (2.11) and (2.13). \square

3. The flashing process. In this section, we construct the flashing process, and state the crucial “uniform hitting property.” We then present a useful equivalent construction in terms of exploration waves. Finally, we explain the coupling of Lemma 1.3, but postpone its proof to the [Appendix](#).

3.1. *Construction of the process.*

Partitioning the lattice. We are given a sequence $\mathcal{H} = \{h_n, n \in \mathbb{N}\}$. We partition the lattice into shells $(\mathcal{S}_j : j \geq 0)$. For an illustration, see Figure 1. For a given parameter $h_0 > 0$, the first shell \mathcal{S}_0 is the ball $\mathbb{B}(0, h_0)$. For $j \geq 1$, shell j is the annulus [see its definition (2.1)]

$$\mathcal{S}_j = \mathbb{A}(r_j - h_j, r_j + h_j),$$

where $\{r_j, j \geq 1\}$ is defined inductively by $r_1 = h_0 + h_1$, and for $j \geq 1$,

$$r_{j+1} - h_{j+1} = r_j + h_j.$$

In Section 4, we need that (o) \mathcal{H} is increasing, (i) $j \mapsto h_j/r_j$ is decreasing and (ii) $h_j = O(r_j^{1/3})$. These properties are a straightforward consequence of our hypothesis $h_j \leq h_{j+1} \leq (1 + \frac{1}{2j})h_j$. Actually we will only need these properties, and our hypothesis is no more than a sufficient condition.

We also define

$$\Sigma_0 = \{0\} \quad \text{and} \quad \Sigma_j = \partial\mathbb{B}(0, r_j), \quad j \geq 1.$$

Flashing times. The key feature we expect from the *flashing process* is that its covering mechanism be simple. More precisely, our construction is guided by property (ii) of the [Introduction](#) which states that *the flashing position, in a shell, is essentially uniform over the cell an explorer first hits upon entering the shell*. Thus, we need to define together *cells* and *flashing times* to realize property (ii). It is important that all sites of a shell can be chosen as *flashing sites* with about the same frequency. In this respect, let us remark that a cell in shell \mathcal{S}_j cannot be a ball of radius h_j centered on Σ_j . Indeed, if this were the case, sites at a distance about h_j would be in much fewer cells than sites of Σ_j , and this would fail to make the covering of a shell uniform. We find it convenient to build a cell with a *mixture* of balls and annuli. A (random) flag Y_j tells the explorers whether it flashes upon

exiting either a sphere or the boundary of an annulus, whose distance from Σ_j is governed with a random radius R_j of appropriate density. Also, to allow for the possibility of flashing on its hitting position on Σ_j , we introduce an additional flag X_j .

More precisely, consider $\{X_j, Y_j, j \geq 0\}$ a sequence of independent Bernoulli variables such that

$$P(X_j = 1) = 1 - P(X_j = 0) = \frac{1}{h_j^d}$$

and

$$P(Y_j = 1) = 1 - P(Y_j = 0) = \begin{cases} 1, & \text{if } j = 0, \\ \frac{1}{2}, & \text{if } j \geq 1. \end{cases}$$

Consider also a sequence of continuous independent variables $\{R_j, j \geq 0\}$ each of which has density $g_j : [0, h_j] \rightarrow \mathbb{R}^+$ with

$$(3.1) \quad g_j(h) = \frac{dh^{d-1}}{h_j^d}.$$

For $j \geq 0$, and z_j in Σ_j , let S be a random walk starting in z_j , and define a stopping time σ as follows. If $R_j = h$ for some $h \leq h_j$, then

$$\sigma = \begin{cases} 0, & \text{if } X_j = 1, \\ H(\mathbb{B}(z_j, h \wedge (r_j + h_j - \|z_j\|))^c), & \text{if } X_j = 0 \text{ and } Y_j = 1, \\ H(\mathbb{A}(r_j - h, r_j + h)^c), & \text{if } X_j = 0 \text{ and } Y_j = 0. \end{cases}$$

We set $H_j = H(\Sigma_j)$, and we define the stopping times $(\sigma_j : j \geq 0)$ as

$$\sigma_j = H_j + \sigma(S \circ \theta_{H_j}),$$

where θ stands for the usual time-shift operator. For $j \geq 0$ we note that, by construction, $S(t) \in S_j$ for all t such that $H_j \leq t < \sigma_j$ and we say that σ_j is a *flashing time* when $S(\sigma_j)$ is contained in the intersection between S_j and the cone with base $B(S(H_j), h_j/2)$. We call such an intersection *a cell centered at $S(H_j)$* , that we denote $\mathcal{C}(S(H_j))$. In other words, for any $z \in \Sigma_j$

$$(3.2) \quad \mathcal{C}(z) = S_j \cap \{x \in \mathbb{R}^d : \exists \lambda \geq 0, \exists y \in B(z, h_j/2), x = \lambda y\}.$$

The uniform hitting property. The main property of the hitting time σ constructed above is the following proposition, which yields property (ii) of the flashing process to be defined soon.

PROPOSITION 3.1. *There are two positive constants $\alpha_1 < \alpha_2$, such that, for h_0 large enough, $j \geq 0$, $z_j \in \Sigma_j$, and $z^* \in \mathcal{C}(z_j)$.*

$$(3.3) \quad \frac{\alpha_1}{h_j^d} \leq \mathbb{P}_{z_j}(S(\sigma) = z^*) \leq \frac{\alpha_2}{h_j^d}.$$

The proof of Proposition 3.1 is given in Section 5.

The flashing process. Consider a family of N independent random walks $(S_i : 1 \leq i \leq N)$ with their stopping times $(H_{i,j}, \sigma_{i,j} : j \geq 0)$. Let also $z_{i,j} = S_i(H_{i,j})$ be the first hitting position of S_i on Σ_j .

We define the cluster inductively. Set $A^*(0) = \emptyset$. For $i \geq 1$, we define τ_i^* as the first flashing time associated with S_i when the explorer stands outside $A^*(i - 1)$. In other words,

$$\tau_i^* = \min\{\sigma_{i,j} : j \geq 0, S_i(\sigma_{i,j}) \in \mathcal{C}(z_{i,j}) \cap A^*(i - 1)^c\}$$

and

$$A^*(i) = A^*(i - 1) \cup \{S_i(\tau_i^*)\}.$$

3.2. Exploration waves. Rather than building $A^*(N)$ following the whole journey of one explorer after another, we can build $A^*(N)$ as an increasing union of clusters formed by stopping explorers on successive shells. Similar wave constructions are introduced in [10] and [9]. We use this alternative construction in the proof of Propositions 1.5 and 1.6.

We denote by $\xi_k \in (\mathbb{Z}^d)^N$ the explorers positions after the k th wave. We denote by $\mathcal{A}_k^*(N)$ the set of sites where settled explorers are after the k th wave. Our inductive construction will be such that

$$\xi_k(i) \notin \Sigma_k \iff \xi_k(i) \in \bigcup_{j < k} S_j \iff \xi_k(i) \in \mathcal{A}_k^*(N).$$

For $k = 0$ we set $\xi_0(i) = 0$, and $\mathcal{A}_0^*(i) = \emptyset$, for $1 \leq i \leq N$. Assume that for $k \geq 0$, $\mathcal{A}_k^*(i)$ is built for $i = 0, \dots, N$. We set $\mathcal{A}_{k+1}^*(0) = \mathcal{A}_k^*(N)$. For i in $\{1, \dots, N\}$, we set the following:

- If $\xi_k(i) \notin \Sigma_k$, then

$$\xi_{k+1}(i) = \xi_k(i) \in \bigcup_{j < k} S_j \quad \text{and} \quad \mathcal{A}_{k+1}^*(i) = \mathcal{A}_{k+1}^*(i - 1).$$

- If $\xi_k(i) \in \Sigma_k$ and $S_i(\sigma_{i,k}) \in \mathcal{C}(z_{i,k}) \cap \mathcal{A}_k^*(i - 1)^c$, then

$$\xi_{k+1}(i) = S_i(\sigma_{i,k}) \in S_k \quad \text{and} \quad \mathcal{A}_{k+1}^*(i) = \mathcal{A}_{k+1}^*(i - 1) \cup \{S_i(\sigma_{i,k})\}.$$

- If $\xi_k(i) \in \Sigma_k$ and $S_i(\sigma_{i,k}) \notin \mathcal{C}(z_{i,k}) \cap \mathcal{A}_k^*(i - 1)^c$, then

$$\xi_{k+1}(i) = S_i(H_{i,k+1}) \in \Sigma_{k+1} \quad \text{and} \quad \mathcal{A}_{k+1}^*(i) = \mathcal{A}_{k+1}^*(i - 1).$$

In words, for each $k \geq 1$, during the k th wave of exploration, the unsettled explorers move one after the other in the order of their labels until either settling in S_{k-1} , or reaching Σ_k where they stop. We then define $\mathcal{A}^*(N)$ by

$$\mathcal{A}^*(N) = \bigcup_{k \geq 1} \mathcal{A}_k^*(N).$$

We explain now why this construction yields the same cluster as our previous definition. An explorer cannot settle inside a shell it has left, and thus cannot settle in any shell \mathcal{S}_j with $j < k$ if it reaches Σ_k . Now, since each wave of exploration is organized according to the label ordering, the fact that an explorer has to wait for the following explorers before proceeding its journey beyond Σ_k does not interfere with the site where it eventually settles.

3.3. Coupling internal DLA and flashing processes.

Proof of Lemma 1.3. For each positive integer N , we build a coupling between $A(N)$ and $A^*(N)$. We first describe the main features of our coupling in words. Its precise definition is postponed to the [Appendix](#).

We launch N independent random walks, and build inductively the associated clusters $A(1), A(2), \dots, A(N)$. In doing so, we use the increments of these random walks to define, step by step, N flashing trajectories S_1^*, \dots, S_N^* up to some times $\bar{t}_1, \dots, \bar{t}_N$. Let us describe informally step $i + 1$ of the induction. Assume that S_1^*, \dots, S_i^* are defined up to some times $t_1 \leq \bar{t}_1, \dots, t_i \leq \bar{t}_i$, and that each site of $A(i)$ is covered by exactly one $S_k^*(t_k)$ with $1 \leq k \leq i$. We can think of $S_1^*(t_1), \dots, S_i^*(t_i)$ as the positions of stopped flashing explorers, some of them stopped at one of their flashing times—say on *blue* sites—some of them not—say on *red* sites. Then, we add the $i + 1$ th explorer and flashing explorer. We set $S_{i+1}^*(0) = S_{i+1}(0) = 0$. We add new increments both to S_{i+1} and to the trajectory of one flashing explorer, say with label j in $\{1; \dots; i + 1\}$, in such a way that the current position of the walker $i + 1$ and that of the flashing explorer j coincide. The label j is defined inductively as follows. Initially, $j = i + 1$. Assume now that the walker $i + 1$ flashes on a red or blue site inside $A(i)$. This site is occupied by exactly two stopped flashing explorers, j and j' [and all other red and blue sites of $A(i)$ are occupied by exactly one flashing explorer]. Since flashing explorers can settle at their flashing times, it makes sense, when j is flashing, to add the next increment to the trajectory of flashing explorer j' rather than j . We do so in two cases, first, when this happens on a red site. In this case, we turn blue that site since j is stopped at a flashing time. Second, when this happens on a blue site, say z , and $j' > j$. Note that in this case, both explorers flash on z , but explorer j reaches z before explorer j' when launched in their label order. Our choice is such that the eventual cluster $A^*(N)$ has the correct law. In all other cases, we keep adding the increments of S_{i+1} to the same flashing trajectory. It is important to note that the value of the increment does not depend on the index of the trajectory we choose to extend. Walker $i + 1$ eventually steps outside $A(i)$, say on z^* , while following a flashing trajectory, say the j th one. We stop the j th flashing trajectory on z^* , and paint z^* blue or red according to whether z^* is one of its flashing sites or not.

When the last walker steps outside $A(N - 1)$, we have

$$(3.4) \quad A(N) = \{S_1^*(\bar{t}_1); \dots; S_N^*(\bar{t}_N)\} \quad \text{with } |A(N)| = N.$$

To define $A^*(N)$ we launch again, in their label's order, the flashing explorers from their current positions (possibly some or none of them since some or all of them can already have reached their settling position). We then get

$$(3.5) \quad A^*(N) = \{S_1^*(\tau_1^*); \dots; S_N^*(\tau_N^*)\}$$

with $|A^*(N)| = N$ and $\tau_k^* \geq \bar{t}_k$ for all k .

Proof of Corollary 1.4. Since a flashing explorer that visited some site beyond a given shell cannot settle in that shell, the one-to-one map

$$(3.6) \quad \psi_N : S_k^*(\bar{t}_k) \in A(N) \mapsto S_k^*(\tau_k^*) \in A^*(N), \quad k = 1, \dots, N,$$

satisfies, for all k and l ,

$$(3.7) \quad S_k^*(\bar{t}_k) \notin \bigcup_{m < l} \mathcal{S}_m \Rightarrow S_k^*(\tau_k^*) = \psi_N(S_k^*(\bar{t}_k)) \notin \bigcup_{m < l} \mathcal{S}_m.$$

Thus, for all $N \geq 0$ there is a coupling and a one-to-one map ψ_N between $A(N)$ and $A^*(N)$ such that for all $k \geq 1$,

$$(3.8) \quad \psi_N(A(N) \cap \mathbb{B}_{r_k+h_k}^c) \subset A^*(N) \cap \mathbb{B}_{r_k+h_k}^c.$$

Inclusion (3.8) has two important consequences:

- (a) If $A^*(N) \subset \mathbb{B}_{r_k+h_k}$, then $A(N) \subset \mathbb{B}_{r_k+h_k}$. Indeed, any site in $A(N)$ outside $\mathbb{B}_{r_k+h_k}$ produces, through ψ_N , a site in $A^*(N)$ outside $\mathbb{B}_{r_k+h_k}$.
- (b) If $\mathbb{B}_{r_k+h_k} \subset A^*(N)$, then $\mathbb{B}_{r_k+h_k} \subset A(N)$. Indeed, those sites in $A(N)$ that are mapped through ψ_N on $A^*(N) \cap \mathbb{B}_{r_k+h_k} = \mathbb{B}_{r_k+h_k}$ are necessarily contained in $\mathbb{B}_{r_k+h_k}$. Since their number is $|\mathbb{B}_{r_k+h_k}|$ and ψ_N is one-to-one, they completely cover $\mathbb{B}_{r_k+h_k}$.

4. Fluctuations. In this section, we prove Propositions 1.5 and 1.6. To do so we use the construction in terms of exploration waves of Section 3.2. Thus, we think of the growing cluster as evolving in discrete time, where time counts the number of exploration waves. The proofs in this section rely on potential theory estimates which we have gathered in Section 5, for the ease of reading.

4.1. *Tiles.* We recall that we have defined a *cell* of \mathcal{S}_j in (3.2), as the intersection of a cone with \mathcal{S}_j . We need also a smaller shape. We define, for any z_j in Σ_j , and for a small ε_0 to be defined later,

$$(4.1) \quad \tilde{\mathcal{C}}(z_j) = \mathcal{S}_j \cap \{x \in \mathbb{R}^d : \exists \lambda \geq 0, \exists y \in B(z_j, \varepsilon_0 h_j), x = \lambda y\}.$$

As in Lemma 12 in [9], concerning locally finite coverings, we claim that, for h_0 large enough, there exist a positive constant K_F , and, for each $j \geq 0$, a subset $\tilde{\Sigma}_j$ of Σ_j such that

$$(4.2) \quad \forall y \in \mathcal{S}_j \quad |\{z \in \tilde{\Sigma}_j : y \in \tilde{\mathcal{C}}(z)\}| \leq K_F \quad \text{and} \quad \mathcal{S}_j = \bigcup_{z_j \in \tilde{\Sigma}_j} \tilde{\mathcal{C}}(z_j).$$

For any $z_j \in \tilde{\Sigma}_j$, we call *tile centered at z_j* , the intersections of $\tilde{\mathcal{C}}(z_j)$ with Σ_j . We denote by $\mathcal{T}(z_j)$ a tile centered at z_j , and by \mathcal{T}_j the set of tiles associated with the shell \mathcal{S}_j .

$$(4.3) \quad \mathcal{T}_j = \{\mathcal{T}(z_j) : z_j \in \tilde{\Sigma}_j\}.$$

We choose ε_0 to satisfy two properties. First, for any $z \in \mathcal{S}_j$, there is $\tilde{z}_j \in \tilde{\Sigma}_j$ such that

$$(4.4) \quad z \in \bigcap_{y \in \mathcal{T}(\tilde{z}_j)} \mathcal{C}(y).$$

This is ensured by the choice of a small enough ε_0 . Indeed, let $z_j \in \Sigma_j$ be a site realizing the minimum of $\{\|z - y\| : y \in \Sigma_j\}$. There is $\lambda > 0$ and $u \in B(z_j, 1)$, such that $z = \lambda u$. Now, there is $\tilde{z}_j \in \tilde{\Sigma}_j$ such that $\|\tilde{z}_j - z_j\| < \varepsilon_0 h_j$, and for any $y \in \mathcal{T}(\tilde{z}_j)$, we have $\|y - z_j\| < 2\varepsilon_0 h_j$. Thus, for ε_0 small enough so that $1 + 2\varepsilon_0 h_j \leq h_j/2$,

$$\forall y \in \mathcal{T}(\tilde{z}_j) \quad \|u - y\| \leq \|u - z_j\| + \|z_j - y\| \leq 1 + 2\varepsilon_0 h_j \leq \frac{h_j}{2},$$

which implies (4.4). Second, the size of a tile should be such that for some $\kappa > 1$, for any $j \geq 1$, and any tile $\mathcal{T} \in \mathcal{T}_j$

$$(4.5) \quad \sup_{z \in \mathbb{B}(0, r_j - h_j)} \mathbb{P}_z(\mathcal{S}(H(\Sigma_j)) \in \mathcal{T}) \leq \frac{\kappa - 1}{\kappa}.$$

Inequality (4.5) follows from Lemma 5(b) of [10] (or Lemma 5.1 below) which for a constant J_d yields

$$\sup_{z \in \mathbb{B}(0, r_j - h_j)} \mathbb{P}_z(\mathcal{S}(H(\Sigma_j)) \in \mathcal{T}) \leq J_d \frac{|\mathcal{T}|}{h_j^{d-1}}.$$

The choice of ε_0 is such that $J_d |\mathcal{T}| \leq \frac{\kappa - 1}{\kappa} h_j^{d-1}$.

4.2. *Bounding inner fluctuations.* For $n \geq 0$, we take $N = |\mathbb{B}_n|$, we recall that $\mathcal{A}_k^*(N) \subset \mathcal{A}_{k+1}^*(N)$ for $k \in \mathbb{N}$, and $\mathcal{A}^*(N) = \bigcup_{k \geq 1} \mathcal{A}_k^*(N)$. We consider

$$(4.6) \quad T^* = \min \left\{ k \geq 1 : \bigcup_{j < k} \mathcal{S}_j \not\subset \mathcal{A}_k^*(N) \right\}.$$

Note that $\mathcal{A}_k^*(N) \subset \bigcup_{j < k} \mathcal{S}_j$, so that T^* is the first time k when the k th wave does not cover all its allowed space. We recall that time counts the number of exploration waves.

For the flashing process if $\bigcup_{j < k} \mathcal{S}_j \not\subset \mathcal{A}_k^*(N)$, then for any $k' > k$, we have $\bigcup_{j < k} \mathcal{S}_j \not\subset \mathcal{A}_{k'}^*(N)$, so that T^* is also the shell label where the first hole of $\mathcal{A}^*(N)$ appears. We have, for l with $r_l < n$,

$$(4.7) \quad P(T^* \leq l) = P(\mathbb{B}(0, r_l + h_l) \not\subset \mathcal{A}^*(N)) \leq \sum_{k \leq l} P(T^* = k + 1).$$

In this section, we estimate from above the probability $P(T^* = k + 1)$ assuming $r_k < n$.

For $k \geq 1$ and $\Lambda \subset \Sigma_k$, we call $W_k(\Lambda)$ the number of unsettled explorers that stand in Λ after the k th wave, that is,

$$(4.8) \quad W_k(\Lambda) = \sum_{i=1}^N \mathbf{1}_\Lambda(\xi_k(i)).$$

We now look at the *crossings* of tiles of \mathcal{T}_k . On the one hand, we will use that if $W_k(\mathcal{T})$ is *large*, then it is unlikely that a hole appears in the cell containing \mathcal{T} during the $k + 1$ th-wave. We use for this purpose the fact that covering for the flashing process is similar to filling an album for a coupon-collector model. On the other hand, if r_k is *small*, it is unlikely that $W_k(\mathcal{T})$ is *small*. We now make precise what we intend by *small* and *large*. For any positive constant ξ , we write

$$(4.9) \quad \begin{aligned} P(T^* = k + 1) &= P(T^* = k + 1, \forall \mathcal{T} \in \mathcal{T}_k, W_k(\mathcal{T}) \geq \xi) \\ &\quad + P(T^* = k + 1, \exists \mathcal{T} \in \mathcal{T}_k, W_k(\mathcal{T}) < \xi) \\ &\leq P(T^* = k + 1 | \forall \mathcal{T} \in \mathcal{T}_k, W_k(\mathcal{T}) \geq \xi) \\ &\quad + P(\exists \mathcal{T} \in \mathcal{T}_k, W_k(\mathcal{T}) < \xi). \end{aligned}$$

A coupon-collector estimate. The first term in the right-hand side of (4.9) is bounded using a simple coupon-collector argument. Indeed, the event $\{T^* = k + 1\}$ implies that there is an uncovered site in \mathcal{S}_k , say z , when explorers stopped in Σ_k are released. By (4.4), there is $z_k \in \tilde{\Sigma}_k$, such that z is a possible settling position of all explorers stopped in $\mathcal{T}(z_k)$. Now, knowing that $\{W_k(\mathcal{T}(z_k)) \geq \xi\}$, Proposition 3.1 tells us that the probability of not covering this site is less than $(1 - \alpha_1/h_k^d)$ to the power ξ . In other words,

$$P(T^* = k + 1 | \forall \mathcal{T} \in \mathcal{T}_k, W_k(\mathcal{T}) \geq \xi) \leq |\mathcal{S}_k| \left(1 - \frac{\alpha_1}{h_k^d}\right)^\xi \leq |\mathcal{S}_k| \exp\left(-\alpha_1 \frac{\xi}{h_k^d}\right).$$

Henceforth, we set

$$(4.10) \quad \xi = Ah^d \log(n) \quad \text{with } h = \sup\{h_k : r_k \leq n\}$$

and A large enough so that

$$(4.11) \quad \begin{aligned} &\sum_{k: r_k < n} P(T^* = k + 1 | \forall \mathcal{T} \in \mathcal{T}_k, W_k(\mathcal{T}) \geq \xi) \\ &\leq |\mathbb{B}_n| \exp(-\alpha_1 A \log n) \leq \frac{1}{n^2}. \end{aligned}$$

Estimating $\{W_k(\mathcal{T}) < \xi\}$. For any $\mathcal{T} \in \mathcal{T}_k$, we consider the counting variable $L_k(\mathcal{T}) = M(\mathbb{B}(0, r_k - h_k), r_k, \mathcal{T})$, and define

$$(4.12) \quad M_k(\mathcal{T}) = W_k(\mathcal{T}) + M(A_k^*, r_k, \mathcal{T})$$

so that $M_k(\mathcal{T}) \stackrel{\text{law}}{=} M(N\mathbf{1}_{\{0\}}, r_k, \mathcal{T})$.

The idea of defining M_k and L_k (for the internal DLA process), and bounding W_k by $M_k - L_k$, is introduced in [10]. Our main observation is that $L_k(\mathcal{T})$ is independent of $W_k(\mathcal{T})$, and

$$W_k(\mathcal{T}) + L_k(\mathcal{T}) \geq M_k(\mathcal{T}).$$

As a consequence, for any positive constants t and ξ (and with the notation $\bar{X} = X - E[X]$),

$$\begin{aligned} P(W_k(\mathcal{T}) < \xi) &\leq e^{t\xi} \times E[\exp(-tW_k(\mathcal{T}))] = e^{t\xi} \frac{E[\exp(-t(W_k(\mathcal{T}) + L_k(\mathcal{T})))]}{E[\exp(-tL_k(\mathcal{T}))]} \\ &\leq \exp(-t(E[M_k(\mathcal{T}) - L_k(\mathcal{T})] - \xi)) \times \frac{E[\exp(-t(\bar{M}_k(\mathcal{T})))]}{E[\exp(-t\bar{L}_k(\mathcal{T}))]}. \end{aligned}$$

Using Lemma 2.3 with condition (4.5), we obtain

$$\begin{aligned} \log P(W_k(\mathcal{T}) < \xi) &\leq -t(E[M_k(\mathcal{T}) - L_k(\mathcal{T})] - \xi) + f(-t)E[M_k(\mathcal{T}) - L_k(\mathcal{T})] \\ &\quad + \frac{\kappa}{2}g(-t) \sum_{y \in \mathbb{B}(0, r_k - h_k)} P_y^2(S(H(\Sigma_k)) \in \mathcal{T}). \end{aligned}$$

We now proceed in two steps. We show in step 1 that for some constant κ' ,

$$E[M_k(\mathcal{T}) - L_k(\mathcal{T})] \geq \kappa'(n^d - (r_k - h_k)^d) \frac{h_k^{d-1}}{r_k^{d-1}}.$$

Since $\{h_k/r_k, k \geq 0\}$ is nonincreasing, it follows that there is a constant $\kappa_1 > 0$ such that, for all $\alpha > 0$, and $k_\alpha := \sup\{j \in \mathbb{N} : r_j < n - \alpha h \log n\}$, where h is defined in (4.10), we have

$$\inf_{k \leq k_\alpha} E[M_k(\mathcal{T}) - L_k(\mathcal{T})] \geq \kappa'(n^d - (n - h)^d) \frac{h^{d-1}}{n^{d-1}} \geq \kappa_1 \alpha h^d \log n.$$

Now, if we choose ξ as in (4.10), with $\alpha = 2A/\kappa_1$ and $k^* = k_\alpha$, that is,

$$k^* := \sup \left\{ j \in \mathbb{N} : r_j \leq n - \frac{2A}{\kappa_1} h \log(n) \right\},$$

then, we get, for all $k \leq k^*$,

$$(4.13) \quad E[M_k(\mathcal{T}) - L_k(\mathcal{T})] \geq 2\xi.$$

We show in step 2, that for a constant C depending on the dimension only

$$(4.14) \quad \sum_{y \in \mathbb{B}(0, r_k - h_k)} \mathbb{P}_y^2(S(H(\Sigma_k)) \in \mathcal{T}) \leq CE[M_k(\mathcal{T}) - L_k(\mathcal{T})].$$

Suppose for a moment that steps 1 and 2 hold. Since, for some $c > 0$, $\max(f(-t), g(-t)) \leq ct^2$ when $t \leq 1$, there is $c' > 0$ such that for $k \leq k^*$

$$(4.15) \quad \begin{aligned} & \log P(W_k(\mathcal{T}) < Ah^d \log(n)) \\ & \leq \inf_{0 \leq t \leq 1} \left(-t + c \left(1 + \frac{C\kappa}{2} \right) t^2 \right) E[M_k(\mathcal{T}) - L_k(\mathcal{T})] \\ & \leq -c' E[M_k(\mathcal{T}) - L_k(\mathcal{T})] \leq -2c' Ah^d \log(n). \end{aligned}$$

Now, using (4.9), (4.11) and (4.15) for A large enough, we have

$$\sum_{k < k^*} P(T^* = k) \leq \frac{2}{n^2}.$$

Borel–Cantelli’s lemma yields then the inner control of Proposition 1.5.

Step 1. We invoke Corollary 5.4, with $n = r_k$, and $\Delta_n = h_k$ [the hypotheses $h_k = O(r_k^{1/3})$ and h_k large enough hold here, as seen in the first paragraph of Section 3.1]. We have for some positive constants κ' , K and for n large enough,

$$(4.16) \quad \begin{aligned} E[M_k(\mathcal{T}) - L_k(\mathcal{T})] &= E[M(|\mathbb{B}_n| - |\mathbb{B}_{r_k - h_k}|)\mathbf{1}_0, r_k, \mathcal{T})] \\ &\quad + E[M(|\mathbb{B}_{r_k - h_k}|\mathbf{1}_0, r_k, \mathcal{T})] - E[M(\mathbb{B}_{r_k - h_k}, r_k, \mathcal{T})] \\ &\geq (|\mathbb{B}_n| - |\mathbb{B}_{r_k - h_k}|)\mathbb{P}_0(S(H_k) \in \mathcal{T}) - Kh_k^{d-1} \\ &\geq 2\kappa'(n^d - (r_k - h_k)^d) \frac{h_k^{d-1}}{r_k^{d-1}} - Kh_k^{d-1} \\ &\geq \kappa'(n^d - (r_k - h_k)^d) \frac{h_k^{d-1}}{r_k^{d-1}} \end{aligned}$$

for $r_k \leq n$ and h_0 large enough.

Step 2. By Lemma 5.1 below, there is a constant κ_G such that, for $y \in \mathbb{B}(0, r_k - h_k)$, and $z \in \tilde{\Sigma}_k$

$$\mathbb{P}_y(S(H(\Sigma_k)) \in \mathcal{T}(z)) \leq \frac{\kappa_G |\mathcal{T}(z)|}{\|z - y\|^{d-1}}.$$

Therefore

$$(4.17) \quad \sum_{y \in \mathbb{B}(0, r_k - h_k)} \mathbb{P}_y^2(S(H(\Sigma_k)) \in \mathcal{T}(z)) \leq \sum_{j: h_k \leq j \leq 2r_k} \sum_{y: j \leq |z - y| < j+1} \frac{\kappa_G^2 |\mathcal{T}(z)|^2}{j^{2(d-1)}}.$$

For a constant C_d , we bound $|\{y : k \leq |z - y| < k + 1\}| \leq C_d k^{d-1}$. Thus,

$$\begin{aligned}
 & \sum_{y \in \mathbb{B}(0, r_k - h_k)} \mathbb{P}_y^2(S(H(\Sigma_k)) \in \mathcal{T}(z)) \\
 (4.18) \quad & \leq \sum_{j : h_k \leq j \leq 2r_k} \frac{C_d \kappa_G^2 |\mathcal{T}(z)|^2}{j^{d-1}} \\
 & \leq C' |\mathcal{T}(z)|^2 \left(\mathbb{1}_{d=2} \log(n) + \mathbb{1}_{d>2} \frac{1}{h_k^{d-2}} \right).
 \end{aligned}$$

Since $|\mathcal{T}(z)|$ is of order h_k^{d-1} , (4.14) holds.

4.3. *Bounding outer fluctuations.* This section follows [9] closely. The features of the flashing process allow for some simplification. We keep the notation of the previous subsection. There, we proved that for some integer k^* , which depends on n ,

$$P(T^* > k^*) = 1 - \varepsilon(n) \quad \text{with} \quad \sum_{n \geq 1} \varepsilon(n) < +\infty.$$

The integer k^* is the largest such that $r_{k^*} \leq n - 2Ah \log(n)/\kappa_1$, for a large constant A and with h defined in (4.10). As a consequence, the following conditional law can be seen as a slight modification of P :

$$(4.19) \quad P^*(\cdot) = P(\cdot | T^* > k^*).$$

We begin by proving that under P^* the probability to find some k with $n \leq r_k < 2n$ and some tile \mathcal{T} in \mathcal{T}_k with $W_k(\mathcal{T})$ larger than or equal to $\xi' = 2A'h^d \log n$ for a large enough A' decreases faster than any given power of n . First, note that on $\{T^* > k^*\}$,

$$(4.20) \quad W_k(\mathcal{T}) + L_k^*(\mathcal{T}) \leq M_k(\mathcal{T}) \quad \text{with} \quad L_k^* = M(\mathbb{B}(0, r_{k^*} - h_{k^*}), r_k, \mathcal{T}).$$

Our key observation is that the pair $(W_k(\mathcal{T}), \mathbb{1}_{\{T^* > k^*\}})$ is independent of L_k^* . Thus, for any $t > 0$,

$$\begin{aligned}
 P(W_k(\mathcal{T}) \geq \xi', T^* > k^*) & \leq e^{-t\xi'} E[e^{tW_k(\mathcal{T})} \mathbb{1}_{\{T^* > k^*\}}] \\
 & = e^{-t\xi'} \frac{E[\exp(t(W_k(\mathcal{T}) + L_k^*)) \mathbb{1}_{\{T^* > k^*\}}]}{E[e^{tL_k^*}]} \\
 & \leq e^{-t\xi'} \frac{E[e^{tM_k(\mathcal{T})}]}{E[e^{tL_k^*}]} \\
 & = \exp(-t(\xi' - E[M_k(\mathcal{T}) - L_k^*])) \times \frac{E[e^{t\bar{M}_k(\mathcal{T})}]}{E[e^{t\bar{L}_k^*}]}.
 \end{aligned}$$

By Lemma 2.3, we have [for $f(t)$ and $g(t)$ quadratic for t small]

$$\begin{aligned}
 & \log P(W_k(\mathcal{T}) \geq \xi', T^* > k^*) \\
 (4.21) \quad & \leq -t(\xi' - E[M_k(\mathcal{T}) - L_k^*]) + f(t) \times E[M_k(\mathcal{T}) - L_k^*] \\
 & \quad + g(t) \times \sum_{y \in \mathbb{B}(0, r_{k^*} - h_{k^*})} \mathbb{P}_y^2(S(H(\Sigma_k)) \in \mathcal{T}).
 \end{aligned}$$

The steps are now similar to the previous proof. We first estimate $E[M_k(\mathcal{T}) - L_k^*]$. By Corollary 5.4, for some positive constant K' and for n large enough,

$$\begin{aligned}
 (4.22) \quad E[M_k(\mathcal{T}) - L_k^*(\mathcal{T})] & \leq K'(n^d - (r_{k^*})^d) \frac{h_k^{d-1}}{r_k^{d-1}} + O(h_k^{d-1}) \\
 & \leq K' d n^{d-1} (n - r_{k^*}) \frac{h_k^{d-1}}{r_k^{d-1}} + O(h_k^{d-1}).
 \end{aligned}$$

Note that since $r_k \leq 2n$, we have $r_k^{d-1} = o(n^{d-1}(n - r_{k^*}))$ so that $O(h_k^{d-1})$ is small compared to the first term in (4.22). Since $k \mapsto h_k/r_k$ is decreasing, we have for some constant K

$$E[M_k(\mathcal{T}) - L_k^*(\mathcal{T})] \leq K h^d \log n.$$

Second, we estimate the sum of $\mathbb{P}_y^2(S(H(\Sigma_k)) \in \mathcal{T})$ which appears on (4.21). We use (4.17) again to obtain as in (4.18), and for a constant C ,

$$\begin{aligned}
 & \sum_{y \in \mathbb{B}(0, r_{k^*} - h_{k^*})} \mathbb{P}_y^2(S(H(\Sigma_k)) \in \mathcal{T}) \\
 & \leq C h_k^{2(d-1)} \left(\mathbb{1}_{d=2} \log(n) + \mathbb{1}_{d>2} \frac{1}{(r_k - (r_{k^*} - h_{k^*}))^{d-2}} \right).
 \end{aligned}$$

Note that $r_k - (r_{k^*} - h_{k^*}) \geq h_k$, and since $k \mapsto h_k/r_k$ is decreasing, we have, for n large enough, $h_k \leq h_{k^*}(r_k/r_{k^*}) \leq h \times (2n)/(n/2)$. Thus, for a constant C ,

$$\sum_{y \in \mathbb{B}(0, r_{k^*} - h_{k^*})} \mathbb{P}_y^2(S(H(\Sigma_k)) \in \mathcal{T}) \leq C(\mathbb{1}_{d=2} h^2 \log(n) + \mathbb{1}_{d>2} h^d) \leq C h^d \log(n).$$

We choose $A' = K$ to obtain, for any $t > 0$

$$\log P(W_k(\mathcal{T}) \geq \xi', T^* > k^*) \leq -(Kt - Kf(t) - Cg(t))h^d \log(n).$$

Since we have $P(T^* > k^*) \geq 1/2$ for A large enough and K can be taken as large as we want, we have that $P^*(W_k(\mathcal{T}) \geq \xi')$ decreases faster than any given power of n .

Now, let F_k denote the event that no tile \mathcal{T} in Σ_k contains more than $\xi' = 2A'h^d \log n$ unsettled explorers after the k th exploration wave. We define, with the notation of Section 3, $\mathcal{G}_k = \sigma(\xi_0, \dots, \xi_k)$, and note that F_k and $\{T^* > k^*\}$ are \mathcal{G}_k -measurable.

For any tile $\mathcal{T} \in \mathcal{T}_k$, let $z_k \in \tilde{\Sigma}_k$ be such that $\mathcal{T} = \mathcal{T}(z_k)$, and denote by $\tilde{\mathcal{C}} = \tilde{\mathcal{C}}(z_k)$. We are entitled, by Proposition 3.1, to use a coupon-collector estimate on the number of settled explorers during the $k + 1$ th exploration wave. On $F_k \cap \{T^* > k^*\}$, and for some positive constant K_1 ,

$$\begin{aligned} E[|\mathcal{A}_{k+1}^* \cap \tilde{\mathcal{C}}| | \mathcal{G}_k] &\geq |\tilde{\mathcal{C}}| \left(1 - \left(1 - \frac{\alpha_1}{h_k^d} \right)^{W_k(\mathcal{T})} \right) \\ &\geq |\tilde{\mathcal{C}}| \left(1 - \exp \left\{ -\alpha_1 \frac{W_k(\mathcal{T})}{h_k^d} \right\} \right) \\ &= \frac{|\tilde{\mathcal{C}}|}{h_k^d} W_k(\mathcal{T}) \frac{h_k^d}{W_k(\mathcal{T})} \left(1 - \exp \left\{ -\alpha_1 \frac{W_k(\mathcal{T})}{h_k^d} \right\} \right) \\ &\geq K_1 W_k(\mathcal{T}) \inf_{x \leq 2A' \log n} \frac{1 - e^{-\alpha_1 x}}{x}. \end{aligned}$$

We now write for some positive constant K_2 ,

$$\begin{aligned} \inf_{x \leq 2A' \log n} \frac{1 - e^{-\alpha_1 x}}{x} &\geq \frac{1}{2A' \log n} \inf_{x \leq 2A' \log n} \frac{1 - e^{-\alpha_1 x / 2A' \log n}}{x / 2A' \log n} \\ &\geq \frac{1}{2A' \log n} \inf_{x \leq 1} \frac{1 - e^{-\alpha_1 x}}{x} \geq \frac{K_2}{\log n}. \end{aligned}$$

We conclude that on $F_k \cap \{T^* > k^*\}$,

$$(4.23) \quad E[|\mathcal{A}_{k+1}^* \cap \tilde{\mathcal{C}}| | \mathcal{G}_k] \geq K_1 K_2 \frac{W_k(\mathcal{T})}{\log n}.$$

Recall now that property (4.2) implies that $K_F |\mathcal{A}_{k+1}^* \cap \mathcal{S}_k| \geq \sum_{z_k \in \tilde{\Sigma}_k} |\mathcal{A}_{k+1}^* \cap \tilde{\mathcal{C}}(z_k)|$. Thus, summing over $z_k \in \tilde{\Sigma}_k$ with $\mathcal{C} = \mathcal{C}(z_k)$ and $\mathcal{T} = \mathcal{T}(z_k)$ in (4.23), we obtain on $F_k \cap \{T^* > k^*\}$,

$$E[|\mathcal{A}_{k+1}^* \cap \mathcal{S}_k| | \mathcal{G}_k] \geq K \frac{W_k(\mathcal{S}_k)}{\log n} \quad \text{where } K = \frac{K_1 K_2}{K_F}.$$

Also, since $W_k(\mathcal{S}_k) \leq |\mathbb{B}(0, n)|$, we have, for n large enough,

$$E[\mathbf{1}_{F_k \cap \{T^* > k^*\}} |\mathcal{A}_{k+1}^* \cap \mathcal{S}_k|] \geq K \frac{E[\mathbf{1}_{\{T^* > k^*\}} W_k(\mathcal{S}_k)]}{\log n} - n^d P(F_k^c).$$

Since $P(T^* > k^*) \geq 1/2$,

$$(4.24) \quad E^*[|\mathcal{A}_{k+1}^* \cap \mathcal{S}_k|] \geq K \frac{E^*[W_k(\mathcal{S}_k)]}{\log n} - 2n^d P(F_k^c).$$

In other words, noting that $|\mathcal{A}_{k+1}^* \cap \mathcal{S}_k| = W_k(\mathcal{S}_k) - W_{k+1}(\mathcal{S}_{k+1})$,

$$(4.25) \quad E^*[W_{k+1}(\mathcal{S}_{k+1})] \leq \left(1 - \frac{K}{\log n} \right) E^*[W_k(\mathcal{S}_k)] + 2n^d P(F_k^c).$$

By iterating (4.25), and using our previous estimate on $P^*(W_k(\mathcal{T}) \geq \xi')$, we obtain that for a large enough ε , $E^*[W_{l_n+\varepsilon \log^2 n}(\mathcal{S}_{l_n+\varepsilon \log^2 n})]$, is summable, when l_n is the lowest index for which $r_{l_n} \geq n$. Also, the probability (under $P!$) of seeing at least one explorer reaching the shell $\mathcal{S}_{l_n+\varepsilon \log^2 n}$ is summable. Using the Borel–Cantelli lemma, this yields the proof of Proposition 1.5.

4.4. Lower bound for the deviations.

4.4.1. Proof of Proposition 1.6: The outer deviation. We denote by K_n the largest index such that $\mathcal{S}_{K_n} \subset \mathbb{B}(0, n)$, and by E_n the event that all explorers stopped on Σ_{K_n} , at time K_n , settle afterward in one of the shells $\{\mathcal{S}_j : K_n \leq j < K_n + b \log(n)\}$ for some positive constant b , and note that $E_n = \{A^*(N) \subset \bigcup_{j < K_n + b \log(n)} \mathcal{S}_j\}$.

We want to find b such that $\sum_{n \geq 1} P(E_n) < \infty$. Using that the flashing times of the different explorers are independent, we have

$$\begin{aligned} P(E_n) &\leq E \left[P \left(\text{all explorers, stopped in } \Sigma_{K_n}, \text{ flash in } \bigcup_{j < K_n + b \log(n)} \mathcal{S}_j | \mathcal{G}_{K_n} \right) \right] \\ &\leq E \left[\left(\sup_{z \in \Sigma_{K_n}} P \left(\text{an explorer, started on } z, \right. \right. \right. \\ &\qquad \qquad \qquad \left. \left. \left. \text{flashes in } \bigcup_{j < K_n + b \log(n)} \mathcal{S}_j | \mathcal{G}_{K_n} \right) \right)^{W_{K_n}(\Sigma_{K_n})} \right]. \end{aligned}$$

Also, there are at least $|\mathcal{S}_{K_n}|$ explorers stopped on Σ_{K_n} , and there is a positive ε_0 such that the probability of crossing a given shell without flashing is larger than ε_0 . Thus

$$\begin{aligned} P(E_n) &\leq E \left[\left(1 - \inf_{z \in \Sigma_{K_n}} P(\text{an explorer started on } z \right. \right. \\ &\qquad \qquad \qquad \left. \left. \text{is unsettled at time } K_n + b \log(n)) \right)^{|\mathcal{S}_{K_n}|} \right] \\ &\leq (1 - \varepsilon_0^{b \log(n)})^{|\mathcal{S}_{K_n}|}. \end{aligned}$$

When choosing b small enough, we reach $\sum_{n \geq 1} P(E_n) < \infty$.

4.4.2. Proof of Proposition 1.6: The inner deviation. We recall that K_n is the largest index such that $\mathcal{S}_{K_n} \subset \mathbb{B}(0, n)$. The rough idea here is that when we stop explorers on Σ_{K_n-1} , there are necessarily tiles (of Σ_{K_n-1}) containing of the order of $h_{K_n-1}^{d-1}$ sites and which receive $h_{K_n-1}^d$ explorers. The number of explorers on these tiles is not enough to cover the associated cells with the coupon collector mechanism. We now make rigorous such an argument for shells with index of order $K_n - \log(K_n)$.

To simplify the notation, let us first define three positive constants c_1, c_2 and c_3 such that for any k with $n/2 \leq r_k \leq n$, we have

$$(4.26) \quad \begin{aligned} |\mathcal{S}_k| &\leq c_1 h_k n^{d-1}, \\ \frac{h_k^d}{\sup_{z \in \Sigma_k} |\mathbb{B}(z, 6h_k) \cap \Sigma_k|} |\Sigma_k| &\geq c_2 n^{d-1} h_k \quad \text{and} \\ \inf_{z \in \Sigma_k} |\tilde{\mathcal{C}}(z)| &\geq c_3 h_k^d. \end{aligned}$$

Using α_2 given in Proposition 3.1, we define

$$(4.27) \quad a_n = \frac{1}{8\alpha_2} \log(h_{K_n}) \quad \text{and} \quad A_n = \left\lceil \frac{c_2 c_3}{4c_1} a_n \right\rceil.$$

Now we assume h_0 large enough to have A_n a strictly positive integer.

We wish now to consider a peel of A_n shells before $\partial\mathbb{B}(0, n)$. Let I_n be the index of the inner shell in this peel, that is, $r_{I_n+A_n} \leq n < r_{I_n+A_n+1}$. Since $T^* \leq I_n + 1$ implies that $\bigcup_{j \leq I_n} \mathcal{S}_j \not\subset A^*(N)$, it is enough to show that $P(T^* > I_n + 1)$ decays faster than any polynomial in n .

Note that the monotonicity of $k \mapsto h_k/r_k$ and $r_{I_n} \geq n/2$, imply that $2h_{I_n} \geq h_k$ for $I_n \leq k \leq I_n + A_n$, and n large enough. Also, on the event $\{T^* > I_n + 1\}$, we have $\mathbb{B}(0, r_{I_n} - h_{I_n}) = A_{I_n}^*(N)$ after the I_n th wave. Thus,

$$W_{I_n}(\Sigma_{I_n}) = |\mathbb{B}(0, n)| - |\mathbb{B}(0, r_{I_n} - h_{I_n})| \leq 2c_1 A_n n^{d-1} \times h_{I_n}.$$

A key feature of the flashing process is that explorers stopped, at time I_n , outside $\mathbb{B}(z, 3h_{I_n}) \cap \Sigma_{I_n}$ cannot settle in $\tilde{\mathcal{C}}(z)$. In other words, knowing \mathcal{G}_k , the covering of a family of cells $\{\tilde{\mathcal{C}}(z_j), j = 1, \dots, \mathcal{N}\}$ are independent events if $\|z_i - z_j\| \geq 6h_{I_n}$ for $i \neq j$. Now, there is an integer \mathcal{N} and sites $\{z_j, j = 1, \dots, \mathcal{N}\}$ with

$$\forall i \neq j \quad \|z_i - z_j\| \geq 6h_{I_n} \quad \text{and} \quad \sum_{j \leq \mathcal{N}} |\mathbb{B}(z_j, 6h_{I_n}) \cap \Sigma_{I_n}| \geq |\Sigma_{I_n}|.$$

We then get using (4.26),

$$(4.28) \quad \mathcal{N} h_{I_n}^d \geq \frac{1}{2} c_2 h_{I_n} n^{d-1}.$$

Let

$$\Gamma = \{j \in [1, \mathcal{N}]: W_{I_n}(\mathbb{B}(z_j, 3h_{I_n}) \cap \Sigma_{I_n}) \leq c_3 a_n h_{I_n}^d\} \quad \text{and} \quad \Gamma^c = [1, \mathcal{N}] \setminus \Gamma.$$

On $\{T^* > I_n + 1\}$,

$$\begin{aligned} 2c_1 A_n n^{d-1} \times h_{I_n} &\geq W_{I_n}(\Sigma_{I_n}) \geq \sum_{j \in \Gamma^c} W_{I_n}(\mathbb{B}(z_j, 3h_{I_n}) \cap \Sigma_{I_n}) \\ &\geq |\Gamma^c| \times (c_3 a_n h_{I_n}^d). \end{aligned}$$

Thus, using the definition of A_n in (4.27), and bound (4.28) on \mathcal{N} , we obtain

$$|\Gamma^c| \leq \frac{2c_1 A_n h_{I_n} n^{d-1}}{c_3 a_n h_{I_n}^d} \leq \frac{c_1 c_2 c_3 a_n \mathcal{N}}{2c_3 c_1 a_n c_2} = \frac{\mathcal{N}}{2}.$$

In other words, we have that $|\Gamma| \geq \mathcal{N}/2$. Now, as already noticed, knowing \mathcal{G}_{I_n} , for any subset $I \subset [1, \mathcal{N}]$, the events $\{\tilde{\mathcal{C}}(z_j) \subset \mathcal{A}_{I_n+1}^*(N), j \in I\}$ are independent. By conditioning on \mathcal{G}_{I_n} , we obtain for h_0 large enough,

$$\begin{aligned} &P(\{T^* > I_n + 1\}) \\ &= E \left[\sum_{I \subset [1, \mathcal{N}], |I| \geq \mathcal{N}/2} \mathbb{1}_{\Gamma=I} \times P(\forall i \in I, \tilde{\mathcal{C}}(z_j) \subset \mathcal{A}_{I_n+1}^*(N) | \mathcal{G}_{I_n}) \right] \\ (4.29) \quad &= E \left[\sum_{I \subset [1, \mathcal{N}], |I| \geq \mathcal{N}/2} \mathbb{1}_{\Gamma=I} \times \prod_{i \in I} P(\tilde{\mathcal{C}}(z_j) \subset \mathcal{A}_{I_n+1}^*(N) | \mathcal{G}_{I_n}) \right] \\ &\leq \sup_{z_j \in \Sigma_{I_n}} P(\mathcal{A}_{I_n+1}^*(N) \supset \tilde{\mathcal{C}}(z_j), W_{I_n}(B(z_j, 3h_{I_n}) \cap \Sigma_{I_n}) \leq c_3 h_{I_n}^d a_n)^{\mathcal{N}/2}. \end{aligned}$$

Considering the probability appearing on the right-hand side of (4.29), we can think of a coupon-collector problem, where an album of size $|\tilde{\mathcal{C}}(z)|$ has to be filled when we collect no more than $c_3 h_{I_n}^d a_n$ coupons. Using inequality (4.31) of Lemma 4.1 below, we show that

$$P(\{T^* > I_n + 1\}) \leq \exp\left(-\frac{\alpha_1}{4} a_n^2 \frac{c_2}{2} h_{I_n}^{1-d} n^{d-1}\right).$$

This concludes the proof.

The result about filling an album, that we just mentioned, is based on the following simple coupon-collector lemma (together with Proposition 3.1), which we did not find in the vast literature on such problems.

LEMMA 4.1. *Consider an album of L items for which are bought independent random coupons, each of them covering one (or possibly none) of the possible L items. If Y_i is the item associated with the i th coupons, we assume that for positive constants α_1, α_2 , such that for any $j = 1, \dots, L$,*

$$(4.30) \quad \frac{\alpha_1}{L} \leq P(Y_i = j) \leq \frac{\alpha_2}{L}.$$

Let τ_L be the number of coupons needed to complete the album. Then, for any $0 < A < \frac{1}{4\alpha_2} \log(L)$, we have

$$(4.31) \quad P(\tau_L < AL) \leq \exp\left(-\frac{\alpha_1^2 A^2 e^{-2\alpha_2 A}}{4} \sqrt{L}\right) \leq \exp\left(-\frac{\alpha_1^2 A^2}{4}\right).$$

PROOF. We denote by σ_i the time needed to collect the i th distinct item after having collected $i - 1$ distinct items. The sequence $\{\sigma_1, \sigma_2, \dots, \sigma_L\}$ is not independent, but if $\mathcal{Y}_k = \sigma(\{Y_1, \dots, Y_k\})$, and $\tau(k) = \sigma_1 + \dots + \sigma_k$, then for $i = 1, \dots, L$,

$$(4.32) \quad \left(1 - \frac{\alpha_1(L - i + 1)}{L}\right)^k \geq P(\sigma_i > k | \mathcal{Y}_{\tau(i-1)}) \geq \left(1 - \frac{\alpha_2(L - i + 1)}{L}\right)^k.$$

Indeed, calling $\mathcal{E}(i - 1)$ the set of the first $i - 1$ collected items,

$$(4.33) \quad \begin{aligned} P(\sigma_i > k | \mathcal{Y}_{\tau(i-1)}) &= P(\{Y_{\tau(i-1)+1}, \dots, Y_{\tau(i-1)+k}\} \subset \mathcal{E}(i - 1) | \mathcal{Y}_{\tau(i-1)}) \\ &= (P(Y \in \mathcal{E}(i - 1) | \mathcal{Y}_{\tau(i-1)}))^k \\ &= (1 - P(Y \notin \mathcal{E}(i - 1) | \mathcal{Y}_{\tau(i-1)}))^k. \end{aligned}$$

Using (4.30) we deduce (4.32) from (4.33). Formula (4.32) gives that

$$\frac{L}{\alpha_1(L - i + 1)} \geq E[\sigma_i | \mathcal{Y}_{\tau(i-1)}] \geq \frac{L}{\alpha_2(L - i + 1)}$$

as well as

$$(4.34) \quad E[\sigma_i^2 | \mathcal{Y}_{\tau(i-1)}] \leq 2 \frac{L^2}{\alpha_1^2(L - i + 1)^2}.$$

Now, we look for $B \leq \sqrt{L}$ such that

$$(4.35) \quad \sum_{i=\sqrt{L}}^{B\sqrt{L}} E[\sigma_{L-i}] \geq 2AL.$$

Note that

$$\sum_{i=\sqrt{L}}^{B\sqrt{L}} E[\sigma_{L-i}] \geq \frac{L}{\alpha_2} \sum_{i=\sqrt{L}}^{B\sqrt{L}} \frac{1}{i + 1} \geq \frac{L}{\alpha_2} \log(B).$$

Thus, condition (4.35) holds for $B \geq \exp(2\alpha_2 A)$, but recall that $B \leq \sqrt{L}$ also, and this gives a bound on A . Finally, note that

$$\max\{E[\sigma_{L-i} | \mathcal{Y}_{\tau(L-i-1)}], i = \sqrt{L}, \dots, B\sqrt{L}\} \leq \frac{\sqrt{L}}{\alpha_1}$$

and set

$$X_i = \frac{E[\sigma_{L-i} | \mathcal{Y}_{\tau(L-i-1)}] - \sigma_{L-i}}{(\sqrt{L}/\alpha_1)} \leq 1.$$

For $x \leq 1$, note that $e^x \leq 1 + x + x^2$ to obtain for $0 \leq \lambda \leq 1$, by successive conditioning,

$$\begin{aligned}
 P\left(\sum_{i=\sqrt{L}}^{B\sqrt{L}} \sigma_{L-i} \leq AL\right) &\leq P\left(\sum_{i=\sqrt{L}}^{B\sqrt{L}} X_i \geq \alpha_1 A\sqrt{L}\right) \\
 (4.36) \qquad &\leq e^{-\lambda\alpha_1 A\sqrt{L}} \prod_{i=\sqrt{L}}^{B\sqrt{L}} (1 + \lambda^2 \sup E[X_i^2 | \mathcal{Y}_{\tau(L-i-1)}]) \\
 &\leq \exp\left(-\lambda\alpha_1 A\sqrt{L} + \lambda^2 \sum_i \sup E[X_i^2 | \mathcal{Y}_{\tau(L-i-1)}]\right).
 \end{aligned}$$

Finally, we have, using (4.34),

$$\sum_{i=\sqrt{L}}^{B\sqrt{L}} \sup E[X_i^2 | \mathcal{Y}_{\tau(L-i-1)}] \leq \sum_{i=\sqrt{L}}^{B\sqrt{L}} \alpha_1^2 \sup \frac{E[\sigma_{L-i}^2 | \mathcal{Y}_{\tau(L-i-1)}]}{L} \leq 2B\sqrt{L}.$$

The results follows as we optimize on $\lambda \leq 1$ in the upper bound in (4.36). \square

5. Potential theory estimates. We collect in this section three technical results. In Corollary 5.4, we estimate the difference between the expected number of independent random walks exiting a ball $\mathbb{B}(0, n)$ at a distinguished site, whether the random walks are initially on the origin or are spread over a sphere $\mathbb{B}(0, r_n)$ with $r_n < n$. Corollary 5.4 is used to bound the mean number of explorers exiting some large ball from a given site, and its proof relies on a discrete mean value property Theorem 5.2, which in turns relies on Blachère’s Proposition B.1 written in the Appendix. Then, Lemma 5.1 improves an estimate of Lawler, Bramson and Griffeath in [10], dealing with the exit site distribution from a sphere when the initial position is not the origin. Indeed, Lemma 5(b) of [10], states that when $d \geq 2$, there is a positive constant J_d such that for any $r > 0$, $z \in \mathbb{B}(0, r)$ and $z^* \in \partial\mathbb{B}(0, r)$, we have

$$(5.1) \qquad \mathbb{P}_z(S(H_r) = z^*) \leq \frac{J_d}{(\|z^*\| - \|z\|)^{d-1}}.$$

Thus, when $\|z^*\| - \|z\|$ is small, (5.1) is useless. Since we need bounds on the sum of squares of $\mathbb{P}_z(S(H_r) = z^*)$ over $z \in \mathbb{B}(0, r - h)$ of order $\log(r)$ in $d = 2$, and of order $1/h^{d-2}$ when $d > 2$, we establish the following.

LEMMA 5.1. *There is a positive constant κ_G such that, for all $r > 0$, if $z \in \mathbb{B}_r$, $z^* \in \partial\mathbb{B}_r$, then*

$$(5.2) \qquad \mathbb{P}_z(S(H_r) = z^*) \leq \frac{\kappa_G}{\|z - z^*\|^{d-1}}.$$

Finally, we prove the *uniform hitting property*, Proposition 3.1, for the boundary of a *cell*. Though the property is natural, the nonspherical nature of a cell, makes its proof tedious.

5.1. *A discrete mean value theorem.* The following result has interest on its own.

THEOREM 5.2. *There are positive constants K_0 and K_a such that for any sequence $\{\Delta_n, n \in \mathbb{N}\}$ with $K_0 \leq \Delta_n \leq n^{1/3}$, for any $z \in \mathbb{B}_n$ with $n - \|z\| \leq 1$, we have, setting $r_n = n - \Delta_n$,*

$$(5.3) \quad \left| |\mathbb{B}_{r_n}| \times G_n(0, z) - \sum_{y \in \mathbb{B}_{r_n}} G_n(y, z) \right| \leq K_a.$$

REMARK 5.3. Note that a related (but distinct) property was also at the heart of [10]. Namely, for $\varepsilon > 0$, and n large enough, if $z \in \mathbb{B}_n$, and $n - \|z\| \geq \varepsilon n$,

$$(5.4) \quad |\mathbb{B}_n| \times G_n(0, z) \geq \sum_{y \in \mathbb{B}_n} G_n(y, z).$$

We start with proving the following useful corollary of Theorem 5.2.

COROLLARY 5.4. *In the setting of Theorem 5.2, and for any $\Lambda \subset \partial \mathbb{B}_n$,*

$$(5.5) \quad |E[M(\mathbb{B}_{r_n} | \mathbf{1}_0, n, \Lambda)] - E[M(\mathbb{B}_{r_n}, n, \Lambda)]| \leq K_a |\Lambda|.$$

PROOF. Note that (5.5) holds if for any $z^* \in \partial \mathbb{B}_n$,

$$(5.6) \quad \left| |\mathbb{B}_{r_n}| \times \mathbb{P}_0(S(H_n) = z^*) - \sum_{y \in \mathbb{B}_{r_n}} \mathbb{P}_y(S(H_n) = z^*) \right| \leq K_a.$$

By a classical decomposition (Lemma 6.3.6 of [11]), we have for a finite subset $B \subset \mathbb{Z}^d$, $y \in B$, and $z^* \in \partial B$

$$(5.7) \quad \mathbb{P}_y(S(H(\partial B)) = z^*) = \frac{1}{2d} \sum_{z \in B, z \sim z^*} G_B(y, z).$$

For $B = \mathbb{B}(0, n)$, we replace in (5.6) the value of $\mathbb{P}_y(S(H(\partial B)) = z^*)$ by the right-hand side in (5.7), and are left with proving that for any $z \in \mathbb{B}_n$ with $n - \|z\| \leq 1$, we have (5.3). \square

PROOF OF THEOREM 5.2. When $d \geq 3$, we express $G_n(0, z)$ in term of Green’s function (Proposition 4.6.2(a) of [11]),

$$G_n(0, z) = G(0, z) - \mathbb{E}_z[G(0, S(H_n))].$$

Now, using Green’s function asymptotics (2.2), there is a constant K_1 (independent on n) such that

$$(5.8) \quad \left| \nu_d G_n(0, z) - 2 \frac{\alpha(z)}{n^{d-1}} \right| \leq \frac{K_1}{n^d} \quad \text{where } \alpha(z) = \mathbb{E}_z[\|S(H_n)\| - \|z\|].$$

In $d = 2$, G_n is expressed in terms of the potential kernel (Proposition 4.6.2(b) of [11])

$$G_n(0, z) = -a(0, z) + \mathbb{E}_z[a(0, S(H_n))].$$

Using (2.3), we have

$$\pi G_n(0, z) = 2\alpha(z)/n + O(1/n^2).$$

Now, $r_n^d = n^d - d\Delta_n n^{d-1} + O(\Delta_n^2 n^{d-2})$, so that using (5.8), and the hypothesis $\Delta_n = O(n^{1/3})$, and $0 \leq n - \|z\| \leq 1$

$$(5.9) \quad \begin{aligned} |\mathbb{B}_{r_n}| G_n(0, z) &= (r_n^d + O(r_n^{d-1})) \left(2 \frac{\alpha(z)}{n^{d-1}} + O\left(\frac{1}{n^d}\right) \right) \\ &= (n^d - d\Delta_n n^{d-1} + O(\Delta_n^2 n^{d-2}) + O(n^{d-1})) \\ &\quad \times \left(2 \frac{\alpha(z)}{n^{d-1}} + O\left(\frac{1}{n^d}\right) \right) \\ &= 2\alpha(z)(n - d\Delta_n) + O(1). \end{aligned}$$

Since $\{\|S_n\|^2 - n, n \in \mathbb{N}\}$ is a martingale (with the natural filtration),

$$\mathbb{E}_z[\|S(H_n)\|^2] - \|z\|^2 = \mathbb{E}_z[H_n] = \sum_{y \in \mathbb{B}_n} G_n(y, z).$$

Using $n - \|z\| \leq 1$, this yields for a constant K_l ,

$$(5.10) \quad \left| \sum_{y \in \mathbb{B}_n} G_n(y, z) - 2\alpha(z)n \right| \leq K_l.$$

We now invoke Proposition B.1 of the Appendix. There is K_b such that for $z \in \mathbb{B}_n$ with $n - \|z\| \leq 1$,

$$(5.11) \quad \left| \sum_{y \in \mathcal{A}(r_n, n)} G_n(y, z) - 2\alpha_0(z)d\Delta_n \right| \leq K_b$$

where $\alpha_0(z) = \mathbb{E}_z[\|S(H_n)\| - \|z\| | H_n < H(B_{r_n})]$.

From (5.10) and (5.11), we obtain

$$(5.12) \quad \left| \sum_{y \in \mathbb{B}_{r_n}} G_n(y, z) - 2n\alpha(z) + 2\alpha_0(z)d\Delta_n \right| \leq K_l + K_b.$$

Now, from (5.9) and (5.12) we obtain for a constant K_2 ,

$$\left| \left(|\mathbb{B}_{r_n}| G_n(0, z) - \sum_{y \in \mathbb{B}_{r_n}} G_n(y, z) \right) - 2(\alpha_0(z) - \alpha(z)) d \Delta_n \right| \leq K_2.$$

Now, from

$$|\alpha_0(z) - \alpha(z)| \leq \mathbb{P}_z(H(B_{r_n}) < H_n) \times (\alpha_0(z) + \mathbb{E}_z[\|S(H_n)\| - \|z\| | H_n > H(B_{r_n})]),$$

and the Gambler’s ruin estimate, for $K_0 > 0$ and $z \in \mathbb{A}(n - 1, n)$,

$$\mathbb{P}_z(H(B_{r_n}) < H_n) \leq \frac{K_0}{\Delta_n},$$

we deduce that

$$\Delta_n |\alpha_0(z) - \alpha(z)| \leq 2 \Delta_n \mathbb{P}_z(H(B_{r_n}) < H_n) \leq 2K_0.$$

The desired result follows. \square

5.2. *Proof of Lemma 5.1.* We follow the proof of Lemma 5(b) of [10]. Set $D := \|z - z^*\|$. Let O' be a closest point to $(1 + \frac{D}{4r})z^*$ in $\mathbb{B}(z^*, \frac{D}{4})$. We define $B'_1 := \mathbb{B}(O', \frac{D}{4})$, $B'_2 := \mathbb{B}(O', \frac{D}{2})$, and we note that

$$\|z - z^*\| \leq \|z - O'\|$$

and, for all x in $\partial B'_2$, the triangle inequality $\|z - z^*\| \leq \|z - x\| + \|x - z^*\|$ implies that

$$\min_{x \in \partial B'_2} \|z - x\| \geq \frac{D}{3}.$$

Now, define

$$\tau := \inf\{t > 0 : S(t) \in \{z\} \cup B_r^c\} \quad \text{and} \quad \tau' := \inf\{t > 0 : S(t) \in B'_1 \cup \partial B'_2\}.$$

By a last exit decomposition, together with the strong Markov property,

$$\begin{aligned} \mathbb{P}_z(S(H_r) = z^*) &= G_r(z, z) \mathbb{P}_{z^*}(S(\tau) = z) \\ &\leq G_r(z, z) \mathbb{P}_{z^*}(S(\tau') \in B'_2) \max_{x \in \partial B'_2} \mathbb{P}_x(S(\tau) = z) \\ (5.13) \quad &= \mathbb{P}_{z^*}(S(\tau') \in B'_2) \max_{x \in \partial B'_2} G_r(x, z) \\ &\leq \mathbb{P}_{z^*}(S(\tau') \in B'_2) \max_{x \in \partial B'_2} G_{r+D}(x, z). \end{aligned}$$

A Gambler’s ruin estimate yields, for some positive constant c ,

$$\mathbb{P}_{z^*}(S(\tau') \in B'_2) \leq \frac{c}{D}.$$

The desired result follows from (5.13) and the previous bound, after we show that for a constant c , such that for all x satisfying $\|x - z\| \geq \frac{D}{3}$,

$$(5.14) \quad G_{r+D}(x, z) \leq \frac{c}{D^{d-2}}.$$

On the set $V := \mathbb{B}(z, \frac{D}{4})$, the map $y \mapsto G(x, y)$ is harmonic. By Harnack's inequality, we have

$$(5.15) \quad G_{r+D}(x, z) \leq \frac{c}{D^d} \sum_{y \in V} G_{r+D}(x, y) = \frac{c}{D^d} \mathbb{E}_x[Y],$$

where c is a positive constant, and Y is the number of visits of V before time H_{r+D} . By taking the supremum over the entering site of V in (5.15),

$$G_{r+D}(x, z) \leq \frac{c}{D^d} \sup_{y \in V} \mathbb{E}_y[Y].$$

It remains to show that $\sup_{y \in V} \mathbb{E}_y[Y] \leq JD^2$, for some positive constant J . This is identical to (2.10) of [10], and we omit this last step.

5.3. Proof of Proposition 3.1. For $j \geq 0$, consider z_j in Σ_j . We show that for positive constants α_1, α_2 , and for all z^* in $\mathcal{C}(z_j)$, we have (3.3). The random walk has initial condition $S(0) = z_j$.

First, when $z^* = z_j$, $S(\sigma_j) = z_j$ if and only if $X_j = 1$. This happens with probability $1/h_j^d$, and gives the result in this case.

Assume $z^* \in \mathcal{C}(z_j) \setminus \{z_j\}$. We recall that the unbiased Bernoulli variable Y_j decides whether the explorer can flash upon exiting either a sphere or an annulus. More precisely, we draw R_j with density g_j given in (3.1), and if $Y_j = 1$ (resp., $Y_j = 0$) the walk flashes upon exiting the ball of center z_j and radius $R_j \wedge (r_j + h_j - \|z_j\|)$ [resp., $\mathbb{A}(r_j - R_j, r_j + R_j)$] provided $S(\sigma_j) \in \mathcal{C}(z_j)$.

Step 1: Flashing when exiting a sphere ($Y_j = 1$). We first prove the upper bound when $Y_j = 1$ and $X_j = 0$. It is obvious that

$$z^* \in \partial\mathbb{B}(z_j, \|z^* - z_j\|) \quad \text{but } z^* \notin \partial\mathbb{B}(z_j, \|z^* - z_j\| - 1).$$

Thus, $R_j \in]\|z^* - z_j\| - 1, \|z^* - z_j\|]$, and there is a constant C such that

$$(5.16) \quad \begin{aligned} P(X_j = 0, Y_j = 1, R_j \in]\|z^* - z_j\| - 1, \|z^* - z_j\|]) \\ \leq C \frac{\|z^* - z_j\|^{d-1}}{h_j^d}. \end{aligned}$$

On the other hand, by (2.4) of Section 2.2,

$$(5.17) \quad \begin{aligned} P(S(\sigma_j) = z^* | X_j = 0, Y_j = 1, R_j \in]\|z^* - z_j\| - 1, \|z^* - z_j\|]) \\ \leq \frac{c_2}{\|z^* - z_j\|^{d-1}}. \end{aligned}$$

The upper bound in the case $\{X_j = 0, Y_j = 1\}$ follows from (5.16) and (5.17).

We now turn to the lower bound when $Y_j = 1$ and $X_j = 0$. Since we want a lower bound, we consider the event that the walk flashes on z^* when exiting a sphere only in the case where $|\|z^*\| - r_j| < h_j/2$. Note that by Lemma 2.2, z^* has a nearest neighbor, say z , which satisfies

$$\|z - z_j\| \leq \|z^* - z_j\| - \frac{1}{4\sqrt{d}}.$$

This means that if $h \in V := [\|z^* - z_j\| - 1/(4\sqrt{d}), \|z^* - z_j\|]$, then $z^* \in \partial\mathbb{B}(z_j, h)$. Thus

$$\begin{aligned} P_{z_j}(S(\sigma_j) = z^*) &\geq P(X_j = 0, Y_j = 1, R_j \in V) \times \inf_{h \in V} \mathbb{P}_{z_j}(S(H(\partial\mathbb{B}(z_j, h))) = z^*) \\ &\geq \frac{ch^{d-1}}{h_j^d} \inf_{h \in V} \mathbb{P}_{z_j}(S(H(\partial\mathbb{B}(z_j, h))) = z^*) \\ &\geq \frac{ch^{d-1}}{h_j^d} \times \frac{c_1}{h^{d-1}} \quad [\text{using (2.4)}]. \end{aligned}$$

The lower bound in the case $Y_j = 1, X_j = 0$, and $|\|z^*\| - r_j| < h_j/2$ is obtained.

Step 2: Flashing when exiting an annulus ($Y_j = 0$). The upper bound for this case is close to the case $Y_j = 1$. It is obvious that

$$\begin{aligned} z^* &\in \partial\mathbb{A}(r_j - \|z^*\| - r_j, r_j + \|z^*\| - r_j) \\ &\text{but } z^* \notin \partial\mathbb{A}(r_j - \|z^*\| - r_j + 1, r_j + \|z^*\| - r_j - 1). \end{aligned}$$

Thus necessarily, $R_j \in]\|z^*\| - r_j| - 1, \|z^*\| - r_j|]$, and

$$P(Y_j = 0, R_j \in]\|z^*\| - r_j| - 1, \|z^*\| - r_j|]) \leq C \frac{\|z^*\| - r_j|^{d-1}}{h_j^d}.$$

For $h > 0$, define $\mathcal{D}_h = \mathbb{A}(r_j - h, r_j + h)$. It is enough to prove that for some constant c , and for any h such that $z^* \in \partial\mathcal{D}_h$ (and $h \in]\|z^*\| - r_j| - 1, \|z^*\| - r_j|]$),

$$(5.18) \quad \mathbb{P}_{z_j}(S(H(\mathcal{D}_h^c)) = z^*) \leq \frac{c}{h^{d-1}}.$$

Note the following fact. If $\|z^*\| > \|z_j\|$, and the walk exits \mathcal{D}_h at z^* , then the walk exits $\mathbb{B}(0, r_j + h)$ at z^* , whereas if $\|z^*\| < \|z_j\|$, and the walk exits \mathcal{D}_h at z^* , then the walk enters $\mathbb{B}(0, r_j - h)$ at z^* . In both cases, Lemma 5(b) of [10] yields (5.18). (Actually, Lemma 5(b) of [10] is formulated to cover only the case $\|z^*\| > \|z_j\|$, but its proof covers both cases.)

We turn now to the lower bound. By Lemma 2.2, z^* has a nearest neighbor, say z ,

$$(5.19) \quad |\|z\| - r_j| \leq \|z^*\| - r_j - \frac{1}{4\sqrt{d}}.$$

This means that if $h \in V := [|\|z^*\| - r_j| - 1/(4\sqrt{d}), |\|z^*\| - r_j|]$, then $z^* \in \partial\mathcal{D}_h$.

We only need to consider the case $|\|z^*\| - r_j| \geq h_j/2$. It is enough to prove, for $h \in V$, $z^* \in \mathcal{C}(z_j) \cap \partial\mathcal{D}_h$, and for some constant c (that depends on d), that

$$(5.20) \quad \mathbb{P}_{z_j}(S(H(\mathcal{D}_h^c)) = z^*) \geq \frac{c}{h^{d-1}}.$$

Let y^* be the closest site of $\partial\mathbb{B}(0, r_j)$ to the segment $[0, z^*]$, and let x^* be in \mathbb{R}^d given by

$$x^* = \left(r_j + \frac{h}{2}\right) \frac{z^*}{\|z^*\|}.$$

Note that if $\tilde{z} \sim z$ and \tilde{z} satisfies (5.19), then $\|x^* - \tilde{z}\| < \inf_{z \in \partial\mathcal{D}_h} \|x^* - z\|$, and we define

$$R^* = \frac{1}{2} \left(\|x^* - \tilde{z}\| + \inf_{z \in \partial\mathcal{D}_h} \|x^* - z\| \right).$$

Define

$$\tilde{\mathcal{D}}_h = \mathbb{A} \left(r_j - \frac{h}{2}, r_j + \frac{h}{2} \right) \quad \text{and set} \quad \Gamma = \mathbb{B}(x^*, R^*) \cap \partial(\tilde{\mathcal{D}}_h^c).$$

Thus, if $\|z^*\| > \|z_j\|$, then Γ is the boundary of the lower hemisphere of the ball $\mathbb{B}(x^*, R^*)$. We need also the time $\tau^+ = \inf\{n \geq 1 : S(n) \in \mathcal{D}_h^c \cup \{z_j\}\}$. By a last exit decomposition, and the strong Markov property, we have

$$(5.21) \quad \begin{aligned} \mathbb{P}_{z_j}(S(H(\partial\mathcal{D}_h)) = z^*) &= G_{\mathcal{D}_h}(z_j, z_j) \mathbb{P}_{z^*}(S(\tau^+) = z_j) \\ &\geq G_{\mathcal{D}_h}(z_j, z_j) \mathbb{P}_{z^*}(H(\Gamma) < \tau^+) \min_{x \in \Gamma} \mathbb{P}_x(S(\tau) = z_j) \\ &\geq \mathbb{P}_{z^*}(H(\Gamma) < \tau^+) \min_{x \in \Gamma} G_{\mathcal{D}_h}(x, z_j). \end{aligned}$$

Since $z^* \in \mathcal{C}(z_j)$, we have $\|y^* - z_j\| \leq h_j/2$, so that y^* and z_j can be connected by 10 overlapping balls of radius $h_j/10$ in such a way that, applying Harnack's inequality 10 times (see Theorem 6.3.9 in [11]) to the harmonic map $y \mapsto G_{\mathcal{D}_h}(x, y)$, we can estimate from below the last factor in (5.21). For any $x \in \Gamma$,

$$G_{\mathcal{D}_h}(x, z_j) \geq c_H^{10} G_{\mathcal{D}_h}(x, y^*).$$

We use again Harnack's inequality on the harmonic functions $x \mapsto G_{\mathcal{D}_h}(x, y^*)$, to obtain

$$\min_{x \in \Gamma} G_{\mathcal{D}_h}(x, y^*) \geq c_H G_{\mathcal{D}_h}(x', y^*),$$

where $x' \in \mathbb{B}(x^*, R^*/2)$ and $\|x' - y^*\| \in [\frac{h}{4} - 1, \frac{h}{4}]$. The purpose of choosing x' is to have $y^* \in \mathbb{B}(x', h/4)$, and $\mathbb{B}(x', h/2) \subset \mathcal{D}_h$ so that $G_{\mathcal{D}_h}(x', y^*) \geq G_{\mathbb{B}(x', h/2)}(x', y^*)$.

When dimension is 2, the classical expansion of $G_{\mathbb{B}(x',h/2)}(x', \cdot)$ (see Proposition 6.3.5 of [11]) gives with a constant K_2 ,

$$(5.22) \quad G_{\mathbb{B}(x',h/2)}(x', y^*) \geq \frac{2}{\pi} \log\left(\frac{h/2}{\|x' - y^*\|}\right) - \frac{K_2}{\|x' - y^*\|} \geq \frac{2}{\pi} \log 2 - \frac{4K_2}{h}.$$

When h is large enough, $G_{\mathbb{B}(x',h/2)}(x', y^*) \geq \log(2)/\pi$.

When dimension is larger than 2, by using (2.2), there is a constant K_d such that, when h is large enough,

$$(5.23) \quad \begin{aligned} G_{\mathcal{D}_h}(x', y^*) &= G(x', y^*) - \mathbb{E}_{x'}[G(S(H(\mathcal{D}_h^c)), y^*)] \\ &\geq \frac{C_d}{h^{d-2}}(4^{d-2} - 1) - \frac{K_g}{h_j^d} \geq \frac{K_d}{h^{d-2}}. \end{aligned}$$

As a consequence of (5.23), we just need to prove that the first factor in (5.21) is of order $1/h$, at least. We realize the event $\{H(\Gamma) < \tau^+\}$ in two moves: the walk first hits the sphere $\mathbb{B}(x^*, R^*/2)$, and then exits from the cap $\partial\mathbb{B}(x^*, R^*) \cap \tilde{\mathcal{D}}_h$,

$$(5.24) \quad \begin{aligned} \mathbb{P}_{z^*}(H(\Gamma) < \tau^+) &\geq \frac{1}{2d} \mathbb{P}_{z^*}(H(\mathbb{B}(x^*, R^*/2)) < H(\mathbb{B}^c(x^*, R^*))) \\ &\quad \times \inf_{y \in \partial\mathbb{B}(x^*, R^*/2)} \mathbb{P}_y(S(H(\mathbb{B}^c(x^*, R^*))) \in \tilde{\mathcal{D}}_h). \end{aligned}$$

The first factor in the right-hand side of (5.24) is of order $1/R^*$, that is, of order $1/h$. To deal with the second factor, we invoke Harnack’s inequality to have for $y \in \partial\mathbb{B}(x^*, R^*/2)$, and for x'' the closest point of \mathbb{Z}^d to x^* ,

$$(5.25) \quad \mathbb{P}_y(S(H(\mathbb{B}^c(x^*, R^*))) \in \tilde{\mathcal{D}}_h) \geq c_H \mathbb{P}_{x''}(S(H(\mathbb{B}^c(x^*, R^*))) \in \tilde{\mathcal{D}}_h).$$

We invoke now (2.4) to obtain for some constant K_3 ,

$$(5.26) \quad \mathbb{P}_{x''}(S(H(\mathbb{B}^c(x^*, R^*))) \in \tilde{\mathcal{D}}_h) \geq c_1 \frac{|\partial\mathbb{B}(x^*, R^*) \cap \tilde{\mathcal{D}}_h|}{|\partial\mathbb{B}(x^*, R^*)|} \geq K_3.$$

We gather (5.24), (5.25) and (5.26) to obtain the desired lower bound.

APPENDIX A: COUPLING AND PROOF OF LEMMA 1.3

We give a precise definition of our coupling. To avoid heavy notation, we write the coupling algorithm as a pseudo-code.

First of all, we draw N independent sequences of independent Bernoulli and continuous random variables $((X_{k,l}, Y_{k,l}, R_{k,l} : l \geq 0) : 1 \leq k \leq N)$ as in Section 3. In addition, we call $(U_k : k \geq 1)$ the sequence of the increments of a generic independent simple random walk on \mathbb{Z}^d . From these two sources of randomness we extract our explorer and flashing explorer trajectories with their associated clusters. The flashing times will be adapted to the flashing explorer trajectories as in Section 3.

Our pseudo-code is made of two loops of size N that make precise the previous description. With the first loop we build our N random walk trajectories $((S_i(t) : 0 \leq t \leq \tau_i) : 1 \leq i \leq N)$ with their associated clusters $A(1), \dots, A(N)$. Step by step, within this first loop, we also define pieces of the flashing explorers trajectories S_1^*, \dots, S_N^* . With the second loop we complete the trajectories of the flashing explorers to build the associated cluster $A^*(N)$. During the algorithm, $t_k \in \mathbb{N}$ stands for the time up to which the trajectory of flashing explorer k has been defined ($k \in \{1; \dots; N\}$). We use the same t for the time governing the evolution of each simple random walk S_i . The index j is updated before adding each random walk increment to the partial sum of S_i and S_j^* . The updating procedure uses the index j' described in Section 3.3, and we denote by $\Delta = U$ the increment. Each encountered U stands for the first unused random variable in the sequence $(U_k : k \geq 1)$.

The main advantage of the pseudo-code formalism is that it allows, through the assignment operator “ \leftarrow ,” expressions of the kind $j \leftarrow \max(j, j')$ or $t_j \leftarrow t_j + 1$ rather than $j(\theta + 1) = \max(j(\theta), j'(\theta))$ and $t_{j(\theta+1)}(\theta + 1) = t_{j(\theta)}(\theta) + 1$ with θ a discrete parameter ordering the sequence of our elementary moves. It makes also implicit identities like $t_k(\theta + 1) = t_k(\theta)$ for any quantity t_k that does not need to be updated. Our following pseudo-code can be re-written in a classical inductive way with θ running through $\{(i, t) \in \{1; \dots; N\} \times \mathbb{N} : t \leq \tau_i\}$ according to lexicographic order. Marks ^(a) and ^(b) refer to remarks (a) and (b) below:

```

A(0) ← ∅;
For i = 1 to N
  {
    j ← i;
    t ← 0;    t_j ← 0;
    S_i(t) ← 0;    S_j^*(t_j) ← 0    [Note that S_i(t) = S_j^*(t_j)]
    While S_i(t) ∈ A(i - 1)
      {
        If t_j is a flashing time for explorer j(a), then
          {
            j' ← unique index(b) k ∈ {1; ...; i} \ {j}
                  such that S_k^*(t_k) = S_j^*(t_j) = S_i(t);
          }
        If t_j is not a flashing time for explorer j(a), then j ← j';
        Otherwise j ← max(j, j');
        Δ ← U; S_i(t + 1) ← S_i(t) + Δ; S_j^*(t_j + 1) ← S_j^*(t_j) + Δ
          [so that S_i(t + 1) = S_j^*(t_j + 1)]
        t ← t + 1; t_j ← t_j + 1;
        A(i) ← A(i - 1) ∪ {S_i(t)};
      }
    i ← i + 1;
  }
A^*(0) ← ∅;
For k = 1 to N
  
```

$$\left\{ \begin{array}{l} \text{While } t_k \text{ is not a flashing time for explorer } k^{(a)} \text{ or } S_k^*(t_k) \in A_{k-1}^* \\ \left\{ \begin{array}{l} S_k^*(t_k + 1) \leftarrow S_k^*(t_k) + U; \\ t_k \leftarrow t_k + 1; \\ A^*(k) \leftarrow A^*(k - 1) \cup \{S_k^*(t_k)\}; \\ k \leftarrow k + 1; \end{array} \right. \end{array} \right.$$

Remarks:

(a) Recall that for $l = j, j'$ or k , S_l^* is defined up to time t_l as well as its associated flashing times.

(b) One checks by induction on i that just after the instruction “ $A(i) \leftarrow A(i - 1) \cup \{S_i(t)\}$,” we have

$$(A.1) \quad A(i) = \{S_1^*(t_1); \dots; S_i^*(t_i)\} \quad \text{and} \quad |A(i)| = i.$$

To do so, one checks by induction on $t < \tau_i$, that

$$(A.2) \quad \begin{aligned} A(i - 1) &= \{S_1^*(t_1); \dots; S_{j-1}^*(t_{j-1}); S_{j+1}^*(t_{j+1}); \dots; S_i^*(t_i)\} \quad \text{and} \\ |A(i - 1)| &= i - 1. \end{aligned}$$

Since we always have $S_j^*(t_j) = S_i(t)$, this proves by induction that j' is well defined.

The key observation is that for each increment U , the index of the explorer that follows this increment depends on the whole previous construction, but the value of U does not depend on it. As a consequence, we build independent random walks S_1, \dots, S_N coupled with independent flashing random walks S_1^*, \dots, S_N^* . Then, one simply checks by induction on i and k that

$$(A.3) \quad A(i) = \{S_1(\tau_1); \dots; S_i(\tau_i)\} \quad \text{and} \quad A^*(k) = \{S_1^*(\tau_1^*); \dots; S_k^*(\tau_k^*)\}$$

for all $1 \leq i, k \leq N$.

Finally, define $(\bar{t}_1, \dots, \bar{t}_N)$ and $(\tau_1^*, \dots, \tau_N^*)$ the values of (t_1, \dots, t_N) at the end of the first and last cycle, respectively. Since t_1, \dots, t_N can only increase during our loops, we have $\tau_k^* \geq \bar{t}_k$ for all k . Then (3.4) and (3.5) follow from (A.1) and (A.3).

APPENDIX B: TIME SPENT IN AN ANNULUS (BY BLACHÈRE)

This section is devoted to an asymptotic expansion of the expected time spent in an annulus $\mathcal{A}(r_n, n)$ for $r_n < n$, when the random walk is started at some point z within the annulus, and before it exits the outer shell.

PROPOSITION B.1. *There are positive constants K_0, K_b , such that for any sequence $\{\Delta_n, n \in \mathbb{N}\}$ with $K_0 \leq \Delta_n \leq n^{1/3}$, for any $z \in \mathcal{A}(r_n, n)$, we have setting $r_n = n - \Delta_n$,*

$$(B.1) \quad \left| \sum_{y \in \mathcal{A}(r_n, n)} G_n(z, y) - (2d\Delta_n\alpha_0(z) - d(n - \|z\|)^2) \right| \leq K_b((n - \|z\|) \vee 1)$$

with

$$\alpha_0(z) = \mathbb{E}_z[\|S(H_n)\| - \|z\| \mathbf{1}_{H(B^c(0, n)) < H(B(0, r_n))}].$$

PROOF. Our strategy is to decompose a path into successive strands lying entirely in the annulus. The first strand is special since the starting point is any $z \in \mathcal{A}(r_n, n)$. The other strands, if any, start all on $\partial\mathbb{B}(0, r_n)$. We estimate the time spent inside the annulus for each strand. Let us remark that we make use of three facts: (i) precise asymptotics for Green’s function, (ii) $(G(0, S(n)), n \in \mathbb{N})$ is a martingale and (iii) $(\|S(n)\|^2 - n, n \in \mathbb{N})$ is a martingale.

Choose $z \in \mathcal{A}(r_n, n)$. We define the following stopping times $(D_i, U_i, i \geq 0)$, corresponding to the i th downward and upward crossings of the sphere of radius r_n . Let $\theta(n)$ act on trajectories by time-translation of n -units. Let $\tau = H(B_{r_n}) \wedge H_n, D_0 = U_0 = 0$ and

$$D_1 = \tau \mathbf{1}_{H(B_{r_n}) < H_n} + \infty \mathbf{1}_{H_n < H(B_{r_n})}.$$

If $D_1 < \infty$, then $U_1 = H_{r_n} \circ \theta(D_1) + D_1$, whereas if $D_1 = \infty$, then we set $U_1 = \infty$. We now proceed by induction, and assume D_i, U_i are defined. If $D_i = \infty$, then $D_{i+1} = \infty$, whereas if $D_i < \infty$ (and necessarily $U_i < \infty$), then

$$D_{i+1} = U_i + (\tau \mathbf{1}_{\tau = H(B_{r_n})} + \infty \mathbf{1}_{\tau = H_n}) \circ \theta(U_i)$$

and

$$U_{i+1} = D_{i+1} + H_{r_n} \circ \theta(D_{i+1}).$$

With this notation, we can write

$$\begin{aligned} \sum_{y \in \mathcal{A}(r_n, n)} G_n(z, y) &= \mathbb{E}_z[\tau] + \sum_{i=1}^{\infty} \mathbb{E}_z[\tau \circ \theta(U_i) \mathbf{1}_{D_i < \infty}] \\ \text{(B.2)} \qquad \qquad \qquad &= \mathbb{E}_z[\tau] + \mathbb{P}_z(D_1 < \infty) \times l(z), \end{aligned}$$

where

$$\text{(B.3)} \quad l(z) = \sum_{i=1}^{\infty} \mathbb{E}_z[\tau \circ \theta(U_i) | D_i < \infty] \prod_{j=1}^{i-1} (1 - \mathbb{P}_z(D_{j+1} = \infty | D_j < \infty)).$$

Now, we compute each term of the right-hand side of (B.2).

We have divided the proof into three steps.

Step 1: First, we show that there is a positive constant K (independent of z and n) such that when $z \in \mathcal{A}(r_n, n)$, then

$$\text{(B.4)} \quad \left| \mathbb{P}_z(D_1 < \infty) - \frac{\alpha_0(z)}{\Delta_n} \right| \leq \frac{K}{\Delta_n^2} ((n - \|z\|) \vee 1).$$

Note that when $z \in \mathbb{B}(0, n)$, and $n - \|z\| \leq 1$, (B.4) yields

$$(B.5) \quad \left| \mathbb{P}_z(D_1 < \infty) - \frac{\mathbb{E}_z[\|S(\tau)\| - \|z\| | D_1 = \infty]}{\Delta_n} \right| \leq \frac{K}{\Delta_n^2}.$$

Second, we show that for $z \in \mathcal{A}(r_n, n)$, and $i \geq 1$,

$$(B.6) \quad \left| \mathbb{P}_z(D_{i+1} = \infty | D_i < \infty) - \frac{\mathbb{E}_z[(\|S(U_i)\| - \|S(D_{i+1})\|)\mathbf{1}_{D_1 \circ \theta(U_i) < \infty} | D_i < \infty]}{\Delta_n} \right| \leq \frac{K}{\Delta_n^2}.$$

Our starting point is the classical Gambler ruin estimate, which in dimension 2, reads with the potential kernel instead of Green’s function,

$$(B.7) \quad \mathbb{P}_z(D_1 < \infty) = \frac{G(0, z) - \mathbb{E}_z[G(0, S(\tau)) | D_1 = \infty]}{\mathbb{E}_z[G(0, S(\tau)) | D_1 < \infty] - \mathbb{E}_z[G(0, S(\tau)) | D_1 = \infty]}.$$

We now expand Green’s function (resp., the potential kernel) using asymptotics (2.2) [resp., (2.3)]. For this purpose, it is convenient to define a random variable

$$X(z) = \frac{1}{\|z\|} (\|S(\tau)\|^2 - \|z\|^2).$$

Note that for any $z \in \mathcal{A}(r_n, n)$, $X(z)/\|z\|$ is small. Indeed,

$$(B.8) \quad \frac{X(z)}{\|z\|} = \frac{(\|S(\tau)\| - \|z\|)(\|S(\tau)\| + \|z\|)}{\|z\|^2}.$$

Since $\Delta_n = n - r_n = O(n^{1/3})$, we have for n large enough,

$$(B.9) \quad \frac{X(z)}{\|z\|} \leq \frac{2(n+1)\Delta_n}{(n-\Delta_n)^2} \leq \frac{8\Delta_n}{n} \quad \text{and} \quad \sup_{z \in \mathcal{A}(r_n, n)} \left(\frac{|X(z)|}{\|z\|} \right)^3 \leq \frac{8^3 \Delta_n^3}{n} \times \frac{1}{n^2}.$$

More precisely, $X(z)$ is of order $2(\|S(\tau)\| - \|z\|)$. Indeed, $\Delta_n^3 \leq n$, and (B.8) yields

$$(B.10) \quad X(z) = 2(\|S(\tau)\| - \|z\|) + \left(\frac{(\|S(\tau)\| - \|z\|)^2}{\|z\|} \right) \implies |X(z) - 2(\|S(\tau)\| - \|z\|)| \leq \frac{1}{\Delta_n}.$$

When dimension $d > 2$, we set $\eta(d) = \frac{d-2}{2}$. In order to use Green’s function asymptotics (2.2), we express $S(\tau)$ in terms of $X(z)$ as follows:

$$(B.11) \quad \frac{1}{\|S(\tau)\|^{d-2}} = \frac{1}{\|z\|^{d-2}} \left(1 + \frac{X(z)}{\|z\|} \right)^{-\eta(d)}.$$

We have a constant K_d such that

$$\begin{aligned}
 & \left| \left(1 + \frac{X(z)}{\|z\|} \right)^{-\eta(d)} \right. \\
 \text{(B.12)} \quad & \left. - \left(1 - \eta(d) \frac{X(z)}{\|z\|} + \eta(d) \frac{\eta(d) + 1}{2} \left(\frac{X(z)}{\|z\|} \right)^2 \right) \right| \\
 & \leq \frac{K_d}{n^2}.
 \end{aligned}$$

For $d > 2$ and any $z \neq 0$, (2.2), (B.11) and (B.12) yield

$$\text{(B.13)} \quad \left| G(0, S(\tau)) - G(0, z) - \eta(d)C_d \left(-\frac{X(z)}{\|z\|^{d-1}} + \frac{\eta(d) + 1}{2} \frac{X(z)^2}{\|z\|^d} \right) \right| \leq \frac{K_d}{n^d}.$$

In dimension 2, the potential kernel asymptotic yields for $K_2 > 0$,

$$\text{(B.14)} \quad \left| a(0, S(\tau)) - a(0, z) - \frac{1}{\pi} \left(\frac{X(z)}{\|z\|} + \frac{1}{2} \frac{X(z)^2}{\|z\|^2} \right) \right| \leq \frac{K_2}{n^2}.$$

In view of (B.14), we assume henceforth that (B.13) holds, but in $d = 2$, we think of $\eta(d)C_d = 1/\pi$, and $\frac{\eta(d)+1}{2} = 1/2$.

Using (B.7) and (B.13), we obtain

$$\begin{aligned}
 & \mathbb{P}_z(D_1 < \infty) \\
 & = \left(\mathbb{E}_z[X(z)|D_1 = \infty] - \bar{C}(z) + O\left(\frac{1}{n}\right) \right) \\
 \text{(B.15)} \quad & \times \left(\mathbb{E}_z[X(z)|D_1 = \infty] - \mathbb{E}_z[X(z)|D_1 < \infty] \right. \\
 & \left. + \underline{C}(z) - \bar{C}(z) + O\left(\frac{1}{n}\right) \right)^{-1},
 \end{aligned}$$

where

$$\begin{aligned}
 \bar{C}(z) &= \frac{\eta(d) + 1}{2} \mathbb{E}_z \left[\frac{X^2(z)}{\|z\|} \middle| D_1 = \infty \right] \quad \text{and} \\
 \underline{C}(z) &= \frac{\eta(d) + 1}{2} \mathbb{E}_z \left[\frac{X^2(z)}{\|z\|} \middle| D_1 < \infty \right].
 \end{aligned}
 \text{(B.16)}$$

Using (B.9), we have some rough estimates on \bar{C} and \underline{C} . For any $z \in \mathcal{A}(r_n, n)$,

$$\text{(B.17)} \quad \bar{C}(z) = O\left(\frac{\Delta_n^2}{n}\right) = O\left(\frac{1}{\Delta_n}\right) \quad \text{and} \quad \underline{C}(z) = O\left(\frac{\Delta_n^2}{n}\right) = O\left(\frac{1}{\Delta_n}\right).$$

Using (B.10), we have better estimates for \bar{C} and \underline{C} .

$$(B.18) \quad \begin{aligned} \bar{C}(z) &= d \frac{(n - \|z\|)^2}{\|z\|} + O\left(\frac{(n - \|z\|) \vee 1}{n}\right), \\ \underline{C}(z) &= d \frac{(\|z\| - r_n)^2}{\|z\|} + O\left(\frac{(\|z\| - r_n) \vee 1}{n}\right). \end{aligned}$$

The rough estimates (B.17) together with (B.10) allow us to derive from (B.15) an estimate for $\mathbb{P}_z(D_1 < \infty)$, for any $z \in \mathcal{A}(r_n, n)$.

$$(B.19) \quad \begin{aligned} &\mathbb{P}_z(D_1 < \infty) \\ &= \frac{\mathbb{E}_z[\|S(\tau)\| - \|z\| | D_1 = \infty] + O(1/\Delta_n)}{\mathbb{E}_z[\|S(\tau)\| - \|z\| | D_1 = \infty] - \mathbb{E}_z[\|S(\tau)\| - \|z\| | D_1 < \infty] + O(1/\Delta_n)} \\ &= \frac{\alpha_0(z) + O(1/\Delta_n)}{\Delta_n(1 + O(1/\Delta_n))}. \end{aligned}$$

This yields (B.4) since $\alpha_0(z) \leq 1 + (n - \|z\|) \vee 1 \leq 2(n - \|z\|) \vee 1$.

Case where $z \in \partial B(0, r_n)$. On $\{D_1 = \infty\}$, we have

$$(B.20) \quad X(z) = 2(\|S(\tau)\| - \|z\|) + O\left(\frac{1}{\Delta_n}\right).$$

On $\{D_1 < \infty\}$, we have

$$X(z) = 2(\|S(\tau)\| - \|z\|) + O\left(\frac{1}{n}\right).$$

This implies

$$(B.21) \quad \bar{C}(z) = d \frac{\Delta_n^2}{\|z\|} + O\left(\frac{\Delta_n}{n}\right) \quad \text{and} \quad \underline{C}(z) = O\left(\frac{1}{n}\right).$$

Thus

$$(B.22) \quad \begin{aligned} &\mathbb{P}_z(D_1 = \infty) \\ &= \frac{2\mathbb{E}_z[\|z\| - \|S(\tau)\| | D_1 < \infty] + \underline{C}(z) + O(1/n)}{\mathbb{E}_z[X(z) | D_1 = \infty] - \mathbb{E}_z[X(z) | D_1 < \infty] + \underline{C}(z) - \bar{C}(z) + O(1/n)} \\ &= \frac{\mathbb{E}_z[\|z\| - \|S(\tau)\| | D_1 < \infty] + O(1/n)}{\Delta_n + O(1)} \\ &= \frac{\mathbb{E}_z[\|z\| - \|S(\tau)\| | D_1 < \infty]}{\Delta_n} + O\left(\frac{1}{\Delta_n^2}\right). \end{aligned}$$

In order to obtain (B.6), we write (B.22) on $\{D_i < \infty\}$, and $z = S(U_i)$ as follows. There is a constant K such that on the event $\{D_i < \infty\}$,

$$(B.23) \quad \left| \mathbb{E}_{S(U_i)}[\mathbf{1}_{D_{i+1}=\infty}] - \frac{\mathbb{E}_{S(U_i)}[(\|S(U_i)\| - \|S(\tau)\|)\mathbf{1}_{D_1 \circ \theta(U_i) < \infty}]}{\Delta_n \times \mathbb{P}_{S(U_i)}(D_1 < \infty)} \right| \leq \frac{K}{\Delta_n^2}.$$

Note that (B.22) implies that $\mathbb{P}_{S(U_i)}(D_1 < \infty) = 1 + O(1/\Delta_n)$, so that (B.23) reads as we integrate over $\{D_i < \infty\}$ with respect to \mathbb{E}_z

$$(B.24) \quad \left| \mathbb{P}_z(D_{i+1} = \infty, D_i < \infty) - \frac{\mathbb{E}_z[\mathbf{1}_{D_i < \infty}(\|S(U_i)\| - \|S(\tau)\|)\mathbf{1}_{D_1 \circ \theta(U_i) < \infty}]}{\Delta_n} \right| \leq \frac{K\mathbb{P}_z(D_i < \infty)}{\Delta_n^2}.$$

We obtain (B.6) as we divide both sides of (B.24) by $\mathbb{P}_z(D_i < \infty)$.

Step 2: We show now that for any $z \in \mathcal{A}(r_n, n)$, we have

$$(B.25) \quad |\mathbb{E}_z[\tau] - (d\Delta_n\alpha_0(z) - d(n - \|z\|)^2)| \leq K((n - \|z\|) \vee 1).$$

When $z \in B_n$ and $n - \|z\| \leq 1$, (B.25) reads

$$(B.26) \quad |\mathbb{E}_z[\tau] - (d\Delta_n\alpha_0(z) - d(n - \|z\|)^2)| \leq K.$$

When $z \in \mathcal{A}(r_n, n)$, and $i \geq 1$, we show that

$$(B.27) \quad \left| \frac{\mathbb{E}_z[\tau \circ \theta(U_i) | D_i < \infty]}{d\Delta_n^2} - \frac{\mathbb{E}_z[(\|S(U_i)\| - \|S(D_{i+1})\|)\mathbf{1}_{D_1 \circ \theta(U_i) < \infty} | D_i < \infty]}{\Delta_n} \right| \leq \frac{K}{\Delta_n^2}.$$

Using that $\{\|S(n)\|^2 - n, n \in \mathbb{N}\}$ is a martingale and the optional sampling theorem (see Lemma 3 of [10]),

$$\begin{aligned} \mathbb{E}_z[\tau] &= \mathbb{E}_z[\|S(\tau)\|^2] - \|z\|^2 = \|z\| \times \mathbb{E}_z[X(z)] \\ &= \|z\| \times (\mathbb{E}_z[X(z) | D_1 = \infty]\mathbb{P}_z(D_1 = \infty) \\ &\quad + \mathbb{E}_z[X(z) | D_1 < \infty]\mathbb{P}_z(D_1 < \infty)). \end{aligned}$$

Thus, using (B.15), simple algebra yields

$$(B.28) \quad \mathbb{E}_z[\tau] = \|z\| \times ((\underline{C}(z) - \bar{C}(z))\mathbb{P}_z(D_1 < \infty) + \bar{C}(z)) + O(1).$$

By recalling (B.18) and (B.4),

$$\begin{aligned} \mathbb{E}_z[\tau] &= d\left(((\|z\| - r_n)^2 - (n - \|z\|)^2 + O(\Delta_n))\left(\frac{\alpha_0(z)}{\Delta_n} + O\left(\frac{(n - \|z\|) \vee 1}{\Delta_n^2}\right)\right) \right) \\ &\quad + d(n - \|z\|)^2 + O((n - \|z\|) \vee 1) \\ &= d(2\|z\| - n - r_n)\alpha_0(z) + d(n - \|z\|)^2 + O((n - \|z\|) \vee 1) \\ &= d\Delta_n\alpha_0(z) - d(n - \|z\|)^2 + O((n - \|z\|) \vee 1). \end{aligned}$$

This yields (B.25).

Assume now that $z \in \partial B(0, r_n)$. From (B.28), we have

$$\mathbb{E}_z[\tau] = \|z\| \times ((\bar{C}(z) - \underline{C}(z))\mathbb{P}_z(D_1 = \infty) + \underline{C}(z)) + O(1).$$

We use (B.6) and (B.21) to obtain

$$\begin{aligned} \mathbb{E}_z[\tau] &= \|z\| \left((d\Delta_n^2 + O(\Delta_n)) \left(\frac{\mathbb{E}_z[\|z\| - \|S(\tau)\| | D_1 < \infty]}{\Delta_n} + O\left(\frac{1}{\Delta_n^2}\right) \right) \right) \\ (B.29) \quad &+ O(1) \\ &= d\Delta_n \mathbb{E}_z[\|z\| - \|S(\tau)\| | D_1 < \infty] + O(1). \end{aligned}$$

Now, write (B.29) as follows. There is a constant K such that for any $z \in \partial \mathbb{B}(0, r_n)$,

$$(B.30) \quad \left| \frac{\mathbb{E}_z[\tau]}{d\Delta_n^2} - \frac{\mathbb{E}_z[\|z\| - \|S(\tau)\| \mathbf{1}_{D_1 < \infty}]}{\Delta_n \mathbb{P}_z(D_1 < \infty)} \right| \leq \frac{K}{\Delta_n^2}.$$

Note that by (B.22) $\Delta_n \mathbb{P}_z(D_1 < \infty) = \Delta_n + O(1)$ and $\|\|z\| - \|S(\tau)\| \mathbf{1}_{D_1 < \infty}\| \leq 1$, thus

$$(B.31) \quad \left| \frac{\mathbb{E}_z[\tau]}{d\Delta_n^2} - \frac{\mathbb{E}_z[\|z\| - \|S(\tau)\| \mathbf{1}_{D_1 < \infty}]}{\Delta_n} \right| \leq \frac{K}{\Delta_n^2}.$$

We replace z by $S(U_i)$ in (B.31) under the event $\{D_i < \infty\}$ to obtain

$$(B.32) \quad \left| \frac{\mathbb{E}_{S(U_i)}[\tau]}{d\Delta_n^2} - \frac{\mathbb{E}_{S(U_i)}[(\|S(U_i)\| - \|S(D_1 \circ \theta(U_i))\|) \mathbf{1}_{D_1 \circ \theta(U_i) < \infty}]}{\Delta_n} \right| \leq \frac{K}{\Delta_n^2}.$$

We multiply both sides of (B.32) by $\mathbf{1}_{D_i < \infty}$, take the expectation on both side of (B.32) and divide by $\mathbb{P}_z(D_i < \infty)$ to obtain (B.27).

Step 3: For $i \geq 1$, we show the following bounds:

$$(B.33) \quad 2 \geq \gamma_i \geq \frac{1}{4d\sqrt{d}}$$

where $\gamma_i = \mathbb{E}_z[(\|S(U_i)\| - \|S(D_{i+1})\|) \mathbf{1}_{D_{i+1} < \infty} | D_i < \infty]$.

The upper bound is obvious. For the lower bound, first we restrict to $\{D_i < \infty\}$, so that $U_i < \infty$. By Lemma 2.2, $S(U_i)$ has a nearest neighbor x , within $\mathbb{B}(0, r_n)$ such that $\|S(U_i)\| - \|x\| \geq 1/(2\sqrt{d})$, and (B.33) is immediate.

Step 4: We show (B.1) using (B.2). For p such that $1 \leq p \leq \infty$, let

$$(B.34) \quad \sigma_p = \sum_{i=1}^p \mathbb{E}_z[\tau \circ \theta(U_i) | D_i < \infty] \prod_{j=1}^{i-1} (1 - \mathbb{P}_z(D_{j+1} = \infty | D_j < \infty)).$$

Now, (B.3) reads $l(z) = \lim_{p \rightarrow \infty} \sigma_p$ (this is the limit of an increasing sequence).

We establish in this step that, for some constant \tilde{K} , any integer n ,

$$(B.35) \quad \lim_{p \rightarrow \infty} \left| 1 - \frac{\sigma_p}{d\Delta_n^2} \right| \leq \frac{\tilde{K}}{\Delta_n}.$$

Once we prove (B.35), we have all the bounds to estimate the right-hand side of (B.2). Indeed, using (B.25), (B.4) and (B.35), we have

$$\begin{aligned}
 & \mathbb{E}_z[\tau] + \mathbb{P}_z(D_1 < \infty) \times l(z) \\
 &= d\Delta_n\alpha_0(z) - d(n - \|z\|)^2 + O((n - \|z\|) \vee 1) \\
 (B.36) \quad &+ \left(\frac{\alpha_0(z)}{\Delta_n} + O\left(\frac{(n - \|z\|) \vee 1}{\Delta_n^2}\right) \right) \times (d\Delta_n^2 + O(\Delta_n)) \\
 &= 2d\Delta_n\alpha_0(z) - 2d(n - \|z\|)^2 + O((n - \|z\|) \vee 1).
 \end{aligned}$$

In order now to prove (B.35), we introduce first some shorthand notation. For p and j positive integers,

$$\begin{aligned}
 (B.37) \quad & a_p = 1 - \frac{\sigma_p}{d\Delta_n^2}, \quad \alpha_j = \mathbb{P}_z(D_{j+1} = \infty | D_j < \infty) \quad \text{and} \\
 & \beta_j = \frac{\mathbb{E}_z[\tau \circ \theta(U_j) | D_j < \infty]}{d\Delta_n^2}.
 \end{aligned}$$

With this notation, (B.6) and (B.27) read as follows:

$$(B.38) \quad \left| \alpha_j - \frac{\gamma_j}{\Delta_n} \right| \leq \frac{K}{\Delta_n^2} \quad \text{and} \quad \left| \beta_j - \frac{\gamma_j}{\Delta_n} \right| \leq \frac{K}{\Delta_n^2} \quad \text{so that } |\alpha_j - \beta_j| \leq \frac{2K}{\Delta_n^2}.$$

Let us rewrite (B.34) as

$$(B.39) \quad a_1 = 1 - \beta_1 \quad \text{and} \quad a_p = a_{p-1} - \beta_p \prod_{j=1}^{p-1} (1 - \alpha_j) \quad \text{for } p > 1.$$

In order to establish (B.35), we show by induction that

$$(B.40) \quad \left| a_p - \prod_{j=1}^p (1 - \alpha_j) \right| \leq \varepsilon_p$$

with for $p > 1$,

$$(B.41) \quad \varepsilon_p = \varepsilon_{p-1} + \frac{2K}{\Delta_n^2} \prod_{j=1}^{p-1} (1 - \alpha_j) \quad \text{and} \quad \varepsilon_1 = \frac{2K}{\Delta_n^2}.$$

Note that it is easy to estimate ε_p from (B.41). By (B.38) and for K_0 large enough there is a constant κ_S such that

$$\begin{aligned}
 \varepsilon_p &\leq \frac{2K}{\Delta_n^2} \left(1 + \sum_{k=1}^p \exp\left(-\sum_{j=1}^k \alpha_j\right) \right) \\
 &\leq \frac{2K}{\Delta_n^2} \left(1 + \sum_{k=1}^p \exp\left(-\sum_{j=1}^k \frac{\gamma_j}{2\Delta_n}\right) \right) \leq \frac{2K}{\Delta_n^2} \kappa_S \Delta_n = \frac{2K\kappa_S}{\Delta_n}.
 \end{aligned}$$

Now, by (B.38), (B.40) holds for $p = 1$, and we assume it holds for $p - 1$. Then

$$(B.42) \quad (1 - \beta_p) \prod_{j=1}^{p-1} (1 - \alpha_j) - \varepsilon_{p-1} \leq a_p \leq (1 - \beta_p) \prod_{j=1}^{p-1} (1 - \alpha_j) + \varepsilon_{p-1}.$$

Then by (B.38), we have (B.40) with ε_p satisfying (B.41).

Now (B.35) follows as we notice that step 3 implies, together with (B.38) and for K_0 large enough, that

$$\lim_{p \rightarrow \infty} \prod_{j=1}^p (1 - \alpha_j) = 0. \quad \square$$

Acknowledgments. The authors thank the CIRM for a friendly atmosphere during their stay as part of a *research in pairs* program.

REFERENCES

- [1] ASSELAH, A. and GAUDILLIÈRE, A. (2010). Sub-logarithmic fluctuations for internal DLA. Preprint. Available at arXiv:1011.4592.
- [2] DIACONIS, P. and FULTON, W. (1991). A growth model, a game, an algebra, Lagrange inversion, and characteristic classes. *Rend. Semin. Mat. Univ. Politec. Torino* **49** 95–119. MR1218674
- [3] FRIEDRICH, T. and LEVINE, L. (2011). Fast simulation of large-scale growth models. In *Approximation, Randomization, and Combinatorial Optimization. Lecture Notes in Computer Science* **6845** 555–566. Springer, Heidelberg. MR2863290
- [4] JERISON, D., LEVINE, L. and SHEFFIELD, S. (2010). Internal DLA in higher dimensions. Preprint. Available at arXiv:1012.3453.
- [5] JERISON, D., LEVINE, L. and SHEFFIELD, S. (2012). Logarithmic fluctuations for internal DLA. *J. Amer. Math. Soc.* **25** 271–301. MR2833484
- [6] KLEBER, M. (2005). Goldbug variations. *Math. Intelligencer* **27** 55–63. MR2164256
- [7] LANDOLT, D. (1987). Fundamental aspects of electropolishing. *Electrochimica Acta* **32** 1–11.
- [8] LAWLER, G. F. (1991). *Intersections of Random Walks*. Birkhäuser, Boston, MA. MR1117680
- [9] LAWLER, G. F. (1995). Subdiffusive fluctuations for internal diffusion limited aggregation. *Ann. Probab.* **23** 71–86. MR1330761
- [10] LAWLER, G. F., BRAMSON, M. and GRIFFEATH, D. (1992). Internal diffusion limited aggregation. *Ann. Probab.* **20** 2117–2140. MR1188055
- [11] LAWLER, G. F. and LIMIC, V. (2010). *Random Walk: A Modern Introduction. Cambridge Studies in Advanced Mathematics* **123**. Cambridge Univ. Press, Cambridge. MR2677157
- [12] LEVINE, L. and PERES, Y. (2009). Strong spherical asymptotics for rotor-router aggregation and the divisible sandpile. *Potential Anal.* **30** 1–27. MR2465710
- [13] MEAKIN, P. and DEUTCH, J. M. (1986). The formation of surfaces by diffusion limited annihilation. *J. Chem. Phys.* **85** 2320–2325.
- [14] MOORE, C. and MACHTA, J. (2000). Internal diffusion-limited aggregation: Parallel algorithms and complexity. *J. Stat. Phys.* **99** 661–690. MR1766912

LAMA
UNIVERSITÉ PARIS-EST
61 AVENUE DU GÉNÉRAL DE GAULLE
94010 CRÉTEIL CEDEX
FRANCE
E-MAIL: amine.asselah@univ-paris12.fr

LATP
UNIVERSITÉ DE PROVENCE
CNRS, 39 RUE F. JOLIOT CURIE
13013 MARSEILLE
FRANCE
E-MAIL: gaudilli@cmi.univ-mrs.fr