

Multivariate Gaussian approximations on Markov chaoses

Simon Campese* Ivan Nourdin[†] Giovanni Peccati[‡]
Guillaume Poly[§]

Abstract

We prove a version of the multidimensional Fourth Moment Theorem for chaotic random vectors, in the general context of diffusion Markov generators. In addition to the usual componentwise convergence and unlike the infinite-dimensional Ornstein-Uhlenbeck generator case, another moment-type condition is required to imply joint convergence of a given sequence of vectors.

Keywords: Markov diffusion generator; Fourth Moment Theorem; multivariate normal approximations.

AMS MSC 2010: 60F05; 60J35; 60J99.

Submitted to ECP on October 8, 2015, final version accepted on May 16, 2016.

1 Introduction

The *Fourth Moment Theorem* (discovered by Nualart and Peccati in [13] and later extended by Nualart and Ortiz-Latorre in [12]) states that, inside a fixed Wiener chaos, a sequence of random variables $(F_n, n \geq 1)$ converges in distribution towards a standard Gaussian random variable if and only if $\mathbb{E}[F_n^2] \rightarrow 1$ and $\mathbb{E}[F_n^4] \rightarrow 3$ as $n \rightarrow \infty$. Recently, in the pathbreaking contribution [5], Ledoux approached this Fourth Moment Phenomenon in the more general context of diffusion Markov generators, and was able to provide a new proof of such a result adopting a purely spectral point of view. Later on, in [1], Azmoodeh, Campese and Poly generalized the concept of chaos originally introduced in [5] and were able not only to obtain a more transparent proof of the classical Fourth Moment Theorem, but also to exhibit many new situations where the Fourth Moment Phenomenon occurs (e.g. the Laguerre or Jacobi chaoses). One should notice that the collection of techniques introduced in [1] have also been successfully applied in other contexts, e.g. for deducing moment conditions in limit theorems (see [2]), or in the study of the so-called *real Gaussian product conjecture* (see [6]).

In this paper, we investigate a multidimensional counterpart of the Fourth Moment Theorem by using the aforementioned approach based on Markov semigroup. In the case of Wiener chaos, the multidimensional version of the Fourth Moment Theorem is due to Peccati and Tudor [14], and is given by the following statement. For the rest of the paper, the symbol ' \xrightarrow{d} ' indicates convergence in distribution.

*Università di Roma Tor Vergata, Italy. E-mail: campese@mat.uniroma2.it

[†]Université du Luxembourg, Luxembourg. E-mail: ivan.nourdin@uni.lu

[‡]Université du Luxembourg, Luxembourg. E-mail: giovanni.peccati@gmail.com

[§]Université de Rennes 1, France. E-mail: guillaume.poly@univ-rennes1.fr

Theorem 1.1 (See [14]). *Let $p_1, \dots, p_d \geq 1$ be fixed integers and $F_n = (F_{1,n}, \dots, F_{d,n})$, $n \geq 1$, be a sequence of vectors such that $F_{i,n}$ belongs to the p_i th Wiener chaos of some Gaussian field, for all $i = 1, \dots, d$ and all n . Furthermore, assume that $\lim_{n \rightarrow \infty} \text{Cov } F_n = C$, and denote by $Z = (Z_1, \dots, Z_d)$ a centered Gaussian random vector with covariance matrix C . Then, the following three assertions are equivalent, as $n \rightarrow \infty$:*

- (i) $F_n \xrightarrow{d} Z$
- (ii) $F_{i,n} \xrightarrow{d} Z_i$ for all $1 \leq i \leq d$.
- (iii) $\mathbb{E}[F_{i,n}^4] \rightarrow \mathbb{E}[Z_i^4]$ for all $1 \leq i \leq d$.

In other words, for sequences of random vectors living inside a fixed system of Wiener chaoses, *componentwise* convergence implies *joint* convergence. Our main result is the following analogue of Theorem 1.1 in the abstract Markov generator framework (unexplained notation and definitions – in particular the notion of a *chaotic vector* – will be formally introduced in the sequel).

Theorem 1.2. *Let L be a diffusion Markov generator acting on $L^2(E, \mu)$ with invariant measure μ and discrete spectrum $0 < \lambda_0 < \lambda_1 < \dots$, fix integers $k_1, \dots, k_d \geq 1$, and let $F_n = (F_{1,n}, \dots, F_{d,n})$, $n \geq 1$, be a sequence of chaotic vectors such that $F_{i,n} \in \ker(L + \lambda_{k_i} \text{Id})$, for $1 \leq i \leq d$ and all n . Furthermore, assume that $\lim_{n \rightarrow \infty} \text{Cov } F_n = C$ and denote by $Z = (Z_1, \dots, Z_d)$ a centered Gaussian random vector with covariance matrix C (defined on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$). Consider the following asymptotic relations (i) and (ii), for $n \rightarrow \infty$:*

- (i) $F_n \xrightarrow{d} Z$,
- (ii) for every $1 \leq i, j \leq d$,

$$\int_E F_{i,n}^2 F_{j,n}^2 d\mu \rightarrow \mathbb{E}[Z_i^2 Z_j^2]. \tag{1.1}$$

Then, (ii) implies (i), and the converse implication (i) \Rightarrow (ii) holds whenever the sequence $(F_{i,n}^2 F_{j,n}^2, n \geq 1)$ is uniformly integrable for every $1 \leq i, j \leq d$.

Remark 1.3. The additional mixed moment condition (ii) has no counterpart in the statement of Theorem 1.1: indeed, whenever the components of the vectors F_n belong to the Wiener chaos of some Gaussian field, we will explain in detail in Section 3 why (1.1) is automatically satisfied for every $1 \leq i \neq j \leq d$ when it holds for $1 \leq i = j \leq d$. We also observe that a sufficient condition for the class

$$(F_{i,n}^2 F_{j,n}^2, n \geq 1),$$

to be uniformly integrable for every i, j , is that, for some $\epsilon > 0$,

$$\sup_n \int_E |F_{i,n}|^{4+\epsilon} d\mu < \infty, \quad \text{for every } i = 1, \dots, d. \tag{1.2}$$

Finally, if the sequence $(F_n, n \geq 1)$ lives in a fixed sum of Gaussian chaoses, then (1.2) is automatically implied by the relation $\lim_{n \rightarrow \infty} \text{Cov } F_n = C$, by virtue of a standard hypercontractivity argument – see e.g. [8, Section 2.8.3]. General sufficient conditions for the semigroup associated with L to be hypercontractive can be found e.g. in [3].

As in the one-dimensional case, it is possible in our abstract framework to provide a proof of Theorem 1.2 that is *not* based on the use of product formulae, and that exploits instead the spectral information embedded into the underlying generator L .

The rest of this paper is organized as follows. In Section 2, we will introduce the abstract Markov generator setting and recall the main one-dimensional findings from [1]. In Section 3, we will define multidimensional chaos and present the proof of Theorem 1.2; we also provide a careful analysis of the additional condition (ii)_b appearing in our Theorem 1.2.

2 Preliminaries

In this section, we introduce the general diffusion Markov generator setting. For a detailed treatment, we refer to the monograph [4].

Throughout the rest of the paper, we fix a probability space (E, \mathcal{F}, μ) and a symmetric Markov generator L with state space E and invariant measure μ . We assume that L has discrete spectrum $S = \{-\lambda_k : k \geq 0\}$ and order its eigenvalues by magnitude, i.e. $0 = \lambda_0 < \lambda_1 < \lambda_2 < \dots$. In the language of functional analysis, L is a self-adjoint, linear operator on $L^2(E, \mu)$ with the property that $L1 = 0$. By standard spectral theory, L is diagonalizable and we have that

$$L^2(E, \mu) = \bigoplus_{k=0}^{\infty} \ker(L + \lambda_k \text{Id}).$$

We denote by L^{-1} the pseudo-inverse of L , defined on $L^2(E, \mu)$ by $L^{-1}1 = 0$ and $L^{-1}F = -\frac{1}{\lambda}F$ for any $F \in \ker(L + \lambda \text{Id})$ such that $\lambda \neq 0$. It is immediate to check that $LL^{-1}F = F - \int_E F d\mu$ for every $F \in L^2(E, \mu)$. The associated bilinear carré du champ operator Γ is defined by

$$\Gamma(F, G) = \frac{1}{2} (L(FG) - FLG - GLF),$$

whenever the right-hand side exists. This is in particular the case when F and G are two jointly chaotic eigenfunctions in the sense of the forthcoming Definition 3.3. Furthermore, as L is self-adjoint and $L1 = 0$, we immediately deduce the integration by parts formula

$$\int_E \Gamma(F, G) d\mu = - \int_E FLG d\mu = - \int_E GLF d\mu. \tag{2.1}$$

A symmetric Markov generator is called *diffusive*, if it satisfies the diffusion property

$$L\phi(F) = \phi'(F)LF + \phi''(F)\Gamma(F, F)$$

for all smooth test functions $\phi: \mathbb{R} \rightarrow \mathbb{R}$ and any $F \in L^2(E, \mu)$. Equivalently, Γ is a derivation i.e.

$$\Gamma(\phi(F), G) = \phi'(F)\Gamma(F, G). \tag{2.2}$$

Considering vectors $F = (F_1, \dots, F_d)$ and test functions $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}$, the diffusion and derivation properties yield that

$$L\phi(F) = L\phi(F_1, \dots, F_d) = \sum_{i=1}^d \frac{\partial \varphi}{\partial x_i}(F) LF_i + \sum_{i,j=1}^d \frac{\partial^2 \varphi}{\partial x_i \partial x_j}(F) \Gamma(F_i, F_j)$$

and

$$\Gamma(\varphi(F), G) = \sum_{i=1}^d \frac{\partial \varphi}{\partial x_i}(F) \Gamma(F_i, G),$$

respectively. In [1], the following definition of chaos was given.

Definition 2.1 (See [1]). *Let L be a symmetric Markov generator with discrete spectrum S , and let $F \in \ker(L + \lambda_p \text{Id})$ be an eigenfunction of L (with eigenvalue $\lambda_p \in S$). We say that F is chaotic, or a chaos eigenfunction, if*

$$F^2 \in \bigoplus_{k: \lambda_k \leq 2\lambda_p} \ker(L + \lambda_k \text{Id}). \tag{2.3}$$

Condition (2.3) means that, if F is a chaos eigenfunction of L with eigenvalue λ_p (say), then the (orthogonal) decomposition of its square along the spectrum of L only contains eigenfunctions associated with eigenvalues that are *less than or equal to twice* λ_p . Note that this property is satisfied by *all* eigenfunctions of L in many crucial instances, e.g. when L is the generator of the Ornstein-Uhlenbeck semigroup, the Laguerre generator or the Jacobi generator (see [1]). Starting from this definition, and by only using the spectral information embedded into the generator L , the authors of [1] were able to deduce Fourth Moment Theorems for sequences of chaotic eigenfunctions and several target distributions, drastically simplifying all known proofs. The analogue in this framework of the classical Fourth Moment Theorem reads as follows.

Theorem 2.2 (Abstract Fourth Moment Theorem, see [1]). *Let L be a symmetric diffusion Markov generator with discrete spectrum S and let $(F_n, n \geq 1)$ be a sequence of chaotic eigenfunctions of L with respect to the same (fixed) eigenvalue, such that $\int_E F_n^2 d\mu \rightarrow 1, n \rightarrow \infty$. Consider the following three conditions, as $n \rightarrow \infty$:*

- (i) $F_n \xrightarrow{d} Z$, where $Z \sim \mathcal{N}(0, 1)$,
- (ii) $\int_E F_n^4 d\mu \rightarrow 3$,
- (iii) $\text{Var}(\Gamma(F_n)) \rightarrow 0$.

Then, the implications (ii) \Rightarrow (iii) \Rightarrow (i) hold. Furthermore, if the sequence $(F_n^4, n \geq 1)$ is uniformly integrable, one has that (i) \Rightarrow (ii).

The next section is devoted to the proof of our main results.

3 Main results

Let us begin by noting the general fact that (under standard regularity assumptions) the distance between the distribution of a random vector $F = (F_1, \dots, F_d)$ and a multivariate Gaussian law is controlled by the expression $\sum_{i,j=1}^d \text{Var} \Gamma(F_i, -L^{-1} F_j)$. This fact can either be shown by using the characteristic function method (see [12]), quantitatively by Stein’s method (see [9]) or by means of the so-called “smart path” technique (see [7]). The proofs carry over to our setting almost verbatim, by replacing the integration by parts formula of Malliavin calculus with the analogous relation (2.1) for the carré du champ operator Γ .

In order to keep our paper as self-contained as possible, instead of using the above mentioned bounds, in the sequel we shall exploit the following estimate involving characteristic functions: the proof is a Fourier-type variation of the smart path method.

Proposition 3.1. *Let $F = (F_1, \dots, F_d)$ be such that $F_j \in L^2(E, \mu)$ and $\int_E F_j d\mu = 0$ for $1 \leq j \leq d$. Assume furthermore the existence of a finite $N \geq 1$ such that $F_j \in \bigoplus_{k=0}^N \ker(L + \lambda_k \text{Id})$ for all $1 \leq j \leq d$. Let γ_d be the law of a d -dimensional centered Gaussian random variable with covariance matrix C . Then, for any $t \in \mathbb{R}^d$ it holds that*

$$\left| \int_E e^{i\langle t, F \rangle_2} d\mu - \int_{\mathbb{R}^d} e^{i\langle t, x \rangle_2} d\gamma_d(x) \right| \leq \|t\|_2^2 \sqrt{\sum_{i,j=1}^d \int_E (C_{ij} - \Gamma(F_i, -L^{-1} F_j))^2 d\mu}.$$

Proof. Fix $t \in \mathbb{R}^d$ and define $\Psi(\theta) = e^{\frac{\theta^2}{2} t^* C t} \int_E e^{i\theta \langle t, F \rangle_2} d\mu$ for $\theta \in [0, 1]$. It is straightforward to verify that Ψ is differentiable on $(0, 1)$, so that

$$\int_E e^{i\langle t, F \rangle_2} d\mu - \int_{\mathbb{R}^d} e^{i\langle t, x \rangle_2} d\gamma_d(x) = e^{-\frac{1}{2} t^* C t} (\Psi(1) - \Psi(0)) = e^{-\frac{1}{2} t^* C t} \int_0^1 \Psi'(\theta) d\theta,$$

and that the derivative of Ψ is given by

$$\Psi'(\theta) = e^{\frac{\theta^2}{2} t^* C t} \left(\theta t^* C t \int_E e^{i\theta \langle t, F \rangle_2} d\mu + i \int_E \langle t, F \rangle_2 e^{i\theta \langle t, F \rangle_2} d\mu \right). \tag{3.1}$$

Using both the integration by parts formula (2.1) and the diffusion property (2.2) for Γ yields that

$$\Psi'(\theta) = \theta e^{\frac{\theta^2}{2} t^* C t} \sum_{i,j=1}^d t_i t_j \int_E (C_{ij} - \Gamma(F_i, -L^{-1} F_j)) e^{i\theta \langle t, F \rangle_2} d\mu. \tag{3.2}$$

The conclusion follows from an application of the Cauchy-Schwarz inequality. □

Remark 3.2. In the previous statement, to require the existence of a finite $N \geq 1$ such that $F_j \in \bigoplus_{k=0}^N \ker(L + \lambda_k \text{Id})$ for all $1 \leq j \leq d$ is for simplicity only. It is enough for our purpose and it prevents us to impose conditions for carrying out the interchange of derivative and integration and the integration-by-parts.

Definition 3.3 (Jointly chaotic eigenfunction).

1. For $1 \leq i, j \leq d$ and $\lambda_{k_i} \in S$, let $F_i \in \ker(L + \lambda_{k_i} \text{Id})$ and $F_j \in \ker(L + \lambda_{k_j} \text{Id})$ be two eigenfunctions of L . We say that F_i and F_j are jointly chaotic, if

$$F_i F_j \in \bigoplus_{r: \lambda_r \leq \lambda_{k_i} + \lambda_{k_j}} \ker(L + \lambda_r \text{Id}). \tag{3.3}$$

2. Let $F = (F_1, \dots, F_d)$ be a vector of eigenfunctions of L , such that $F_i \in \ker(L + \lambda_{k_i} \text{Id})$ for $1 \leq i \leq d$. Whenever any two of its components (possibly the same) are jointly chaotic, we say that F is chaotic. In particular, this implies that each component of a chaotic vector is chaotic in the sense of Definition 2.1.

Note that if a vector $F = (F_1, \dots, F_d)$ is chaotic in the sense of Definition 3.3, then in particular each component F_i , $1 \leq i \leq d$, is chaotic in the sense of Definition 2.1. Indeed, as F_i is by definition jointly chaotic with itself, statement (3.3) of Definition 3.3 exactly becomes statement (2.3) of Definition 2.1:

$$F_i^2 \in \bigoplus_{r: \lambda_r \leq 2\lambda_{k_i}} \ker(L + \lambda_r \text{Id}).,$$

We also observe that, in many important examples, all vectors of eigenfunctions are chaotic. In particular, this is the case for Wiener, Laguerre and Jacobi chaos (see [1]). A crucial ingredient for the proof of our main result is the following result, whose short proof can be found in [1].

Theorem 3.4 ([1, Thm. 2.1]). Fix an eigenvalue $-\lambda = -\lambda_n \in S$ and assume that $F \in \bigoplus_{k=0}^n \ker(L + \lambda_k \text{Id})$. Then it holds that, for any $\eta \geq \lambda_n$,

$$\int_E F (L + \eta \text{Id})^2 (F) d\mu \leq \eta \int_E F (L + \eta \text{Id}) (F) d\mu.$$

We are now ready to prove our main Theorem 1.2.

Proof of Theorem 1.2. Since the remaining parts of the statement are straightforward, we will only prove the implication (ii) \Rightarrow (i). Throughout the proof, we will denote the covariance matrix of F_n by $C_n = (C_{ij,n})_{i,j=1}^n$, so that thus $C_n \rightarrow C$ as $n \rightarrow \infty$. According to Proposition 3.1, it suffices to show that, if (ii) is satisfied then, for $1 \leq i, j \leq d$,

$$\int_E (\Gamma(F_{i,n}, -L^{-1} F_{j,n}) - C_{ij})^2 d\mu \rightarrow 0, \tag{3.4}$$

as $n \rightarrow \infty$. It holds that $\Gamma(F_{i,n}, -L^{-1}F_{j,n}) = \frac{1}{\lambda_{k_j}}\Gamma(F_{i,n}, F_{j,n})$ and $\int_E \Gamma(F_{i,n}, F_{j,n}) d\mu = \lambda_{k_j}C_{ij,n}$. The integral in (3.4) can thus be rewritten as

$$\frac{1}{\lambda_{k_j}^2} \text{Var} \Gamma(F_{i,n}, F_{j,n}) + (C_{ij} - C_{ij,n})^2. \tag{3.5}$$

For $i = j$, the quantity in (3.5) tends to zero thanks to the one-dimensional Fourth Moment Theorem 2.2.

Let us thus assume that $i \neq j$. By definition of Γ , we can write

$$\Gamma(F_{i,n}, F_{j,n}) - \lambda_{k_j}C_{ij,n} = \frac{1}{2} (L + (\lambda_{k_i} + \lambda_{k_j}) \text{Id}) (F_{i,n}F_{j,n} - a_{ij}C_{ij,n}),$$

where $a_{ij} = \frac{2\lambda_{k_j}}{\lambda_{k_i} + \lambda_{k_j}}$. Inserting the definition of the carré du champ operator, we get that

$$\begin{aligned} & \int_E (\Gamma(F_{i,n}, -L^{-1}F_{j,n}) - C_{ij,n})^2 d\mu \\ &= \frac{1}{\lambda_{k_j}^2} \int_E (\Gamma(F_{i,n}, F_{j,n}) - \lambda_{k_j}C_{ij,n})^2 d\mu \\ &= \frac{1}{4\lambda_{k_j}^2} \int_E ((L + (\lambda_{k_i} + \lambda_{k_j}) \text{Id}) (F_{i,n}F_{j,n} - a_{ij}C_{ij,n}))^2 d\mu \\ &= \frac{1}{4\lambda_{k_j}^2} \int_E (F_{i,n}F_{j,n} - a_{ij}C_{ij,n}) (L + (\lambda_{k_i} + \lambda_{k_j}) \text{Id})^2 (F_{i,n}F_{j,n} - a_{ij}C_{ij,n}) d\mu. \end{aligned}$$

Therefore, by Theorem 3.4 and the fact that $\int_E \Gamma(F_{i,n}, F_{j,n}) d\mu = \lambda_{k_j}C_{ij,n}$, it holds that

$$\begin{aligned} & \int_E (\Gamma(F_{i,n}, -L^{-1}F_{j,n}) - C_{ij,n})^2 d\mu \\ & \leq \frac{\lambda_{k_i} + \lambda_{k_j}}{4\lambda_{k_j}^2} \int_E (F_{i,n}F_{j,n} - a_{ij}C_{ij,n}) (L + (\lambda_{k_i} + \lambda_{k_j}) \text{Id}) (F_{i,n}F_{j,n} - a_{ij}C_{ij,n}) d\mu \\ &= \frac{1}{a_{ij}\lambda_{k_j}} \int_E (F_{i,n}F_{j,n} - a_{ij}C_{ij,n}) (\Gamma(F_{i,n}, F_{j,n}) - \lambda_{k_j}C_{ij,n}) d\mu \\ &= \frac{1}{a_{ij}\lambda_{k_j}} \left(\int_E F_{i,n}F_{j,n}\Gamma(F_{i,n}, F_{j,n}) d\mu - \lambda_{k_j}C_{ij,n}^2 \right) \tag{3.6} \end{aligned}$$

Now, by the diffusion property (2.2), the integration by parts formula (2.1) and the Cauchy-Schwarz inequality, we have

$$\begin{aligned} & \int_E F_{i,n}F_{j,n}\Gamma(F_{i,n}, F_{j,n}) d\mu = \frac{1}{4} \int_E \Gamma(F_{i,n}^2, F_{j,n}^2) d\mu = -\frac{1}{4} \int_E F_{i,n}^2 L(F_{j,n}^2) d\mu \\ &= -\frac{1}{2} \int_E F_{i,n}^2 (F_{j,n} L F_{j,n} + \Gamma(F_{j,n}, F_{j,n})) d\mu \\ &= \frac{\lambda_{k_j}}{2} \int_E F_{i,n}^2 F_{j,n}^2 d\mu - \frac{1}{2} \int_E F_{i,n}^2 \Gamma(F_{j,n}, F_{j,n}) d\mu \\ &= \frac{\lambda_{k_j}}{2} \int_E F_{i,n}^2 (F_{j,n}^2 - C_{jj,n}) d\mu - \frac{1}{2} \int_E F_{i,n}^2 (\Gamma(F_{j,n}, F_{j,n}) - \lambda_{k_j}C_{jj,n}) d\mu \\ &\leq \frac{\lambda_{k_j}}{2} \int_E F_{i,n}^2 (F_{j,n}^2 - C_{jj,n}) d\mu + \frac{1}{2} \sqrt{m_4(F_{i,n})} \sqrt{\text{Var}(\Gamma(F_{j,n}, F_{j,n}))} \\ &= \frac{\lambda_{k_j}}{2} \left(\int_E F_{i,n}^2 F_{j,n}^2 d\mu - C_{ii,n}C_{jj,n} \right) + \frac{1}{2} \sqrt{m_4(F_{i,n})} \sqrt{\text{Var}(\Gamma(F_{j,n}, F_{j,n}))}. \end{aligned}$$

Plugging such an estimate into (3.6), we can thus write

$$\int_E (\Gamma(F_{i,n}, -L^{-1}F_{j,n}) - C_{ij,n})^2 d\mu \leq \frac{1}{2\lambda_{k_j} a_{ij}} \left(\sqrt{m_4(F_{i,n})} \sqrt{\text{Var}(\Gamma(F_{j,n}, F_{j,n}))} + \lambda_{k_j} R_{i,j}(n) \right),$$

where

$$R_{ij}(n) = \int_E F_{i,n}^2 F_{j,n}^2 d\mu - C_{ii,n} C_{jj,n} - 2C_{ij,n}^2.$$

By virtue of Theorem 2.2 and (3.5), we are now left to show that $R_{ij}(n) \rightarrow 0$, as $n \rightarrow \infty$. This follows immediately from $\mathbb{E}[Z_i^2 Z_j^2] = \mathbb{E}[Z_i^2] \mathbb{E}[Z_j^2] + 2\mathbb{E}[Z_{ij}]^2$ and our assumption $C_n \rightarrow C$. \square

Remark 3.5. In the framework of the classical Theorem 1.1, where L is the infinite-dimensional Ornstein-Uhlenbeck generator and its eigenfunctions are multiple Wiener-Itô integrals, componentwise Gaussian convergence in distribution always yields joint Gaussian convergence. From our abstract point of view, we can explain (and generalize) this phenomenon by means of the next result.

Proposition 3.6. *In the setting and with the notation of Theorem 1.2, assume in addition that*

1. L is ergodic, i.e. its kernel only consists of constant functions.
2. For $1 \leq i, j \leq d$ such that $\lambda_{k_i} = \lambda_{k_j}$, it holds that

$$\int_E \pi_{2\lambda_{k_i}}(F_{i,n}^2) \pi_{2\lambda_{k_j}}(F_{j,n}^2) d\mu - 2 \left(\int_E F_{i,n} F_{j,n} d\mu \right)^2 \rightarrow 0, \tag{3.7}$$

where π_λ denotes the orthogonal projection onto $\ker(L + \lambda \text{Id})$.

Then, the following two assertions are equivalent.

- (i) $F_n \xrightarrow{d} Z$
- (ii) For $1 \leq i \leq d$ it holds that $\int_E F_{i,n}^4 d\mu \rightarrow \mathbb{E}[Z_i^4]$.

Proof. In view of Theorem 1.2, we have only to show that condition (ii) implies

$$\int_E F_{i,n}^2 F_{j,n}^2 d\mu \rightarrow \mathbb{E}[Z_i^2 Z_j^2]$$

or equivalently (as $\mathbb{E}[Z_i^2 Z_j^2] = \mathbb{E}[Z_i^2] \mathbb{E}[Z_j^2] + 2\mathbb{E}[Z_{ij}]^2$) that

$$\int_E F_{i,n}^2 F_{j,n}^2 d\mu - \int_E F_{i,n}^2 d\mu \int_E F_{j,n}^2 d\mu - 2 \left(\int_E F_{i,n} F_{j,n} d\mu \right)^2 \rightarrow 0$$

for $1 \leq i, j \leq d$. To do so, we first note that, by the definition of Γ and the chaotic property of $F_{i,n}$, it holds that

$$\Gamma(F_{i,n}, F_{i,n}) = \frac{1}{2} (L + 2\lambda_{k_i} \text{Id})(F_{i,n}^2) = \frac{1}{2} \sum_{\substack{r \in \mathbb{N} \\ \lambda_r < 2\lambda_{k_i}}} (2\lambda_{k_i} - \lambda_r) \pi_{\lambda_r}(F_{i,n}^2).$$

Therefore, by the orthogonality of the projections corresponding to different eigenvalues and by the one-dimensional Fourth Moment Theorem 2.2, $\int_E F_{i,n}^4 d\mu \rightarrow \mathbb{E}[Z_i^4]$ implies that

$$\int_E \pi_{\lambda_r}(F_{i,n}^2)^2 d\mu \rightarrow 0, \quad \text{for all } r \text{ such that } 0 < \lambda_r < 2\lambda_{k_i}. \tag{3.8}$$

We exploit this fact by writing

$$\int_E F_{i,n}^2 F_{j,n}^2 \, d\mu = \int_E \pi_0(F_{i,n}^2) F_{j,n}^2 \, d\mu + \sum_{\substack{r \in \mathbb{N} \\ 0 < \lambda_r < 2\lambda_{k_i}}} \int_E \pi_{\lambda_r}(F_{i,n}^2) F_{j,n}^2 \, d\mu + \int_E \pi_{2\lambda_{k_i}}(F_{i,n}^2) F_{j,n}^2 \, d\mu.$$

Assume without loss of generality that $\lambda_{k_j} \leq \lambda_{k_i}$. The ergodicity assumption on L forces $\pi_0(F_{i,n}^2)$ to be constant and thus

$$\int_E \pi_0(F_{i,n}^2) F_{j,n}^2 \, d\mu = \pi_0(F_{i,n}^2) \int_E F_{j,n}^2 \, d\mu = \int_E F_{i,n}^2 \, d\mu \int_E F_{j,n}^2 \, d\mu.$$

By Cauchy-Schwarz and (3.8), all integrals $\int_E \pi_{\lambda_r}(F_{i,n}^2) F_{j,n}^2 \, d\mu$ inside the sum in the middle vanish in the limit. Finally, assumption (3.7) ensures that the third term (which is zero if $\lambda_{k_i} \neq \lambda_{k_j}$), exhibits the wanted asymptotic behaviour. \square

Remark 3.7. As already mentioned, both of these additional assumptions are always verified in the case of the infinite-dimensional Ornstein-Uhlenbeck generator (see for example [8] or [11] for any unexplained notation). While the ergodicity is immediate, more effort is needed to show (3.7). For a multiple integral $I_p(f)$ with $f \in \mathfrak{H}^{\odot p}$, the well known product formula yields that $\pi_{2p}(I_p(f)^2) = I_{2p}(f \otimes f)$. If $I_p(g)$, $g \in \mathfrak{H}^{\odot p}$ is another multiple integral, one can show (see for example [10, Lemma 2.2(2)]) that

$$\mathbb{E}[I_{2p}(f \otimes f) I_{2p}(g \otimes g)] = 2\mathbb{E}[I_p(f) I_p(g)]^2 + \sum_{r=1}^{p-1} p!^2 \binom{p}{r}^2 \langle f \otimes_r f, g \otimes_r g \rangle_{\mathfrak{H}^{\otimes 2p}}.$$

Replacing the kernels f and g by two sequences (f_n) and (g_n) , the classical Fourth Moment Theorem implies that scalar products inside the sum vanish in the limit if (at least one of) the two sequences $I_p(f_n)$ and $I_p(g_n)$ converges in distribution towards a Gaussian.

Acknowledgments. We would like to thank two anonymous referees for their thorough reading of an earlier version of this manuscript and for comments and suggestions which improved the exposition of our results. S. Campese was partially supported by ERC grant 277742 PASCAL. I. Nourdin was partially supported by the Grant F1R-MTH-PUL-15CONF (CONFLUENT) at Luxembourg University. G. Peccati was partially supported by the Grant F1R-MTH-PUL-15STAR (STARS) at Luxembourg University.

References

- [1] Ehsan Azmoodeh, Simon Campese, and Guillaume Poly. Fourth Moment Theorems for Markov diffusion generators. *J. Funct. Anal.*, 266(4):2341–2359, 2014. MR-3150163
- [2] Ehsan Azmoodeh, Dominique Malicet, Guillaume Mijoule, and Guillaume Poly. Generalization of the Nualart-Peccati criterion. *Ann. Probab.*, 44(2):924–954, 2016. MR-3474463
- [3] Dominique Bakry. L’hypercontractivité et son utilisation en théorie des semigroupes. In *Lectures on probability theory (Saint-Flour, 1992)*, volume 1581 of *Lecture Notes in Math.*, pages 1–114. Springer, Berlin, 1994. MR-1307413
- [4] Dominique Bakry, Ivan Gentil, and Michel Ledoux. *Analysis and geometry of Markov diffusion operators*, volume 348 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Cham, 2014. MR-3155209
- [5] M. Ledoux. Chaos of a Markov operator and the fourth moment condition. *Ann. Probab.*, 40(6):2439–2459, 2012. MR-3050508

- [6] Dominique Malicet, Ivan Nourdin, Giovanni Peccati, and Guillaume Poly. Squared chaotic random variables: new moment inequalities with applications. *J. Funct. Anal.*, 270(2):649–670, 2016. MR-3425898
- [7] Ivan Nourdin and Giovanni Peccati. Stein’s method meets Malliavin calculus: a short survey with new estimates. In *Recent development in stochastic dynamics and stochastic analysis*, volume 8 of *Interdiscip. Math. Sci.*, pages 207–236. World Sci. Publ., Hackensack, NJ, 2010. MR-2807823
- [8] Ivan Nourdin and Giovanni Peccati. *Normal approximations with Malliavin calculus*, volume 192 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2012. From Stein’s method to universality. MR-2962301
- [9] Ivan Nourdin, Giovanni Peccati, and Anthony Réveillac. Multivariate normal approximation using Stein’s method and Malliavin calculus. *Ann. Inst. Henri Poincaré Probab. Stat.*, 46(1):45–58, 2010. MR-2641769
- [10] Ivan Nourdin and Jan Rosiński. Asymptotic independence of multiple Wiener-Itô integrals and the resulting limit laws. *Ann. Probab.*, 42(2):497–526, 2014. MR-3178465
- [11] David Nualart. *The Malliavin calculus and related topics*. Probability and its Applications (New York). Springer-Verlag, Berlin, second edition, 2006. MR-2200233
- [12] D. Nualart and S. Ortiz-Latorre. Central limit theorems for multiple stochastic integrals and Malliavin calculus. *Stochastic Process. Appl.*, 118(4):614–628, 2008. MR-2394845
- [13] David Nualart and Giovanni Peccati. Central limit theorems for sequences of multiple stochastic integrals. *Ann. Probab.*, 33(1):177–193, 2005. MR-2118863
- [14] Giovanni Peccati and Ciprian A. Tudor. Gaussian limits for vector-valued multiple stochastic integrals. In *Séminaire de Probabilités XXXVIII*, volume 1857 of *Lecture Notes in Math.*, pages 247–262. Springer, Berlin, 2005. MR-2126978

Electronic Journal of Probability

Electronic Communications in Probability

Advantages of publishing in EJP-ECP

- Very high standards
- Free for authors, free for readers
- Quick publication (no backlog)
- Secure publication (LOCKSS¹)
- Easy interface (EJMS²)

Economical model of EJP-ECP

- Non profit, sponsored by IMS³, BS⁴, ProjectEuclid⁵
- Purely electronic

Help keep the journal free and vigorous

- Donate to the IMS open access fund⁶ (click here to donate!)
- Submit your best articles to EJP-ECP
- Choose EJP-ECP over for-profit journals

¹LOCKSS: Lots of Copies Keep Stuff Safe <http://www.lockss.org/>

²EJMS: Electronic Journal Management System <http://www.vtex.lt/en/ejms.html>

³IMS: Institute of Mathematical Statistics <http://www.imstat.org/>

⁴BS: Bernoulli Society <http://www.bernoulli-society.org/>

⁵Project Euclid: <https://projecteuclid.org/>

⁶IMS Open Access Fund: <http://www.imstat.org/publications/open.htm>