

Percolation in the Boolean model with convex grains in high dimension

Jean-Baptiste Gouéré* Florestan Labéy†

Abstract

We investigate percolation in the Boolean model with convex grains in high dimension. For each dimension d , one fixes a compact, convex and symmetric set $K \subset \mathbb{R}^d$ with non empty interior. In a first setting, the Boolean model is a reunion of translates of K . In a second setting, the Boolean model is a reunion of translates of K or ρK for a further parameter $\rho \in (1, 2)$. We give the asymptotic behavior of the percolation probability and of the percolation threshold in the two settings.

Keywords: continuum percolation; Boolean model; high dimension.

MSC2020 subject classifications: 60K35; 60D05; 82B43.

Submitted to EJP on March 9, 2021, final version accepted on July 24, 2023.

Supersedes arXiv:2012.04370.

Supersedes HAL : hal-03044355.

1 Introduction and main results

1.1 Setting

Boolean model. For any $d \geq 1$, we denote by $\mathcal{K}(d)$ the set of subsets K of \mathbb{R}^d such that:

- The set K is compact, convex and symmetric (that is, for all $x \in K, -x \in K$).
- The Lebesgue measure of K satisfies $|K| = 1$.

Note that for all $K \in \mathcal{K}(d)$, the interior set $\text{Int}(K)$ is non empty. Otherwise K would be included in a proper affine subspace of \mathbb{R}^d and its Lebesgue measure would be 0.

Let $d \geq 1$ and $K \in \mathcal{K}(d)$. Let ν be a finite measure on $(0, +\infty)$ with positive mass. For simplicity we assume that the support of ν is bounded. Let $\lambda > 0$. Let ξ be a Poisson point process on $\mathbb{R}^d \times (0, +\infty)$ with intensity measure $dx \otimes \lambda\nu$. Consider

$$\Sigma = \Sigma(\lambda, \nu, d, K) = \bigcup_{(c,r) \in \xi} c + rK.$$

*IDP. Université de Tours. France. E-mail: jean-baptiste.gouere@univ-tours.fr

†IDP. Université de Tours. France. E-mail: florestan.labey@univ-tours.fr

This is the Boolean model with parameters λ, ν, d and K .

We call the sets $c + rK$ the *grains* of the Boolean model. We call c the *center* and r the *radius* of the grain $c + rK$.

We can write

$$\xi = \{(c, r(c)), c \in \chi\} \tag{1.1}$$

where χ is a Poisson point process on \mathbb{R}^d with intensity measure $\lambda\nu[(0, \infty)] dx$. Given χ , $(r(c))_{c \in \chi}$ is a family of i.i.d.r.v. with common distribution $\nu[(0, \infty)]^{-1}\nu$. As we shall not need this result, we do not give a more formal statement. We nevertheless think that it can provide some intuition.

We denote by $B(d)$ the Euclidean closed ball of \mathbb{R}^d centered at the origin and such that $|B(d)| = 1$. Note that $B(d)$ belongs to $\mathcal{K}(d)$. Most of the times, we will write simply B . We refer to $\Sigma(\lambda, \nu, d, B)$ as a Euclidean Boolean model. Note that, with our choice of terminology, B is a grain of radius 1 whereas this is not a Euclidean ball of radius 1. This should create no confusion as we will use the term “radius” only in the above sense.

We refer to the books by Kingman [8] and Last and Penrose [10] for background on Poisson point processes.

Percolation in the Boolean model. Fix $r > 0$. Recall (1.1) and set $\chi^0 = \chi \cup \{0\}$ and $r(0) = r$. We define an unoriented graph structure on χ^0 by putting an edge between x and y if $x + r(x)K$ touches $y + r(y)K$. As K is convex and symmetric,

$$x + r(x)K \text{ touches } y + r(y)K \iff x - y \in [r(x) + r(y)]K \iff y - x \in [r(x) + r(y)]K.$$

Note that (see (A.1) in Appendix A) for all $\lambda > 0$,

$$\begin{aligned} & \mathbb{P}[\text{the connected component of the graph } \chi^0 \text{ that contains } 0 \text{ is infinite}] \\ &= \mathbb{P}[\text{the connected component of } \Sigma \cup rK \text{ that contains } 0 \text{ is unbounded}]. \end{aligned} \tag{1.2}$$

We define the percolation threshold as usual (see Appendix A for details):

$$\begin{aligned} \lambda_c(\nu, d, K) &= \inf\{\lambda > 0 : \mathbb{P}[\text{the connected comp. of the graph } \chi^0 \text{ that contains } 0 \text{ is infinite}] > 0\} \end{aligned} \tag{1.3}$$

$$= \inf\{\lambda > 0 : \mathbb{P}[\text{the connected comp. of } \Sigma \cup rK \text{ that contains } 0 \text{ is unbounded}] > 0\} \tag{1.4}$$

$$= \inf\{\lambda > 0 : \mathbb{P}[\text{one of the connected comp. of } \Sigma \text{ is unbounded}] = 1\}. \tag{1.5}$$

In particular this does not depend on the choice of $r > 0$. We refer to the book by Meester and Roy [13] for background on percolation in the Boolean model.

As K is symmetric and as the interior set $\text{Int}(K)$ is non empty, there exists $r^- > 0$ such that $r^-B \subset K$. As K is compact, there exists $r^+ > 0$ such that $K \subset r^+B$. Therefore, by a natural coupling (and with an obvious generalization in our notations as $r^\pm B$ may not belong to $\mathcal{K}(d)$),

$$\Sigma(\lambda, \nu, d, r^-B) \subset \Sigma(\lambda, \nu, d, K) \subset \Sigma(\lambda, \nu, d, r^+B).$$

By standard results on percolation in the Euclidean Boolean model, we deduce that λ_c is always positive and finite. Actually, we will provide lower and upper bounds on λ_c with independent arguments.

Scale invariant thresholds. As in [6], we define ¹

$$c_c(\nu, d, K) = 1 - \exp\left(-\lambda_c(\nu, d, K) \int_{(0,+\infty)} r^d \nu(dr)\right)$$

and

$$\tilde{\lambda}_c(\nu, d, K) = -2^d \ln(1 - c_c(\nu, d, K)) = \lambda_c(\nu, d, K) \int_{(0,+\infty)} (2r)^d \nu(dr).$$

The quantity $c_c(\nu, d, K)$ is the probability that a given point (say 0) belongs to the critical Boolean model $\Sigma(\lambda_c, \nu, d, K)$. By ergodicity, this is also the density of the critical Boolean model. It has therefore a clear geometrical meaning. It is moreover invariant by scaling: if all radii are multiplied by a constant, λ_c changes but c_c remains unchanged. For these reasons c_c – and the related quantity $\tilde{\lambda}_c$ – are more convenient when comparing the threshold for different measures ν . We refer to [6] for a more detailed discussion on these topics.

1.2 Known results in the Euclidean setting

1.2.1 The case of a constant radius

Framework. We are interested here in the case $\nu = \delta_{1/2}$ and $K = B$. In other words, each grain is a translate of $\frac{1}{2}B$. For $\lambda > 0$ and $d \geq 1$, let

$$C^0 = C^0(\lambda, \delta_{1/2}, d, B)$$

be the connected component of

$$\Sigma(\lambda, \delta_{1/2}, d, B) \cup \frac{1}{2}B$$

that contains the origin. As mentioned in (1.4),

$$\lambda_c(\delta_{1/2}, d, B) = \inf\{\lambda > 0 : \mathbb{P}[C^0(\lambda, \delta_{1/2}, d, B) \text{ is unbounded}] > 0\}.$$

Link with a Galton-Watson process. In order to provide some intuition, let us describe the link between C^0 and some Galton-Watson process. A more precise description is given later.

Fix $\lambda > 0$ and $d \geq 1$. Denote by N_d the number of balls that belong to C^0 . The set C^0 is unbounded if and only if N_d is infinite. Call $\frac{1}{2}B$ the ball of generation 0. Define the balls of generation 1 as the random balls of Σ that touch $\frac{1}{2}B$. Define the balls of generation 2 as the random balls of Σ that touch at least one ball of generation 1 without being a ball of generation 0 or 1 and so on.

Note that $x + \frac{1}{2}B$ touches $x' + \frac{1}{2}B$ if and only if $x' \in x + B$. Therefore the number of random balls that intersect a given deterministic ball $x + \frac{1}{2}B$ is a Poisson random variable with parameter λ . Indeed, this is the number of points of a Poisson point process on \mathbb{R}^d with intensity λdx that belongs to $x + B$ and moreover $|B| = 1$. If there were no geometrical interference between children (a ill defined term) of different balls, then N_d would be the total population Z_d of a Galton-Watson process with offspring distribution $\text{Poisson}(\lambda)$. See Section 3.2.1 for details. Taking into account interactions, one can see that N_d is actually stochastically dominated by Z_d .

Let $S(\lambda)$ denote the survival probability of a Galton-Watson process with $\text{Poisson}(\lambda)$ offspring. Turning the above intuition into a proof, one can easily prove

$$\mathbb{P}[C^0(\lambda, \delta_{1/2}, d, B) \text{ is unbounded}] \leq S(\lambda)$$

¹In [6] there is an extra factor v_d which is the volume of the unit Euclidean ball. This is due to the fact that, in [6], the unit grain is the unit Euclidean ball whereas here the unit grain is a set K of volume 1.

and therefore

$$\lambda_c(\delta_{1/2}, d, B) \geq 1.$$

Result. In [14], Penrose proves that these bounds are asymptotically sharp when d tends to ∞ .

Theorem 1.1 ([14]). *For any $\lambda > 0$,*

$$\lim_{d \rightarrow \infty} \mathbb{P}[C^0(\lambda, \delta_{1/2}, d, B) \text{ is unbounded}] = S(\lambda).$$

Moreover,

$$\lim_{d \rightarrow \infty} \lambda_c(\delta_{1/2}, d, B) = 1.$$

The idea is to prove that, when d tends to infinity, the geometrical interference vanishes and the genealogy of the balls becomes closer and closer to the genealogy of the Galton-Watson process.

1.2.2 The case of radii that can take two values 1 and ρ with $1 < \rho < 2$

Framework. Let $\rho > 1$ and $d \geq 1$. We consider here the case

$$\nu = \nu_{d,\rho} = \delta_{1/2} + \frac{1}{\rho^d} \delta_{\rho/2} \tag{1.6}$$

and $K = B$. We refer to [6] for the motivation of the choice of $\nu_{d,\rho}$. Basically, the idea is to keep constant in d the relative influence of grains of different radii.

For all $\lambda > 0$, let

$$C^0 = C^0(\lambda, \nu_{d,\rho}, d, B)$$

be the connected component of

$$\Sigma(\lambda, \nu_{d,\rho}, d, B) \cup \frac{\rho}{2}B$$

that contains the origin. As before, we are interested in the percolation probability, that is the probability that C^0 is unbounded, and the percolation threshold λ_c .

Link with a two-type Galton-Watson process. As in the constant radius case, there is a natural genealogy for the balls of C^0 and this genealogical structure is stochastically bounded from above by a Galton-Watson process. In this setting, the Galton-Watson is a two-type Galton-Watson process: there are $\frac{1}{2}B$ grains and $\frac{\rho}{2}B$ grains. One can check easily that the probability that C^0 is unbounded is bounded from above by the survival probability of this Galton-Watson process. This gives a lower bound on $\lambda_c(\nu_{d,\rho}, d, B)$ which is equivalent to $\kappa_c(\rho)^d$, when d tends to ∞ , where

$$\kappa_c(\rho) = \frac{2\sqrt{\rho}}{1+\rho} < 1. \tag{1.7}$$

Result. In [6], Gou  r   and Marchand prove that if $1 < \rho < 2$, when d tends to infinity, geometrical interference vanishes and the behavior of the percolation threshold is roughly given by the behavior of the critical parameter of the two-type Galton-Watson process.

Theorem 1.2 ([6]). *For all $\rho \in (1, 2)$,*

$$\lim_{d \rightarrow \infty} \frac{1}{d} \ln(\lambda_c(\nu_{d,\rho}, d, B)) = \ln(\kappa_c(\rho)). \tag{1.8}$$

Note $\tilde{\lambda}_c(\delta_{1/2}, d, B) = \lambda_c(\delta_{1/2}, d, B)$ and $\tilde{\lambda}_c(\nu_{d,\rho}, d, B) = 2\lambda_c(\nu_{d,\rho}, d, B)$. Therefore, by Theorems 1.1 and 1.2, for d large enough,

$$\tilde{\lambda}_c(\nu_{d,\rho}, d, B) < \tilde{\lambda}_c(\delta_{1/2}, d, B)$$

and then

$$c_c(\nu_{d,\rho}, d, B) < c_c(\delta_{1/2}, d, B).$$

This result, which refutes a conjecture based on numerical estimations in low dimension and some heuristics in any dimension, was one of the motivations of [6]. We refer to [6] for more details.

1.2.3 The case of radii that can take two values 1 and ρ with $\rho > 2$: geometry still plays a role in high dimension

In [7], Gou er e and Marchand prove that (1.8) does not hold when $\rho > 2$. In that case, when d tends to infinity, geometrical interference does not vanish and the behavior of the percolation threshold is given by a competition between geometrical effects (dependencies due to the lack of space in \mathbb{R}^d) and genealogical aspects (given by the associated Galton-Watson process).

To sum up, geometrical interference does not always vanish in high dimension and geometry can still play a role. This is one of the motivations of this work where we investigate percolation in the Boolean model in high dimensions beyond the Euclidean case.

1.3 Main results

1.3.1 A further notation

Our main results deal with quantities $f(d, K) \in \mathbb{R}$ defined for all $d \geq 1$ and $K \in \mathcal{K}(d)$. For $\ell \in \mathbb{R}$, we write

$$\lim_{d \rightarrow \infty, K \in \mathcal{K}(d)} f(d, K) = \ell \quad \text{unif. in } K \in \mathcal{K}(d)$$

when the following holds

$$\forall \varepsilon > 0, \exists d_0 \geq 1, \forall d \geq d_0, \forall K \in \mathcal{K}(d), |f(d, K) - \ell| \leq \varepsilon.$$

1.3.2 The case of a constant radius

We consider the case $\nu = \delta_{1/2}$. For $d \geq 1$, $K \in \mathcal{K}(d)$ and $\lambda > 0$, let

$$C^0 = C^0(\lambda, \delta_{1/2}, d, K)$$

be the connected component of

$$\Sigma(\lambda, \delta_{1/2}, d, K) \cup \frac{1}{2}K$$

that contains the origin. As mentioned in (1.4),

$$\lambda_c(\delta_{1/2}, d, K) = \inf\{\lambda > 0 : \mathbb{P}[C^0(\lambda, \delta_{1/2}, d, K) \text{ is unbounded}] > 0\}.$$

Exactly as in the Euclidean case, one can check that the probability that $C^0(\lambda, \delta_{1/2}, d, K)$ is unbounded is bounded from above by $S(\lambda)$, the survival probability of a Galton-Watson process with progeny $\text{Poisson}(\lambda)$. This yields $\lambda_c(\delta_{1/2}, d, K) \geq 1$. This is stated as Proposition 3.1.

Our first main result is a generalization of Theorem 1.1.

Theorem 1.3. • For any $\lambda > 0$,

$$\lim_{d \rightarrow \infty, K \in \mathcal{K}(d)} \mathbb{P}[C^0(\lambda, \delta_{1/2}, d, K) \text{ is unbounded}] = S(\lambda) \quad \text{unif. in } K \in \mathcal{K}(d).$$

• Moreover,

$$\lim_{d \rightarrow \infty, K \in \mathcal{K}(d)} \lambda_c(\delta_{1/2}, d, K) = 1 \quad \text{unif. in } K \in \mathcal{K}(d).$$

1.3.3 The case of radii that can take two values 1 and ρ with $1 < \rho < 2$

Framework. Let $\rho > 1$ and $d \geq 1$. We are interested in the case $\nu = \nu_{d,\rho}$ defined in (1.6). With $\beta > 0$ we associate the intensity

$$\lambda = \beta \kappa_c(\rho)^d$$

where $\kappa_c(\rho)$ is defined in (1.7). This choice is natural because of Theorem 1.2. Let

$$C^0 = C^0(\beta \kappa_c(\rho)^d, \nu_{d,\rho}, d, K)$$

be the connected component of

$$\Sigma(\beta \kappa_c(\rho)^d, \nu_{d,\rho}, d, K) \cup \frac{\rho}{2}K$$

that contains the origin. As before, we are interested in the percolation probability, that is the probability that C^0 is unbounded, and the percolation threshold

$$\beta_c(\rho, d, K) = \inf \{ \beta > 0 : \mathbb{P} [C^0(\beta \kappa_c(\rho)^d, \nu_{d,\rho}, d, K) \text{ is unbounded}] > 0 \}.$$

Link with a two-type Galton-Watson process. As explained above, there is a natural genealogy for the balls of C^0 . This genealogical structure is bounded from above by a Galton-Watson process. In this setting, the Galton-Watson is a two-type Galton-Watson process: there are $\frac{1}{2}K$ grains and $\frac{\rho}{2}K$ grains. We will call them 1-particles and ρ -particles. Let

$$M = M(\beta, \rho, d) = \beta \begin{pmatrix} \left(\frac{2\sqrt{\rho}}{1+\rho}\right)^d & \left(\frac{1}{\sqrt{\rho}}\right)^d \\ (\sqrt{\rho})^d & \left(\frac{2\sqrt{\rho}}{1+\rho}\right)^d \end{pmatrix}. \tag{1.9}$$

This is the mean matrix of the Galton-Watson process. Denote by N_1 the random number of children of a 1-particle which are 1-particles. Denote by N_ρ the random number of children of a 1-particle which are ρ -particles. Then N_1 and N_ρ are independent Poisson random variables with parameters given (in that order) by the first row of M . The children of a ρ -particle behave in a similar way described by the second row of M . Note that the Galton-Watson process does not depend on K . However, its link with the cluster of the origin depends on K . The Galton-Watson process starts with one ρ -particle. Denote by $S(\beta, \rho, d)$ its survival probability. As before, one can easily check that the probability that C^0 is unbounded is bounded from above by $S(\beta, \rho, d)$. This gives a lower bound on $\beta_c(\rho, d, K)$. This is formalized in Proposition 4.3.

Result. In our second main result we prove that, if $1 < \rho < 2$, the above mentioned inequalities are asymptotically sharp when d tends to ∞ . As above, $S(\beta^2)$ denotes the survival probability of Galton-Watson process with Poisson(β^2) progeny. The appearance of $S(\beta^2)$ is due to the fact that $S(\beta, \rho, d)$ converges to $S(\beta^2)$ when d tends to ∞ . See Lemma 4.1.

Theorem 1.4. • Let $\beta > 0$ and $\rho \in (1, 2)$. Then

$$\lim_{d \rightarrow \infty, K \in \mathcal{K}(d)} \mathbb{P} [C^0(\beta \kappa_c(\rho)^d, \nu_{d,\rho}, d, K) \text{ is unbounded}] = S(\beta^2) \text{ unif. in } K \in \mathcal{K}(d).$$

• Let $\rho \in (1, 2)$.

$$\lim_{d \rightarrow \infty, K \in \mathcal{K}(d)} \beta_c(\rho, d, K) = 1 \text{ unif. in } K \in \mathcal{K}(d).$$

The second item can be rephrased as follows:

$$\lambda_c(\rho, d, K) \sim \kappa_c(\rho)^d \text{ as } d \rightarrow \infty.$$

This is a strengthening and a generalization of Theorem 1.2 which only provides a logarithmic equivalent of $\lambda_c(\rho, d, B)$. The first item is new. Theorem 1.4 is the most delicate result of the article.

1.3.4 Further comments

Two-radii case with $\rho > 2$. There is no hope to generalize Theorem 1.4 in this form for any ρ . Indeed, by [7], we know that the behavior of the critical threshold of percolation is not given by the critical threshold of the associated Galton-Watson process when $\rho > 2$ and $K = B$. However, for K belonging to some families of convex sets (for example ℓ^p balls), the result of Theorem 1.4 could hold for ρ in a larger interval.

About the proofs. In previous works in the Euclidean framework, the proofs rely on the isotropy of the model and on explicit estimates of volumes of intersections of balls. These tools are no longer available in the convex case and therefore we had to develop more robust strategies. These strategies provide a deeper understanding of the interference phenomena and allow to obtain more accurate results in the case where the radii take two values 1 and $\rho \in (1, 2)$ (compare Theorem 1.2 and Theorem 1.4).

Moreover our proofs suggest a couple of conjectures. They suggest that it is in the Euclidean case that the link between the Galton-Watson process and the cluster of the origin is the weakest. This is therefore in the Euclidean case that the percolation threshold could be the highest. The proofs may provide some tools our ideas toward a proof of this conjecture. We give more details in Section 3.2.2. We also state a related conjecture in Section 2.6.

Describing the plans of the proofs needs some tools and notations. Therefore we postpone the presentation of the proofs (and their comparison with proofs in the Euclidean setting) to Sections 3.4.4 and 4.5.4. However we tried to provide a pointer to the definition of each object used in these descriptions. Therefore the interested and hurried reader may try to jump directly to Sections 3.4.4 and 4.5.4.

Organization of the paper. In Section 2 we gather some notations and some results from analysis and high dimension geometry. In Section 3 we prove Theorem 1.3. In Section 4 we prove Theorem 1.4. In both cases, the difficult part is to establish the lower bound on percolation probabilities. The plans of the proofs are given in Section 3.4.4 and 4.5.4 once the objects are defined.

2 Some tools and notations

2.1 A couple of notations for random variables

When $d \geq 1$ and $K \in \mathcal{K}(d)$ are given, we shall denote by X_K, X'_K, X''_K independent random variables with uniform distribution on K . In the whole of this paper, \mathcal{N} will denote a standard Gaussian random vector in \mathbb{R}^2 .

2.2 Log-concavity

In this section, we gather some well known facts about log-concave functions. A map $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ is log-concave if, for all $x, y \in \mathbb{R}^d$ and all $\lambda \in (0, 1)$, the following inequality holds:

$$f(\lambda x + (1 - \lambda)y) \geq f(x)^\lambda f(y)^{1-\lambda}.$$

For example, the indicator of a convex set is log-concave.

If $g, h : \mathbb{R}^d \rightarrow \mathbb{R}_+$ are log-concave and measurable and if their convolution is defined everywhere, then $g * h$ is log-concave. This is a consequence of Prékopa-Leindler inequality (see for example [5] or [15]).

In particular, for all $d \geq 1$ and $K \in \mathcal{K}(d)$, X_K and $X_K + X'_K$ (see Section 2.1 for notations) have a log-concave density.

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ be log-concave. Let us also assume that f is symmetric, that is, for all $x \in \mathbb{R}^d$, $f(x) = f(-x)$. Then, for all $x \in \mathbb{R}^d$, $f(x) \leq f(0)$. Indeed, for all $x \in \mathbb{R}^d$,

$$\begin{aligned} f(0) &\geq f(x)^{1/2} f(-x)^{1/2} \text{ by log-concavity} \\ &= f(x) \text{ by symmetry.} \end{aligned}$$

2.3 A central limit theorem for random variables with log-concave density

The total variation distance between two random variables X and Y with values in \mathbb{R}^2 is

$$d_{TV}(X, Y) = \sup_{A \in \mathcal{B}(\mathbb{R}^2)} |\mathbb{P}[X \in A] - \mathbb{P}[Y \in A]|.$$

Recall (see for example Theorem 5.2 in the book by Lindvall [12]) that there exists (X', Y') such that X' has the same distribution as X , Y' has the same distribution as Y and

$$d_{TV}(X, Y) = \mathbb{P}[X' \neq Y'].$$

The following result is a rewriting of a weak version of Theorem 1.3 in [9].

Theorem 2.1. *There exists a sequence $(\varepsilon_{CLT}(d))_d$ which tends to 0 such that, for any centered random variable X in \mathbb{R}^d with log-concave density, there exists a linear map $L : \mathbb{R}^d \rightarrow \mathbb{R}^2$ such that*

$$d_{TV}(L(X), \mathcal{N}) \leq \varepsilon_{CLT}(d).$$

We will apply this result to random variables X_K or $X_K + X'_K$ (see Section 2.1 for notations).

Proof. In [9], the result is stated for an isotropic random vector X with log-concave density. One says that X is isotropic if $\mathbb{E}(X) = 0$ and $\text{var}(X) = I_d$. The random variable X may not be isotropic. But as X has a density, $\text{var}(X)$ is positive definite. As moreover X is centered, there exists an invertible linear map $T : \mathbb{R}^d \rightarrow \mathbb{R}^d$ such that $T(X)$ is isotropic. Moreover $T(X)$ still has a log-concave density. Therefore Theorem 1.3 of [9] applies to $T(X)$ and provides, for any $\varepsilon > 0$ and any d large enough depending only on ε , an orthogonal projection π on a plane $P \subset \mathbb{R}^d$ such that $\pi(T(X))$ is close in total variation distance to a standard Gaussian random vector \mathcal{N}_P on P : $d_{TV}(\pi(T(X)), \mathcal{N}_P) \leq \varepsilon$. From this result, one gets Theorem 2.1 with $L = \varphi \circ \pi \circ T$ where φ is an isometry between P and \mathbb{R}^2 . Note that $\varphi(\mathcal{N}_P)$ has the same distribution as \mathcal{N} . \square

2.4 A concentration result for random variables with uniform distribution on convex sets

Let $d \geq 1$. One says that a random variable on \mathbb{R}^d is isotropic if $\mathbb{E}(X) = 0$ and $\text{var}(X) = I_d$. Let $K \in \mathcal{K}(d)$. Let X_K be a random variable with uniform distribution on

K . We will need the following definition:

$$T \text{ is adapted to } X_K \text{ if } T \text{ is an invertible linear map from } \mathbb{R}^d \text{ to } \mathbb{R}^d \text{ such that } T(X_K) \text{ is isotropic.} \tag{2.1}$$

There always exist such maps. The following result is a rewriting of a weak version of Theorem 1.4 in [9].

Theorem 2.2. *Let $\varepsilon > 0$. Let $\ell \geq 1$. There exists d_0 such that, for all $d \geq d_0$, all $K \in \mathcal{K}(d)$ and all T adapted to X_K ,*

$$\mathbb{P} \left[\left| \frac{\|T(X_K^1 + \dots + X_K^\ell)\|_2}{\sqrt{d}} - \sqrt{\ell} \right| \geq \varepsilon \right] \leq \varepsilon$$

where X_K^1, \dots, X_K^ℓ are independent random variables with uniform distribution on K .

Proof. By Theorem 1.4 in [9], there exist absolute constants $c, C > 0$ such that the following holds. For all $d \geq 1$, all isotropic random variable X on \mathbb{R}^d with a log-concave density and all $\eta \in [0, 1]$,

$$\mathbb{P} \left[\left| \frac{\|X\|_2}{\sqrt{d}} - 1 \right| \geq \eta \right] \leq Cd^{-c\eta^2}.$$

Let $d \geq 1, \ell \geq 1, K \in \mathcal{K}(d)$ and T a map adapted to X_K . The random variable

$$X = \frac{T(X_K^1 + \dots + X_K^\ell)}{\sqrt{\ell}}$$

is isotropic. Furthermore, as K is convex, the common density of the X_K^i is log-concave. Therefore the common density of the $T(X_K^i)$ is log-concave. By stability by convolution (see Section 2.2), the density of $T(X_K^1 + \dots + X_K^\ell)$ is log-concave. Therefore, the density of X is log-concave.

Let $\varepsilon \in (0, \sqrt{\ell})$. We apply the result stated at the beginning of the proof with X defined as above and $\eta = \varepsilon/\sqrt{\ell}$. We get

$$\mathbb{P} \left[\left| \frac{\|T(X_K^1 + \dots + X_K^\ell)\|_2}{\sqrt{d}\sqrt{\ell}} - 1 \right| \geq \frac{\varepsilon}{\sqrt{\ell}} \right] \leq Cd^{-c\varepsilon^2\ell^{-1}}.$$

The result follows. □

2.5 Rearrangement inequalities

We will need the following version of Riesz’s rearrangement inequality. This is Theorem 3.7 in [11] in the simple setting of indicator function. When A is a Borel subset of \mathbb{R}^d with finite Lebesgue measure, we denote by A^* the Euclidean ball centered at the origin such that $|A| = |A^*|$.

Theorem 2.3. *Let $d \geq 1$. Let A_1, A_2, A_3 be Borel subsets of \mathbb{R}^d with finite Lebesgue measure. Then*

$$\int_{\mathbb{R}^d} \mathbb{1}_{A_1} * \mathbb{1}_{A_2}(x) \mathbb{1}_{A_3}(x) \, dx \leq \int_{\mathbb{R}^d} \mathbb{1}_{A_1^*} * \mathbb{1}_{A_2^*}(x) \mathbb{1}_{A_3^*}(x) \, dx.$$

2.6 A conjecture

This section is not necessary for the main results of the article. Let $d \geq 1$ and $K \in \mathcal{K}(d)$. Let $\|\cdot\|_K$ be the norm defined by $\|x\|_K = \inf\{r > 0 : x \in rK\}$. In particular

$\|\cdot\|_B = c_d \|\cdot\|_2$ for some constant c_d depending on the dimension d . For any $r > 0$, by Theorem 2.3,

$$\begin{aligned} \mathbb{P}[\|X_K + X'_K\|_K \leq r] &= \int_{\mathbb{R}^d} \mathbf{1}_K * \mathbf{1}_K(s) \mathbf{1}_{rK}(s) \, ds \\ &\leq \int_{\mathbb{R}^d} \mathbf{1}_B * \mathbf{1}_B(s) \mathbf{1}_{rB}(s) \, ds \\ &= \mathbb{P}[\|X_B + X'_B\|_B \leq r]. \end{aligned}$$

In other words, $\|X_B + X'_B\|_B$ is stochastically dominated by $\|X_K + X'_K\|_K$. Similar results hold for the sum of more copies of X_B or X_K . Numerical simulations suggest the following related conjecture. For any $p \in [1, +\infty]$ we denote by $\|\cdot\|_p$ the usual ℓ^p norm on \mathbb{R}^d and by $B_p \in \mathcal{K}(d)$ the associated ball of volume 1 centered at 0. In particular, $B_2 = B$.

Conjecture 2.4. For any $r > 0$, the map from $[1, +\infty]$ to $[0, 1]$ defined by

$$p \mapsto \mathbb{P} \left[\|X_{B_p} + X'_{B_p}\|_{B_p} \leq r \right]$$

is increasing on $[1, 2]$ and decreasing on $[2, +\infty]$.

Note that $\|X_{B_p} + X'_{B_p}\|_{B_p}$ has the same law as $\|X_p + X'_p\|_p$ where X_p and X'_p are two independent random variables with uniform distribution on the unit ball for the norm $\|\cdot\|_p$. When d tends to ∞ , $\|X_{B_p} + X'_{B_p}\|_{B_p}$ converges in probability to a constant $N(p)$. Using the representation of uniform random variables on B_p given in [3] one easily gets, for $p \in [1, \infty)$,

$$N(p) = \left[\frac{p}{4\Gamma^2(1+p^{-1})} \int_{\mathbb{R}^d \times \mathbb{R}^d} |x+y|^p e^{-|x|^p - |y|^p} \, dx \, dy \right]^{\frac{1}{p}}$$

where Γ is the Gamma function. One also easily checks that $N(\infty) = 2$. Here is a related easy looking conjecture: N is decreasing on $[1, 2]$ and increasing on $[2, \infty]$. We have not been able to prove this result. However, as a consequence of the above discussion, one can check that one always has $N(p) \geq N(2) = \sqrt{2}$.

We formulate similar conjecture for the sum of a higher number of copies of X_{B_p} .

2.7 Sum of independent random variables uniformly distributed on an Euclidean ball

We state and prove here a well known result for which we have no ready reference. Recall that $B(d)$ denotes the Euclidean closed ball of \mathbb{R}^d centered at the origin such that $|B(d)| = 1$.

Lemma 2.5. For all $\varepsilon > 0$, $\mathbb{P}[X_{B(d)} + X'_{B(d)} \in (\sqrt{2} - \varepsilon)B(d)] \rightarrow 0$ as $d \rightarrow \infty$.

Proof. Let $X(d)$ and $X'(d)$ be independent random variables with uniform distribution on the unit ball $\tilde{B}(d)$ of \mathbb{R}^d . We will prove that $\|X(d) + X'(d)\|_2$ tends to $\sqrt{2}$ in probability when d tends to ∞ . By a scaling argument, this yields the required result.

By independence and isotropy,

$$\mathbb{E}[|(X(d), X'(d))|] = \mathbb{E}[|(X(d), \|X'(d)\|_2 e_1)|] \leq \mathbb{E}[|(X(d), e_1)|] \tag{2.2}$$

where (\cdot, \cdot) denotes the scalar product in \mathbb{R}^d and e_1 the first vector of the canonical basis. But

$$(X(d), e_1)^2 + \dots + (X(d), e_d)^2 = \|X(d)\|_2^2 \leq 1.$$

Therefore, by symmetry, $\mathbb{E}[(X(d), e_1)^2] \leq d^{-1} \rightarrow 0$ and then $\mathbb{E}[|(X(d), e_1)|] \rightarrow 0$. By (2.2) and Markov inequality, we deduce from the above limit that $(X(d), X'(d))$ tends to 0 in probability. As moreover $\|X(d)\|_2^2$ and $\|X'(d)\|_2^2$ tend to 1 in probability, we get that $\|X(d) + X'(d)\|_2^2$ tends to 2 in probability as d tends to ∞ . \square

2.8 Notations for branching random walks (BRW)

Let $\lambda > 0$ and $d \geq 1$. Let Δ be a random variable on \mathbb{R}^d . Let S be a finite subset of \mathbb{R}^d . Let k be an integer. Let $M > 0$.

- τ^λ denotes a Galton-Watson tree with $\text{Poisson}(\lambda)$ progeny. The generation $|x|$ of a particle x of τ^λ is the graph distance from x to the root. If x is not the root we denote by \bar{x} the parent of x , that is the unique particle of generation $|x| - 1$ on the shortest path from the root to x .
- We see a BRW as a random tree (or forest) where each node x is called a particle and possesses a location $V(x) \in \mathbb{R}^d$. The notation $\bar{\tau}^{\lambda,d,\Delta;S}$ denotes a BRW such that:
 - The process starts with one particle located at each point of S .
 - Each particle x has $N(x)$ children located at

$$V(x) + \Delta(x, 1), \dots, V(x) + \Delta(x, N(x))$$

where the distribution of $N(x)$ is $\text{Poisson}(\lambda)$, where the $\Delta(x, \cdot)$ have the same distribution as Δ and where all variables $N(\cdot)$ and $\Delta(\cdot, \cdot)$ are independent.

If A is a subset of \mathbb{R}^d , we write

$$\bar{\tau}^{\lambda,d,\Delta;S}(A)$$

as a short notation for

$$\sum_{x \in \bar{\tau}^{\lambda,d,\Delta;S}} \mathbb{1}_A(V(x)),$$

that is the number of particles of the BRW located in A .

- $\bar{\tau}^{\lambda,d,\Delta}$ is a short notation for $\bar{\tau}^{\lambda,d,\Delta;\{0\}}$.
- If $\bar{\tau}$ is a BRW, then $\bar{\tau}_{\leq k}$ denotes the restriction of $\bar{\tau}$ to the k first generations and $\bar{\tau}_k$ denotes its restriction to the k -th generation.

Let moreover $\beta > 0$ and $\rho > 1$. In Section 4, we will use a BRW which will alternate ρ -particles for even generations and 1-particles for odd generations.

- $\bar{\tau}^{\beta,\rho,d,\Delta;S}$ denotes a two-types BRW. There are 1-particles and ρ -particles. It starts with one ρ -particle located at each point of S .
 - Each 1-particle x has $N(x)$ children which are ρ -particles located at

$$V(x) + \Delta(x, 1), \dots, V(x) + \Delta(x, N(x))$$

where the distribution of $N(x)$ is $\text{Poisson}(\beta\sqrt{\rho}^{-d})$, where the $\Delta(x, \cdot)$ have the same distribution as Δ and where all variables $N(\cdot)$ and $\Delta(\cdot, \cdot)$ are independent.

- Each ρ -particle x has $N(x)$ children which are 1-particles located at

$$V(x) + \Delta(x, 1), \dots, V(x) + \Delta(x, N(x))$$

where the distribution of $N(x)$ is $\text{Poisson}(\beta\sqrt{\rho}^d)$, where the $\Delta(x, \cdot)$ have the same distribution as Δ and where all variables $N(\cdot)$ and $\Delta(\cdot, \cdot)$ are independent.

- As above we will omit S when $S = \{0\}$.

- We define the event

$$\text{Small}_\rho \left(\bar{\tau}_{\leq k}^{\beta, \rho, d, \Delta; S}, M \right) = \{ \text{the number of } \rho \text{ particles of } \bar{\tau}_{\leq k}^{\beta, \rho, d, \Delta; S} \text{ is at most } M \\ \text{and the number of children of any } \rho \text{ particle is at most } M\sqrt{\rho^d} \}.$$

We will need some ordering on the nodes of a tree. We can for example formalize trees using Neveu formalism. In this formalism, nodes are finite sequences of positive integers (the idea is that $(4, 2)$ is the second child of the fourth child of the root \emptyset). See for example Section 2.2 of [16]. We can then order the nodes by lexicographic order. We will refer to this order as Neveu order.

3 Proof of Theorem 1.3

3.1 Framework and \widehat{C}^0

Let $\lambda > 0$, $d \geq 1$ and $K \in \mathcal{K}(d)$. Let χ be a Poisson point process on \mathbb{R}^d with intensity measure λdx . Note that $\xi = \{(c, 1/2), c \in \chi\}$ is a Poisson point process on $\mathbb{R}^d \times (0, +\infty)$ with intensity measure $dx \otimes \lambda \delta_{1/2}$. Set $\chi^0 = \chi \cup \{0\}$. As in Section 1.1, we define an unoriented graph structure on χ^0 by putting an edge between $x, y \in \chi^0$ if $y - x \in K$. Let \widehat{C}^0 be the connected component of the graph χ^0 that contains 0. We are interested (see (1.2) with $r = 1/2$) in

$$\mathbb{P}[\#\widehat{C}^0 = \infty]$$

and in

$$\lambda_c(\delta_{1/2}, d, K) = \inf\{\lambda > 0 : \mathbb{P}[\#\widehat{C}^0 = \infty] > 0\}.$$

3.2 Branching random walk and cluster of the origin

The content of this section (with the exception of Section 3.2.2) is essentially contained in Section 5 of [14]. Roughly, the aim is to explain that \widehat{C}^0 , or a subset of \widehat{C}^0 , can be seen as the set of positions of a pruned BRW. The framework is the same as in Section 3.1.

3.2.1 Basic construction

Exploring a subset of \widehat{C}^0 . We explore some part of \widehat{C}^0 by revealing successively parts of the point process χ . We will define inductively a tree with root 0 and where each node is a point of \widehat{C}^0 . We will call the graph distance from a node x to the root 0 the generation of x . Start with $\mathbb{A} = \{0\}$ and $\mathbb{B} = \emptyset$. At each stage of the algorithm, perform the following steps.

1. Select one of the elements of \mathbb{A} according to any given rule and call it x .
2. Add each point of

$$\chi \cap ((x + K) \setminus (\mathbb{B} + K))$$

to \mathbb{A} where

$$\mathbb{B} + K = \bigcup_{b \in \mathbb{B}} b + K.$$

Put an arrow from x to each of the points y we have just added to \mathbb{A} . We think about each such y as a child of x .

The idea is the following. Before performing Step 2:

- The points $y \in \chi$ such that there is an edge (“edge” refers to the graph structure defined on χ^0 in the introduction) between x and y are the points of $\chi \cap (x + K)$.

- All the points of $\chi \cap (\mathbb{B} + K)$ have been revealed. All of them are in $\mathbb{A} \cup \mathbb{B}$ and in \widehat{C}^0 .
 - None of the points of $\chi \setminus (\mathbb{B} + K)$ have been revealed.
3. Move x from \mathbb{A} to \mathbb{B} .
 4. If \mathbb{A} is non-empty, go back to Step 1.

There are two cases.

- The algorithm terminates. In that case it provides a tree whose set of nodes is \mathbb{B} . We have $\mathbb{B} = \widehat{C}^0$.
- The algorithm does not terminate. In that case it provides a sequence of growing trees. We can consider the limit tree. Denote its set of nodes by \mathbb{B} . We have $\mathbb{B} \subset \widehat{C}^0$ and $\#\mathbb{B} = \#\widehat{C}^0 = \infty$.

In all cases, we have

$$\mathbb{B} \subset \widehat{C}^0 \text{ and } \#\mathbb{B} = \#\widehat{C}^0.$$

Note that, when \widehat{C}^0 is infinite, it may happen that $\mathbb{B} \neq \widehat{C}^0$. This depends on the rule used to select x in Step 1. If we use the rule “Select one of the elements of \mathbb{A} of minimal generation according to any given rule” or the rule “Select the element of \mathbb{A} of minimal Euclidean distance to 0”, then $\mathbb{B} = \widehat{C}^0$. We shall not need this result and, on the contrary, we will find it convenient to have some freedom in the choice of the point x in Step 1.

Building a subset of \widehat{C}^0 by pruning a BRW. The key is the following. Before Step 2 in the above exploration process, conditioned to what we have revealed,

$$\chi \cap ((x + K) \setminus (\mathbb{B} + K))$$

is a Poisson point process of intensity

$$\lambda \mathbb{1}_{(x+K) \setminus (\mathbb{B}+K)}(y) \, dy.$$

Furthermore, generating such a Poisson point process can be done as follows.

- Generate N a random variable with Poisson(λ) distribution. Note that $\lambda = \lambda|x + K|$ as $|K| = 1$.
- Generate Y_1, \dots, Y_N i.i.d. random variables with uniform distribution on $x + K$.
- The random set $\{Y_i : Y_i \notin \mathbb{B} + K\}$ has the required distribution.

This enables us to build a set which, under a suitable coupling (which we will assume implicitly henceforth) is a subset of \widehat{C}^0 . Consider the BRW

$$\bar{\tau} = \bar{\tau}^{\lambda, d, X_K}$$

where X_K is uniformly distributed on K (see Section 2.8 for notations on BRW). Start with $\mathbb{A} = \{\varnothing\}$ where \varnothing is the root of $\bar{\tau}$ (which is located at 0) and $\mathbb{B} = \emptyset$. At each stage of the algorithm, perform the following steps.

1. Select one of the elements of \mathbb{A} according to any given rule and call it x .
2. Consider successively the children (in the BRW) y of x in any order and for each of them, do the following:

if there does not exist $x' \in \mathbb{B}$ such that $V(y) \in V(x') + K$, then add y to \mathbb{A} .

Let us introduce some vocabulary for future reference. We say that the other children of x are rejected

$$\text{because of } \textit{interference} \text{ between } x \text{ and } x' \text{ or because of } \textit{interference} \text{ with } x'. \tag{3.1}$$

We say that

$$V(x') + K \text{ is the } \textit{region of interference} \text{ of } x'. \tag{3.2}$$

In other words, we reject y because of interference with x' when y belongs to the region of interference of x' . A large part of our proof will be devoted to establish the fact that, in high dimension, there is not too much interference.

3. Move x from \mathbb{A} to \mathbb{B} .
4. If \mathbb{A} is non-empty, go back to Step 1.

If the algorithm terminates it provides a set \mathbb{B} . Otherwise, it provides an increasing sequence of \mathbb{B} (one \mathbb{B} for each stage) and we define \mathbb{B} as the union of all those \mathbb{B} at different stages. In any case,

$$\{V(x), x \in \mathbb{B}\} \subset \widehat{C}^0 \text{ and } \#\mathbb{B} = \#\widehat{C}^0.$$

Note that $\{V(x), x \in \mathbb{B}\}$ is the set of positions of a pruned version of $\bar{\tau}$.

3.2.2 An open question

This section is not necessary for the main result of the article. Let $d \geq 1$ and $K \in \mathcal{K}(d)$. By Theorem 2.3,

$$\mathbb{P}[X_K + X'_K \in K] = \int_{\mathbb{R}^d} \mathbb{1}_K * \mathbb{1}_K(s) \mathbb{1}_K(s) \, ds \leq \int_{\mathbb{R}^d} \mathbb{1}_B * \mathbb{1}_B(s) \mathbb{1}_B(s) \, ds = \mathbb{P}[X_B + X'_B \in B].$$

We now use the setting and notations of Section 3.2.1. To construct \widehat{C}^0 , we have to reject because of interference the grandchildren of the root of $\bar{\tau}^{\lambda, d, X_K}$ that belong to K . If we condition on the underlying tree τ^{λ, d, X_K} , the probability that a given grandchildren is rejected for this reason is $\mathbb{P}[X_K + X'_K \in K]$. As seen above, this probability is maximal when $K = B$. This remark and further similar considerations may suggest that the percolation probability could be minimal when $K = B$ and therefore the percolation threshold λ_c could be maximal when $K = B$. This would be coherent with numerical simulations in low dimension. In particular, by reducing percolation to a local criteria and then using Monte Carlo methods, Balister, Bollobás and Walters provided in [2] the following 99.99% confidence intervals for $\lambda_c(\delta_{1/2}, K, 2)$: $[4.508, 4.515]$ when $K = B$ and $[4.392, 4.398]$ when $K = [-1/2, 1/2]^2$.

3.2.3 Constructing a smaller set by over-pruning

We introduce here a variant of the previous constructions. The general idea is that, by rejecting more than necessary at some stage, we may be able to have a better control on interference at a later stage.

Exploring a smaller subset of \widehat{C}^0 . Start with $\mathbb{A} = \{0\}$ and $\mathbb{B} = \emptyset$. At each stage of the algorithm, perform the following steps.

1. Select one of the elements of \mathbb{A} according to any given rule and call it x .

- Let $K(x)$ be a subset of $x + K$ defined according to any given rule. It can depend on \mathbb{A}, \mathbb{B} and x . Consider successively the points of

$$\chi \cap \left(K(x) \setminus \bigcup_{b \in \mathbb{B}} K(b) \right)$$

in any order and add some of them to \mathbb{A} according to any given rule. Put an arrow from x to each of the points we have just added.

With the vocabulary introduced above, the region of interference of x is $K(x)$.

- Move x from \mathbb{A} to \mathbb{B} .
- If \mathbb{A} is non-empty, go back to Step 1.

We get in the end some set that we denote by \mathbb{B}^- . We have

$$\mathbb{B}^- \subset \widehat{C}^0 \text{ and } \#\mathbb{B}^- \leq \widehat{C}^0.$$

Building a smaller subset of \widehat{C}^0 by over-pruning a BRW. As in Section 3.2.1, consider the BRW $\bar{\tau} = \bar{\tau}^{\lambda, d, X_K}$ and start with $\mathbb{A} = \{\emptyset\}$ and $\mathbb{B} = \emptyset$. At each stage of the algorithm, perform the following steps.

- Select one of the elements of \mathbb{A} according to any given rule and call it x .
- Let $K(x)$ be a subset of $V(x) + K$ defined according to any given rule. We say that this is the *region of interference* of x . It can depend on \mathbb{A}, \mathbb{B} and x . Consider successively in any given order the children y of x whose position belongs to

$$K(x) \setminus \left(\bigcup_{b \in \mathbb{B}} K(b) \right) \tag{3.3}$$

and, for each of them, decide according to any given rule whether one adds it to \mathbb{A} or not.

We thus distinguish (somehow artificially) two kinds of over-pruning:

- By taking $K(x)$ smaller than $V(x) + K$ we may reduce the number of children of x in the pruned BRW. However, as we reduced the region of interference of x , we may reject less points at a later stage of the algorithm.
- Similarly, by adding to \mathbb{A} only some of the children of x which belongs to (3.3), we further reduce the number of children of x in the pruned BRW. However, we can gain some properties on the positions of the points of the pruned BRW which may help us controlling the interference at a later stage of the algorithm.

Each kind of over-pruning amounts to rejecting some of the children y of x belonging to

$$(V(x) + K) \setminus \bigcup_{b \in \mathbb{B}} K(b)$$

according to some given rule. In this work, we will describe over-pruning in that way, that is by rejecting some points according to some specific rules.

- Move x from \mathbb{A} to \mathbb{B} .
- If \mathbb{A} is non-empty, go back to Step 1.

If the algorithm terminates it provides a set \mathbb{B} . Otherwise, it provides an increasing sequence of \mathbb{B} (one \mathbb{B} for each stage) and we define \mathbb{B} as the union of all those \mathbb{B} at different stages. In any case,

$$\{V(x), x \in \mathbb{B}\} \subset \widehat{C}^0 \text{ and } \#\mathbb{B} \leq \#\widehat{C}^0.$$

The set $\{V(x), x \in \mathbb{B}\}$ is the set of positions of an over-pruned version of $\bar{\tau}$.

Intuitive rephrasing. Let us rephrase one of the key ideas at a more intuitive level. When we look for the children of a particle x , we reveal the relevant Poisson point process in a (subset of) the interference region $K(x)$. Therefore at any later stage of the construction we have to reject any particle which falls in $K(x)$. However, we do not have to care about any rejected particle or about any particle whose children we never consider: they generate no interference.

3.3 Proof of the upper bound on percolation probability

The aim is to prove the following proposition, which is the easy part of Theorem 1.3. We refer to Section 1.3.2 for notations.

Proposition 3.1.

- For all $\lambda > 0, d \geq 1$ and $K \in \mathcal{K}(d)$, $\mathbb{P}[C^0(\lambda, \delta_{1/2}, d, K) \text{ is unbounded}] \leq S(\lambda)$.
- For all $d \geq 1$ and $K \in \mathcal{K}(d)$, $\lambda_c(\delta_{1/2}, d, K) \geq 1$.

Proof. We use the framework of Section 3.1 and the BRW $\bar{\tau} = \bar{\tau}^{\lambda, d, X_K}$ of Section 3.2. By the discussion in Section 3.2.1, we know that $\#\widehat{C}^0$ has the same distribution as the total population of a pruned version of $\bar{\tau}$. Therefore $\#\widehat{C}^0$ is stochastically dominated by the total population of $\bar{\tau}$, that is by the population of a Galton-Watson tree τ^λ . Using (1.2) for the first equality, we thus get

$$\mathbb{P}[C^0(\lambda, \delta_{1/2}, d, K) \text{ is unbounded}] = \mathbb{P}[\#\widehat{C}^0(\lambda, \delta_{1/2}, d, K) = \infty] \leq \mathbb{P}[\#\tau^\lambda = \infty] = S(\lambda).$$

This is the first part of Proposition 3.1. The second part follows as $S(\lambda) = 0$ for $\lambda \leq 1$ and as $\lambda_c(\delta_{1/2}, d, K)$ is the infimum of all $\lambda > 0$ such that $\mathbb{P}[C^0(\lambda, \delta_{1/2}, d, K) \text{ is unbounded}]$ is positive. \square

3.4 Proof of the lower bound on percolation probability

We use the framework of Subsection 3.1. Our aim is to prove the following result.

Theorem 3.2. *Let $\lambda > 0$ and $\varepsilon > 0$. There exists $d_0 \geq 1$ such that, for all $d \geq d_0$ and all $K \in \mathcal{K}(d)$,*

$$\mathbb{P}[\#\widehat{C}^0(\lambda, \delta_{1/2}, d, K) = \infty] \geq S(\lambda) - \varepsilon.$$

Proof of Theorem 1.3 using Theorem 3.2. Let $\lambda > 0$ and $\varepsilon > 0$. Let d_0 be as given by Theorem 3.2. Let $d \geq d_0$ and $K \in \mathcal{K}(d)$. We then have,

$$S(\lambda) \geq \mathbb{P}[C^0(\lambda, \delta_{1/2}, d, K) \text{ is unbounded}] = \mathbb{P}[\#\widehat{C}^0(\lambda, \delta_{1/2}, d, K) = \infty] \geq S(\lambda) - \varepsilon.$$

The first inequality is Proposition 3.1. The equality is (1.2). The second inequality is due to our choice of d_0 . We have proved the first part of Theorem 1.3.

Let $\lambda > 1$. Then $S(\lambda) > 0$. Set $\varepsilon = S(\lambda)/2 > 0$. By the first part of Theorem 1.3, there exists d_0 such that, for all $d \geq d_0$ and $K \in \mathcal{K}(d)$,

$$\mathbb{P}[C^0(\lambda, \delta_{1/2}, d, K) \text{ is unbounded}] \geq S(\lambda) - \varepsilon = S(\lambda)/2 > 0.$$

Therefore, for all $d \geq d_0$ and all $K \in \mathcal{K}(d)$, $\lambda_c(\delta_{1/2}, d, K) \leq \lambda$. Combined with the second part of Proposition 3.1, this gives the second part of Theorem 1.3. \square

3.4.1 Good Gaps

Result. Let $d \geq 1$. For any $K \in \mathcal{K}(d)$, denote as usual by X_K, X'_K i.i.d.r.v. uniformly distributed on K . For all $\eta > 0$, set

$$G(d, K, \eta) = \{z \in \mathbb{R}^d : \mathbb{P}(z + X'_K \notin K) \geq 1 - \eta\}$$

where G stands for “Good Gap”. Note that G is symmetric because K is symmetric.

Lemma 3.3. *There exists a sequence $(\varepsilon_G(d))_d$ that tends to 0 such that for all $d \geq 1$, all $K \in \mathcal{K}(d)$, all $a \in \mathbb{R}^d$ and all $\eta > 0$:*

$$\mathbb{P}[a + X_K \notin G(d, K, \eta)] \leq \eta^{-1} \varepsilon_G(d).$$

Proof. Let $d \geq 1$. For all $K \in \mathcal{K}(d)$ and all $a \in \mathbb{R}^d$ we have, using rearrangement inequality,

$$\begin{aligned} \mathbb{P}(a + X_K + X'_K \in K) &= \int_{\mathbb{R}^d} \mathbb{1}_K * \mathbb{1}_K(x) \mathbb{1}_{K-a}(x) \, dx \\ &\leq \int_{\mathbb{R}^d} \mathbb{1}_B * \mathbb{1}_B(x) \mathbb{1}_B(x) \, dx \quad \text{by Theorem 2.3} \\ &= \mathbb{P}(X_B + X'_B \in B). \end{aligned}$$

So, for all $\eta > 0$,

$$\begin{aligned} \mathbb{P}[a + X_K \notin G(d, K, \eta)] &= \mathbb{P}[\mathbb{P}(a + X_K + X'_K \notin K | X_K) < 1 - \eta] \\ &= \mathbb{P}[\mathbb{P}(a + X_K + X'_K \in K | X_K) > \eta] \\ &\leq \eta^{-1} \mathbb{E}[\mathbb{P}(a + X_K + X'_K \in K | X_K)] \\ &= \eta^{-1} \mathbb{P}(a + X_K + X'_K \in K) \\ &\leq \eta^{-1} \mathbb{P}(X_B + X'_B \in B) \quad \text{by the above discussion.} \end{aligned}$$

But $\mathbb{P}(X_B + X'_B \in B) \rightarrow 0$ when d tends to ∞ . This is (up to a scaling) (21) of Lemma 3 in [14]. This is also a consequence of Lemma 2.5. This concludes the proof. \square

3.4.2 Embedding of a two-dimensional lattice in \mathbb{R}^d - oriented percolation

Embedding of a two-dimensional lattice in \mathbb{R}^d . Set $\mathcal{L} = \{(i, j) \in \mathbb{N} \times \mathbb{Z} : i + j \text{ odd } |j| < i\}$ and $\bar{\mathcal{L}} = \mathcal{L} \cup \{(0, 0)\}$. We see $\bar{\mathcal{L}}$ as an oriented graph by putting an edge from $(0, 0)$ to $(1, 0)$ and, for every $(i, j) \in \mathcal{L}$, one edge from (i, j) to $(i + 1, j + 1)$ and one from (i, j) to $(i + 1, j - 1)$. We consider on $\bar{\mathcal{L}}$ the lexicographical order. Thus, the first vertices of $\bar{\mathcal{L}}$ are $(0, 0), (1, 0), (2, -1), (2, 1), (3, -2), \dots$

When $d \geq 1$ and $K \in \mathcal{K}(d)$ are given, one fixes a linear map $L : \mathbb{R}^d \rightarrow \mathbb{R}^2$ given by Theorem 2.1 for X_K . With each $(i, j) \in \bar{\mathcal{L}}$ we associate the sets $A(i, j) \subset \mathbb{R}^2$ and $A_L(i, j) \subset \mathbb{R}^d$ defined by

$$A(i, j) = (i, j) + 4^{-1}D \text{ and } A_L(i, j) = L^{-1}(A(i, j))$$

where D the Euclidean unit ball of \mathbb{R}^2 (not to be confused with $B = B(d)$, which is the Euclidean ball of \mathbb{R}^d of volume 1). The sets $A_L(i, j)$ are pairwise disjoint. Moreover 0 belongs to $A_L(0, 0)$.

Using this embedding, we will compare the cluster of the origin to a supercritical percolation process on $\bar{\mathcal{L}}$.

Oriented percolation on \mathcal{L} . Let $\theta(u)$ be the probability that there exists an infinite open path originating from $(1, 0)$ in a Bernoulli site percolation on the graph \mathcal{L} with parameter u . We will need²:

$$\lim_{u \rightarrow 1} \theta(u) = 1. \tag{3.4}$$

We refer to [4] for background on oriented percolation.

3.4.3 An estimate about BRW

The aim of this section is to prove Lemma 3.5 and its consequence Lemma 3.4. Recall the notations from Section 2.8. Recall in particular that $S(\lambda)$ is defined as the survival probability of a Galton-Watson process with progeny $\text{Poisson}(\lambda)$.

Lemma 3.4. *Let $\lambda > 1$ and $\varepsilon > 0$. There exists $m, d_0, k, M \geq 1$ such that, for all $d \geq d_0$ and all centered random variable X in \mathbb{R}^d with log-concave density, the following properties hold where L is any map given by Theorem 2.1 for X .*

- For all $z \in A_L(0, 0)$,

$$\mathbb{P} \left[\bar{\tau}_k^{\lambda, d, X; \{z\}}(A_L(1, 0)) \geq m \text{ and } \bar{\tau}_{\leq k}^{\lambda, d, X; \{z\}}(\mathbb{R}^d) \leq M \right] \geq S(\lambda) - \varepsilon.$$

- For all $(i, j) \in \mathcal{L}$ and all subsets $\mathcal{S} \subset A_L(i, j)$ of cardinality m ,

$$\mathbb{P} \left[\bar{\tau}_k^{\lambda, d, X; \mathcal{S}}(A_L(i + 1, j + 1)) \geq m \text{ and } \bar{\tau}_k^{\lambda, d, X; \mathcal{S}}(A_L(i + 1, j - 1)) \geq m \text{ and } \bar{\tau}_{\leq k}^{\lambda, d, X; \mathcal{S}}(\mathbb{R}^d) \leq M \right] \geq 1 - \varepsilon.$$

Lemma 3.4 is a consequence of the following result (the proof is given below). In Section 3 we will prove and use another consequence of Lemma 3.5. Recall that \mathcal{N} denotes a standard Gaussian random vector in \mathbb{R}^2 .

Lemma 3.5. *Let $\lambda > 1$ and $\varepsilon > 0$. There exists $m, k, M \geq 1$ such that the following properties hold.*

- For all $z \in A(0, 0)$,

$$\mathbb{P} \left[\bar{\tau}_k^{\lambda, 2, \mathcal{N}; \{z\}}(A(1, 0)) \geq m \text{ and } \bar{\tau}_{\leq k}^{\lambda, 2, \mathcal{N}; \{z\}}(\mathbb{R}^2) \leq M \right] \geq S(\lambda) - \varepsilon.$$

- For all $(i, j) \in \mathcal{L}$ and all subset $\mathcal{S} \subset A(i, j)$ of cardinality m ,

$$\mathbb{P} \left[\bar{\tau}_k^{\lambda, 2, \mathcal{N}; \mathcal{S}}(A(i + 1, j + 1)) \geq m \text{ and } \bar{\tau}_k^{\lambda, 2, \mathcal{N}; \mathcal{S}}(A(i + 1, j - 1)) \geq m \text{ and } \bar{\tau}_{\leq k}^{\lambda, 2, \mathcal{N}; \mathcal{S}}(\mathbb{R}^2) \leq M \right] \geq 1 - \varepsilon.$$

Let us start by the following lemmas.

Lemma 3.6. *The map S is continuous on $[0, +\infty)$.*

Proof. For any $\lambda > 1$, $1 - S(\lambda)$ is the only real $u \in (0, 1)$ such that $u = \exp(\lambda(u - 1))$ which we write

$$\frac{\ln(u)}{u - 1} = \lambda.$$

But $u \mapsto \ln(u)/(u - 1)$ defines a decreasing homeomorphism f from $(0, 1)$ to $(1, +\infty)$ ³. As moreover S vanishes on $[0, 1]$, the result follows. \square

²By the first inequality of (1) of Section 10 of [4] with $N = 0$, one has (thanks to a contour argument): $1 - \theta(u) \leq \sum_{m \geq 4} 3^m (1 - u)^{m/4}$.

³This is for example a consequence of the following facts: $f(0+) = +\infty$; $f(1-) = 1$; for all $v \in (0, 1)$, $f(1 - v) = 1 + v/2 + v^2/3 + v^3/4 + \dots$

Lemma 3.7. *Let $\lambda > 1$ and $\alpha, \beta, \varepsilon > 0$. Let $m \geq 1$. There exists $k \geq 1$ such that, for all $x \in \alpha D$,*

$$\mathbb{P} \left[\bar{\tau}_k^{\lambda, 2, \mathcal{N}}(x + \beta D) \geq m \right] \geq S(\lambda) - \varepsilon.$$

Proof. Let us first define a few constants. Let $\lambda_1 \in]1, \lambda[$ be such that

$$S(\lambda_1) \geq S(\lambda) - \varepsilon.$$

Such a λ_1 exists by continuity of S , see Lemma 3.6. Let $\Lambda_1 > 0$ be such that

$$\lambda_1 := \lambda \mathbb{P}[\|\mathcal{N}\|_2 \leq \Lambda_1].$$

Let $C > 0$ be such that, for all $n \geq 1$ and all $z \in (\alpha + \Lambda_1 n)D$,

$$\mathbb{P} \left[\mathcal{N} \in \frac{1}{n}z + \frac{\beta}{n}D \right] \geq \frac{C}{n^2}. \tag{3.5}$$

Let us show the existence of $\eta > 0$ and $n_0 \geq 1$ such that,

$$\forall n \geq n_0, \mathbb{P} \left[\bar{\tau}_n^{\lambda, 2, \mathcal{N}}(n\Lambda_1 D) \geq \eta \lambda_1^n \right] \geq S(\lambda) - 3\varepsilon. \tag{3.6}$$

Denote by $\bar{\tau}^{\lambda, 2, \mathcal{N}, \Lambda_1}$ the BRW obtained from $\bar{\tau}^{\lambda, 2, \mathcal{N}}$ by pruning one particle and its progeny as soon as it makes a step whose Euclidean norm is larger than Λ_1 . The new BRW has then the same distribution as $\bar{\tau}^{\lambda_1, 2, \mathcal{N}_1}$ where \mathcal{N}_1 has the distribution of \mathcal{N} conditioned on $\|\mathcal{N}\|_2 \leq \Lambda_1$. We have (see for example [1] page 9)

$$\frac{\bar{\tau}_n^{\lambda_1, 2, \mathcal{N}_1}(\mathbb{R}^2)}{\lambda_1^n} \rightarrow W \text{ a.s.}$$

where W is a random variable which is positive on the event $\{\bar{\tau}^{\lambda_1, 2, \mathcal{N}_1} \text{ survives}\}$ whose probability is $S(\lambda_1)$. We can then chose $\eta > 0$ such that $\mathbb{P}[W \geq 2\eta] \geq S(\lambda_1) - \varepsilon$. We can now fix $n_0 \geq 1$ such that, for all $n \geq n_0$,

$$\mathbb{P} \left[\bar{\tau}_n^{\lambda_1, 2, \mathcal{N}_1}(\mathbb{R}^2) \geq \eta \lambda_1^n \right] \geq S(\lambda_1) - 2\varepsilon \geq S(\lambda) - 3\varepsilon$$

and then

$$\mathbb{P} \left[\bar{\tau}_n^{\lambda, 2, \mathcal{N}_1}(n\Lambda_1 D) \geq \eta \lambda_1^n \right] \geq S(\lambda) - 3\varepsilon.$$

One deduces (3.6).

Now, let us show

$$\forall n \geq 1, \forall x \in \alpha D, \forall y \in n\Lambda_1 D, \mathbb{P} \left[\bar{\tau}_{n^2}^{\lambda, 2, \mathcal{N}; \{y\}}(x + \beta D) \geq 1 \right] \geq \frac{S(\lambda)C}{n^2}. \tag{3.7}$$

Let $n \geq 1, x \in \alpha D$ and $y \in n\Lambda_1 D$. We have

$$\mathbb{P} \left[\bar{\tau}_{n^2}^{\lambda, 2, \mathcal{N}; \{y\}}(x + \beta D) \geq 1 \right] \geq S(\lambda) \mathbb{P} \left[y + \sum_{i=1}^{n^2} \mathcal{N}'_i \in x + \beta D \right]$$

where the \mathcal{N}'_i are independent copies of \mathcal{N} . To prove this, it is sufficient to consider, on the event $\{\bar{\tau}^{\lambda, 2, \mathcal{N}; \{y\}} \text{ survives}\}$, the position of a given particle of generation n^2 . As $\sum_{i=1}^{n^2} \mathcal{N}'_i$ has the same distribution as $n\mathcal{N}$, we deduce

$$\mathbb{P} \left[\bar{\tau}_{n^2}^{\lambda, 2, \mathcal{N}; \{y\}}(x + \beta D) \geq 1 \right] \geq S(\lambda) \mathbb{P} \left[\mathcal{N} \in \frac{1}{n}(x - y) + \frac{\beta}{n}D \right].$$

Thanks to (3.5) we deduce (3.7).

We now combine (3.6) and (3.7) and get, for all $x \in \alpha D$ and all $n \geq n_0$,

$$\mathbb{P} \left[\bar{\tau}_{n+n^2}^{\lambda, 2, \mathcal{N}}(x + \beta D) \geq m \right] \geq (\mathbf{S}(\lambda) - 3\varepsilon) \mathbb{P} \left[\text{binomial} \left(\lfloor \eta \lambda_1^n \rfloor, \frac{\mathbf{S}(\lambda)C}{n^2} \right) \geq m \right].$$

This can be proven by first conditioning with respect to the n first generations of the BRW, working on the event $\{\bar{\tau}_n^{\lambda, 2, \mathcal{N}}(n\Lambda_1 D) \geq \eta \lambda_1^n\}$ (whose probability is controlled by (3.6)) and using (3.7) with $\lfloor \eta \lambda_1^n \rfloor$ independent BRW originating from different positions of $\bar{\tau}_n^{\lambda, 2, \mathcal{N}}$ in $n\Lambda_1 D$. But for n large enough we have

$$\lfloor \eta \lambda_1^n \rfloor \frac{\mathbf{S}(\lambda)C}{n^2} - m \geq \frac{1}{2} \eta \lambda_1^n \frac{\mathbf{S}(\lambda)C}{n^2}$$

and then, by Chebyshev's inequality,

$$\mathbb{P} \left[\text{binomial} \left(\lfloor \eta \lambda_1^n \rfloor, \frac{\mathbf{S}(\lambda)C}{n^2} \right) \leq m \right] \leq \eta \lambda_1^n \left(\frac{1}{2} \eta \lambda_1^n \frac{\mathbf{S}(\lambda)C}{n^2} \right)^{-2} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

As a consequence, there exists $n \geq n_0$ such that, for all $x \in \alpha D$,

$$\mathbb{P} \left[\bar{\tau}_{n+n^2}^{\lambda, 2, \mathcal{N}}(x + \beta D) \geq m \right] \geq \mathbf{S}(\lambda) - 4\varepsilon.$$

We fix such a n . The lemma is proven with $k = n + n^2$. □

Proof of Lemma 3.5. Fix m such that

$$(1 - \mathbf{S}(\lambda) + \varepsilon)^m \leq \varepsilon.$$

This is possible if $\varepsilon > 0$ is small enough, which we can assume. We apply Lemma 3.7 with $\alpha = 3$ and $\beta = 1/4$. We get k such that, for all $x \in 3D$,

$$\mathbb{P} \left[\bar{\tau}_k^{\lambda, 2, \mathcal{N}}(x + 4^{-1}D) \geq m \right] \geq \mathbf{S}(\lambda) - \varepsilon.$$

By natural couplings between the involved BRW we get, for all $z \in \mathbb{R}^2$;

$$\begin{aligned} \bar{\tau}_k^{\lambda, 2, \mathcal{N}; \{z\}}(A(1, 0)) &= \bar{\tau}_k^{\lambda, 2, \mathcal{N}; \{z\}}((1, 0) + 4^{-1}D) \\ &= \bar{\tau}_k^{\lambda, 2, \mathcal{N}}((1, 0) - z + 4^{-1}D). \end{aligned}$$

But if $z \in A(0, 0)$, then $(1, 0) - z \in 3D$. Therefore, for all $z \in A(0, 0)$,

$$\mathbb{P} \left[\bar{\tau}_k^{\lambda, 2, \mathcal{N}; \{z\}}(A(1, 0)) \geq m \right] \geq \mathbf{S}(\lambda) - \varepsilon. \tag{3.8}$$

Let $(i, j) \in \mathcal{L}$. Let $\mathcal{S} \subset A(i, j)$ such that $\#\mathcal{S} = m$. Using the independence between the BRW originating from the different points of \mathcal{S} , an argument similar to the above one and the definition of m , we get

$$\mathbb{P} \left[\bar{\tau}_k^{\lambda, 2, \mathcal{N}; \mathcal{S}}(A(i + 1, j + 1)) \geq m \right] \geq 1 - (1 - \mathbf{S}(\lambda) + \varepsilon)^m \geq 1 - \varepsilon.$$

With the same argument, we get

$$\mathbb{P} \left[\bar{\tau}_k^{\lambda, 2, \mathcal{N}; \mathcal{S}}(A(i + 1, j - 1)) \geq m \right] \geq 1 - (1 - \mathbf{S}(\lambda) + \varepsilon)^m \geq 1 - \varepsilon.$$

Therefore

$$\mathbb{P} \left[\bar{\tau}_k^{\lambda, 2, \mathcal{N}; \mathcal{S}}(A(i + 1, j + 1)) \geq m \text{ and } \bar{\tau}_k^{\lambda, 2, \mathcal{N}; \mathcal{S}}(A(i + 1, j - 1)) \geq m \right] \geq 1 - 2\varepsilon. \tag{3.9}$$

Let M be large enough to ensure $\mathbb{P} \left[\bar{\tau}_{\leq k}^{\lambda, 2, \mathcal{N}}(\mathbb{R}^2) \geq M/m \right] \leq \varepsilon/m$. By a natural coupling, we get, for all $\mathcal{S} \subset \mathbb{R}^2$ of cardinality at most m ,

$$\mathbb{P} \left[\bar{\tau}_{\leq k}^{\lambda, 2, \mathcal{N}, \mathcal{S}}(\mathbb{R}^2) \geq M \right] \leq m\varepsilon/m = \varepsilon. \tag{3.10}$$

The lemma follows from (3.8), (3.9) and (3.10). □

Proof of Lemma 3.4. Let m, k, M be given by Lemma 3.5. Let d_0 be such that $\varepsilon_{CLT}(d) \leq \varepsilon/M$ for all $d \geq d_0$ where ε_{CLT} appears in Theorem 2.1. Let $d \geq d_0$, X be a centered random variable in \mathbb{R}^d with log-concave density and L be any map given by Theorem 2.1. Let \mathcal{S} be a finite subset of \mathbb{R}^2 . With an appropriate coupling,

$$\mathbb{P} \left[\left\{ \bar{\tau}_{\leq k}^{\lambda, 2, \mathcal{N}; \mathcal{S}}(\mathbb{R}^2) \leq M \right\} \setminus \left\{ \bar{\tau}_{\leq k}^{\lambda, 2, L(X); \mathcal{S}} = \bar{\tau}_{\leq k}^{\lambda, 2, \mathcal{N}; \mathcal{S}} \right\} \right] \leq M\mathbb{P}[L(X) \neq \mathcal{N}] \leq M\varepsilon_{CLT}(d) \leq \varepsilon.$$

Therefore, for any $z \in A_L(0, 0)$,

$$\begin{aligned} & \mathbb{P} \left[\bar{\tau}_k^{\lambda, d, X; \{z\}}(A_L(1, 0)) \geq m \text{ and } \bar{\tau}_{\leq k}^{\lambda, d, X; \{z\}}(\mathbb{R}^d) \leq M \right] \\ &= \mathbb{P} \left[\bar{\tau}_k^{\lambda, 2, L(X); \{L(z)\}}(A(1, 0)) \geq m \text{ and } \bar{\tau}_{\leq k}^{\lambda, 2, L(X); \{L(z)\}}(\mathbb{R}^2) \leq M \right] \\ &\geq \mathbb{P} \left[\bar{\tau}_k^{\lambda, 2, \mathcal{N}; \{L(z)\}}(A(1, 0)) \geq m \text{ and } \bar{\tau}_{\leq k}^{\lambda, 2, \mathcal{N}; \{L(z)\}}(\mathbb{R}^2) \leq M \right. \\ &\quad \left. \text{and } \left\{ \bar{\tau}_{\leq k}^{\lambda, 2, L(X); \{L(z)\}} = \bar{\tau}_{\leq k}^{\lambda, 2, \mathcal{N}; \{L(z)\}} \right\} \right] \\ &\geq \mathbb{P} \left[\bar{\tau}_k^{\lambda, 2, \mathcal{N}; \{L(z)\}}(A(1, 0)) \geq m \text{ and } \bar{\tau}_{\leq k}^{\lambda, 2, \mathcal{N}; \{L(z)\}}(\mathbb{R}^2) \leq M \right] - \varepsilon \\ &\geq S(\lambda) - 2\varepsilon. \end{aligned}$$

This gives the first item. The second item is proven in exactly the same way. □

3.4.4 Plan and intuition

Setup. Let $\lambda > 0$, $\varepsilon > 0$, $d \geq 1$ and $K \in \mathcal{K}(d)$. Recall the definition of $\widehat{C}^0 = \widehat{C}^0(\lambda, \delta_{1/2}, d, K)$ in Section 3.1 and the notation $S(\lambda)$ for the survival probability of a Poisson(λ) offspring Galton-Watson process. The aim is to prove that the inequality

$$\mathbb{P}[\#\widehat{C}^0 = \infty] \geq S(\lambda) - \varepsilon$$

holds for any d large enough, uniformly in $K \in \mathcal{K}(d)$. Recall that \widehat{C}^0 can be built as the set of positions of a pruned version of the BRW $\bar{\tau}^{\lambda, d, X_K}$ where X_K denotes a random variable with uniform distribution on X_K . Recall in particular the notion of interference defined in (3.1). See Section 2.8 for notations on BRW and Section 3.2.1 for the construction of \widehat{C}^0 from $\bar{\tau}^{\lambda, d, X_K}$. The basic idea is that, up to an event whose probability vanishes when d tends to ∞ , \widehat{C}^0 is infinite when $\bar{\tau}^{\lambda, d, X_K}$ is infinite.

The underlying Galton-Watson tree. The underlying Galton-Watson tree τ of the BRW $\bar{\tau}^{\lambda, d, X_K}$ does not depend on d nor on K . It only depends on λ . This is a Galton-Watson process with Poisson(λ) offspring starting from one particle.

Control of the interference up to a given generation. Consider the case where the root \emptyset has a child x which itself has a child y . In the construction of \widehat{C}^0 we have to reject y because of interference with the root \emptyset if $V(y) \in V(\emptyset) + K^4$ that is if

$$[V(x) - V(\emptyset)] + [V(y) - V(x)] \in K. \tag{3.11}$$

⁴Actually $V(\emptyset) = 0$ but the argument is clearer if we keep writing $V(\emptyset)$.

Recall that X_K and X'_K are independent random variables with uniform distribution on K . Conditioned on the tree τ , the probability of (3.11) is $\mathbb{P}[X_K + X'_K \in K]$. By the rearrangement inequality (Theorem 2.3) this probability is at most $\mathbb{P}[X_B + X'_B \in B]$ where, as usual, B is the Euclidean ball of unit volume. It is moreover easy to check that $\mathbb{P}[X_B + X'_B \in B]$ tends to 0 when d tends to infinity (see Lemma 2.5). Therefore $\mathbb{P}[X_K + X'_K \in K]$ tends to 0 uniformly in K as d tends to infinity. With these ideas it is quite easy to show that for any given generation $k \geq 1$,

$$\mathbb{P}[\text{no particle of } \bar{\tau}_{\leq k}^{\lambda, d, X_K} \text{ is rejected because of interference}] \rightarrow 1 \text{ as } d \rightarrow \infty \text{ unif. in } K.$$

Control of the position of the particles. For various reasons, we need to control the position of the particles. Fix a linear map $L : \mathbb{R}^d \rightarrow \mathbb{R}^2$ given by Theorem 2.1 for X_K . Recall that \mathcal{N} denotes a standard Gaussian random vector \mathcal{N} on \mathbb{R}^2 . The map L (which depends on X_K and thus on d) fulfills the following property. The total variation distance between $L(X_K)$ and \mathcal{N} tends to 0 when d tends to ∞ , uniformly in K . Thanks to this property, we can control the value of the image by L of the position of the particles, uniformly in K when d tends to ∞ . This will be sufficient for our purpose.

Comparison with a super-critical oriented two-dimension percolation process.

The difficulty is to get a control over all generations of $\bar{\tau}_{\leq k}^{\lambda, d, X_K}$. Recall the definition of the sets $A_L(i, j)$, $(i, j) \in \bar{\mathcal{L}}$ in Section 3.4.2. Combining the previous ideas, we can prove the following results which are the basic steps of a renormalization scheme. Here m (number of seeds) and k (number of generations) are to be suitably chosen. See Lemma 3.4. The following results hold for d large enough, uniformly in K .

- With a probability close to $S(\lambda)$, no particle of $\bar{\tau}_{\leq k}^{\lambda, d, X_K}$ is rejected by interference (and thus all of them belong to the cluster \widehat{C}_0 in our construction) and m particles of $\bar{\tau}_k^{\lambda, d, X_K}$ are located in $A_L(1, 0)$. If this is the case, we say that stage $(0, 0)$ is a success.
- With a probability close to 1, for any $(i, j) \in \mathcal{L}$, if we start with a set of m particles whose set of positions is $\mathcal{S}(i, j) \subset A_L(i, j)$, then no particle of $\bar{\tau}_{\leq k}^{\lambda, d, X_K; \mathcal{S}(i, j)}$ (a BRW with initial set of particles located at $\mathcal{S}(i, j)$) is rejected by interference (when considering only interference within $\bar{\tau}_{\leq k}^{\lambda, d, X_K; \mathcal{S}(i, j)}$), m particles of $\bar{\tau}_k^{\lambda, d, X_K; \mathcal{S}(i, j)}$ belong to $A_L(i + 1, j + 1)$ and m particles of $\bar{\tau}_k^{\lambda, d, X_K; \mathcal{S}(i, j)}$ belong to $A_L(i + 1, j - 1)$. If this is the case, we say that stage (i, j) is a success (this is not well defined for the moment as it depends on $\mathcal{S}(i, j)$).

If stage $(0, 0)$ is a success (which occurs with probability close to $S(\lambda)$), then we can use the position of m particles of $\bar{\tau}_k^{\lambda, d, X_K}$ located in $A_L(1, 0)$ (recall that all of them belong to \widehat{C}_0 as none of them was rejected by interference) as a set of seeds $\mathcal{S}(1, 0)$ for stage $(1, 0)$. If stage $(1, 0)$ is a success (which occurs with probability close to 1) we can use the position of m particles of $\bar{\tau}_k^{\lambda, d, X_K; \mathcal{S}(1, 0)}$ located in $A_L(2, \pm 1)$ as a set of seeds $\mathcal{S}(2, \pm 1)$ for stage $(2, \pm 1)$ and so on. With the exception of stage $(0, 0)$, we thus have a natural coupling with a super-critical oriented percolation process on \mathcal{L} .

If stage $(0, 0)$ is a success, if the oriented percolation process percolates and if there were no interference between BRW of different stages, then \widehat{C}_0 would be infinite and the proof would be over. It remains to deal with interference between the BRW of different stages.

Control of the interference between BRW of different stages. This is actually the main difficulty of the proof. Let us mention that in the actual proof we will handle

interference between particles of a given BRW and particles of different BRW in a unified way, based on the following ideas. We perform over-pruning (see Section 3.2.3) to build a subset of \widehat{C}^0 . Concretely this means that, when exploring the BRW, if a particle does not fulfill one of the required properties, we reject the particle and its progeny. This depends on the order in which we explore the BRW, but this is not an issue for our purpose.

1. We fix a large M and do not explore more than M particles at each stage. If M is large enough, this does not modify significantly the probability of success at each stage. Thus there is no drawbacks. However, this gives a bound on the number of particles at each stage that can generate interference at a later stage.
2. We reject a particle and its progeny if it makes a step whose image by L is too large. More precisely, for a large Λ , if y is a child of x , we reject y and its progeny if $\|L(V(y) - V(x))\|_2 \geq \Lambda$. As above, if Λ is large enough, there is no drawbacks. However, there are two advantages:
 - (a) This reduces the interference region. Recall that D denotes the unit disk of \mathbb{R}^2 . The interference of x is $x + K \cap L^{-1}(\Lambda D)$ instead of $x + K^5$.
 - (b) This allows to localize the particles at each stage and then to identify the particles that can interfere at a later stage. As the seeds are in $(i, j) + D$ and as we explore k generations, all the particles considered at stage (i, j) belong to $(i, j) + (1 + \Lambda k)D$.
3. We reject a particle x and its progeny if we previously examined without rejecting a particle x' such that the following condition does *not* hold:

$$\|L(V(x) - V(x'))\|_2 \geq 2\Lambda \text{ or } V(x) - V(x') \in G(d, K, \eta). \quad (3.12)$$

Here $\eta > 0$ is a suitable parameter and $G(d, K, \eta)$ is defined in Section 3.4.1. By the localization properties, if x is examined at stage (i, j) and if x' has been examined at stage (i', j') , then $\|L(V(x) - V(x'))\|_2 \geq 2\Lambda$ holds as soon as (i, j) and (i', j') are far enough from each other. Moreover the number of particles examined at each stage is bounded. Therefore, for a given x , we just have to check the property $V(x) - V(x') \in G(d, K, \eta)$ for a bounded number of particles x' . By Lemma 3.3, conditioned on everything but $V(x) - V(\overleftarrow{x})$ ⁶ (recall that \overleftarrow{x} denotes the parent of x), this holds with high probability for all d large enough, uniformly in K .

Thanks to (3.12), the probability of interference is small. Let us explain this point. Let x, y, x' be three distinct particles where y is a child of x . We have examined and not rejected x and x' . We are examining y and we already now that $\|L(V(x) - V(y))\|_2 \leq \Lambda$ holds. We wonder whether y has to be rejected because of interference with x' . As the interference region of x' is $V(x') + K \cap L^{-1}(\Lambda D)$, we wonder whether $V(y)$ belongs to $V(x') + K \cap L^{-1}(\Lambda D)$. We condition on everything but $V(y) - V(x)$, which is a random variable uniformly distributed on K .

- If $\|L(V(x) - V(x'))\|_2 \geq 2\Lambda$, then y can not be rejected because of interference with x' . Indeed y belongs to $V(x) + K \cap L^{-1}(\Lambda D)$ and therefore y can not belong to $V(x') + K \cap L^{-1}(\Lambda D)$.

⁵Equivalently, we could have worked from the beginning with the BRW

$$\overleftarrow{\tau}^{\lambda P}[\|L(X_K)\|_2 \leq \Lambda], d, X_K^\Lambda$$

where X_K^Λ is distributed as X_K conditioned on $\|L(X_K)\|_2 \leq \Lambda$.

⁶In the proof, we will indeed consider conditional probabilities. We therefore have to be careful with these aspects.

- Otherwise, by (3.12), we have $V(x) - V(x') \in G(d, K, \eta)$. The probability that $V(y) - V(x') = (V(x) - V(x')) + (V(y) - V(x))$ (because of our conditioning, this amounts to consider $V(x) - V(x')$ as constant and $V(y) - V(x)$ as random with its usual law) belongs to K is at most η (by definition of $G(d, K, \eta)$). Thus the probability that y is rejected because of interference with x' is at most η .

Because of our control on the location and number of particles at each stage, this is sufficient.

A two-step approach to handle interference. Let us emphasize that, in addition to the control of the number of particles and the length of the projections of the steps, the main ingredient is thus the following two-step approach:

1. First we ensure that all relevant relative positions are good (this is the content of (3.12)).
2. Then we use this control to control the interference (the set $G(d, K, \eta)$ is designed for this task).

Comparison with the proof by Penrose in the Euclidean case. The plan of the proof is the same. Thanks to results of analysis and high dimension geometry (see Section 2) most parts of the proof of Penrose can actually be adapted to the non Euclidean case. One of the main difference is due to the lack of isotropy in our setting. In particular, the equivalent of the set of Good Gaps $G(d, K, \eta)$ in the Euclidean setting is simply $\mathbb{R}^d \setminus \frac{3}{4}B$. (Penrose actually does not use an exact equivalent of (3.12); see [14] page 542, condition (iii).) In our setting we had to provide an alternative description of this set of Good Gaps. Fortunately, an abstract definition was sufficient thanks to rearrangement inequalities (see Theorem 2.3).

3.4.5 Construction of a subset of \widehat{C}^0 related to an oriented percolation on \mathcal{L}

Parameters. Fix $\lambda > 1$ and $\varepsilon > 0$. Fix $m, d_1, k, M \geq 1$ as provided by Lemma 3.4 for the parameters λ and ε . Fix $\Lambda \geq 1$ such that $\mathbb{P}[\|\mathcal{N}\|_2 \geq \Lambda] \leq \varepsilon/M$. Let d_2 be such that, for all $d \geq d_2$, $\varepsilon_{CLT}(d) \leq \varepsilon/M$ where ε_{CLT} appears in Theorem 2.1. Fix $\eta > 0$ such that

$$400k^2\Lambda^2M^2\eta \leq \varepsilon. \tag{3.13}$$

Let d_3 be such that, for all $d \geq d_3$,

$$400k^2\Lambda^2M^2\eta^{-1}\varepsilon_G(d) \leq \varepsilon/3. \tag{3.14}$$

Set $d_0 = \max(d_1, d_2, d_3)$.

Setting and aim. Let $d \geq d_0$ and $K \in \mathcal{K}(d)$. Fix $L : \mathbb{R}^d \rightarrow \mathbb{R}^2$ given by Theorem 2.1 for X_K . By definition of Λ and as $d \geq d_0 \geq d_2$, we get (under an appropriate coupling)

$$\mathbb{P}[\|L(X_K)\|_2 \geq \Lambda] \leq \mathbb{P}[L(X_K) \neq \mathcal{N}] + \mathbb{P}[\|\mathcal{N}\|_2 \geq \Lambda] \leq 2\varepsilon/M. \tag{3.15}$$

We aim at proving

$$\mathbb{P}[\#\widehat{C}^0 = \infty] \geq (S(\lambda) - 5\varepsilon)\theta(1 - 5\varepsilon)$$

where θ is defined in Section 3.4.2. Theorem 3.2 will follow easily. Thanks to the choice of parameters, the following properties hold.

- For all $z \in A_L(0, 0)$,

$$\mathbb{P} \left[\bar{\tau}_k^{\lambda, d, X_K; \{z\}}(A_L(1, 0)) \geq m \text{ and } \bar{\tau}_{\leq k}^{\lambda, d, X_K; \{z\}}(\mathbb{R}^d) \leq M \right] \geq S(\lambda) - \varepsilon. \quad (3.16)$$

- For all $(i, j) \in \mathcal{L}$ and all $S \subset A_L(i, j)$ with cardinality m ,

$$\mathbb{P} \left[\bar{\tau}_k^{\lambda, d, X_K; S}(A_L(i + 1, j + 1)) \geq m \text{ and } \bar{\tau}_k^{\lambda, d, X_K; S}(A_L(i + 1, j - 1)) \geq m \text{ and } \bar{\tau}_{\leq k}^{\lambda, d, X_K; S}(\mathbb{R}^d) \leq M \right] \geq 1 - \varepsilon. \quad (3.17)$$

- (3.13), (3.14), (3.15).

Randomness and σ -fields. Recall that X_K stands for a random variable with uniform distribution on K . Let

$$(\bar{\tau}^{i, j, n})_{(i, j, n) \in \bar{\mathcal{L}} \times \{1, \dots, m\}}$$

be a family of independent copies of $\bar{\tau}^{\lambda, d, X_K}$. Let $(\alpha^{i, j})_{(i, j) \in \bar{\mathcal{L}}}$ be a family of i.i.d. Bernoulli random variables with parameter $1 - \varepsilon$. For all $(i, j) \in \bar{\mathcal{L}}$ we denote by $\mathcal{F}_{i, j}$ (resp. $\mathcal{F}_{i, j}^-$) the σ -field generated by the $\bar{\tau}^{i', j', n}$ and the $\alpha^{i', j'}$ for $(i', j', n) \in \bar{\mathcal{L}} \times \{1, \dots, m\}$ such that (i', j') is smaller (resp. strictly smaller) than (i, j) for the lexicographic order.

We will not formalize it but we will merge appropriately pruned versions of the BRW $\bar{\tau}^{i, j, n}$ to get a unique BRW starting from one particle located at 0. This latter BRW is the BRW explored in order to build a subset of \hat{C}^0 .

Further notations and remarks. At the beginning, the site $(0, 0)$ is active and each site $(i, j) \in \mathcal{L}$ is inactive. Moreover, with $(0, 0)$ is associated the singleton $\mathcal{S}(0, 0) = \{0_{\mathbb{R}^d}\} \subset A_L(0, 0)$. We then enumerate the sites (i, j) of $\bar{\mathcal{L}}$ by lexicographic order. Some sites $(i, j) \in \mathcal{L}$ will be activated. With each active site (i, j) will be associated a subset $\mathcal{S}(i, j) \subset A_L(i, j)$ of cardinality m . This set of m points will always be, in a coupling with the Boolean model, a subset of \hat{C}^0 . If at the end of the construction there exists in the graph $\bar{\mathcal{L}}$ an infinite path of active sites, then \hat{C}^0 is infinite and percolation occurs in the Boolean model.

To simplify some arguments, we will also associate with each site of $\bar{\mathcal{L}}$ a state: open or closed. It will be done in such a way that if π is an infinite open path in $\bar{\mathcal{L}}$ from $(0, 0)$, then π only contains active sites and therefore percolation occurs in the Boolean model.

We will use the over-pruning algorithm described in Section 3.2.3. We will often use over-pruning implicitly by rejecting more particles than necessary and by not considering the children of some particles (which amount to define their interference region as the empty set).

Set

$$G = G(d, K, \eta)$$

where $G(d, K, \eta)$ is introduced in Section 3.4.1.

Stage $(0, 0)$. We consider the BRW $\bar{\tau} = \bar{\tau}^{0, 0, 1}$.

1. We examine successively the particles of generation at most k of this BRW in any admissible order. The last requirement means that:

- Children are examined after their parents.

- Children of a given parent are examined in a row: once we start examining one of them, we then examine all the children of this parent.

We stop as soon as we have examined all the particles or as soon as the particle x under examination fulfills one of the following conditions:

- (a) Overpopulation. The particle is the M -th particle examined.
- (b) Bad Gap. This occurs in any of the following conditions.
 - x is not the root and $V(x) \notin V(\overleftarrow{x}) + K \cap L^{-1}(\Lambda D)$. This ensures that the interference region of \overleftarrow{x} is indeed $V(\overleftarrow{x}) + K \cap L^{-1}(\Lambda D)$ (see Section 3.2.3 and the remarks in the paragraph above).
 - There exists a particle y examined strictly before x such that $V(x) \notin V(y) + G$.
- (c) Interference. There exists a particle y examined strictly before x which is not the parent \overleftarrow{x} nor a sibling of x and which is such that $V(x) \in V(y) + K \cap L^{-1}(\Lambda D)$ ⁷.

One defines the following two sets.

- The set $\mathcal{G}(0, 0)$ (\mathcal{G} stands for generated) of all the particles examined except the last one if it caused Overpopulation or Bad Gap or Interference. By some abuse of notation, we will sometimes see $\mathcal{G}(0, 0)$ (and other similar sets) as the set of positions of the particles. We will call them the particles *generated* at stage $(0, 0)$. Note that each particle whose children have been examined is a generated particle (this is due to the fact that we examine the particles in an admissible order). This is a key remark when considering interference later in the construction.
 - In the coupling with the Boolean model, all the points of $\mathcal{G}(0, 0)$ (seen as the set of positions of the particles) belong to \widehat{C}^0 . Indeed, none of them caused a stop by Interference.
 - All the gaps between any two distinct particles of $\mathcal{G}(0, 0)$ are good:

$$\forall x, y \in \mathcal{G}(0, 0), x \neq y \implies V(y) \in V(y) + G.$$

Indeed, none of them caused a stop by Bad Gap.

–

$$\text{The image by } L \text{ of } \mathcal{G}(0, 0) \text{ is included in } k\Lambda D. \tag{3.18}$$

Here we see $\mathcal{G}(0, 0)$ as the set of positions of the particles. This is due to the fact that the position of the root is 0, the fact that the image by L of each step belongs to ΛD and the fact that we did not explore the BRW beyond generation k .

- $\mathcal{G}(0, 0)$ contains at most M points.
- The set $\mathcal{G}_k(0, 0) \subset \mathcal{G}(0, 0)$ of particles of generation k examined except the last one if it caused Overpopulation or Bad Gap or Interference.

⁷With the notations of Section 3.2.3, we want to reject any particle which belongs to

$$\bigcup_{y \in \mathcal{B}} V(y) + K \cap L^{-1}(\Lambda D).$$

Let \mathcal{B}' be the set of particles examined strictly before x and which are neither \overleftarrow{x} nor a sibling of x . As $\mathcal{B} \subset \mathcal{B}'$, we are performing over-pruning at this step.

2. If

$$\#(\mathcal{G}_k(0,0) \cap A_L(1,0)) \geq m \tag{3.19}$$

we say that the site $(0,0)$ is open, that the site $(1,0)$ is active and we define $\mathcal{S}(1,0)$, where \mathcal{S} stands for seeds, as the m first points of

$$\mathcal{G}_k(0,0) \cap A_L(1,0)$$

in Neveu ordering (see Section 2.8). In the coupling with the Boolean model, all the points of $\mathcal{S}(1,0)$ belong to \widehat{C}^0 . This holds because $\mathcal{S}(1,0) \subset \mathcal{G}_k(0,0) \subset \mathcal{G}(0,0)$ and all points of $\mathcal{G}(0,0)$ belong to \widehat{C}^0 . If (3.19) does not hold, we say that the site $(0,0)$ is closed and $(1,0)$ remains inactive.

Stage (i,j) . Recall that we now consider successively each $(i,j) \in \mathcal{L}$ by lexicographic order. If (i,j) is inactive, then we decide as follows: it is open if $\alpha^{i,j} = 1$; it is closed otherwise. Therefore, in this case, it is open independently of everything else with probability $1 - \varepsilon$.

Thereafter, we consider the case where (i,j) is active. The set $\mathcal{S}(i,j)$ is well defined. It's a subset of cardinal m of $A_L(i,j)$. List the points of $\mathcal{S}(i,j)$ in an arbitrary order: $\mathcal{S}(i,j) = \{x^1, \dots, x^m\}$. We consider the m BRW $\bar{\tau}^n = x^n + \bar{\tau}^{i,j,n}$ where $x^n + \bar{\tau}^{i,j,n}$ designates the BRW $\bar{\tau}^{i,j,n}$ in which x^n was added to the position of all the particles. We gather these m BRW into a single BRW originating from $\mathcal{S}(i,j)$. We denote it by $\bar{\tau}^{\mathcal{S}(i,j)}$.

1. We examine successively the particles of generation between 1 and k of $\bar{\tau}^{\mathcal{S}(i,j)}$ in any admissible order (see Stage $(0,0)$). In particular, we never examine the roots of this BRW (note that the positions of the roots are also positions of particles of BRW examined during one of the previous stages). We stop as soon as we have examined all the particles or as soon as the particle x under examination fulfills one of the following conditions:

- (a) Overpopulation. The particle is the M -th particle examined during this stage (i,j) .
- (b) Bad Gap. One of the following conditions occurs.
 - $V(x) \notin V(\overleftarrow{x}) + K \cap L^{-1}(\Lambda D)$. This ensure that the interference region of \overleftarrow{x} is indeed $V(\overleftarrow{x}) + K \cap L^{-1}(\Lambda D)$ (see stage $(0,0)$ for more explanations).
 - There exists a particle y examined strictly before x during this stage (i,j) or generated during one of the previous stages (its position then belongs to $\mathcal{G}(i',j')$ for some $(i',j') < (i,j)$ for the lexicographic order) such that:

$$\|L(V(x)) - L(V(y))\|_2 \leq 2\Lambda \text{ and } V(x) \notin V(y) + G.$$

- (c) Interference. There exists a particle y examined strictly before x during this stage (i,j) or generated during one of the previous stages (its position then belongs to $\mathcal{G}(i',j')$ for some $(i',j') < (i,j)$ for the lexicographic order) such that: this is not the parent \overleftarrow{x} nor a sibling of x and we have:

$$V(x) \in V(y) + K \cap L^{-1}(\Lambda D).$$

One defines the following two sets.

- The set $\mathcal{G}(i,j)$ of all particles examined except the last one if it caused Overpopulation or Bad Gap or Interference. We will call them the particles *generated* at stage (i,j) . As before, each particle whose children have been examined is a generated particle.

- In the coupling with the Boolean model, all the points of $\mathcal{G}(i, j)$ belong to \widehat{C}^0 . Indeed, none of them caused a stop by Interference.
- All the gaps between two distinct points x and y of

$$\bigcup_{(i', j') \leq (i, j)} \mathcal{G}(i', j')$$

satisfy

$$\|L(V(x)) - L(V(y))\|_2 > 2\Lambda \text{ or } V(x) \in V(y) + G. \quad (3.20)$$

Indeed, none of them caused a stop by Bad Gap.

-

$$\text{The image by } L \text{ of } \mathcal{G}(i, j) \text{ is included in } (i, j) + 2k\Lambda D. \quad (3.21)$$

This is due to the fact that the positions of the roots belong to $(i, j) + D$, the fact that the image by L of each step belongs to ΛD , the fact that we only explored the first k generations and the fact that $1 + k\Lambda \leq 2k\Lambda$.

- $\mathcal{G}(i, j)$ contains at most M points.
- The set $\mathcal{G}_k(i, j) \subset \mathcal{G}(i, j)$ of particles of generation k examined, except the last one if it caused a stop by Overpopulation or Bad Gap or Interference.

2. If

$$\#(\mathcal{G}_k(i, j) \cap A_L(i + 1, j + 1)) \geq m \text{ and } \#(\mathcal{G}_k(i, j) \cap A_L(i + 1, j - 1)) \geq m \quad (3.22)$$

then:

- We say that the site (i, j) is open.
- If the site $(i + 1, j + 1)$ is inactive then we say that it is henceforth active and we define $\mathcal{S}(i + 1, j + 1)$ as the m first point by Neveu ordering (see Section 2.8) of $\mathcal{G}_k(i, j) \cap A_L(i + 1, j + 1)$. In the coupling with the Boolean model, all the points of $\mathcal{S}(i + 1, j + 1)$ belong to \widehat{C}^0 .
- If the site $(i + 1, j - 1)$ is inactive then we say that it is henceforth active and we define $\mathcal{S}(i + 1, j - 1)$ as the m first point by Neveu ordering of $\mathcal{G}_k(i, j) \cap A_L(i + 1, j - 1)$. In the coupling with the Boolean model, all the points of $\mathcal{S}(i + 1, j - 1)$ belong to \widehat{C}^0 .

Otherwise, we say that the site (i, j) is closed.

3.4.6 Bounds on conditional probabilities

The aim of this section is to prove the following lemmas. Parameters have been fixed in Section 3.4.5.

Lemma 3.8. For all $(i, j) \in \overline{\mathcal{L}}$, $\{(i, j) \text{ active}\} \in \mathcal{F}(i, j)^-$ and $\{(i, j) \text{ open}\} \in \mathcal{F}(i, j)$.

Lemma 3.9. We have $\mathbb{P}[(0, 0) \text{ open}] \geq S(\lambda) - 5\varepsilon$.

Lemma 3.10. For all $(i, j) \in \mathcal{L}$, $\mathbb{P}[(i, j) \text{ open} | \mathcal{F}^-(i, j)] \geq 1 - 5\varepsilon$.

Proof of Lemma 3.8. This is straightforward by construction. □

Proof of Lemma 3.9. We have

$$\{(0, 0) \text{ closed}\} \setminus (\text{Overpopulation} \cup \text{BadGap} \cup \text{Interference}) \subset \text{Else} \quad (3.23)$$

where

$$\begin{aligned} \text{Overpopulation} &= \{\text{the enumeration stops because of Overpopulation}\}, \\ \text{BadGap} &= \{\text{the enumeration stops because of Bad Gap} \\ &\quad \text{and not Overpopulation}\}, \\ \text{Interference} &= \{\text{the enumeration stops because of Interference} \\ &\quad \text{and not Overpopulation or Bad Gap}\}, \\ \text{Else} &= \{\bar{\tau}_k(A_L(1, 0)) < m\} \end{aligned}$$

and where $\bar{\tau}$ is the BRW used in Stage $(0, 0)$. Indeed, if the event on the left-hand side of (3.23) occurs, then (3.19) does not hold and $\mathcal{G}_k(0, 0) = \{V(x), x \in \bar{\tau}_k\}$ hence $\bar{\tau}_k(A_L(1, 0)) < m$. As a consequence

$$\{(0, 0) \text{ closed}\} \subset \text{BadGap} \cup \text{Interference} \cup \text{Overpopulation} \cup \text{Else}. \quad (3.24)$$

Let us give an upper bound for the probability of BadGap. We have $\text{BadGap} \subset \text{BadGap}_1 \cup \text{BadGap}_2$ with

$$\text{BadGap}_1 = \bigcup_x \{V(x) \notin V(\bar{x}) + K \cap L^{-1}(\Lambda D)\}$$

and

$$\text{BadGap}_2 = \bigcup_{x \neq y} \{V(x) \notin V(y) + G\} \setminus \text{BadGap}_1$$

where for BadGap_1 the union is over all x in the M first particles (otherwise this is the event Overpopulation which occurs) of the tree except the root and where for BadGap_2 the union is over the same set of x and over all y strictly preceding x (for the order used for the examination).

The event BadGap_2 is then the union of at most M^2 events. Rewriting this events, we get

$$\text{BadGap}_2 \subset \bigcup_{x \neq y} \{V(\bar{x}) - V(y) + V(x) - V(\bar{x}) \notin G\}.$$

But, conditioned on the underlying Galton-Watson tree, $V(x) - V(\bar{x})$ is independent of $(V(\bar{x}), V(y))$ (actually it is independent of the positions of all previously examined particles) and then of $V(\bar{x}) - V(y)$. Moreover $V(x) - V(\bar{x})$ has the same distribution as X_K . Therefore, conditioning on the underlying tree τ we have, for all x, y as above,

$$\mathbb{P}[V(\bar{x}) - V(y) + V(x) - V(\bar{x}) \notin G | \tau] \leq \sup_{z \in \mathbb{R}^d} \mathbb{P}[z + X_K \notin G] \leq \eta^{-1} \varepsilon_G(d)$$

where $\varepsilon_G(d)$ is defined in Lemma 3.3. We then get

$$\mathbb{P}[\text{BadGap}_2] \leq M^2 \eta^{-1} \varepsilon_G(d).$$

Hence $\mathbb{P}[\text{BadGap}_2] \leq \varepsilon$ by (3.14). (We also use the fact that several of our constants are greater than 1.)

The event BadGap_1 is the union of at most M event. Arguing as above and using (3.15), we get $\mathbb{P}[\text{BadGap}_1] \leq M2\varepsilon/M = 2\varepsilon$. Thus

$$\mathbb{P}[\text{BadGap}] \leq 3\varepsilon.$$

Let us now provide an upper bound for the probability of the event Interference. Denote x and y the two particles which cause the event Interference. In particular:

- x is one of the first M particles (otherwise this is the event Overpopulation which occurs) of the tree except the root.
- y is a particle of $\bar{\tau}$ strictly preceding x (for the order of enumeration).
- $V(x) \in V(y) + K$.
- $V(\bar{x}) \in V(y) + G$ because otherwise BadGap would occur instead of Interference (recall also $G = -G$).

Thus,

$$\begin{aligned} \text{Interference} &\subset \bigcup_{x,y} \{V(x) \in V(y) + K\} \cap \{V(\bar{x}) \in V(y) + G\} \\ &\subset \bigcup_{x,y} \{V(\bar{x}) + (V(x) - V(\bar{x})) \in V(y) + K\} \cap \{V(\bar{x}) \in V(y) + G\} \end{aligned}$$

where the union is over all x in the first M particles of $\bar{\tau}$ except the root and y in the particles of $\bar{\tau}$ strictly preceding x . The right-hand side is then the union of at most M^2 events. By definition of $G = G(d, k, \eta)$ and using a reasoning similar to the one used for the event Bad, we then deduce

$$\mathbb{P}[\text{Interference}] \leq M^2 \eta \leq \varepsilon$$

by (3.13). Finally, by (3.16), we have

$$\mathbb{P}[\text{Overpopulation} \cup \text{Else}] \leq 1 - S(\lambda) + \varepsilon.$$

As a consequence, $\mathbb{P}[(0, 0) \text{ open}] \geq S(\lambda) - 5\varepsilon$. □

Proof of Lemma 3.10. Let $(i, j) \in \mathcal{L}$. The event $\{(i, j) \text{ is active}\}$ is $\mathcal{F}^-(i, j)$ measurable. Therefore, we have to show

$$\mathbb{P}[(i, j) \text{ open} | \mathcal{F}^-(i, j)] \geq 1 - \varepsilon \text{ on the event } \{(i, j) \text{ is active}\}$$

and

$$\mathbb{P}[(i, j) \text{ open} | \mathcal{F}^-(i, j)] \geq 1 - \varepsilon \text{ on the event } \{(i, j) \text{ is inactive}\}.$$

The second property is straightforward. Indeed, when (i, j) is inactive, (i, j) has been defined as open independently of everything else with probability $1 - \varepsilon$. Let us prove the first property.

We now work on the event $\{(i, j) \text{ is active}\}$.

On the event $\{(i, j) \text{ is active}\}$ there is a well defined set $\mathcal{S}(i, j)$ with cardinality m whose points are the starting points of m BRW. This set is measurable with respect to $\mathcal{F}^-(i, j)$. We also have the BRW $\bar{\tau}^{\mathcal{S}(i, j)}$ which has been used in Stage (i, j) . We have, as in the proof of Lemma 3.9,

$$\{(i, j) \text{ is closed}\} \cap \{(i, j) \text{ is active}\} \subset \text{BadGap} \cup \text{Interference} \cup \text{Overpopulation} \cup \text{Else}$$

where the events BadGap, Interference and Overpopulation are defined as in the proof of Lemma 3.9 and where

$$\text{Else} = \{\bar{\tau}_k^{\mathcal{S}(i, j)}(A_L(i + 1, j + 1)) < m \text{ or } \bar{\tau}_k^{\mathcal{S}(i, j)}(A_L(i + 1, j - 1)) < m\}.$$

Let us provide an upper bound for the conditional probability of the event BadGap. The proof is similar to the one in Lemma 3.9. The proof is identical for BadGap₁. For BadGap₂, the combinatorial factor is not M^2 anymore. There are at most M choices for x . Let us fix some x . There are at most M choices for y if y has been examined during

Stage (i, j) . If $y \in \mathcal{G}(i', j')$ for $(i', j') < (i, j)$, then $L(V(y)) \in (i', j') + 2k\Lambda D$ (see (3.18) and (3.21)). Similarly, $L(V(x)) \in (i, j) + 2k\Lambda D$ (otherwise BadGap_1 would have occurred and not BadGap_2). As a consequence, the condition $\|L(V(x)) - L(V(y))\|_2 \leq 2\Lambda$ yields

$$\|(i, j) - (i', j')\|_2 \leq (4k + 2)\Lambda \leq 6k\Lambda.$$

Therefore there are at most $(12k\Lambda + 1)^2$ choice for (i', j') and therefore at most

$$(12k\Lambda + 1)^2 M + M \leq 200k^2 \Lambda^2 M$$

choices for y . Finally, the number of choices for (x, y) is at most $200k^2 \Lambda^2 M^2$. As a consequence, arguing as in Lemma 3.9,

$$\begin{aligned} \mathbb{P}[\text{BadGap} | \mathcal{F}^-(i, j)] &\leq 200k^2 \Lambda^2 M^2 \sup_{z \in \mathbb{R}^d} \mathbb{P}[z + X_K \notin G] \\ &\leq 200k^2 \Lambda^2 M^2 \eta^{-1} \varepsilon_G(d). \end{aligned}$$

where $\varepsilon_G(d)$ is defined in Lemma 3.3. Therefore,

$$\mathbb{P}[\text{BadGap} | \mathcal{F}^-(i, j)] \leq \mathbb{P}[\text{BadGap}_1 | \mathcal{F}^-(i, j)] + \mathbb{P}[\text{BadGap}_2 | \mathcal{F}^-(i, j)] \leq 3\varepsilon$$

by (3.14).

Let us now give an upper bound for the probability of the event Interference. Denote x and y the two particles which cause the event Interference. In particular,

- x is one of the first M particles (otherwise the event Overpopulation would occur) of $\bar{\tau}$ not belonging to generation 0.
- y is a particle examined strictly before x during this stage (i, j) or generated during one of the previous stage (its position belongs in this case to $\mathcal{G}(i', j')$ for some $(i', j') < (i, j)$ for the lexicographical order).
- y is not \overleftarrow{x} .
- $V(x) \in V(y) + K \cap L^{-1}(\Lambda D)$. This property yields $\|L(V(x)) - L(V(y))\|_2 \leq \Lambda$. By an argument already used to give an upper bound for the event BadGap , we get that there are at most $400k^2 \Lambda^2 M^2$ choices for (x, y) . By $\|L(V(x)) - L(V(y))\|_2 \leq \Lambda$ we also get

$$\|L(V(\overleftarrow{x})) - L(V(y))\|_2 \leq 2\Lambda. \tag{3.25}$$

Indeed, as BadGap does not occur, $\|L(V(x)) - L(V(\overleftarrow{x}))\|_2 \leq \Lambda$.

- $V(\overleftarrow{x}) \in V(y) + G$. Indeed, we have

$$V(\overleftarrow{x}), V(y) \in \bigcup_{(i', j') \leq (i, j)} \mathcal{G}(i', j'). \tag{3.26}$$

This is clear for y if y has been generated during some Stage $(i', j') < (i, j)$. This is also true if y has been examined during the current stage. Indeed, y has been examined before x . Therefore y did not stopped the enumeration of the particles and therefore $V(y) \in \mathcal{G}(i, j)$. With the same argument we check that this is also true for $V(\overleftarrow{x})$. The required result is then a consequence of (3.20), (3.25), (3.26) and of $\overleftarrow{x} \neq y$.

We then have

$$\text{Interference} \subset \bigcup_{x, y} \{V(x) \in V(y) + K\} \cap \{V(\overleftarrow{x}) \in V(y) + G\} \tag{3.27}$$

where x and y are as above. In particular, as already mentioned, there are at most $400k^2\Lambda^2M^2$ choices for (x, y) . By definition of Good Gaps we then get, with the same arguments as before (we only use the randomness on $V(x) - V(\bar{x})$),

$$\mathbb{P}[\text{Interference}|\mathcal{F}^-(i, j)] \leq 400k^2\Lambda^2M^2\eta \leq \varepsilon$$

by (3.13).

Finally, by (3.17), we get

$$\mathbb{P}[\text{Overpopulation} \cup \text{Else}|\mathcal{F}^-(i, j)] \leq \varepsilon.$$

As a consequence, (we are still working on the $\mathcal{F}_{i,j}^-$ measurable event $\{(i, j) \text{ is active}\}$),

$$\mathbb{P}[(i, j) \text{ open}|\mathcal{F}^-(i, j)] \geq 1 - 5\varepsilon. \quad \square$$

3.4.7 Proof of Theorem 3.2

In Section 3.4.5 we fixed $\lambda > 1$ and $\varepsilon > 0$. We then got some integer d_0 and several other parameters satisfying various properties. We then let $d \geq d_0$ and $K \in \mathcal{K}(d)$ and built some process in Section 3.4.5. We studied some properties of this process in Section 3.4.6. We now conclude the proof of Theorem 3.2.

Let $(i, j), (i', j') \in \bar{\mathcal{L}}$ be such that $(i, j) \rightarrow (i', j')$, that is there is an arrow from (i, j) to (i', j') in the graph $\bar{\mathcal{L}}$. If (i, j) is active and open, then (i', j') is active and $\mathcal{S}(i', j')$ is a well defined subset of \widehat{C}^0 .

Recall that $(0, 0)$ is active. If there exists an infinite path π in $\bar{\mathcal{L}}$ originating from $(0, 0)$ and containing only open sites, by the previous discussion, we get that \widehat{C}^0 is infinite. Therefore

$$\mathbb{P}[\#\widehat{C}^0 = \infty] \geq \mathbb{P}[\text{there exists an infinite open path from } (0, 0)].$$

By Lemmas 3.8, 3.9 and 3.10 we get

$$\mathbb{P}[\text{there exists an infinite open path from } (0, 0)] \geq (S(\lambda) - 5\varepsilon)\theta(1 - 5\varepsilon)$$

where $\theta(1 - 5\varepsilon)$ is the probability that there exists an infinite open path originating from $(1, 0)$ in a Bernoulli site percolation on \mathcal{L} with parameter $1 - 5\varepsilon$. Therefore

$$\mathbb{P}[\#\widehat{C}^0 = \infty] \geq (S(\lambda) - 5\varepsilon)\theta(1 - 5\varepsilon).$$

But $\theta(1 - 5\varepsilon)$ tends to 1 as ε tends to 0 (this is stated as (3.4)). This proves Theorem 3.2 in the case $\lambda > 1$. When $\lambda \leq 1$, $S(\lambda) = 0$ and the required result is straightforward.

4 Proof of Theorem 1.4

4.1 Framework and the clusters \widehat{C}^0 and \widehat{A}^0

Framework. Let $\beta > 0, \rho > 1, d \geq 1$ and $K \in \mathcal{K}(d)$. Set

$$\lambda = \beta\kappa_c(\rho)^d.$$

Let χ_1 and χ_ρ be independent Poisson point processes on \mathbb{R}^d with intensity λdx and $\lambda\rho^{-d} dx$. Set $r(c) = 1/2$ for $c \in \chi_1$ and $r(c) = \rho/2$ for $c \in \chi_\rho$. Write $\chi = \chi_1 \cup \chi_\rho$. The process $\{(c, r(c)), c \in \chi\}$ is a Poisson point process on $\mathbb{R}^d \times (0, +\infty)$ with intensity measure $dx \otimes \lambda\nu_{d,\rho}(dr)$.

Write $\chi^0 = \chi \cup \{0\}$ and set $r(0) = \rho/2$.

The cluster \widehat{C}^0 . As in the constant radius case, we can define a relevant undirected graph structure on χ^0 as follows. We put an edge between x and y if the associated grains $x + r(x)K$ and $y + r(y)K$ touch each other, that is if

$$y \in x + (r(y) + r(x))K.$$

As in the constant radius case, $\widehat{C}^0 = \widehat{C}^0(\beta, \rho, d, K)$ (we simplify the notations for the parameters) is the connected component of the graph χ^0 which contains 0. We are interested in the probability that \widehat{C}^0 is unbounded (see (1.2)) and in

$$\beta_c(\rho, d, K) = \inf \left\{ \beta > 0 : \mathbb{P}[\widehat{C}^0 \text{ is unbounded}] > 0 \right\}.$$

The cluster \widehat{A}^0 . We define a new undirected graph on χ^0 . In this new graph, we put an edge between points x and y if $r(x) \neq r(y)$ and if the associated grains touch each other. In other words, we put an edge between points x and y if⁸

$$r(x) \neq r(y) \text{ and } y \in x + \frac{1 + \rho}{2}K.$$

We denote by $\widehat{A}^0 = \widehat{A}^0(\beta, \rho, d, K)$ (A stands for “alternate” as we alternate between 1-grains and ρ -grains) the connected component of 0 in this new graph. Clearly,

$$\widehat{A}^0 \subset \widehat{C}^0.$$

4.2 Two related BRW

We will not formalize the algorithms of exploration of \widehat{C}^0 and \widehat{A}^0 . This can be done as in Section 3.2 to which we refer for more details. We will give directly the relation with two BRW and provide some intuition. Let $\beta > 0, \rho > 1, d \geq 1$ and $K \in \mathcal{K}(d)$.

Recall that \widehat{A}^0 is a subset of \widehat{C}^0 and that we are interested in whether \widehat{C}^0 is infinite or not. Therefore, instead of investigating \widehat{C}^0 or \widehat{A}^0 , we can investigate

$$\frac{2}{1 + \rho}\widehat{C}^0 \text{ or } \frac{2}{1 + \rho}\widehat{A}^0.$$

More specifically, the BRW we will study will provide subsets of $\frac{2}{1 + \rho}\widehat{C}^0$ or $\frac{2}{1 + \rho}\widehat{A}^0$. This is a very tiny change but it will simplify formulas by removing numerous $\frac{1 + \rho}{2}$ factors.

4.2.1 A BRW for the upper bound on percolation probability

The BRW. Consider for example a deterministic 1-grain

$$x + \frac{1}{2}K.$$

The random ρ -grains which touch the previous grain are the set of grains centered at points of

$$\left(x + \frac{1 + \rho}{2}K \right) \cap \chi_\rho.$$

Therefore, this is a Poisson random variable with parameter

$$\beta \kappa_c(\rho)^d \rho^{-d} \left| \frac{1 + \rho}{2}K \right| = \beta \left(\frac{1}{\sqrt{\rho}} \right)^d$$

⁸Note that one of the two radii equals $1/2$ and the other equals $\rho/2$, therefore the sum is always $(\rho + 1)/2$.

as $|K| = 1$ and by definition of $\kappa_c(\rho)$ (see (1.7)). Similar results hold for the other cases. We can therefore see that \widehat{C}^0 is the set of points of some pruned two-type BRW. We refer to Section 3.2 for details on the pruning but mention one important difference below. It is a two-type BRW. There are 1-particles and ρ -particles. It starts with one ρ -particle located at 0. The progeny are independent (between the two types) and Poisson distributed. The matrix of mean is the matrix $M(\beta, \rho, d)$ defined in (1.9). The steps of the BRW (which we will actually not use) are i.i.d. with uniform distribution on K for type 1 to type 1 progeny, on ρK for type ρ to type ρ progeny, on $\frac{\rho+1}{2}K$ otherwise.

Let us conclude with the important difference in the pruning process. Recall the vocabulary introduced in (3.1). Here we are revealing points of two independent Poisson point processes: χ_1 and χ_ρ . Therefore there can only be interference between two 1-particles (when we are revealing χ_ρ) or between two ρ -particles (when are revealing χ_1). In other words, a child y of ρ -particle x can only be rejected because of interference with another ρ -particle x' and a child y of 1-particle x can only be rejected because of interference with another 1-particle x' .

The plan. The upper bound for the percolation probability will follow from a simple analysis of the survival probability of the underlying Galton-Watson process.

4.2.2 A BRW for the lower bound on percolation probability

The BRW. Recall

$$\frac{2}{1+\rho} \widehat{A}^0 \subset \frac{2}{1+\rho} \widehat{C}^0.$$

We will prove a lower bound on the probability that the set of the left is infinite. This will give a lower bound on the probability that the set of the right is infinite.

The set $\frac{2}{1+\rho} \widehat{A}^0$ is the set of positions of a pruned BRW. We refer to Section 3.2 for details on the pruning. Here we only describe the BRW. It is a two-type BRW. There are 1-particles and ρ -particles. It starts with one ρ -particle located at 0. The progeny are independent (between the two types) and Poisson distributed. The matrix of mean is

$$M'(\beta, \rho, d) = \beta \begin{pmatrix} 0 & \left(\frac{1}{\sqrt{\rho}}\right)^d \\ (\sqrt{\rho})^d & 0 \end{pmatrix}. \tag{4.1}$$

The steps of the BRW are i.i.d. with uniform distribution on K . (Recall that we are interested in $\frac{2}{1+\rho} \widehat{A}^0$.) Such BRW will be denoted by

$$\overline{\tau}^{\beta, \rho, d, X_K}.$$

As in Section 4.2.1, there can only be interference between two 1-particles (when we are revealing χ_ρ) or between two ρ -particles (when are revealing χ_1).

4.3 A result on survival probability of Galton-Watson processes

Let $\beta > 0, \rho > 1$ and $d \geq 1$. Recall the matrices $M(\beta, \rho, d)$ and $M'(\beta, \rho, d)$ defined by (1.9) and (4.1). Let $\tau(\beta, \rho, d)$ be a two-type Galton-Watson process with independent Poisson progeny with mean matrix $M(\beta, \rho, d)$ and starting with one ρ -particle. Denote by $S(\beta, \rho, d)$ its survival probability. Let $\tau'(\beta, \rho, d)$ be a two-type Galton-Watson process with independent Poisson progeny with mean matrix $M'(\beta, \rho, d)$ and starting with one ρ -particle. Denote by $S'(\beta, \rho, d)$ its survival probability. Let $\tau''(\beta^2)$ be a one type Galton-Watson process with Poisson(β^2) progeny starting with one particle. As usual we denote by $S(\beta^2)$ its survival probability.

The aim of this section is to prove the following lemma.

Lemma 4.1. *Let $\beta > 0$ and $\rho > 1$. There exists a sequence $(\varepsilon_{GW}(d))_d$ of positive real numbers which tends to 0 such that,*

$$\forall d \geq 1, S(\beta^2) - \varepsilon_{GW}(d) \leq S'(\beta, \rho, d) \leq S(\beta, \rho, d) \leq S(\beta^2) + \varepsilon_{GW}(d).$$

This is a straightforward consequence of the following lemma. We keep the notations given at the beginning of Section 4.3.

Lemma 4.2. *Let $\beta > 0$ and $\rho > 1$.*

1. *For all $n \geq 1$,*

$$\limsup_{d \rightarrow \infty} \mathbb{P}[\tau(\beta, \rho, d) \text{ survives}] - \mathbb{P}[\tau'(\beta, \rho, d) \text{ survives up to step } n] \leq 0.$$

2. *For all $n \geq 1$,*

$$\limsup_{d \rightarrow \infty} \mathbb{P}[\tau'(\beta, \rho, d) \text{ survives up to step } 2n] \leq \mathbb{P}[\tau''(\beta^2) \text{ survives up to step } n].$$

3. $\lim_{n \rightarrow \infty} \mathbb{P}[\tau''(\beta^2) \text{ survives up to step } n] = \mathbb{P}[\tau''(\beta^2) \text{ survives}].$

4. *For all $d \geq 1$, $\mathbb{P}[\tau'(\beta, \rho, d) \text{ survives}] \leq \mathbb{P}[\tau(\beta, \rho, d) \text{ survives}].$*

5. $\liminf_{d \rightarrow \infty} \mathbb{P}[\tau'(\beta, \rho, d) \text{ survives}] \geq \mathbb{P}[\tau''(\beta^2) \text{ survives}].$

Proof of Lemma 4.1 using Lemma 4.2. Fix $\beta > 0$ and $\rho > 1$. Let $\varepsilon > 0$. By Item 3 we fix n_0 such that

$$\mathbb{P}[\tau''(\beta^2) \text{ survives up to step } n_0] \leq \mathbb{P}[\tau''(\beta^2) \text{ survives}] + \varepsilon.$$

Combining Item 1 (with $n = 2n_0$) and Item 2 (with $n = n_0$), we get the existence of d_2 such that,

$$\forall d \geq d_2, \mathbb{P}[\tau(\beta, \rho, d) \text{ survives}] \leq \mathbb{P}[\tau''(\beta^2) \text{ survives up to step } n_0] + \varepsilon$$

and then

$$\forall d \geq d_2, \mathbb{P}[\tau(\beta, \rho, d) \text{ survives}] \leq \mathbb{P}[\tau''(\beta^2) \text{ survives}] + 2\varepsilon. \tag{4.2}$$

By Item 5 of Lemma 4.2, there exists d_1 such that,

$$\forall d \geq d_1, \mathbb{P}[\tau''(\beta^2) \text{ survives}] \leq \mathbb{P}[\tau'(\beta, \rho, d) \text{ survives}] + \varepsilon. \tag{4.3}$$

The result follows from (4.3), Item 4 of Lemma 4.2 and (4.2). \square

Proof of Item 1 of Lemma 4.2. Let $\beta > 0$ and $\rho > 1$. Hereafter, we assume that d is large enough to ensure

$$\beta \left(\frac{2\sqrt{\rho}}{1+\rho} \right)^d \leq \frac{1}{2}. \tag{4.4}$$

We couple in a natural way $\tau(\beta, \rho, d)$ and $\tau'(\beta, \rho, d)$. If $\tau(\beta, \rho, d)$ survives, then there exists an infinite branch $0 = x_0, x_1, x_2, \dots$ in $\tau(\beta, \rho, d)$. If moreover $\tau'(\beta, \rho, d)$ does not survive at least up to step n , then there exists $k \in \{1, \dots, n\}$ such that x_k and x_{k-1} are two particles of the same type. By (4.4), there exists no infinite branch of 1-particles. Therefore there exists $\ell \geq n$ such that x_ℓ is a ρ -particle. Let ℓ be the smallest such integer. The probability of the event

$$\{\tau(\beta, \rho, d) \text{ survives}\} \setminus \{\tau'(\beta, \rho, d) \text{ survives up to step } n\}$$

is thus bounded from above by the expected number of such paths (x_0, \dots, x_ℓ) .

In order to bound this expected number of paths, we will associate a type with each such path. For each $k \in \{0, \dots, \ell\}$ we set $t_k = 1$ if x_k is a 1-particle and $t_k = \rho$ otherwise. We thus get a sequence $T = (t_0, \dots, t_\ell)$ of types. This sequence belongs to the set \mathcal{T} of finite sequences $(t'_0, \dots, t'_{\ell'})$ such that

1. $\ell' \geq n$.
2. $t'_0 = t'_{\ell'} = \rho$. As a consequence, the sets $\{k \in \{1, \dots, \ell'\} : t'_{k-1} = 1 \text{ and } t'_k = \rho\}$ and $\{k \in \{1, \dots, \ell'\} : t'_{k-1} = \rho \text{ and } t'_k = 1\}$ have the same cardinality. This is a crucial property which will cancel a large factor later in the proof.
3. There exists $k \in \{1, \dots, n\}$ such that $t'_k = t'_{k-1}$.
4. If $\ell' > n$, then $t'_n = t'_{n+1} = \dots = t'_{\ell'-1} = 1$. This is due to the fact that we stop, after step n , with the first ρ -particle.

Let us fix $T = (t_0, \dots, t_\ell) \in \mathcal{T}$. The expected number of paths of the tree $\tau(\beta, \rho, d)$ whose type is T is $\beta^\ell \Delta^{I(T)}$ where

$$\Delta = \left(\frac{2\sqrt{\rho}}{1+\rho} \right)^d < 1$$

and

$$I(T) = \#\{k \in \{1, \dots, \ell\} : t_k = t_{k-1}\}.$$

We have used the second property of types which yields some cancellations in antidiagonal coefficients of M . By the third property and as $\ell \geq n$ we have $I(T) \geq 1$. By the fourth one we have $I(T) \geq \ell - 1 - n$. Therefore $I(T) \geq \max(1, \ell - 1 - n)$ and the expected number of paths of τ whose type is T is at most

$$\beta^\ell \Delta^{\max(1, \ell - 1 - n)}.$$

Summing over types, we get

$$\begin{aligned} & \mathbb{P}[\{\tau(\beta, \rho, d) \text{ survives}\} \setminus \{\tau'(\beta, \rho, d) \text{ survives at least up to step } n\}] \\ & \leq \sum_{T \in \mathcal{T}} \beta^\ell \Delta^{\max(1, \ell - 1 - n)}. \end{aligned}$$

The contribution of types of length $\ell = n$ is at most $2^n \beta^n \Delta$. The contribution of types of length $\ell = n + 1$ is at most $2^n \beta^{n+1} \Delta$. The contribution of types of length $\ell \geq n + 2$ is at most (we use $\beta \Delta \leq 1/2$, see (4.4)),

$$2^n \sum_{\ell \geq n+2} \beta^\ell \Delta^{\ell-1-n} = 2^n \beta^{n+1} \frac{\beta \Delta}{1 - \beta \Delta} \leq 2^{n+1} \beta^{n+1} \beta \Delta.$$

Therefore

$$\begin{aligned} & \mathbb{P}[\{\tau(\beta, \rho, d) \text{ survives}\} \setminus \{\tau'(\beta, \rho, d) \text{ survives at least up to step } n\}] \\ & \leq 2^n \beta^n (1 + \beta + 2\beta^2) \left(\frac{2\sqrt{\rho}}{1+\rho} \right)^d \end{aligned}$$

and then

$$\begin{aligned} & \mathbb{P}[\tau(\beta, \rho, d) \text{ survives}] \\ & \leq \mathbb{P}[\tau'(\beta, \rho, d) \text{ survives at least up to step } n] + 2^n \beta^n (1 + \beta + 2\beta^2) \left(\frac{2\sqrt{\rho}}{1+\rho} \right)^d. \end{aligned}$$

This yields the result. □

Proof of Item 2 of Lemma 4.2. Let $\beta > 0$, $\rho > 1$ and $n_0 \geq 1$. Let $d \geq 1$. Write

$$\mu = \beta \sqrt{\rho}^d \text{ and } \mu^* = \beta \frac{1}{\sqrt{\rho}^d}. \tag{4.5}$$

Let $N(\mu)$ be a Poisson(μ) random variable and $N_1(\mu^*), N_2(\mu^*), \dots$ be Poisson(μ^*) random variables. Assume that these random variables are independent. Set

$$X(\beta, \rho, d) = \sum_{i=1}^{N(\mu)} N_i(\mu^*). \tag{4.6}$$

The process $(\tau'_{2n}(\beta, \rho, d))_n$ is a Galton-Watson process with progeny distributed as $X(\beta, \rho, d)$. When d converges to ∞ , $X(\beta, \rho, d)$ converges in distribution to a Poisson(β^2) random variable $N(\beta^2)$. This can for example be shown by computing the characteristic functions: for all $t \in \mathbb{R}$,

$$\begin{aligned} \mathbb{E}[e^{itX(\beta, \rho, d)}] &= \exp \left[-\beta\sqrt{\rho}^d \left(1 - \exp \left(-\beta\sqrt{\rho}^{-d} (1 - e^{it}) \right) \right) \right] \\ &\rightarrow_{d \rightarrow \infty} \exp \left[-\beta^2 (1 - e^{it}) \right] \\ &= \mathbb{E}[e^{itN(\beta^2)}]. \end{aligned}$$

Thus, there exists a sequence of random variables $\tilde{X}(d)$, each of which has the same distribution of $X(\beta, \rho, d)$, and a random variable $\tilde{N}(\beta^2)$ with Poisson(β^2) distribution such that $\tilde{X}(d)$ converges almost surely to \tilde{N} . Let us use such coupling for all variables defining our Galton-Watson processes. We thus get a new version of $(\tau'_{2n}(\beta, \rho, d))_n$ – which we denote by $(\tilde{\tau}'_{2n}(\beta, \rho, d))_n$ – and a new version of $\tau''(\beta^2)$ – which we denote by $\tilde{\tau}''(\beta^2)$ – such that $(\tilde{\tau}'_{2n}(\beta, \rho, d))_{n \leq n_0}$ converges almost surely to $(\tilde{\tau}''(\beta^2))_{n \leq n_0}$ when d tends to ∞ . The almost sure convergence is due to the fact that, due to the coupling, the progeny of the root in the first process converges almost surely (and thus is almost surely equal for d large enough) to the progeny of the root in the second process, the progeny of each child of the root in the first process converges almost surely (and thus is almost surely equal for d large enough) to the progeny of the corresponding child in the second process, and so on. Therefore

$$\begin{aligned} &\mathbb{P}[\tau'(\beta, \rho, d) \text{ survives up to step } 2n] \\ &= \mathbb{P}[(\tau'_{2n}(\beta, \rho, d))_n \text{ survives up to step } n] \\ &= \mathbb{P}[(\tilde{\tau}'_{2n}(\beta, \rho, d))_n \text{ survives up to step } n] \quad \text{as they have the same distribution} \\ &\rightarrow \mathbb{P}[\tilde{\tau}''(\beta^2) \text{ survives up to step } n] \quad \text{as } d \rightarrow \infty \text{ by the discussion above} \\ &= \mathbb{P}[\tau''(\beta^2) \text{ survives up to step } n] \quad \text{as they have the same distribution.} \end{aligned}$$

This yields the result. □

Proof of Item 3 of Lemma 4.2. This is straightforward as the sequence of events is non-increasing. □

Proof of Item 4 of Lemma 4.2. This is straightforward by a natural coupling. □

Proof of Item 5 of Lemma 4.2. Let $\beta > 0$, $\rho > 1$ and $d \geq 1$. We will use some remarks and notations from the proof of Item 2, in particular (4.5) and (4.6). The process $(\tau'(\beta, \rho, d)_{2n})_n$ is a Galton-Watson process with progeny distributed as $X(\beta, \rho, d)$. But

$$X(\beta, \rho, d) \geq \sum_{i=1}^{N(\mu)} \mathbb{1}_{N_i(\mu^*) \geq 1}$$

which is $\text{Poisson}(\mu(1 - \exp(-\mu^*)))$ distributed. Therefore

$$\begin{aligned} & \mathbb{P}[\tau'(\beta, \rho, d) \text{ survives}] \\ &= \mathbb{P}[(\tau'(\beta, \rho, d)_{2n})_n \text{ survives}] \\ &= \mathbb{P}[\text{a Galton-Watson process with progeny distributed as } X(\beta, \rho, d) \text{ survives}] \\ &\geq \mathbb{P}[\text{a Galton-Watson process with Poisson}(\mu(1 - \exp(-\mu^*))) \text{ progeny survives}] \\ &= S(\mu(1 - \exp(-\mu^*))) \\ &\rightarrow S(\beta^2) \text{ as } d \rightarrow \infty \end{aligned}$$

as $\mu(1 - \exp(-\mu^*)) \rightarrow \beta^2$ and as S is continuous by Lemma 3.6. □

4.4 Proof of the upper bound on percolation probability

We use the framework of Section 4.1 and the first BRW defined in Section 4.2. The aim of this section is to prove the following result. This is the easy part in the control of percolation probability and percolation threshold.

Proposition 4.3. • Let $\beta > 0$. For any $\rho > 1$, any $d \geq 1$ and any $K \in \mathcal{K}(d)$,

$$\mathbb{P} \left[\#\widehat{C}^0(\beta, \rho, d, K) = \infty \right] \leq S(\beta, \rho, d). \tag{4.7}$$

• For any $\rho > 1$, any $d \geq 1$ and any $K \in \mathcal{K}(d)$,

$$\beta_c(\rho, d, K) \geq \frac{1}{1 + \left(\frac{2\sqrt{\rho}}{1+\rho}\right)^d}. \tag{4.8}$$

• Let $\beta > 0$ and $\rho > 1$. There exists a sequence $(\varepsilon(d))_d$ of positive real numbers which tends to 0 such that,

$$\forall d \geq 1, \forall K \in \mathcal{K}(d), \mathbb{P} \left[\#\widehat{C}^0(\beta, \rho, d, K) = \infty \right] \leq S(\beta^2) + \varepsilon(d).$$

• Let $\rho > 1$. There exists a sequence $(\varepsilon'(d))_d$ of positive real numbers which tends to 0 such that,

$$\forall d \geq 1, \forall K \in \mathcal{K}(d), \beta_c(\rho, d, K) \geq 1 - \varepsilon'(d).$$

Proof. Let $\beta > 0, \rho > 0, d \geq 1$ and $K \in \mathcal{K}(d)$. Denote by $\bar{\tau}$ be the associated BRW introduced in Section 4.2.1. In particular, the mean matrix of the associated two-type Galton-Watson process is $M(\beta, \rho, d)$ which is defined in (1.9). By the discussion of Section 4.2.1,

$$\mathbb{P} \left[\#\widehat{C}^0(\beta, \rho, d, K) = \infty \right] \leq \mathbb{P}[\#\bar{\tau} = \infty] = S(\beta, \rho, d).$$

This gives (4.7). The largest eigenvalue of $M(\beta, \rho, d)$ is

$$r = \beta \left[1 + \left(\frac{2\sqrt{\rho}}{1+\rho} \right)^d \right].$$

Therefore $S(\beta, \rho, d)$ equals 0 when $r \leq 1$ (see [1] page 186). Combined with the upper bound on the percolation probability, this yields (4.8).

Lemma 4.1 and (4.7) yield Item 3. Item 4 is a straightforward consequence of (4.8). □

4.5 Proof of the lower bound on percolation probability

This is the hard part in the control of percolation probability and percolation threshold. Our aim is to prove the following theorem.

Theorem 4.4. • Let $\beta > 0$ and $1 < \rho < 2$. There exists a sequence $(\varepsilon(d))_d$ of positive real numbers which tends to 0 such that,

$$\forall d \geq 1, \forall K \in \mathcal{K}(d), \mathbb{P} \left[\#\widehat{C}^0(\beta, \rho, d, K) = \infty \right] \geq S(\beta^2) - \varepsilon(d).$$

• Let $1 < \rho < 2$. There exists a sequence $(\varepsilon'(d))_d$ of positive real numbers which tends to 0 such that,

$$\forall d \geq 1, \forall K \in \mathcal{K}(d), \beta_c(\rho, d, K) \leq 1 + \varepsilon'(d).$$

Proof of Theorem 1.4 using Theorem 4.4. Recall Section 4.1 for notations. In particular, for all $\beta > 0, \rho > 1, d \geq 1$ and $K \in \mathcal{K}(d)$, by (1.2) and by our choice of notations (which we simplified in Section 4):

$$\mathbb{P}[\#\widehat{C}^0(\beta, \rho, d, K) = \infty] = \mathbb{P}[C^0(\beta\kappa_c(\rho)^d, \nu_{d,\rho}, d, K) \text{ is unbounded}].$$

Item 1 of Theorem 4.4, Item 1 of Proposition 4.3 and Lemma 4.1 yield Item 1 of Theorem 1.4. Item 2 of Theorem 4.4 and Item 4 of Proposition 4.3 yield Item 2 of Theorem 1.4. □

4.5.1 Good Gaps

Let $d \geq 1$ and $K \in \mathcal{K}(d)$. As usual, denote by X_K, X'_K, X''_K i.i.d.r.v. uniformly distributed on K . Let L any map provided by Theorem 2.1 for X_K and define \widehat{L} by $\widehat{L} = 2^{-1/2}L$. Let T be any map adapted to X_K (see (2.1)).

For all $\eta > 0$ smaller than $\sqrt{2}$ set

$$\begin{aligned} G(d, K, \eta) &= \{z \in \mathbb{R}^d : \mathbb{P}[z + X_K \notin K] \geq 1 - \eta\}, \\ H'(d, K, \eta) &= \{z \in \mathbb{R}^d : \|T(z)\|_2 d^{-1/2} < 3/2\}, \\ G'(d, K, \eta) &= \{z \in \mathbb{R}^d : \mathbb{P}[z + X_K + X'_K \notin H'(d, K, \eta)] \geq 1 - \eta\}, \\ H(d, K, \eta) &= \{z \in \mathbb{R}^d : \mathbb{P}[z + X_K \in K] \leq (\sqrt{2} - \eta)^{-d}\}. \end{aligned}$$

The family of sets G was already introduced in Section 3.4.1 with a similar purpose. The only property of $3/2$ we are interested in is that $\sqrt{2} < 3/2 < \sqrt{3}$. When the parameters are clear from the context, we will write G, G' and so on.

Lemma 4.5. Let $\eta \in (0, \sqrt{2})$. There exists a sequence $(\varepsilon'_G(d, \eta))_d$ that tends to 0 such that for all $d \geq 1$, all $K \in \mathcal{K}(d)$, all $a \in \mathbb{R}^d$:

$$\begin{aligned} \mathbb{P}[a + X_K \in G(d, K, \eta)] &\geq 1 - \varepsilon'_G(d, \eta), \\ \mathbb{P}[X_K + X'_K \in H'(d, K, \eta)] &\geq 1 - \varepsilon'_G(d, \eta), \\ \mathbb{P}[a + X_K \in G'(d, K, \eta)] &\geq 1 - \varepsilon'_G(d, \eta), \\ \mathbb{P}[X_K + X'_K \in H(d, K, \eta)] &\geq 1 - \varepsilon'_G(d, \eta). \end{aligned}$$

Proof. We have to prove that four probabilities tends to 1, uniformly in $K \in \mathcal{K}(d)$ and $a \in \mathbb{R}^d$, as d tends to ∞ . This is the content of Lemma 3.3 for $\mathbb{P}[a + X_K \in G(d, K, \eta)]$. By Theorem 2.2, we have the following convergence in probability when d tends to ∞ .

$$\frac{\|T(X_K + X'_K)\|_2}{\sqrt{d}} \rightarrow \sqrt{2} \text{ and } \frac{\|T(X_K + X'_K + X''_K)\|_2}{\sqrt{d}} \rightarrow \sqrt{3}.$$

The convergences are uniform in $K \in \mathcal{K}(d)$. The first convergence gives immediately the required result for $\mathbb{P}[X_K + X'_K \in H'(d, K, \eta)]$ as $\sqrt{2} < 3/2$. With some further work, the second convergence gives the required result for $\mathbb{P}[a + X_K \in G'(d, K, \eta)]$.

Indeed, for all $d \geq 1, a \in \mathbb{R}^d$ and $K \in \mathcal{K}(d)$,

$$\begin{aligned} \mathbb{P}[a + X_K \notin G'] &= \mathbb{P}[\mathbb{P}[a + X_K + X'_K + X''_K \in H' | X_K] \geq \eta] \\ &\leq \eta^{-1} \mathbb{P}[a + X_K + X'_K + X''_K \in H'] \quad \text{by Markov inequality} \\ &= \eta^{-1} \int_{\mathbb{R}^d} \mathbf{1}_{H'}(a + s) \mathbf{1}_K * \mathbf{1}_K * \mathbf{1}_K(s) \, ds \\ &= \eta^{-1} \mathbf{1}_{H'} * \mathbf{1}_K * \mathbf{1}_K * \mathbf{1}_K(a) \quad \text{by symmetry of } K. \end{aligned}$$

But H' and K are convex and symmetric. Therefore $\mathbf{1}'_{H'}$ and $\mathbf{1}_K$ are log-concave and symmetric. Hence, $\mathbf{1}_{H'} * \mathbf{1}_K * \mathbf{1}_K * \mathbf{1}_K$ is log-concave (see Section 2.2) and symmetric. As a consequence,

$$\mathbf{1}_{H'} * \mathbf{1}_K * \mathbf{1}_K * \mathbf{1}_K(a) \leq \mathbf{1}_{H'} * \mathbf{1}_K * \mathbf{1}_K * \mathbf{1}_K(0).$$

Therefore,

$$\begin{aligned} \mathbb{P}[a + X_K \notin G'] &\leq \eta^{-1} \mathbf{1}_{H'} * \mathbf{1}_K * \mathbf{1}_K * \mathbf{1}_K(0) \\ &= \eta^{-1} \mathbb{P}[X_K + X'_K + X''_K \in H'] \quad \text{by the same arguments} \\ &= \eta^{-1} \mathbb{P}[\|T(X_K + X'_K + X''_K)\|_2 d^{-1/2} < 3/2] \quad \text{by definition of } H'. \end{aligned}$$

With the above mentioned consequence of Theorem 2.2, we then get the required result about $\mathbb{P}[a + X_K \in G'(d, K, \eta)]$.

Let us now consider $\mathbb{P}[X_K + X'_K \in H(d, K, \eta)]$. For all $z \in \mathbb{R}^d$ we have, using symmetry of K ,

$$\mathbb{P}[z + X_K \in K] = \int_{\mathbb{R}^d} \mathbf{1}_K(z + x) \mathbf{1}_K(x) \, dx = \mathbf{1}_K * \mathbf{1}_K(z).$$

Therefore,

$$|\mathbb{R}^d \setminus H| = |\{z \in \mathbb{R}^d : \mathbf{1}_K * \mathbf{1}_K(z) \geq (\sqrt{2} - \eta)^{-d}\}| \leq (\sqrt{2} - \eta)^d \int_{\mathbb{R}^d} \mathbf{1}_K * \mathbf{1}_K = (\sqrt{2} - \eta)^d.$$

But then, by Theorem 2.3 (rearrangement inequality),

$$\begin{aligned} \mathbb{P}[X_K + X'_K \notin H] &= \int_{\mathbb{R}^d} \mathbf{1}_{\mathbb{R}^d \setminus H}(s) \mathbf{1}_K * \mathbf{1}_K(s) \, ds \\ &\leq \int_{\mathbb{R}^d} \mathbf{1}_{(\sqrt{2} - \eta)B}(s) \mathbf{1}_B * \mathbf{1}_B(s) \, ds \\ &= \mathbb{P}[\|X_B + X'_B\|_2 \leq (\sqrt{2} - \eta)] \end{aligned}$$

which tends to 0 by Lemma 2.5. □

4.5.2 Embedding of a two-dimensional lattice in \mathbb{R}^d

Let $d \geq 1$ and $K \in \mathcal{K}(d)$. Let L be any map given by Theorem 2.1 for X_K . Define \widehat{L} by $\widehat{L}(x) = 2^{-1/2}L(x)$. Let \mathcal{N} and \mathcal{N}' be two independent standard Gaussian random vectors in \mathbb{R}^2 . Under an appropriate coupling,

$$\mathbb{P}[L(X_K) \neq \mathcal{N}] \leq \varepsilon_{CLT}(d)$$

where ε_{CLT} is the sequence which appears in the statement of Theorem 2.1. Therefore

$$\mathbb{P}[\widehat{L}(X_K) \neq 2^{-1/2}\mathcal{N}] \leq \varepsilon_{CLT}(d). \tag{4.9}$$

Under an appropriate coupling,

$$\mathbb{P}[L(X_K + X'_K) \neq \mathcal{N} + \mathcal{N}'] \leq 2\varepsilon_{CLT}(d).$$

As $2^{-1/2}(\mathcal{N} + \mathcal{N}')$ has the same distribution as \mathcal{N} we then have, under a new coupling,

$$\mathbb{P}[\widehat{L}(X_K + X'_K) \neq \mathcal{N}] \leq 2\varepsilon_{CLT}(d). \tag{4.10}$$

Recall the definition of $\overline{\mathcal{L}}$ and $A(\cdot, \cdot)$ in Section 3.4.2. For any $(i, j) \in \overline{\mathcal{L}}$, we set

$$A_{\widehat{\mathcal{L}}}(i, j) = \widehat{L}^{-1}(A(i, j)).$$

The sets $A_{\widehat{\mathcal{L}}}(i, j)$ are pairwise disjoint. Moreover 0 belongs to $A_{\widehat{\mathcal{L}}}(0, 0)$.

As in the proof of Theorem 1.3, we will use this embedding of $\overline{\mathcal{L}}$ to compare the cluster of the origin to a supercritical percolation process on $\overline{\mathcal{L}}$.

4.5.3 An estimate about BRW

The aim of this section is to prove the following result. This is a consequence of Lemma 3.5. Recall the notations about BRW and related events such as $\text{Small}_\rho(\cdot, \cdot)$ in Section 2.8. In particular, note that the underlying Galton-Watson process of $\overline{\tau}^{\beta, \rho, d, X_K; \mathcal{S}}$ is a two-type Galton-Watson with matrix mean $M'(\beta, \rho, d)$ and not $M(\beta, \rho, d)$.

Lemma 4.6. *Let $\beta, \rho > 1$ and $\varepsilon > 0$. There exist $m, d_0, k, M \geq 1$ with k even such that, for all $d \geq d_0$ and all $K \in \mathcal{K}(d)$, the following properties hold where L is any map given by Theorem 2.1 for X_K and where $\widehat{L} = 2^{-1/2}L$,*

- For all $z \in A_{\widehat{\mathcal{L}}}(0, 0)$,

$$\mathbb{P} \left[\overline{\tau}_k^{\beta, \rho, d, X_K; \{z\}}(A_{\widehat{\mathcal{L}}}(1, 0)) \geq m \text{ and } \text{Small}_\rho \left(\overline{\tau}_{\leq k}^{\beta, \rho, d, X_K; \{z\}}, M \right) \right] \geq S(\beta^2) - \varepsilon.$$

- For all $(i, j) \in \overline{\mathcal{L}}$ and all subset $\mathcal{S} \subset A_{\widehat{\mathcal{L}}}(i, j)$ of cardinality m ,

$$\mathbb{P} \left[\overline{\tau}_k^{\beta, \rho, d, X_K; \mathcal{S}}(A_{\widehat{\mathcal{L}}}(i+1, j+1)) \geq m \text{ and } \overline{\tau}_k^{\beta, \rho, d, X_K; \mathcal{S}}(A_{\widehat{\mathcal{L}}}(i+1, j-1)) \geq m \right. \\ \left. \text{and } \text{Small}_\rho \left(\overline{\tau}_{\leq k}^{\beta, \rho, d, X_K; \mathcal{S}}, M \right) \right] \geq 1 - \varepsilon.$$

Proof. Let $\beta, \rho > 1$ and $\varepsilon > 0$. We can and will assume that ε small enough to ensure $S(\beta^2) - \varepsilon > 0$. By Lemma 3.6, we can choose $\eta > 0$ such that

$$S(\beta^2 - \eta) \geq S(\beta^2) - \varepsilon.$$

Applying Lemma 3.5 with $\lambda = \beta^2 - \eta$ and ε we get constants that we denote by m_2, k_2, M_2 . For all $d \geq 1$ write

$$\mu = \beta\sqrt{\rho}^d \text{ and } \mu^* = \beta \frac{1}{\sqrt{\rho}^d}.$$

Let d_1 be such that, for all $d \geq d_1$,

$$\mu(1 - \exp(-\mu^*)) \geq \beta^2 - \eta.$$

Let d_2 be such that, for all $d \geq d_2$, $\varepsilon_{CLT}(d) \leq \varepsilon/M_2$ where $(\varepsilon_{CLT}(d))_d$ is the sequence given by Theorem 2.1. Let M_3 be such that, for all $d \geq 1$, all $K \in \mathcal{K}(d)$ and all $\mathcal{S} \subset \mathbb{R}^d$ of cardinality at most m_2 ,

$$\mathbb{P}[\text{Small}_\rho \left(\overline{\tau}_{\leq 2k_2}^{\beta, \rho, d, X_K; \mathcal{S}}, M_3 \right)] \geq 1 - \varepsilon. \tag{4.11}$$

Let us check that this is possible. Denote by N_ρ the number of ρ -particles of $\bar{\tau}_{\leq 2k_2}^{\beta, \rho, d, X_K; \mathcal{S}}$. Denote by N_1 the number of 1-particles of $\bar{\tau}_{\leq 2k_2}^{\beta, \rho, d, X_K; \mathcal{S}}$. Then, for any M_3 ,

$$\mathbb{P}[N_\rho \geq M_3] \leq \frac{\mathbb{E}[N_\rho]}{M_3} \leq m_2 \frac{\sum_{i=0}^{k_2} (\mu\mu^*)^i}{M_3} = m_2 \frac{\sum_{i=0}^{k_2} \beta^{2i}}{M_3}$$

and

$$\mathbb{P}[N_1 \geq M_3\sqrt{\rho^d}] \leq \frac{\mathbb{E}[N_1]}{M_3\sqrt{\rho^d}} \leq m_2\mu \frac{\sum_{i=0}^{k_2-1} (\mu^*\mu)^i}{M_3\sqrt{\rho^d}} = m_2\beta\sqrt{\rho^d} \frac{\sum_{i=0}^{k_2-1} \beta^{2i}}{M_3\sqrt{\rho^d}} = m_2\beta \frac{\sum_{i=0}^{k_2-1} \beta^{2i}}{M_3}.$$

We can thus fix M_3 as stated above.

Finally set $d_0 = \max(d_1, d_2)$.

We now prove that the conclusion of the lemma holds with $m_2, d_0, 2k_2, M_3$. Let $d \geq d_0, K \in \mathcal{K}(d)$. Let L be any map associated with X_K by Theorem 2.1 and set $\widehat{L} = 2^{-1/2}L$. Let \mathcal{S} be a finite subset of \mathbb{R}^d .

Consider the BRW $(\bar{\tau}_{2n}^{\beta, \rho, d, X_K; \mathcal{S}})_n$ where we sample at even steps. Let us prove the following stochastic domination between two BRW⁹:

$$\left(\bar{\tau}_{2n}^{\beta, \rho, d, X_K; \mathcal{S}}\right)_n \geq \left(\bar{\tau}_n^{\beta^2 - \eta, X_K + X'_K; \mathcal{S}}\right)_n. \tag{4.12}$$

Prune $\bar{\tau}^{\beta, \rho, d, X_K; \mathcal{S}}$ in the following way. If a 1-particle has strictly more than one children, keep the first one in Neveu ordering (see Section 2.8) and remove all the other ones and their progeny. Denote by $\bar{\tau}^-$ the new BRW. We have the following stochastic domination:

$$\left(\bar{\tau}_n^{\beta, \rho, d, X_K; \mathcal{S}}\right)_n \geq \left(\bar{\tau}_n^-\right)_n.$$

and then

$$\left(\bar{\tau}_{2n}^{\beta, \rho, d, X_K; \mathcal{S}}\right)_n \geq \left(\bar{\tau}_{2n}^-\right)_n. \tag{4.13}$$

Consider the BRW on the right. The steps are independent copies of $X_K + X'_K$. The progeny is distributed as the sum of a Poisson(μ) number of independent Bernoulli($1 - \exp(-\mu^*)$) random variables. Therefore, the progeny is Poisson($\mu(1 - \exp(-\mu^*))$) distributed. As $d \geq d_0 \geq d_1$, the progeny stochastically dominates a Poisson($\beta^2 - \eta$) random variable. As a consequence $(\bar{\tau}_{2n}^-)_n$ stochastically dominates the BRW on the right of (4.12). With (4.13), this yields (4.12).

As $d \geq d_0 \geq d_2$ we have, under an appropriate coupling,

$$\mathbb{P}[\widehat{L}(X_K + X'_K) \neq \mathcal{N}] \leq 2\varepsilon/M_2. \tag{4.14}$$

Assume that $\mathcal{S} \subset \mathbb{R}^d$ is a set of cardinality at most m_2 . Let $(i', j') \in \widehat{\mathcal{L}}$. We have

$$\begin{aligned} & \mathbb{P} \left[\bar{\tau}_{k_2}^{\beta^2 - \eta, \mathcal{N}; \widehat{L}(\mathcal{S})}(A(i', j')) \geq m_2 \text{ and there are at most } M_2 \text{ particles in } \bar{\tau}_{\leq k_2}^{\beta^2 - \eta, \mathcal{N}; \widehat{L}(\mathcal{S})} \right] \\ & \leq \mathbb{P} \left[\bar{\tau}_{k_2}^{\beta^2 - \eta, \widehat{L}(X_K + X'_K); \widehat{L}(\mathcal{S})}(A(i', j')) \geq m_2 \right] + M_2 \frac{2\varepsilon}{M_2} \quad \text{by (4.14)} \\ & = \mathbb{P} \left[\bar{\tau}_{k_2}^{\beta^2 - \eta, X_K + X'_K; \mathcal{S}}(A_{\widehat{L}}(i', j')) \geq m_2 \right] + 2\varepsilon \\ & \leq \mathbb{P} \left[\bar{\tau}_{2k_2}^{\beta, \rho, d, X_K; \mathcal{S}}(A_{\widehat{L}}(i', j')) \geq m_2 \right] + 2\varepsilon \quad \text{by (4.12)}. \end{aligned} \tag{4.15}$$

⁹Note that we sample at even times on the left and at all times on the right. In particular, even on the left this is a one type BRW(the type is always ρ).

Therefore,

$$\begin{aligned} & \mathbb{P} \left[\bar{\tau}_{2k_2}^{\beta, \rho, d, X_K; \mathcal{S}}(A_{\widehat{L}}(i', j')) \geq m_2 \text{ and } \text{Small}_\rho \left(\bar{\tau}_{\leq 2k_2}^{\beta, \rho, d, X_K; \mathcal{S}}, M_3 \right) \right] \\ & \geq \mathbb{P} \left[\bar{\tau}_{2k_2}^{\beta, \rho, d, X_K; \mathcal{S}}(A_{\widehat{L}}(i', j')) \geq m_2 \right] - \varepsilon \quad \text{by (4.11)} \\ & \geq \mathbb{P} \left[\bar{\tau}_{k_2}^{\beta^2 - \eta, \mathcal{N}; \widehat{L}(\mathcal{S})}(A(i', j')) \geq m_2 \right. \\ & \quad \left. \text{and there are at most } M_2 \text{ particles in } \bar{\tau}_{\leq k_2}^{\beta^2 - \eta, \mathcal{N}; \widehat{L}(\mathcal{S})} \right] - 3\varepsilon \end{aligned}$$

where we used (4.15) in the last step.

We can now conclude using the definition of m_2, k_2, M_2 (which comes from Lemma 3.5 with $\lambda = \beta^2 - \eta$ and ε). If $\mathcal{S} = \{z\} \subset A_{\widehat{L}}(0, 0)$ we get, with $(i', j') = (1, 0)$ and the inequality obtained above,

$$\begin{aligned} & \mathbb{P} \left[\bar{\tau}_{2k_2}^{\beta, \rho, d, X_K; \{z\}}(A_{\widehat{L}}(1, 0)) \geq m_2 \text{ and } \text{Small}_\rho \left(\bar{\tau}_{\leq 2k_2}^{\beta, \rho, d, X_K; \{z\}}, M_3 \right) \right] \\ & \geq \mathbb{P} \left[\bar{\tau}_{k_2}^{\beta^2 - \eta, \mathcal{N}; \{\widehat{L}(z)\}}(A(1, 0)) \geq m_2 \right. \\ & \quad \left. \text{and there are at most } M_2 \text{ particles in } \bar{\tau}_{\leq k_2}^{\beta^2 - \eta, \mathcal{N}; \{\widehat{L}(z)\}} \right] - 3\varepsilon \\ & \geq S(\beta^2 - \eta) - 4\varepsilon \quad \text{by the choice of } m_2, k_2, M_2 \\ & \geq S(\beta^2) - 5\varepsilon \quad \text{by the choice of } \eta. \end{aligned}$$

Let $(i, j) \in \mathcal{L}$. If \mathcal{S} is a subset of cardinality m_2 of $A_{\widehat{L}}(i, j)$ we get similarly, with $(i', j') = (i + 1, j \pm 1)$,

$$\mathbb{P} \left[\bar{\tau}_{2k_2}^{\beta, \rho, d, X_K; \mathcal{S}}(A_{\widehat{L}}(i + 1, j \pm 1)) \geq m_2 \text{ and } \text{Small}_\rho \left(\bar{\tau}_{\leq 2k_2}^{\beta, \rho, d, X_K; \mathcal{S}}, M_3 \right) \right] \geq 1 - 4\varepsilon.$$

Therefore,

$$\begin{aligned} & \mathbb{P} \left[\bar{\tau}_{2k_2}^{\beta, \rho, d, X_K; \mathcal{S}}(A_{\widehat{L}}(i + 1, j + 1)) \geq m_2 \text{ and } \bar{\tau}_{2k_2}^{\beta, \rho, d, X_K; \mathcal{S}}(A_{\widehat{L}}(i + 1, j - 1)) \geq m_2 \right. \\ & \quad \left. \text{and } \text{Small}_\rho \left(\bar{\tau}_{\leq 2k_2}^{\beta, \rho, d, X_K; \mathcal{S}}, M_3 \right) \right] \geq 1 - 8\varepsilon. \end{aligned}$$

Items 1 and 2 hold with the choice of parameters $m_2, d_0, 2k_2, M_3$. □

4.5.4 Plan and intuition

The aim of this section is to present in an informal way the plan of the proof. This a refinement of the plan given in Section 3.4.4 in the constant radius case. We assume that the reader is familiar with the content of Section 3.4.4.

Setup. Let $\beta > 0, 1 < \rho < 2, \varepsilon > 0, d \geq 1$ and $K \in \mathcal{K}(d)$. As usual, we denote by X_K and X'_K two independent random variables with uniform distribution on K . Recall the definition of $\widehat{C}^0 = \widehat{C}^0(\beta, \rho, d, K)$ and $\widehat{A}^0 = \widehat{A}^0(\beta, \rho, d, K)$ in Section 4.1. Recall that $S(\beta^2)$ denotes the survival probability of a Poisson(β^2) offspring Galton-Watson process. The aim is to prove that the inequality

$$\mathbb{P}[\#\widehat{C}^0(\beta, \rho, d, K) = \infty] \geq S(\beta^2) - \varepsilon$$

holds for any d large enough, uniformly in $K \in \mathcal{K}(d)$. We actually prove that for any d large enough, uniformly in $K \in \mathcal{K}(d)$,

$$\mathbb{P}[\#\widehat{A}^0(\beta, \rho, d, K) = \infty] \geq S(\beta^2) - \varepsilon.$$

This is a stronger result as \widehat{A}^0 is a subset of \widehat{C}^0 . The advantage is that the structure of \widehat{A}^0 is easier. The idea of focusing on \widehat{A}^0 instead of \widehat{C}^0 was already used in [6]. However, it was used in a cruder way. Recall that the main result in [6] is a logarithmic equivalent of the critical parameter in the Euclidean case, stated here as Theorem 1.2.

The set \widehat{A}^0 can be built from the BRW $\overline{\tau}^{\beta, \rho, d, X_K}$. See Section 2.8 for notations on BRW and Section 4.2.2 for the construction of \widehat{A}^0 from $\overline{\tau}^{\beta, \rho, d, X_K}$. Note in particular that, in $\overline{\tau}^{\beta, \rho, d, X_K}$, children of ρ -particles are 1-particles and children of 1-particles are ρ -particles. The basic intuition is that, up to an event whose probability vanishes when d tends to ∞ , \widehat{A}^0 is infinite when $\overline{\tau}^{\beta, \rho, d, X_K}$ is infinite.

The underlying Galton-Watson tree. The underlying Galton-Watson tree of the BRW $\overline{\tau}^{\beta, \rho, d, X_K}$ does not depend on K but it depends on d . The number of children of a ρ -particle tends to ∞ when d tends to ∞ . Indeed, this is a Poisson random variable with parameter $\beta\sqrt{\rho}^d$. Thus, the Galton-Watson process underlying $\overline{\tau}^{\beta, \rho, d, X_K}$ degenerates when d tends to ∞ . This is a major difference with respect to the constant radius case and this is why the non constant case is much more involved. It is also for this reason that Theorem 1.4 does not hold for $\rho > 2$ (the explosion of the number of children of ρ -particles increases when ρ increases; see [7] where a logarithmic equivalent for the critical probability is given in the Euclidean case for every ρ : the behavior is not given by the sole underlying Galton-Watson process when $\rho > 2$). There is however a nice feature. When d tends to ∞ ,

$$\left(\overline{\tau}_{2n}^{\beta, \rho, d, X_K}\right)_n \approx \left(\overline{\tau}_n^{\beta^2, d, X_K + X'_K}\right)_n. \tag{4.16}$$

(See (4.12).) Note that the first BRW is sampled at even times. The second BRW is similar to the BRW studied in Section 3. We simply replaced λ by β^2 and X_K by $X_K + X'_K$. We can therefore use many results from Section 3. The underlying Galton-Watson process of the second BRW only depends on β^2 . This is a key property. More precisely this is a Poisson(β^2) offspring Galton-Watson process.

The plan is to prove that, asymptotically when d tends to ∞ , if the second BRW survives, which occurs with probability $S(\beta^2)$, then \widehat{A}^0 is infinite.

Control of the interference between BRW. The basic plan is the same as in the constant radius case. In particular, we use a two-step approach (see Section 3.4.4: make sure that the relevant gaps are good and then use this fact to control interference). However, the number of 1-particles of the BRW explodes when d tends to ∞ and each of them generates a region of interference. Moreover, many of them (and their progeny) have to be rejected because of interference but this point is not a real issue as most of them have no progeny at all. As the number of ρ -particles is well behaved when d tends to ∞ , the basic idea is to focus as much as possible on ρ -particles.

We perform over-pruning (see Section 3.2.3) in order to control rejection by interference. This means that we reject particles and their progeny if they break one the following rules. The constants M and Λ are large and the constant η is small. The map \widehat{L} is associated with X_K . See Section 4.5.2.

1. If y is a children of x , then $\|\widehat{L}(V(y) - V(x))\|_2 \leq \Lambda$. (Recall that this means that we reject y and its progeny if $\|\widehat{L}(V(y) - V(x))\|_2 > \Lambda$. Similar remarks apply below.)
2. There are no more than M ρ -particles at each stage and each ρ -particle has no more than $M\sqrt{\rho}^d$ children (which are 1-particles).
3. If x and x' are two distinct ρ -particles, then

$$\|\widehat{L}(V(x') - V(x))\|_2 \geq 4\Lambda \text{ or } V(x') - V(x) \in G(d, K, \eta) \cap G'(d, K, \eta).$$

Good Gaps are defined in Section 4.5.1.

4. If z is a ρ -particle and if x is his grandparent, then

$$V(z) - V(x) \in H(d, K, \eta) \cap H'(d, K, \eta).$$

Note that if y is the parent of z , the above property implies that the interference region of y is included in

$$V(x) + H'(d, K, \eta).$$

Note that Properties 3 and 4 depend only on ρ -particles which are not too numerous. This is why Lemma 4.5 enables us to prove that the extra-pruning required to get these properties is harmless. However these properties freely give some further properties on 1-particles (see for example the comment after the statement of Property 4).

Thanks to the above properties, the probability of interference will be small. Let us explain the general ideas. Recall that there can be interference only between a 1-particle and a ρ -particle and that we want to avoid rejection of ρ -particles (and thus of 1-particles with at least one child). Note that the number of 1-particles with at least one child is bounded by the number of ρ -particles which itself is controlled uniformly in d .

- Rejection of a 1-particle y . Denote by x its parent. This is a ρ -particle. Let x' be another ρ -particle. Let us consider rejection of y because of interference with x' .
 - If $\|\widehat{L}(V(x') - V(x))\|_2 \geq 2\Lambda$, then $x + K \cap \widehat{L}^{-1}(\Lambda D)$ and $x' + K \cap \widehat{L}^{-1}(\Lambda D)$ are disjoint. But y belongs to $x + K \cap \widehat{L}^{-1}(\Lambda D)$ and the interference region of x' is included in $x' + K \cap \widehat{L}^{-1}(\Lambda D)$. Therefore y can not be rejected because of interference with x' .
 - Otherwise $V(x) - V(x') \in G(d, K, \eta)$. Write $V(y) - V(x') = (V(y) - V(x)) + (V(x) - V(x'))$. By definition of $G(d, K, \eta)$, we see that the probability, conditioned on everything but $V(y) - V(x)$, that y is rejected because of interference with x' is at most η .
- Rejection of a ρ -particle z . Let y be its parent (it is a 1-particle) and x its grandparent (it is a ρ -particle). Let x' be a ρ -particle different from z but which can be x . We are wondering if z can be rejected because of interference with a child y' of x' . The number of 1-children of a ρ -individual diverges as d tends to ∞ . Therefore we want, as much as possible, to avoid considering each child y' of x' individually. In other words we want, as much as possible, to consider only ρ -particles.
 - If $\|\widehat{L}(V(x') - V(x))\|_2 \geq 4\Lambda$, then for any child y' of x' , $\|\widehat{L}(V(y') - V(z))\|_2 \geq \Lambda$ and therefore z can not be rejected because of y' .
 - Otherwise and if x and x' are distinct, then $V(x) - V(x') \in G'(d, K, \eta)$. Write $V(z) - V(x') = (V(z) - V(y)) + (V(y) - V(x)) + (V(x) - V(x'))$. Recall that the interference region of the 1-particle y' is included in $V(x') + H'(d, K, \eta)$. Therefore, by definition of $G'(d, K, \eta)$, the probability, conditioned on everything but $(V(z) - V(y))$ and $(V(y) - V(x))$, that *there exists a child y' of x' such that z is rejected because of interference with y'* is at most η . Note that we do not have to take into account the number of children of x' as we only use the fact that the interference region of *any child y' of x'* is included in $V(x') + H'(d, K, \eta)$. This is a key fact as the number of 1-children explodes when d tends to ∞ .
 - Otherwise, $x = x'$. This is the crucial case where the assumption $\rho < 2$ is needed. We have $V(z) - V(x) \in H(d, K, \eta)$ and thus $V(x) - V(z) \in H(d, K, \eta)$. Therefore, by definition of $H(d, K, \eta)$, the probability, conditioned on everything but $V(y') - V(x')$, that a given child y' of x (with $y' \neq y$) is responsible

for the rejection of z is at most $(\sqrt{2} - \eta)^{-d}$. But x has no more than $M\sqrt{\rho}^d$ children. Therefore the probability that there exists a child y' of $x' = x$ such that z is rejected because of interference with y' is at most $M\sqrt{\rho}^d(\sqrt{2} - \eta)^{-d}$ which is smaller than η for d large enough, provided that $\sqrt{\rho} < \sqrt{2} - \eta$.

4.5.5 Construction of a subset of $\frac{2}{1+\rho}\widehat{A}^0$ related to an oriented percolation on \mathcal{L}

Choice of parameters. Fix $\beta > 1$, $2 > \rho > 1$ and $\varepsilon > 0$. Fix $m, d_1, k, M \geq 1$ with k even as provided by Lemma 4.6 for the parameters β, ρ and ε . Let $\Lambda \geq 1$ be such that $\mathbb{P}[\|\mathcal{N}\|_2 > 2^{1/2}\Lambda] \leq \varepsilon/(2M)$. Fix $\eta \in (0, \sqrt{2})$ such that

$$400k^2\Lambda^2M^2\eta \leq \varepsilon \tag{4.17}$$

and

$$\frac{\sqrt{\rho}}{\sqrt{2} - \eta} < 1. \tag{4.18}$$

Recall the definition of $\varepsilon'_G(d, \eta)$ in Lemma 4.5. Let d_2 be such that, for all $d \geq d_2$,

$$400k^2\Lambda^2M^2\varepsilon'_G(d, \eta) \leq \varepsilon \tag{4.19}$$

and

$$M^2\sqrt{\rho}^d(\sqrt{2} - \eta)^{-d} \leq \varepsilon. \tag{4.20}$$

Let d_3 be such that, for all $d \geq d_3$, for all $K \in \mathcal{K}(d)$, for any L associated with X_K by Theorem 2.1, $\mathbb{P}[L(X_K) \neq \mathcal{N}] \leq \varepsilon/(2M)$. Then, with $\widehat{L} = 2^{-1/2}L$,

$$\begin{aligned} \mathbb{P}[\|\widehat{L}(X_K)\|_2 > \Lambda] &\leq \mathbb{P}[L(X_K) \neq \mathcal{N}] + \mathbb{P}[\|\mathcal{N}\|_2 > 2^{1/2}\Lambda] \\ &\leq \varepsilon/M. \end{aligned} \tag{4.21}$$

Let $d_0 = \max(d_1, d_2, d_3)$.

Setting and aim. Let $d \geq d_0$, $K \in \mathcal{K}(d)$. As usual, we denote by X_K and X'_K independent random variables with uniform distribution on K . Let L be any map associated with X_K by Theorem 2.1 for X_K . Define \widehat{L} by $\widehat{L}(x) = 2^{-1/2}L(x)$ as above. We aim at proving

$$\mathbb{P}[\#\widehat{A}^0 = \infty] \geq (\mathbf{S}(\beta^2) - 8\varepsilon)\theta(1 - \varepsilon)$$

where θ is defined in Section 3.4.2. From this result and Proposition 4.3, Theorem 4.4 will follow easily.

Thanks to the choice of parameters, the following properties hold (in particular the first ones are due to the choice of $m, d_1, k, M \geq 1$ using Lemma 4.6):

- For all $z \in \widehat{A}_L(0, 0)$,

$$\mathbb{P}\left[\overline{\tau}_k^{\beta, \rho, d, X_K; \{z\}}(\widehat{A}_L(1, 0)) \geq m \text{ and } \text{Small}_\rho\left(\overline{\tau}_{\leq k}^{\beta, \rho, d, X_K; \{z\}}, M\right)\right] \geq \mathbf{S}(\beta^2) - \varepsilon. \tag{4.22}$$

- For all $(i, j) \in \mathcal{L}$ and all subset $\mathcal{S} \subset \widehat{A}_L(i, j)$ of cardinality m ,

$$\begin{aligned} \mathbb{P}\left[\overline{\tau}_k^{\beta, \rho, d, X_K; \mathcal{S}}(\widehat{A}_L(i+1, j+1)) \geq m \text{ and } \overline{\tau}_k^{\beta, \rho, d, X_K; \mathcal{S}}(\widehat{A}_L(i+1, j-1)) \geq m \right. \\ \left. \text{and } \text{Small}_\rho\left(\overline{\tau}_{\leq k}^{\beta, \rho, d, X_K; \mathcal{S}}, M\right)\right] \geq 1 - \varepsilon. \end{aligned} \tag{4.23}$$

- (4.17), (4.19), (4.20) and (4.21).

Randomness and σ -fields. Let

$$(\bar{\tau}^{i,j,n})_{(i,j,n) \in \bar{\mathcal{L}} \times \{1, \dots, m\}}$$

be a family of independent copies of $\bar{\tau}^{\beta, \rho, d, X_K}$. Let $(\alpha^{i,j})_{(i,j) \in \bar{\mathcal{L}}}$ be a family of i.i.d. Bernoulli random variables with parameter $1 - \varepsilon$. For all $(i, j) \in \bar{\mathcal{L}}$ we denote by $\mathcal{F}_{i,j}$ (resp. $\mathcal{F}_{i,j}^-$) the σ -field generated by the $\bar{\tau}^{i',j',n}$ and the $\alpha^{i',j'}$ for $(i', j', n) \in \bar{\mathcal{L}} \times \{1, \dots, m\}$ such that (i', j') is smaller (resp. strictly smaller) than (i, j) for the lexicographic order. Initially, the site $(0, 0)$ is active and all other sites are inactive.

Stage $(0, 0)$. This stage is slightly different and slightly less involved than the next stages. The difference is similar to the difference in the corresponding construction given in Section 3. Here, in order to avoid lengthy repetitions, we only give the construction for sites $(i, j) \in \mathcal{L}$ and quickly explain by footnotes the modifications needed for the site $(0, 0)$. We hope that this is clear but, if this is not the case, we refer the reader to Section 3.

Stage (i, j) for $(i, j) \in \mathcal{L}$. Recall that we now consider successively each $(i, j) \in \mathcal{L}$ by lexicographic order. If (i, j) is inactive we decide, independently of everything else, that it is open with probability $1 - \varepsilon$ and closed otherwise. More precisely, we decide that (i, j) is open when $\alpha^{i,j} = 1$ and closed otherwise.

Thereafter, we consider the case where (i, j) is active. The set $\mathcal{S}(i, j)$ is well defined. It's a subset of cardinal m of $A_{\bar{\mathcal{L}}}(i, j)$. List the points of $\mathcal{S}(i, j)$ in an arbitrary order: $\mathcal{S}(i, j) = \{x^1, \dots, x^m\}$. We consider the m BRW $\bar{\tau}^n = x^n + \bar{\tau}^{i,j,n}$ where $x^n + \bar{\tau}^{i,j,n}$ designates the BRW $\bar{\tau}^{i,j,n}$ in which x^n has been added to the position of all the particles. We gather these m BRW into a single BRW originating from $\mathcal{S}(i, j)$. We denote it by $\bar{\tau}^{\mathcal{S}(i,j)}$ ¹⁰.

1. We examine successively the particles of generation between 1 ¹¹ and k of $\bar{\tau}^{\mathcal{S}(i,j)}$ in any admissible order. The last requirement means that:

- Children are examined after their parents.
- Children of a given parent are examined in a row: once we start examining the children of one parent, we then examine all the children of this parent.

Some of the particles will be rejected. We will reject more particles than necessary, thus performing some over-pruning as explained in Section 4.2. Some particles can be rejected before examination (if we have examined and rejected one of its ancestor). If we have examined a particle and not rejected it, we say that the particle has been “generated”. If we are examining a particle x and talk about a particle y generated before x , it means that one of the following two properties occur:

- y has been examined at stage (i, j) strictly before x and y has not been rejected.
- y has been examined at a previous stage (i', j') (thus $(i', j') < (i, j)$ in lexicographical order) and y has not been rejected ¹².

¹⁰In Stage $(0, 0)$, $(0, 0)$ is always active and we simply consider the BRW $\bar{\tau}^{0,0,1}$ originating from $\{0\}$.

¹¹In Stage $(0, 0)$ we also examine the particle of generation 0.

¹²This property never occurs if $(i, j) = 0$.

We never examine the roots of this BRW (note that the positions of the roots are also positions of particles of BRW examined during one of the previous stages¹³). We stop as soon as we have examined all the particles or as soon as the particle examined causes a stop by “Overpopulation” (equivalently, if “Overpopulation” occurs, we reject all the remaining particles). When examining a particle x , several things can occur:

- (a) Overpopulation. This occurs if x is the M -th ρ -particle examined during this stage (i, j) or if x is the $\lceil M\sqrt{\rho^d} \rceil$ -th children examined of a given ρ -particle. If “Overpopulation” occurs, we reject x and all the particles of $\bar{\tau}^{S(i,j)}$ which have not been examined yet. This stops the examination for this stage.
- (b) Bad Gap. This occurs if one of the following three conditions occurs:
 - x is a ρ -particle and there exists a ρ -particle x' generated before x such that:

$$\|\widehat{L}(V(x') - V(x))\|_2 < 4\Lambda \text{ and } V(x) - V(x') \notin G(d, K, \eta) \cap G'(d, K, \eta).$$

- x is a ρ -particle and the grand parent x' of x ¹⁴ is such that

$$V(x) - V(x') \notin H(d, K, \eta) \cap H'(d, K, \eta).$$

- The following condition holds¹⁵:

$$\|\widehat{L}(V(x) - V(\bar{x}))\|_2 > \Lambda.$$

In this case we do *not* stop the examination. However, we reject x and all its progeny¹⁶ and we do not consider rejection of x by interference (see below). This over-pruning will help us controlling the interference. Let us note a few consequences right now, as this is necessary to explain the paragraph “interference” below.

- If x' and y' are two generated particles such that x' is the parent of y' , then $\|\widehat{L}(V(y') - V(x'))\|_2 \leq \Lambda$.
- The interference region (see (3.2)) of a generated particle y' is included in

$$V(y') + K \cap \widehat{L}^{-1}(\Lambda D)$$

where D is the unit disk of \mathbb{R}^2 . If moreover y' is a 1-particle with parent x' , then the interference region of y' is included in

$$V(x') + H'(d, K, \eta). \tag{4.24}$$

- (c) Interference. It occurs if there exists a particle y' generated before x , such that y' is not the parent \bar{x} of x and such that one of the following conditions occurs (see the remark on the interference region in the paragraph “Bad Gap”):

¹³In Stage $(0, 0)$ we examine the root, even if there is nothing to examine. This is just a convention which ensures that the root is generated.

¹⁴In Stage $(0, 0)$ the condition is empty for the root 0.

¹⁵In Stage $(0, 0)$ the condition is empty for the root 0.

¹⁶In Section 3 we stopped the examination as soon as we saw a “Bad Gap”. Here we can not avoid some particles to be rejected because of “Bad Gap”. However, as we will see, we can avoid (with high probability) rejection of ρ -particle because of “Bad Gap” (note that we do not care about rejection of 1-particles without children). It will be sufficient.

- x is a 1-particle, y' is a ρ -particle and

$$V(x) \in V(y') + K \cap \widehat{L}^{-1}(\Lambda D).$$

- x is a ρ -particle, y' is a 1-particle and

$$V(x) \in V(y') + K \cap \widehat{L}^{-1}(\Lambda D) \text{ and } V(x) \in V(y') + H'(d, K, \eta).$$

In this case we do *not* stop the examination. However, we reject the particle x and all its progeny¹⁷.

One defines the following sets.

- The set $\mathcal{G}(i, j)$ of positions of all particles examined and not rejected. We call them the particles generated at stage (i, j) . By an abuse of notation, we see $\mathcal{G}(i, j)$ as inheriting the genealogical structure of the BRW.
 - In the coupling with the Boolean model, all the points of $\mathcal{G}(i, j)$ belong to $\frac{2}{1+\rho}A^0$. Indeed, none of them was rejected (see also the definition and properties of the seeds $\mathcal{S}(\cdot, \cdot)$ below¹⁸).

$$\text{The image by } \widehat{L} \text{ of } \mathcal{G}(i, j) \text{ is included in } (i, j) + 2k\Lambda D. \tag{4.25}$$

This is due to the fact that the images by \widehat{L} of the positions of the roots belong to $(i, j) + D$ (see definition of $\mathcal{S}(\cdot, \cdot)$ below¹⁹), the fact that the image by \widehat{L} of each step leading to a non rejected particle belongs to ΛD , the fact that we only explored the first k generations and the fact that $1 + k\Lambda \leq 2k\Lambda$.

- The subset $\mathcal{G}^\rho(i, j) \subset \mathcal{G}(i, j)$ of positions of ρ -particles generated and the subset $\mathcal{G}^1(i, j) \subset \mathcal{G}(i, j)$ of positions of 1-particles generated.
 - All the gaps between two distinct points x and x' of

$$\bigcup_{(i', j') \leq (i, j)} \mathcal{G}^\rho(i', j')$$

satisfy

$$\|\widehat{L}(V(x)) - \widehat{L}(V(x'))\|_2 \geq 4\Lambda \text{ or } V(x) \in V(x') + G(d, K, \eta) \cap G'(d, K, \eta). \tag{4.26}$$

- If moreover x' is the grand-parent of x , then

$$V(x) - V(x') \in H(d, K, \eta) \cap H'(d, K, \eta). \tag{4.27}$$

- $\mathcal{G}^\rho(i, j)$ contains at most M points.
- Any point in $\mathcal{G}^\rho(i, j)$ has at most $M\sqrt{\rho}^d$ children in $\mathcal{G}^1(i, j)$.

Indeed, none of them was rejected because of “Bad Gap” or “Overpopulation”.

- The set $\mathcal{G}_k(i, j) \subset \mathcal{G}^\rho(i, j)$ of positions of generated particles of generation k . Note that since k is even and since we started with ρ -particles, the particles of $\mathcal{G}_k(i, j)$ are ρ -particles.

¹⁷This is again a difference with respect to Section 3. The reason is the same as for “Bad Gap”.

¹⁸In stage $(0, 0)$, the set of seeds is $\{0\}$ which belong to $\frac{2}{1+\rho}A^0$.

¹⁹In stage $(0, 0)$, the root is located at 0.

2. If

$$\#(\mathcal{G}_k(i, j) \cap A_{\widehat{\mathcal{L}}}(i + 1, j + 1)) \geq m \text{ and } \#(\mathcal{G}_k(i, j) \cap A_{\widehat{\mathcal{L}}}(i + 1, j - 1)) \geq m \quad (4.28)$$

then²⁰:

- We say that the site (i, j) is open.
- If the site $(i + 1, j + 1)$ is inactive then we say that it is henceforth active and we define $\mathcal{S}(i + 1, j + 1)$ as the m first particles of $\mathcal{G}_k(i, j) \cap A_{\widehat{\mathcal{L}}}(i + 1, j + 1)$ in an arbitrary order. In the coupling with the Boolean model, all the points of $\mathcal{S}(i + 1, j + 1)$ belong to $\frac{2}{1+\rho} \widehat{A}^0$ ²¹.
- If the site $(i + 1, j - 1)$ is inactive then we say that it is henceforth active and we define $\mathcal{S}(i + 1, j - 1)$ as the m first particles of $\mathcal{G}_k(i, j) \cap A_{\widehat{\mathcal{L}}}(i + 1, j - 1)$ in an arbitrary order. In the coupling with the Boolean model, all the points of $\mathcal{S}(i + 1, j - 1)$ belong to $\frac{2}{1+\rho} \widehat{A}^0$ ²².

Otherwise, we say that the site (i, j) is closed.

4.5.6 Bounds on conditional probabilities

The aim of this section is to prove the following lemmas. Parameters have been fixed in Section 4.5.5.

Lemma 4.7. For all $(i, j) \in \overline{\mathcal{L}}$, $\{(i, j) \text{ is active}\} \in \mathcal{F}^-(i, j)$ and $\{(i, j) \text{ is open}\} \in \mathcal{F}(i, j)$.

Lemma 4.8. We have $\mathbb{P}[(0, 0) \text{ is open}] \geq S(\beta^2) - 8\varepsilon$.

Lemma 4.9. For all $(i, j) \in \mathcal{L}$, $\mathbb{P}[(i, j) \text{ is open} | \mathcal{F}^-(i, j)] \geq 1 - 8\varepsilon$.

Proof of Lemma 4.7. This is straightforward by construction. □

The proof of Lemma 4.8 is less involved than the proof of Lemma 4.9. The difference is similar to the difference in the proofs of Lemmas 3.9 and 3.10. There are moreover similarities in the proofs of the four lemmas. Therefore, in order to avoid lengthy repetitions, we only give a somewhat sketchy proof of Lemma 4.9 and quickly explain in footnotes the modifications needed for the proof of Lemma 4.8. For more details, we refer the reader to the proofs of Lemmas 3.9 (which is the more detailed) and Lemma 3.10.

Proof of Lemma 4.9. Let $(i, j) \in \mathcal{L}$. The event $\{(i, j) \text{ is active}\}$ is $\mathcal{F}^-(i, j)$ measurable. Therefore, we have to show

$$\mathbb{P}[(i, j) \text{ open} | \mathcal{F}^-(i, j)] \geq 1 - 8\varepsilon \text{ on the event } \{(i, j) \text{ is active}\}$$

and

$$\mathbb{P}[(i, j) \text{ open} | \mathcal{F}^-(i, j)] \geq 1 - 8\varepsilon \text{ on the event } \{(i, j) \text{ is inactive}\}$$

The second property is straightforward. Indeed, when (i, j) is inactive, (i, j) has been defined as open independently of everything else with probability $1 - \varepsilon$. Let us prove the first property.

²⁰In Stage $(0, 0)$, the condition is

$$\#(\mathcal{G}_k(0, 0) \cap A_{\widehat{\mathcal{L}}}(1, 0)) \geq m. \quad (4.29)$$

²¹In Stage $(0, 0)$, we say that the site $(1, 0)$ is active and we define $\mathcal{S}(1, 0)$ as the m first particles of $\mathcal{G}_k(0, 0) \cap A_{\widehat{\mathcal{L}}}(1, 0)$ in an arbitrary order. In the coupling with the Boolean model, all the points of $\mathcal{S}(1, 0)$ belong to $\frac{2}{1+\rho} \widehat{A}^0$.

²²In Stage $(0, 0)$ this item does not exist.

Below, we implicitly work on the event $\{(i, j) \text{ is active}\}$ and probabilities are always conditional to $\mathcal{F}^-(i, j)$. Therefore, there is a well defined set $\mathcal{S}(i, j)$ with cardinality m whose points are the starting points of m BRW. This set is measurable with respect to $\mathcal{F}^-(i, j)$. We also have the BRW $\bar{\tau}^{\mathcal{S}(i, j)}$ which has been used in Stage (i, j) ²³. Moreover

$$\{(i, j) \text{ is closed}\} \subset \text{BadGap}_\rho \cup \text{Interference}_\rho \cup \text{Small}_\rho \left(\bar{\tau}^{\mathcal{S}(i, j)}, M \right)^c \cup \text{Else}$$

where

$$\begin{aligned} \text{BadGap}_\rho &= \{\text{a } \rho\text{-particle has been rejected directly or indirectly because of Bad Gap}\} \\ &\cap \text{Small}_\rho \left(\bar{\tau}^{\mathcal{S}(i, j)}, M \right), \end{aligned}$$

and

$$\begin{aligned} \text{Interference}_\rho &= \{\text{a } \rho\text{-particle has been rejected directly or indirectly because of Interference}\} \\ &\cap \text{Small}_\rho \left(\bar{\tau}^{\mathcal{S}(i, j)}, M \right) \end{aligned}$$

and where²⁴

$$\text{Else} = \{ \bar{\tau}_k^{\mathcal{S}(i, j)}(A_{\hat{L}}(i+1, j+1)) < m \text{ or } \bar{\tau}_k^{\mathcal{S}(i, j)}(A_{\hat{L}}(i+1, j-1)) < m \}.$$

By “a ρ -particle x has been rejected directly or indirectly” we means that x has been rejected or that one of its ancestors has been rejected. The inclusion is due to the fact that, if (i, j) is active but $\text{BadGap}_\rho \cup \text{Interference}_\rho \cup \text{Small}_\rho \left(\bar{\tau}^{\mathcal{S}(i, j)}, M \right)^c$ does not occur, then $\mathcal{G}_k(i, j)$ is the set of positions of the particles of $\bar{\tau}_k^{\mathcal{S}(i, j)}$ (which are ρ particles).

Let us provide an upper bound for the probability of the event BadGap_ρ . Denote by x the first particle (in the order of enumeration) that causes BadGap_ρ . This is either a ρ -particle rejected because of Bad Gap or a 1-particle with at least one child in $\bar{\tau}_k^{\mathcal{S}(i, j)}$ (in other word, a parent in $\bar{\tau}_k^{\mathcal{S}(i, j)}$ of a ρ particle) rejected because of Bad Gap. On $\text{Small}_\rho \left(\bar{\tau}^{\mathcal{S}(i, j)}, M \right)$ there are at most M ρ -particles and therefore at most M 1-particles with at least one child. This gives us a bound on the number of particles that can be the cause of BadGap_ρ .

We further distinguish according to the three different types of BadGap_ρ .

1. Suppose that x satisfies the third condition of rejection by Bad Gap. There are at most $2M$ choices for x (which is either a ρ -particle or a parent of a ρ -particle). By (4.21) one deduces (by first conditioning by everything but $V(x) - V(\bar{x})$) that the probability that BadGap_ρ occurs because of the third type of Bad Gap is at most

$$2M\varepsilon/M = 2\varepsilon.$$

In what follows we assume that x does not satisfy the third condition of rejection by Bad Gap. In particular, arguing as in (4.25),

$$\hat{L}(V(x)) \in (i, j) + 2\Lambda kD. \tag{4.30}$$

²³In the proof of Lemma 4.8, there is no need to discuss about conditional probabilities. Moreover, the BRW is simply $\bar{\tau}^{0,0,1}$.

²⁴In the proof of Lemma 4.8,

$$\text{Else} = \{ \bar{\tau}_k^{0,0,1}(A_{\hat{L}}(1, 0)) < m \}.$$

2. Consider the first case. There at most M choices for the ρ -particle x . As $\|\widehat{L}(V(x') - V(x))\|_2 \leq 4\Lambda$, there are at most $200k^2\Lambda^2M$ choices for x' . Let us prove this fact. Either x' is one of the at most M ρ -particles generated at stage (i, j) or x' is one of the at most M ρ -particles generated at an earlier stage (i', j') . In the latter case, $\widehat{L}(V(x)) \in (i, j) + 2k\Lambda D$ (this is (4.30)) and $\widehat{L}(V(x')) \in (i', j') + 2k\Lambda D$ (see (4.25)). As $\|\widehat{L}(V(x') - V(x))\|_2 \leq 4\Lambda$ this implies

$$\|(i, j) - (i', j')\|_2 \leq (4k + 4)\Lambda \leq 6k\Lambda.$$

(Recall that k is even, so $k \geq 2$). Therefore there are at most $(12k\Lambda + 1)^2$ choices for (i', j') . Combining the previous properties, the number of choice for x' is at most

$$M + M(12k\Lambda + 1)^2 \leq 200k^2\Lambda^2M$$

as announced. We used $k, \Lambda \geq 1$ to simplify the upper bound²⁵. Finally, the number of choices for (x, x') is at most $200k^2\Lambda^2M^2$.

By Lemma 4.5 one deduces (by first conditioning on everything but $V(x) - V(\overleftarrow{x})$) that the probability that BadGap_ρ occurs because of the first type of Bad Gap is at most

$$400k^2\Lambda^2M^2\varepsilon'_G(d, \eta).$$

By (4.19), this is at most ε .

3. Consider the second case. There are at most M choice for the ρ -particle x . Then x' is the grandparent of x . By Lemma 4.5 one deduces (by first conditioning on everything but $V(x) - V(\overleftarrow{x})$ and $V(\overleftarrow{x}) - V(x')$) that the probability that BadGap_ρ occurs because of the second type of Bad Gap is at most

$$2M\varepsilon'_G(d, \eta).$$

By (4.19), as $k, \Lambda, M \geq 1$, this is at most ε .

Finally,

$$\mathbb{P}[\text{BadGap}_\rho] \leq 4\varepsilon. \tag{4.31}$$

Let us now give an upper bound for the probability of the event Interference_ρ . Denote by x the first particle (in the order of enumeration) that causes Interference_ρ . This is either a ρ -particle rejected because of Interference or a 1-particle with at least one child in $\overleftarrow{\tau}_k^{S(i,j)}$ rejected because of Interference . On $\text{Small}_\rho(\overleftarrow{\tau}^{S(i,j)}, M)$ there are at most M ρ -particles and therefore at most M 1-particles with at least one child. We distinguish the two cases of rejection by interference.

1. In the first case, a 1-particle x is rejected because of a ρ -particle y' which is not the parent \overleftarrow{x} of x . Note that x has not been rejected because of Bad Gap, otherwise we would not consider x for rejection by interference.

There are at most M choices for the 1-particle x (as it has as least one child which is a ρ -particle). The ρ -particle y' is such that $V(x) \in V(y') + K \cap \widehat{L}^{-1}(\Lambda D)$. Therefore $\|\widehat{L}(V(x) - V(y'))\|_2 \leq \Lambda$. As above, we conclude that there are at most $200k^2\Lambda^2M$ choices for y' (this is a crude bound). Therefore, there are at most $200k^2\Lambda^2M^2$ choices for (x, y') .

We have $\|\widehat{L}(V(\overleftarrow{x}) - V(x))\|_2 \leq \Lambda$ (otherwise x would have been rejected for Bad Gap) and therefore $\|\widehat{L}(V(\overleftarrow{x}) - V(y'))\|_2 \leq 2\Lambda$. As a consequence $V(\overleftarrow{x}) - V(y') \in G(d, K, \eta)$

²⁵In the proof of Lemma 4.8, there are actually at most M choices for x' . There are therefore also at most $200k^2\Lambda^2M$ choices for x' . Similar remarks apply several times below.

(otherwise y' or \overleftarrow{x} , which are distinct ρ -particles, would have been rejected for Bad Gap). Write $V(x) - V(y') = (V(\overleftarrow{x}) - V(y')) + (V(x) - V(\overleftarrow{x}))$. By definition of $G(d, K, \eta)$, conditioned on everything but $V(x) - V(\overleftarrow{x})$, the probability that x is rejected because of interference with y' is at most η .

Therefore, the probability that Interference_ρ occurs because of the first type of Rejection is at most

$$200k^2\Lambda^2M^2\eta.$$

By (4.17), this is at most ε .

2. In the second case, Interference_ρ is caused by the rejection of a ρ -particle z – with parent y and grand parent x – due to interference with a 1-particle y' – with parent x' – where y' is different from y . Note that x and x' are ρ -particles, that y and y' are 1-particles and that z is a ρ -particle.

As z is rejected because of interference with y' , at some point in the examination process:

- z is being examined and has not been rejected for Bad Gap (otherwise we would not even consider rejection of z due to interference).
- y' has been examined before and has not been rejected (otherwise it would not be a generated particle and we would not consider the interference caused by y').

This also implies that y, x and x' have been examined (parents are examined before their children) and have not been rejected (because when a particle is rejected, its progeny is rejected). As we examine in an admissible order, x' can not be a progeny of y (because we already examined the child y' of x' and we are currently examining z) or z . As z is a grandchild of x , z is different from x . To sum up, among all particles x, y, z, x', y' the only possible equality is $x = x'$.

We further distinguish according whether $x = x'$ or not.

- (a) Case $x \neq x'$.

As z is a grand-child of x we have $\|\widehat{L}(V(z) - V(x))\|_2 \leq 2\Lambda$ (otherwise y or z would have been rejected for Bad Gap). As z interferes with y' we have $\|\widehat{L}(V(z) - V(y'))\|_2 \leq \Lambda$. As y' is a child of x' we have $\|\widehat{L}(V(y') - V(x'))\|_2 \leq \Lambda$ (otherwise y' would have been rejected for Bad Gap). Therefore

$$\|\widehat{L}(V(z) - V(x'))\|_2 \leq 2\Lambda \tag{4.32}$$

and $\|\widehat{L}(V(x') - V(x))\|_2 \leq 4\Lambda$. As the ρ -particles x and x' have not been rejected by Bad Gap,

$$V(x) - V(x') \in G'(d, K, \eta). \tag{4.33}$$

This is where we use the assumption $x \neq x'$. Note that rejection of z because of y' implies (see (4.24))

$$V(z) - V(x') \in H'(d, K, \eta). \tag{4.34}$$

We thus see that if Interference_ρ occurs because of this type of case, then there exists a ρ -particle z with grandparent x and a ρ -particle x' distinct of x – which is not a progeny of y or z – such that (4.32), (4.33) and (4.34) occurs. Note that there is no mention of y' anymore. This is a key point as it allows us to control the number of particles involved. As z and x' are ρ -particles,

by (4.32) and arguments already used above, there are at most $200k^2\Lambda^2M^2$ choices for (z, x') .

Write $V(z) - V(x') = (V(z) - V(y)) + (V(y) - V(x)) + (V(x) - V(x'))$. By definition of $G'(d, K, \eta)$ and $H'(d, H, \eta)$, for each (z, x') , conditioned on everything but $V(z) - V(y)$ and $V(y) - V(x)$, on the event $\{(4.33) \text{ occurs}\}$ (which is measurable with respect to the conditioning σ -field because x' is not a progeny of y), the probability of (4.34) is at most η . To sum up, the probability that Interference_ρ occurs because of this type of case is at most

$$200k^2\Lambda^2M^2\eta.$$

By (4.17), this is at most ε .

- (b) Case $x = x'$. There are at most M choices for the ρ -particle z . As the enumeration has not been stopped by Overpopulation, given z , there are at most $M\sqrt{\rho^d}$ choices for y' which is a child of the grandfather x of z . Thus, there are at most $M^2\sqrt{\rho^d}$ choices for (z, y') . As z has not been rejected because of Bad Gap,

$$V(x) - V(z) \in H(d, K, \eta). \tag{4.35}$$

As z is rejected by interference with y' , $V(z) \in V(y') + K$ and therefore (using also symmetry of K)

$$(V(x) - V(z)) + (V(y') - V(x)) \in K. \tag{4.36}$$

But by definition of $H(d, H, \eta)$, given z and y' , conditioned on everything but $V(y') - V(x)$, on $\{(4.35) \text{ occurs}\}$, the probability of (4.36) is at most $(\sqrt{2} - \eta)^{-d}$. Therefore the probability that Interference_ρ occurs because of this type of case is at most

$$M^2\sqrt{\rho^d}(\sqrt{2} - \eta)^{-d}.$$

By (4.20) this is at most ε .

Finally,

$$\mathbb{P}[\text{Interference}_\rho] \leq 3\varepsilon. \tag{4.37}$$

By (4.23)²⁶

$$\mathbb{P} \left[\text{Small}_\rho \left(\bar{\tau}^{S(i,j)}, M \right)^c \cup \text{Else} \right] \leq \varepsilon.$$

This concludes the proof. □

4.5.7 Proof of Theorem 4.4

Proof of Item 1 of Theorem 4.4. In Section 4.5.5 we fixed $\beta > 1$, $2 > \rho > 1$ and $\varepsilon > 0$. We then got some integer d_0 (depending only on β, ρ and ε) and several other parameters satisfying various properties. We then let $d \geq d_0$ and $K \in \mathcal{K}(d)$ and built some process in Section 4.5.5. We studied some properties of this process in Section 4.5.6.

As in Section 3.4.7, we have

$$\mathbb{P}[\#\hat{A}^0 = \infty] \geq \mathbb{P}[\text{there exists an infinite open path from } (0, 0)].$$

²⁶In the proof of Lemma 4.8, we apply (4.22) to get

$$\mathbb{P} \left[\text{Small}_\rho \left(\bar{\tau}^{S(i,j)}, M \right)^c \cup \text{Else} \right] \leq 1 - (S(\beta^2) - \varepsilon).$$

By Lemmas 4.7, 4.8 and 4.9 we get

$$\mathbb{P}[\text{there exists an infinite open path from } (0, 0)] \geq (S(\lambda) - 8\varepsilon)\theta(1 - 8\varepsilon)$$

where $\theta(1 - 8\varepsilon)$ is the probability that there exists an infinite open path originating from $(1, 0)$ in a Bernoulli site percolation on \mathcal{L} with parameter $1 - 8\varepsilon$. Therefore

$$\mathbb{P}[\#\widehat{C}^0 = \infty] \geq \mathbb{P}[\#\widehat{A}^0 = \infty] \geq (S(\beta^2) - 8\varepsilon)\theta(1 - 8\varepsilon).$$

But $\theta(1 - 8\varepsilon)$ tends to 1 as ε tends to 0 (see (3.4)). This proves Item 1 of Theorem 4.4 in the case $\beta > 1$. The case $\beta \leq 1$ is trivial as, in this case, $S(\beta^2) = 0$.

Proof of Item 2 of Theorem 4.4. Let $\beta > 1$, $2 > \rho > 1$. Let $\varepsilon > 0$ such that $S(\beta^2) - \varepsilon > 0$. By Item 1, there exists d_0 such that, for all $d \geq d_0$ and all $K \in \mathcal{K}(d)$,

$$\mathbb{P}[\#\widehat{C}^0 = \infty] \geq S(\beta^2) - \varepsilon > 0.$$

Therefore, for all such ρ, d and K , $\beta_c(\rho, d, K) \leq \beta$.

A Some details on the definition of λ_c

All the results in this section are very standard and simple, but we have no ready reference for them. We provide a few technical details related to the definition of λ_c . We refer to the notations used in (1.3), (1.4) and (1.5).

Measurability. We prove in this paragraph the measurability of

$$\{\text{the connected component of the graph } \chi^0 \text{ that contains } 0 \text{ is unbounded}\}.$$

This is a consequence of the following facts:

- There exists a sequence of random variables $(C_n, R_n)_n$ such that, on the full event $\{\xi \text{ is infinite}\}$,

$$\xi = \{(C_n, R_n), n \in \mathbb{N}\}.$$

In other words, $\chi = \{(C_n), n \in \mathbb{N}\}$ and for all n , $r(C_n) = R_n$.

- The map defined by $(c_1, r_1, c_2, r_2) \mapsto \mathbb{1}_{\{c_1+r_1K \cap c_2+r_2K \neq \emptyset\}}$ is measurable because

$$\mathbb{1}_{\{c_1+r_1K \cap c_2+r_2K \neq \emptyset\}} = \mathbb{1}_K \left(\frac{1}{r_1 + r_2} (c_2 - c_1) \right).$$

The above equality is a consequence of $r_2K - r_1K = (r_2 + r_1)K$ which, in turn, is a consequence of the fact that K is symmetric and convex.

Equality (1.4). This is a consequence of

$$\begin{aligned} & \{\text{the connected component of the graph } \chi^0 \text{ that contains } 0 \text{ is unbounded}\} \\ &= \{\text{the connected component of } \Sigma \cup rK \text{ that contains } 0 \text{ is unbounded}\}. \end{aligned} \tag{A.1}$$

Let us prove this equality. Let \widehat{C}^0 be the connected component of the origin in the graph χ^0 . For all $x, y \in \chi^0$, there is an edge between x and y if and only if $x + r(x)K$ touches $y + r(y)K$. For all $x \in \chi^0$, $x + r(x)K$ is connected. Using these two facts, we get that

$$\bigcup_{x \in \widehat{C}^0} x + r(x)K \text{ is connected} \tag{A.2}$$

and that

$$\bigcup_{x \in \widehat{C}^0} x + r(x)K \text{ and } \bigcup_{x \in \chi^0 \setminus \widehat{C}^0} x + r(x)K \text{ are disjoint.} \quad (\text{A.3})$$

For all $x \in \chi^0$, $x + r(x)K$ is compact. The number of grains that touches a given bounded region is finite. This is a simple consequence of the fact that ν has bounded support, that K is bounded that χ^0 is locally finite. Using these two facts, we get that the two sets appearing in (A.3) are closed subsets of \mathbb{R}^d and therefore of $\Sigma \cup rK$.

Using (A.2), (A.3) and the closedness of the sets we get that the set which appears in (A.2) is the connected component of $\Sigma \cup rK$ that contains the origin. This yields (A.1) as K is bounded.

Equality (1.5). We define on χ a graph structure by putting an edge between x and y in χ when $x + r(x)K$ touches $y + r(y)K$. In other words, this is the natural restriction of the graph χ^0 to χ . With the same ideas as in the previous paragraph, we can check that “one of the connected component of Σ is unbounded” if and only if “one of the connected component of the graph χ is unbounded”. Then, with the same ideas as in the first paragraph, we can check that this is measurable.

Define

$$\begin{aligned} \Lambda_r &= \{\lambda > 0 : \mathbb{P}[\text{the connected component of } \Sigma \cup rK \text{ containing } 0 \text{ is unbounded}] > 0\}, \\ \Lambda' &= \{\lambda > 0 : \mathbb{P}[\text{one of the connected components of } \Sigma \text{ is unbounded}] > 0\}, \\ \Lambda &= \{\lambda > 0 : \mathbb{P}[\text{one of the connected components of } \Sigma \text{ is unbounded}] = 1\}. \end{aligned}$$

Let us show that the three sets are equal. As K is connected with non empty interior, by stationarity we get $\Lambda \subset \Lambda_r$. The connected component of $\Sigma \cup rK$ that contains the origin is (A.2). The connected components of Σ are in similar correspondence with connected components of the graph χ . As moreover the degree of 0 in the graph χ^0 is finite we get $\Lambda_r \subset \Lambda'$. Finally, $\Lambda' \subset \Lambda$ by ergodicity. Thus the three sets are equal. In particular, $\Lambda_r = \Lambda$ and this proves (1.5).

References

- [1] K. B. Athreya and P. E. Ney, *Branching processes*, Dover Publications, Inc., Mineola, NY, 2004, Reprint of the 1972 original [Springer, New York; MR0373040]. MR2047480
- [2] Paul Balister, Béla Bollobás, and Mark Walters, *Continuum percolation with steps in the square or the disc*, *Random Structures Algorithms* **26** (2005), no. 4, 392–403. MR2139873
- [3] Franck Barthe, Olivier Guédon, Shahar Mendelson, and Assaf Naor, *A probabilistic approach to the geometry of the l_p^n -ball*, *Ann. Probab.* **33** (2005), no. 2, 480–513. MR2123199
- [4] Richard Durrett, *Oriented percolation in two dimensions*, *Ann. Probab.* **12** (1984), no. 4, 999–1040. MR757768
- [5] R. J. Gardner, *The Brunn-Minkowski inequality*, *Bull. Amer. Math. Soc. (N.S.)* **39** (2002), no. 3, 355–405. MR1898210
- [6] Jean-Baptiste Gouéré and Régine Marchand, *Nonoptimality of constant radii in high dimensional continuum percolation*, *Ann. Probab.* **44** (2016), no. 1, 307–323. MR3456339
- [7] Jean-Baptiste Gouéré and Régine Marchand, *Continuum percolation in high dimensions*, *Ann. Inst. Henri Poincaré Probab. Stat.* **54** (2018), no. 4, 1778–1804. MR3865657
- [8] J. F. C. Kingman, *Poisson processes*, *Oxford Studies in Probability*, vol. 3, The Clarendon Press, Oxford University Press, New York, 1993, Oxford Science Publications. MR1207584
- [9] B. Klartag, *A central limit theorem for convex sets*, *Invent. Math.* **168** (2007), no. 1, 91–131. MR2285748
- [10] Günter Last and Mathew Penrose, *Lectures on the Poisson process*, *Institute of Mathematical Statistics Textbooks*, vol. 7, Cambridge University Press, Cambridge, 2018. MR3791470

- [11] Elliott H. Lieb and Michael Loss, *Analysis*, second ed., Graduate Studies in Mathematics, vol. 14, American Mathematical Society, Providence, RI, 2001. MR1817225
- [12] Torgny Lindvall, *Lectures on the coupling method*, Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics, John Wiley & Sons, Inc., New York, 1992, A Wiley-Interscience Publication. MR1180522
- [13] Ronald Meester and Rahul Roy, *Continuum percolation*, Cambridge Tracts in Mathematics, vol. 119, Cambridge University Press, Cambridge, 1996. MRMR1409145
- [14] Mathew D. Penrose, *Continuum percolation and Euclidean minimal spanning trees in high dimensions*, *Ann. Appl. Probab.* **6** (1996), no. 2, 528–544. MR1398056
- [15] András Prékopa, *On logarithmic concave measures and functions*, *Acta Sci. Math. (Szeged)* **34** (1973), 335–343. MR404557
- [16] Zhan Shi, *Branching random walks*, Lecture Notes in Mathematics, vol. 2151, Springer, Cham, 2015, Lecture notes from the 42nd Probability Summer School held in Saint Flour, 2012, École d'Été de Probabilités de Saint-Flour. [Saint-Flour Probability Summer School]. MR3444654

Acknowledgments. The authors are grateful to an anonymous referee for his very careful reading and for his detailed list of useful comments.

Electronic Journal of Probability

Electronic Communications in Probability

Advantages of publishing in EJP-ECP

- Very high standards
- Free for authors, free for readers
- Quick publication (no backlog)
- Secure publication (LOCKSS¹)
- Easy interface (EJMS²)

Economical model of EJP-ECP

- Non profit, sponsored by IMS³, BS⁴, ProjectEuclid⁵
- Purely electronic

Help keep the journal free and vigorous

- Donate to the IMS open access fund⁶ (click here to donate!)
- Submit your best articles to EJP-ECP
- Choose EJP-ECP over for-profit journals

¹LOCKSS: Lots of Copies Keep Stuff Safe <http://www.lockss.org/>

²EJMS: Electronic Journal Management System: <https://vtex.lt/services/ejms-peer-review/>

³IMS: Institute of Mathematical Statistics <http://www.imstat.org/>

⁴BS: Bernoulli Society <http://www.bernoulli-society.org/>

⁵Project Euclid: <https://projecteuclid.org/>

⁶IMS Open Access Fund: <https://imstat.org/shop/donation/>