

## THE TIME CONSTANT AND CRITICAL PROBABILITIES IN PERCOLATION MODELS

LEANDRO P. R. PIMENTEL<sup>1</sup>*Institut de Mathématiques, École Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland*

email: leandro.pimentel@epfl.ch

*Submitted 3 February 2005 , accepted in final form 19 July 2006*

AMS 2000 Subject classification: 60K35, 82D30

Keywords: Percolation, time constant, critical probabilities, Delaunay triangulations

*Abstract*

We consider a first-passage percolation (FPP) model on a Delaunay triangulation  $\mathcal{D}$  of the plane. In this model each edge  $e$  of  $\mathcal{D}$  is independently equipped with a nonnegative random variable  $\tau_e$ , with distribution function  $\mathbb{F}$ , which is interpreted as the time it takes to traverse the edge. Vahidi-Asl and Wierman [9] have shown that, under a suitable moment condition on  $\mathbb{F}$ , the minimum time taken to reach a point  $\mathbf{x}$  from the origin  $\mathbf{0}$  is asymptotically  $\mu(\mathbb{F})|\mathbf{x}|$ , where  $\mu(\mathbb{F})$  is a nonnegative finite constant. However the exact value of the time constant  $\mu(\mathbb{F})$  still a fundamental problem in percolation theory. Here we prove that if  $\mathbb{F}(0) < 1 - p_c^*$  then  $\mu(\mathbb{F}) > 0$ , where  $p_c^*$  is a critical probability for bond percolation on the dual graph  $\mathcal{D}^*$ .

**Introduction**

First-passage percolation theory on periodic graphs was presented by Hammersley and Welsh [4] to model the spread of a fluid through a porous medium. In this paper we continue a study of planar first-passage percolation models on random graphs, initiated by Vahidi-Asl and Wierman [9], as follows. Let  $\mathcal{P}$  denote the set of points realized in a two-dimensional homogeneous Poisson point process with intensity 1. To each  $\mathbf{v} \in \mathcal{P}$  corresponds an open polygonal region  $\mathbf{C}_{\mathbf{v}} = \mathbf{C}_{\mathbf{v}}(\mathcal{P})$ , the Voronoi tile at  $\mathbf{v}$ , consisting of the set of points of  $\mathbb{R}^2$  which are closer to  $\mathbf{v}$  than to any other  $\mathbf{v}' \in \mathcal{P}$ . Given  $\mathbf{x} \in \mathbb{R}^2$  we denote by  $\mathbf{v}_{\mathbf{x}}$  the almost surely unique point in  $\mathcal{P}$  such that  $\mathbf{x} \in \mathbf{C}_{\mathbf{v}_{\mathbf{x}}}$ . The collection  $\{\mathbf{C}_{\mathbf{v}} : \mathbf{v} \in \mathcal{P}\}$  is called the Voronoi Tiling of the plane based on  $\mathcal{P}$ .

The Delaunay Triangulation  $\mathcal{D}$  is the graph where the vertex set  $\mathcal{D}_v$  equals  $\mathcal{P}$  and the edge set  $\mathcal{D}_e$  consists of non-oriented pairs  $(\mathbf{v}, \mathbf{v}')$  such that  $\mathbf{C}_{\mathbf{v}}$  and  $\mathbf{C}_{\mathbf{v}'}$  share a one-dimensional edge (Figure 1). One can see that almost surely each Voronoi tile is a convex and bounded polygon, and the graph  $\mathcal{D}$  is a triangulation of the plane [7]. The Voronoi Tessellation  $\mathcal{V}$  is the graph where the vertex set  $\mathcal{V}_v$  is the set of vertices of the Voronoi tiles and the edge set  $\mathcal{V}_e$  is the set

<sup>1</sup>RESEARCH SUPPORTED BY SWISS NATIONAL SCIENCE FOUNDATION GRANT 510767




Figure 1: The Delaunay Triangulation and the Voronoi Tessellation.

of edges of the Voronoi tiles. The edges of  $\mathcal{V}$  are segments of the perpendicular bisectors of the edges of  $\mathcal{D}$ . This establishes duality of  $\mathcal{D}$  and  $\mathcal{V}$  as planar graphs:  $\mathcal{V} = \mathcal{D}^*$ .

To each edge  $\mathbf{e} \in \mathcal{D}_e$  is independently assigned a nonnegative random variable  $\tau_{\mathbf{e}}$  from a common distribution  $\mathbb{F}$ , which is also independent of the Poisson point process that generates  $\mathcal{P}$ . From now on we denote  $(\Omega, \mathcal{F}, \mathbb{P})$  the probability space induced by the Poisson point process  $\mathcal{P}$  and the passage times  $(\tau_{\mathbf{e}})_{\mathbf{e} \in \mathcal{D}_e}$ . The passage time  $t(\gamma)$  of a path  $\gamma$  in the Delaunay Triangulation is the sum of the passage times of the edges in  $\gamma$ . The first-passage time between two vertices  $\mathbf{v}$  and  $\mathbf{v}'$  is defined by

$$T(\mathbf{v}, \mathbf{v}') := \inf\{t(\gamma) ; \gamma \in \mathcal{C}(\mathbf{v}, \mathbf{v}')\},$$

where  $\mathcal{C}(\mathbf{v}, \mathbf{v}')$  the set of all paths connecting  $\mathbf{v}$  to  $\mathbf{v}'$ . Given  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$  we define  $T(\mathbf{x}, \mathbf{y}) := T(\mathbf{v}_{\mathbf{x}}, \mathbf{v}_{\mathbf{y}})$ .

To state the main result of this work we require some definitions involving a bond percolation model on the Voronoi Tessellation  $\mathcal{V}$ . Such a model is constructed by choosing each edge of  $\mathcal{V}$  to be open independently with probability  $p$ . An open path is a path composed of open edges. We denote  $\mathbb{P}_p^*$  the law induced by the Poisson point process and the random state (open or not) of an edge. Given a planar graph  $\mathcal{G}$  and  $\mathbf{A}, \mathbf{B} \subseteq \mathbb{R}^2$  we say that a self-avoiding path  $\gamma = (\mathbf{v}_1, \dots, \mathbf{v}_k)$  is a path connecting  $\mathbf{A}$  to  $\mathbf{B}$  if  $[\mathbf{v}_1, \mathbf{v}_2] \cap \mathbf{A} \neq \emptyset$  and  $[\mathbf{v}_{k-1}, \mathbf{v}_k] \cap \mathbf{B} \neq \emptyset$  ( $[\mathbf{x}, \mathbf{y}]$  denotes the line segment connecting  $\mathbf{x}$  to  $\mathbf{y}$ ). For  $L > 0$  let  $A_L$  be the event that there exists an open path  $\gamma = (\mathbf{v}_j)_{1 \leq j \leq h}$  in  $\mathcal{V}$ , connecting  $\{0\} \times [0, L]$  to  $\{3L\} \times [0, L]$ , and with  $\mathbf{v}_j \in [0, 3L] \times [0, L]$  for all  $j = 2, \dots, h-1$ . In this case we also say that  $\gamma$  crosses the rectangle  $[0, 3L] \times [0, L]$ . Define the function

$$\eta^*(p) := \liminf_{L \rightarrow \infty} \mathbb{P}_p^*(A_L),$$

and consider the percolation threshold,

$$p_c^* := \inf\{p > 0 : \eta^*(p) = 1\}. \quad (1)$$

We have that  $p_c^* \in (0, 1)$ , which follows by standard arguments in percolation theory. For more in percolation thresholds on Voronoi tilings we refer to [1, 2, 11].

**Theorem 1** *If  $\mathbb{F}(0) < 1 - p_c^*$  then there exist constants  $c_j > 0$  such that for all  $n \geq 1$*

$$\mathbb{P}(T(\mathbf{0}, \mathbf{n}) < c_1 n) \leq c_2 \exp(-c_3 n), \quad (2)$$

where  $\mathbf{0} := (0, 0)$  and  $\mathbf{n} := (n, 0)$ .

To show the importance of Theorem 1 we recall two fundamental results proved by Vahidi-Asl and Wierman [9, 10]. Consider the growth process

$$\mathbf{B}_{\mathbf{x}}(t) := \{\mathbf{y} \in \mathbb{R}^2 : \mathbf{y} \in c(\mathbf{C}_{\mathbf{v}}) \text{ with } \mathbf{v} \in \mathcal{D}_v \text{ and } T(\mathbf{v}_{\mathbf{x}}, \mathbf{v}) \leq t\}.$$

where  $c(\mathbf{C})$  denotes the closure of  $\mathbf{C} \in \mathbb{R}^2$ . Set

$$\mu(\mathbb{F}) := \inf_{n>0} \frac{\mathbb{E}T(\mathbf{0}, \mathbf{n})}{n} \in [0, \infty].$$

and let  $\tau_1, \tau_2, \tau_3$  be independent random variables with distribution  $\mathbb{F}$ . If

$$\mathbb{E}\left(\min_{j=1,2,3} \{\tau_j\}\right) < \infty \quad (3)$$

then  $\mu(\mathbb{F}) < \infty$  and for all unit vectors  $\vec{\mathbf{x}} \in S^1$  ( $|\vec{\mathbf{x}}| = 1$ )  $\mathbb{P}$ -a.s.

$$\lim_{n \rightarrow \infty} \frac{T(\mathbf{0}, n\vec{\mathbf{x}})}{n} = \lim_{n \rightarrow \infty} \frac{\mathbb{E}T(\mathbf{0}, \mathbf{n})}{n} = \mu(\mathbb{F}). \quad (4)$$

Further, if

$$\mathbb{E}\left(\min_{j=1,2,3} \{\tau_j\}^2\right) < \infty \quad (5)$$

and  $\mu(\mathbb{F}) > 0$  then for all  $\epsilon > 0$   $\mathbb{P}$ -a.s. there exists  $t_0 > 0$  such that for all  $t > t_0$

$$(1 - \epsilon)t\mathbf{D}(1/\mu) \subseteq \mathbf{B}_{\mathbf{0}}(t) \subseteq (1 + \epsilon)t\mathbf{D}(1/\mu), \quad (6)$$

where  $\mathbf{D}(r) := \{\mathbf{x} \in \mathbb{R}^2 : |\mathbf{x}| \leq r\}$ .

We note here that the asymptotic shape is an Euclidean ball due to the statistical invariance of the Poisson point process. Unfortunately the exact value of the time constant  $\mu(\mathbb{F})$ , as a functional of  $\mathbb{F}$ , still a basic problem in first-passage percolation theory. Our result provides a sufficient condition on  $\mathbb{F}$  to ensure  $\mu(\mathbb{F}) > 0$ .

**Corollary 1** *Under assumption (3), if  $\mathbb{F}(0) < 1 - p_c^*$  then  $\mu(\mathbb{F}) \in (0, \infty)$ .*

PROOF. Together with the Borel-Cantelli Lemma, Theorem 1 and (4) imply

$$0 < c_1 \leq \liminf_{n \rightarrow \infty} \frac{T(\mathbf{0}, \mathbf{n})}{n} = \lim_{n \rightarrow \infty} \frac{T(\mathbf{0}, \mathbf{n})}{n} = \mu(\mathbb{F}) < \infty,$$

which is the desired result.  $\square$

For FPP models on the  $\mathbb{Z}^2$  lattice Kesten (1986) has shown that  $\mathbb{F}(0) < 1/2 = p_c(\mathbb{Z}^2)$  (the critical probability for bond percolation on  $\mathbb{Z}^2$ ) is a sufficient condition to get (2) by using a stronger version of the BK-inequality. Here we follow a different method and we apply a simple renormalization argument to obtain a similar result. We expect that our condition to get (2) is equivalent to

$$\mathbb{F}(0) < p_c := \inf\{p > 0; \theta(p) = 1\},$$

where  $\theta(p)$  is the probability that bond percolation on  $\mathcal{D}$  occurs with density  $p$ , since it is conjectured that  $p_c + p_c^* = 1$  (duality) for many planar graphs. In fact, by combining Corollary 1 with (6) we have:

**Corollary 2**

$$1 \leq p_c + p_c^* .$$

PROOF. To see this assume we have a first-passage percolation model on  $\mathcal{D}$  with

$$\mathbb{P}(\tau_{\mathbf{e}} = 0) = 1 - \mathbb{P}(\tau_{\mathbf{e}} = 1) = \mathbb{F}(0) = 1 - p > p_c^* . \tag{7}$$

Then  $\mathbb{P}$ -a.s. there exists an infinite cluster  $\mathcal{W} \subseteq \mathcal{D}$  composed by edges  $\mathbf{e}$  with  $\tau_{\mathbf{e}} = 0$ . Denote by  $T(\mathbf{0}, \mathcal{W})$  the first-passage time from  $\mathbf{0}$  to  $\mathcal{W}$ . Then for all  $t > T(\mathbf{0}, \mathcal{W})$  we have that  $\mathbf{B}_{\mathbf{0}}(t)$  is an unbounded set. By (6) (since such a distribution satisfies (3) and (5)), this implies that  $\mu(\mathbb{F}) = \mu(p) = 0$  if  $1 - p > p_c$ . On the other hand, by Corollary 1,  $\mu(p) > 0$  if  $1 - p < 1 - p_c^*$ , and so (2) must hold.  $\square$

Other passage times have been considered in the literature such as  $T(\mathbf{0}, \mathbf{H}_n)$ , where  $\mathbf{H}_n$  is the hyperplane consisting of points  $\mathbf{x} = (x^1, x^2)$  so that  $x_1 = n$ , and  $T(\mathbf{0}, \partial[-n, n]^2)$ . The arguments in this article can be used to prove the analog of Theorem 1 when  $T(\mathbf{0}, \mathbf{n})$  is replaced by  $T(\mathbf{0}, \mathbf{H}_n)$  or  $T(\mathbf{0}, \partial[-n, n]^2)$ . For site versions of FPP models the method works as well if we change the condition on  $\mathbb{F}$  to  $\mathbb{F}(0) < 1 - \bar{p}_c$ , where now  $\bar{p}_c$  is the critical probability for site percolation. Similarly to Corollary 2, in this case one can also obtain the inequality  $1/2 \leq \bar{p}_c$ . For more details we refer to [8].

## 1 Renormalization

For the moment we assume that  $\mathbb{F}$  is Bernoulli with parameter  $p$ . Let  $L \geq 1$  be a parameter whose value will be specified later. Let  $\mathbf{z} = (z^1, z^2) \in \mathbb{Z}^2$  and

$$|\mathbf{z}|_{\infty} := \max_{j=1,2} \{|z^j|\} .$$

Denote  $\mathbf{C}_{\mathbf{z}}$  the circuit composed by sites  $\mathbf{z}' \in \mathbb{Z}^2$  with  $|\mathbf{z} - \mathbf{z}'|_{\infty} = 2$ . For each  $\mathbf{A} \subseteq \mathbb{R}^2$ , we denote by  $\partial\mathbf{A}$  its boundary. For each  $\mathbf{z} \in \mathbb{Z}^2$  and  $r \in \{j/2 : j \in \mathbb{N}\}$  consider the box

$$\mathbf{B}_{\mathbf{z}}^{rL} := L\mathbf{z} + [-rL, rL]^2 .$$

Divide  $\mathbf{B}_{\mathbf{z}}^{L/2}$  into thirty-six sub-boxes with the same size and declare that  $B_{\mathbf{z}}^{L/2}$  is a **full box** if all these thirty-six sub-boxes contain at least one point of  $\mathcal{P}$ . Let

$$H_{\mathbf{z}}^L := [\mathbf{B}_{\mathbf{z}'}^{L/2} \text{ is a full box } \forall \mathbf{z}' \in \mathbf{C}_{\mathbf{z}}] .$$

Let  $\mathcal{C}_L$  be the set of all self-avoiding paths  $\gamma = (\mathbf{v}_j)_{1 \leq j \leq h}$  in  $\mathcal{D}$ , connecting  $\partial\mathbf{B}_{\mathbf{z}}^{L/2}$  to  $\partial\mathbf{B}_{\mathbf{z}}^{3L/2}$  and with  $\mathbf{C}_{\mathbf{v}_j} \cap \mathbf{B}_{\mathbf{z}}^{3L/2}$  for all  $j = 2, \dots, h - 1$ . Let

$$G_{\mathbf{z}}^L := [t(\gamma) \geq 1 \forall \gamma \in \mathcal{C}_L] .$$

We say that  $\mathbf{B}_{\mathbf{z}}^{L/2}$  is a **good box** (or that  $\mathbf{z}$  is a good point) if

$$Y_{\mathbf{z}}^L := \mathbb{I}(H_{\mathbf{z}}^L \cap G_{\mathbf{z}}^L) = 1 ,$$

where  $\mathbb{I}(E)$  denotes the indicator function of the event  $E$ .




Figure 2: Renormalization

**Lemma 1** *If  $\mathbb{P}(\tau_{\mathbf{e}} = 0) = 1 - p < 1 - p_c^*$  then*

$$\lim_{L \rightarrow \infty} \mathbb{P}(Y_{\mathbf{0}}^L = 1) = 1.$$

PROOF. First notice that

$$\mathbb{P}(Y_{\mathbf{0}}^L = 0) \leq \mathbb{P}((H_{\mathbf{0}}^L)^c) + \mathbb{P}((G_{\mathbf{0}}^L)^c). \tag{8}$$

By the definition of a two-dimensional homogeneous Poisson point process,

$$\lim_{L \rightarrow \infty} \mathbb{P}((H_{\mathbf{0}}^L)^c) = 0. \tag{9}$$

Now, let  $X_{\mathbf{e}^*} := \tau_{\mathbf{e}}$ , where  $\mathbf{e}^*$  is the edge in  $\mathcal{V}_{\mathbf{e}}$  (the Voronoi tessellation) dual to  $\mathbf{e}$ . Then  $\{X_{\mathbf{e}^*} ; \mathbf{e}^* \in \mathcal{V}_{\mathbf{e}}\}$  defines a bond percolation model on  $\mathcal{V}$  with law  $\mathbb{P}_p^*$ . Consider the rectangles

$$R_L^1 := [L/2, 3L/2] \times [-3L/2, 3L/2], \quad R_L^2 := [-3L/2, 3L/2] \times [L/2, 3L/2]$$

$$R_L^3 := [-3L/2, -L/2] \times [-3L/2, 3L/2] \text{ and } R_L^4 := [-3L/2, 3L/2] \times [-3L/2, -L/2].$$

We denote by  $A_L^i$  the event  $A_L$  (recall the definition of  $p_c^*$ ) but now translate to the rectangle  $R_L^i$ , and by  $F_L$  the event that an open circuit  $\sigma^*$  in  $\mathcal{V}$  which surrounds  $B_{\mathbf{0}}^{L/2}$  and lies inside  $B_{\mathbf{0}}^{3L/2}$  does not exist. Thus one can easily see that

$$\bigcap_{i=1}^4 A_L^i \subseteq (F_L)^c.$$

Notice that if there exists an open circuit  $\sigma^*$  in  $\mathcal{V}$  which surrounds  $B_{\mathbf{0}}^{L/2}$  and lies inside  $B_{\mathbf{0}}^{3L/2}$ , then every path  $\gamma$  in  $\mathcal{C}_L$  has an edge crossing with  $\sigma^*$  and thus  $t(\gamma) \geq 1$ . Therefore,

$$\mathbb{P}((G_{\mathbf{0}}^L)^c) \leq \mathbb{P}_p^*(F_L) \leq 4(1 - \mathbb{P}_p^*(A_L)). \tag{10}$$

Since  $p > p_c^*$ , by using (8), (9), (10) and the definition of  $p_c^*$ , we get Lemma 1. □

To obtain some sort of independence between the random variables  $Y_{\mathbf{z}}^L$  we shall study some geometrical aspects of Voronoi tilings. Given  $\mathbf{A} \subseteq \mathbb{R}^2$ , let  $\mathcal{I}_{\mathcal{P}}(\mathbf{A})$  be the sub-graph of  $\mathcal{D}$  composed of vertices  $\mathbf{v}_1$  in  $\mathcal{D}_v$  and edges  $(\mathbf{v}_2, \mathbf{v}_3)$  in  $\mathcal{D}_e$  so that  $\mathbf{C}_{\mathbf{v}_i} \cap \mathbf{A} \neq \emptyset$  for all  $i = 1, 2, 3$ .

**Lemma 2** *Let  $L > 0$  and  $\mathbf{z} \in \mathbb{Z}^2$ . Assume that  $\mathcal{P}$  and  $\mathcal{P}'$  are two configurations of points so that  $\mathcal{P} \cap \mathbf{B}_{\mathbf{z}}^{5L/2} = \mathcal{P}' \cap \mathbf{B}_{\mathbf{z}}^{5L/2}$  and that  $\mathbf{B}_{\mathbf{z}'}^{L/2}$  is a full box with respect to  $\mathcal{P}$ , for all  $\mathbf{z}' \in \mathbf{C}_{\mathbf{z}}$ . Then  $\mathcal{I}_{\mathcal{P}}(\mathbf{B}_{\mathbf{z}}^{3L/2}) = \mathcal{I}_{\mathcal{P}'}(\mathbf{B}_{\mathbf{z}}^{3L/2})$ .*

PROOF. By the definition of the Delaunay Triangulation, Lemma 2 holds if we prove that

$$\mathbf{C}_{\mathbf{v}}(\mathcal{P}) \cap \mathbf{B}_{\mathbf{z}}^{3L/2} \neq \emptyset \Rightarrow \mathbf{C}_{\mathbf{v}}(\mathcal{P}) = \mathbf{C}_{\mathbf{v}}(\mathcal{P}'). \quad (11)$$

To prove this we claim that

$$\mathbf{C}_{\mathbf{v}}(\mathcal{P}) \cap \mathbf{B}_{\mathbf{z}}^{3L/2} \neq \emptyset \Rightarrow \mathbf{C}_{\mathbf{v}}(\mathcal{P}) \subseteq \mathbf{B}_{\mathbf{z}}^{2L}. \quad (12)$$

If (12) does not hold then there exist  $\mathbf{x}_1 \in \partial \mathbf{B}_{\mathbf{z}}^{3L/2} \cap \mathbf{C}_{\mathbf{v}}(\mathcal{P})$  and  $\mathbf{x}_2 \in \partial \mathbf{B}_{\mathbf{z}}^{2L} \cap \mathbf{C}_{\mathbf{v}}(\mathcal{P})$  (by convexity of Voronoi tilings). Since every box  $\mathbf{B}_{\mathbf{z}'}^{L/2}$  with  $|\mathbf{z} - \mathbf{z}'|_{\infty} = 2$  is a full box, there exist  $\mathbf{v}_1, \mathbf{v}_2 \in \mathcal{P}$  so that

$$|\mathbf{v}_1 - \mathbf{x}_1| \leq \sqrt{2}L/6 \text{ and } |\mathbf{v}_2 - \mathbf{x}_2| \leq \sqrt{2}L/6.$$

Although,  $\mathbf{x}_1$  and  $\mathbf{x}_2$  belong to  $\mathbf{C}_{\mathbf{v}}(\mathcal{P})$  and so

$$|\mathbf{v} - \mathbf{x}_1| \leq |\mathbf{v}_1 - \mathbf{x}_1| \text{ and } |\mathbf{v} - \mathbf{x}_2| \leq |\mathbf{v}_2 - \mathbf{x}_2|.$$

Thus,

$$L/2 \leq |\mathbf{x}_1 - \mathbf{x}_2| \leq |\mathbf{x}_1 - \mathbf{v}| + |\mathbf{x}_2 - \mathbf{v}| \leq \sqrt{2}L/3,$$

which leads to a contradiction since  $\sqrt{2}/3 < 1/2$ . By an analogous argument, one can prove that

$$\mathbf{C}_{\mathbf{v}'}(\mathcal{P}') \cap (\mathbf{B}_{\mathbf{z}}^{5L/2})^c \neq \emptyset \Rightarrow \mathbf{C}_{\mathbf{v}'}(\mathcal{P}') \subseteq (\mathbf{B}_{\mathbf{z}}^{2L})^c. \quad (13)$$

Now suppose (11) does not hold. Without loss of generality, we may assume that there exists  $\mathbf{v} \in \mathcal{P}$  with  $\mathbf{C}_{\mathbf{v}}(\mathcal{P}) \cap \mathbf{B}_{\mathbf{z}}^{3L/2} \neq \emptyset$  and  $\mathbf{x} \in \mathbf{C}_{\mathbf{v}}(\mathcal{P})$  with  $\mathbf{x} \notin \mathbf{C}_{\mathbf{v}'}(\mathcal{P}')$ . So  $\mathbf{x} \in \mathbf{C}_{\mathbf{v}'}(\mathcal{P}')$  for some  $\mathbf{v}' \in \mathcal{P}'$ . Although,  $\mathcal{P} \cap \mathbf{B}_{\mathbf{z}}^{5L/2} = \mathcal{P}' \cap \mathbf{B}_{\mathbf{z}}^{5L/2}$  and then  $\mathbf{v}' \in (\mathbf{B}_{\mathbf{z}}^{5L/2})^c$ , which is a contradiction with (12) and (13).  $\square$

For each  $l \geq 1$ , we say that the collection of random variables  $\{Y_{\mathbf{z}} : \mathbf{z} \in \mathbb{Z}^2\}$  is  $l$ -dependent if  $\{Y_{\mathbf{z}} : \mathbf{z} \in \mathbf{A}\}$  and  $\{Y_{\mathbf{z}} : \mathbf{z} \in \mathbf{B}\}$  are independent whenever

$$l < d_{\infty}(\mathbf{A}, \mathbf{B}) := \min\{|\mathbf{z} - \mathbf{z}'|_{\infty} : \mathbf{z} \in \mathbf{A} \text{ and } \mathbf{z}' \in \mathbf{B}\}.$$

Combining Lemma 2 with the translation invariance and the independence property of the Poisson point process we obtain:

**Lemma 3** *For all  $L > 0$ ,  $\{Y_{\mathbf{z}}^L : \mathbf{z} \in \mathbb{Z}^2\}$  is a 5-dependent collection of identically distributed Bernoulli random variables.*

Denote  $Y^L := \{Y_{\mathbf{z}}^L ; \mathbf{z} \in \mathbb{Z}^2\}$  and let  $M_m(Y^L)$  be the maximum number of pairwise disjoint good circuits in  $\mathbb{Z}^2$ , surrounding the origin and lying inside the box  $[-m, m]^2$ .

**Lemma 4** *If  $\mathbb{F}(0) < 1 - p_c^*$  then there exists  $L_0 > 0$  and  $c_j = c_j(L_0) > 0$  such that*

$$\mathbb{P}(M_m(Y^{L_0}) \leq c_1 m) \leq \exp(-c_2 m).$$

PROOF. Combining Lemmas 1 and 3 with and Theorem 0.0 of Ligget, Schonman and Stacey [6], one gets that  $Y^L$  is dominated from below by a collection  $X^L := \{X_z^L; z \in \mathbb{Z}^2\}$  of i.i.d. Bernoulli random variables with parameter  $\rho(L) \rightarrow 1$  when  $L \rightarrow \infty$ . But for  $\rho_L$  sufficiently close to 1, we can chose  $c > 0$  sufficiently small, so that the probability of the event that  $M_m(X^L) < cm$  decays exponentially fast with  $m$  (see Chapter 3 of Grimmett [3]). Together with domination, this proves Lemma 4.  $\square$

The connection between the variable  $M_m(Y^L)$  and the first-passage time  $T(\mathbf{0}, \mathbf{n})$  is summarize by the following:

**Lemma 5**

$$\frac{M_{nL-1}^L}{6} \leq T(\mathbf{0}, \mathbf{n}).$$

PROOF. We say that  $(B_{\mathbf{z}_j}^{L/2})_{1 \leq j \leq h}$  is a circuit of good boxes if  $(\mathbf{z}_j)_{1 \leq j \leq h}$  is a good circuit in  $\mathbb{Z}^2$ , and that  $(B_{\mathbf{z}_j}^{L/2})_{1 \leq j \leq h}$  and  $(B_{\mathbf{z}'_j}^{L/2})_{1 \leq j \leq h'}$  are  $l$ -distant if

$$d_\infty((\mathbf{z}_j)_{1 \leq j \leq h}, (\mathbf{z}'_j)_{1 \leq j \leq h'}) > l.$$

Denote  $M_m^L := M_m(Y^L)$ . Notice that there exist at least  $(M_{nL-1}^L/6)$  pairwise 5-distant circuits of good boxes surrounding the origin and lying inside  $[-n, n]^2 \subseteq \mathbb{R}^2$ . Therefore, every path  $\gamma$  between the origin and any point outside  $[-n, n]^2$  must cross at least  $(M_{nL-1}^L/6)$  5-distant circuits of good boxes. We claim this yields

$$\frac{M_{nL-1}^L}{6} \leq t(\gamma). \quad (14)$$

Indeed, assume we take two 5-distant good boxes, say  $\mathbf{B}_{\mathbf{z}_1}^{L/2}$  and  $\mathbf{B}_{\mathbf{z}_2}^{L/2}$ , connected by a path  $\gamma$  in  $\mathcal{D}$ . Then  $\gamma$  must contain two sub-paths in  $\mathcal{D}$ , say  $\tilde{\gamma}_i = (\mathbf{v}_j^i)_{1 \leq j \leq h_i}$  for  $i = 1, 2$ , connecting  $\partial \mathbf{B}_{\mathbf{z}_i}^{3L/2}$  to  $\partial \mathbf{B}_{\mathbf{z}_i}^{5L/2}$  and with  $\mathbf{C}_{\mathbf{v}_j^i} \cap \mathbf{B}_{\mathbf{z}_i}^{3L/2}$  for all  $j = 2, \dots, h_i - 1$ . Since  $B_{\mathbf{z}_1}^{L/2}$  and  $B_{\mathbf{z}_2}^{L/2}$  are 5-distant good boxes, by Lemma 2, these sub-paths must be edge disjoint. By the definition of a good box,  $t(\tilde{\gamma}_1) \geq 1$  and  $t(\tilde{\gamma}_2) \geq 1$ , which yields

$$2 \leq t(\tilde{\gamma}_1) + t(\tilde{\gamma}_2) \leq t(\gamma).$$

By repeating this argument inductively (on the number of good boxes which are crossed by  $\gamma$ ) one can get (14). Lemma 5 follows directly from (14).  $\square$

Now we are ready to prove Theorem 1.

PROOF. Together with Lemma 5, Lemma 4 implies Theorem 1 under (7). For the general case, assume  $\mathbb{F}(0) = \mathbb{P}(\tau_{\mathbf{e}} = 0) < 1 - p_1$ . Fix  $\epsilon > 0$  so that  $\mathbb{F}(\epsilon) < 1 - p_c^*$  (we can do so since  $\mathbb{F}$  is right-continuous). Define the auxiliary process  $\tau_{\mathbf{e}}^\epsilon := \mathbb{I}(\tau_{\mathbf{e}} > \epsilon)$  and denote by  $T^\epsilon$  the first-passage time associated to the collection  $\{\tau_{\mathbf{e}}^\epsilon : \mathbf{e} \in \mathcal{D}_e\}$ . Thus  $T^\epsilon(\mathbf{0}, \mathbf{n}) \leq \epsilon^{-1}T(\mathbf{0}, \mathbf{n})$ . Since  $\tau_{\mathbf{e}}^\epsilon$  has a Bernoulli distribution with parameter  $\mathbb{P}(\tau_{\mathbf{e}}^\epsilon = 0) = \mathbb{F}(\epsilon) < 1 - p_c^*$ , together with the previous case this yields Theorem 1.  $\square$

## Acknowledgment

This work was develop during my doctoral studies at Impa and I would like to thank my adviser, Prof. Vladas Sidoravicius, for his dedication and encouragement during this period. I also thank the whole administrative staff of IMPA for their assistance and CNPQ for financing my doctoral studies, without which this work would have not been possible.

## References

- [1] B. Bollobas and O. Riordan. The critical probability for random Voronoi percolation in the plane is  $1/2$ . Preprint available from arXiv.org:math/0410336.
- [2] B. Bollobas and O. Riordan. Sharp thresholds and percolation in the plane. Preprint available from arXiv.org:math/0412510.
- [3] G. Grimmett. Percolation (second edition). Springer (1999).
- [4] J.M. Hammersley, D.J.A. Welsh. First-passage percolation, sub-additive process, stochastic network and generalized renewal theory. Springer-Verlag (1965), 61-110.
- [5] H. Kesten. Aspects of first-passage percolation. *Lectures Notes in Math.* **1180**, Springer-Verlag (1986), 125-264.
- [6] T.M. Ligget, R.H. Schonmann and A.M. Stacey. Domination by product measures. *Ann. Probab.* **25** (1997), 71-95.
- [7] J. Moller. Lectures on random Voronoi tessellations. *Lectures Notes in Stat.* **87**, Springer-Verlag (1991).
- [8] L.P.R. Pimentel. Competing growth, interfaces and geodesics in first-passage percolation on Voronoi tilings. Phd Thesis, IMPA, Rio de Janeiro (2004).
- [9] M.Q. Vahidi-Asl and J.C. Wierman. First-passage percolation on the Voronoi tessellation and Delaunay triangulation. Random Graphs 87 (M. Karonske, J. Jaworski and A. Rucinski, eds.) Wiley (1990), 341-359.
- [10] M.C. Vahidi-Asl and J.C. Wierman. A shape result for first-passage percolation on the Voronoi tessellation and Delaunay triangulation. Random Graphs 89 (A. Frieze and T. Luczak, eds.), Wiley (1992), 247-262.
- [11] A. Zvavitch. The critical probability for Voronoi percolation. MSc. thesis, Weizmann Institute of Science (1996).