

Stein approximation for Itô and Skorohod integrals by Edgeworth type expansions

Nicolas Privault*

Abstract

We derive Edgeworth-type expansions for Skorohod and Itô integrals with respect to Brownian motion, based on cumulant operators defined by the Malliavin calculus. As a consequence we obtain Stein approximation bounds for stochastic integrals, which apply to SDE solutions and to multiple stochastic integrals.

Keywords: Stein method; cumulants; Malliavin calculus; Wiener space; Edgeworth expansions; Itô integral; Skorohod integral.

AMS MSC 2010: 60H07; 62E17; 60G15; 60H05.

Submitted to ECP on September 30, 2014, final version accepted on April 23, 2015.

1 Introduction

Classical Edgeworth series around the Gaussian cumulative distribution function $\Phi(x)$ take the form

$$\Phi(x) + c_1\phi(x)H_1(x) + \cdots + c_m\phi(x)H_m(x) + \cdots,$$

where $\phi(x)$, $x \in \mathbb{R}$, is the standard Gaussian density, $H_k(x)$ is the Hermite polynomial of degree $k \geq 1$, and c_k is a coefficient depending on the sequence of cumulants $(\kappa_n)_{n \geq 1}$ of a random variable F , cf. Chapter 5 of [6] and § A.4 of [11]. Edgeworth expansions are used in particular as asymptotic expansions for the cumulative distribution function $P(F \leq x)$ (or, in more general forms, as asymptotic expansions for expectations of the type $\mathbb{E}[h(F)]$, where h is some test function - see [6]), when F is centered with unit variance $\mathbb{E}[F^2] = 1$, for example when F is a renormalized sum of independent random variables that can be approximated by the central limit theorem, cf. Chapter 2 of [5].

In [1], Edgeworth type expansions of the form

$$\mathbb{E}[Ff(F) - f'(F)] = \sum_{l=2}^{\infty} \frac{\kappa_{l+1}}{l!} E[f^{(l)}(F)], \quad f \in \mathcal{C}^{\infty}(\mathbb{R}),$$

have been derived and connected to classical Edgeworth series for $\mathbb{E}[h(F)]$ by the Stein equation

$$h(F) = \mathbb{E}[h(X)] + Ff(F) - f'(F),$$

*School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore. E-mail: nprivault@ntu.edu.sg

where h is some adequate test function and $X \simeq \mathcal{N}(0, 1)$ is a standard Gaussian random variable. Recently, Edgeworth type expansions with exact remainder term of the form

$$\mathbb{E}[Ff(F) - f'(F)] = \sum_{l=2}^n \frac{\kappa_{l+1}}{l!} \mathbb{E}[f^{(l)}(F)] + \mathbb{E}[f^{(n+1)}(F)\Gamma_{n+1}F], \quad (1.1)$$

have been obtained by the Malliavin calculus in [8], [2], [3], written here for F a centered random variable with unit variance, where Γ_{n+1} is a cumulant type operator on the Wiener space satisfying the relation $n!\mathbb{E}[\Gamma_n F] = \kappa_{n+1}$, $n \in \mathbb{N}$, cf. [10]. This approach refines and extends the application of the Malliavin calculus to Stein approximation, Berry-Esseen bounds and the fourth moment theorem initiated in [9], see also [12], and [7] for a review.

The approaches of [2], [8], [9] and the cumulant operators of [10] rely on covariance identities based on the number (or Ornstein-Uhlenbeck) operator L and its inverse on the Wiener space, and they are particularly well suited to the study of multiple stochastic integrals.

In this paper we derive a Edgeworth type expansions for random variables represented as the Itô or Skorohod integral $F = \delta(u)$ of a process u on the Wiener space. Our expansions rely on properties of the operator δ , which coincides with the Itô stochastic integral with respect to d -dimensional Brownian motion on the square-integrable adapted processes, and are applied to Stein approximation bounds. Although this approach does not rely on the operator L , it nevertheless also covers the case of multiple stochastic integrals.

In Section 2 we derive expansions of the form (1.1) for $\mathbb{E}[\delta(u)f(\delta(u)) - f'(\delta(u))]$, based on a family of cumulant operators that are associated to the process u and specially defined for the Skorohod integral operator δ . In Section 3 we derive Stein type approximation bounds for stochastic integrals, and we apply them to the solutions of stochastic differential equations. In Section 4 we also provide an alternative approach to the results of [9] on multiple stochastic integrals.

Notation and cumulant operators for the Skorohod integral

Consider a standard d -dimensional Brownian motion $(B_t)_{t \in \mathbb{R}_+}$ generating the filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ on the Wiener space Ω . Letting $H = L^2(\mathbb{R}_+; \mathbb{R}^d)$, we consider the standard Sobolev spaces of real-valued, resp. H -valued, functionals $\mathbb{D}_{p,k}$, resp. $\mathbb{D}_{p,k}(H)$, $p, k \geq 1$, for the Malliavin gradient D on the Wiener space, cf. [13] for definitions. Recall that the Skorohod operator δ is the adjoint of the gradient D through the duality relation

$$\mathbb{E}[F\delta(v)] = \mathbb{E}[\langle DF, v \rangle_H], \quad F \in \text{Dom}(D), \quad v \in \text{Dom}(\delta), \quad (1.2)$$

and we have the commutation relation

$$D_t \delta(u) = u(t) + \delta(D_t u), \quad t \in \mathbb{R}_+, \quad (1.3)$$

provided $u \in \mathbb{D}_{2,1}(H)$ and $D_t u \in \text{Dom}(\delta)$, dt -a.e., cf. Proposition 1.3.2 of [13].

Next we define an operator composition $(Du)^k$ and its adjoint D^* in the sense of matrix powers with continuous indices. Namely, given $u \in \mathbb{D}_{2,1}(H)$ and $k \geq 1$, we let $(Du)^k$ denote the random operator on H almost surely defined by

$$(Du)^k h_s = \int_0^\infty \cdots \int_0^\infty (D_{t_k} u_s D_{t_{k-1}} u_{t_k} \cdots D_{t_1} u_{t_2}) h_{t_1} dt_1 \cdots dt_k, \quad s \in \mathbb{R}_+, \quad h \in H, \quad (1.4)$$

cf. e.g. § 7 of [17], [16], [15] for details. In the sequel we will simply denote $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_H$. The adjoint D^*u of Du on H satisfies

$$\langle (Du)v, h \rangle = \langle v, (D^*u)h \rangle, \quad h, v \in H,$$

and is given by

$$(D^*u)v_s = \int_0^\infty (D_s u_t)v_t dt, \quad s \in \mathbb{R}_+, \quad v \in L^2(W; H).$$

Given $u \in \mathbb{D}_{k,2}(H)$, our results will be based on a family of cumulant operators

$$\Gamma_k^u : \mathbb{D}_{2,1} \longrightarrow L^2(\Omega), \quad k \geq 1,$$

defined by $\Gamma_1^u F := \langle u, DF \rangle$ and

$$\Gamma_k^u F := F \langle (Du)^{k-2}u, u \rangle + F \langle D^*u, D((Du)^{k-2}u) \rangle + \langle (Du)^{k-1}u, DF \rangle, \quad k \geq 2.$$

Note that the operator Γ^u is directly relevant to the integrand u in the stochastic integral representation $\delta(u)$ and as such it differs from the Γ operator of [10] appearing in (1.1), in addition, those operators are not directly related to the Bakry-Émery-Ledoux Γ and Γ_2 operators.

Recall that by the proof of Lemma 3.1 in [14] we have

$$\langle D^*u, D((Du)^k v) \rangle_{H \otimes H} = \text{trace}((Du)^{k+1} Dv) + \sum_{i=2}^{k+1} \frac{1}{i} \langle (Du)^{k+1-i} v, D \text{trace}(Du)^i \rangle, \quad (1.5)$$

$u \in \mathbb{D}_{2,2}(H)$, $v \in \mathbb{D}_{2,1}(H)$, $k \in \mathbb{N}$, hence by the relation

$$\langle (Du)^k h, u \rangle = \langle (D^*u)^k u, h \rangle = \frac{1}{2} \langle (D^*u)^{k-1} D \langle u, u \rangle, h \rangle = \frac{1}{2} \langle (Du)^{k-1} h, D \langle u, u \rangle \rangle, \quad (1.6)$$

$h \in H$, $k \geq 1$, $u \in \mathbb{D}_{2,1}(H)$, which follows from $D \langle u, u \rangle = 2(D^*u)u$, for any $u \in \mathbb{D}_{2,2}(H)$ we have

$$\Gamma_k^u \mathbf{1} = \frac{1}{2} \langle (Du)^{k-3}u, D \langle u, u \rangle \rangle + \text{trace}(Du)^k + \sum_{i=2}^{k-1} \frac{1}{i} \langle (Du)^{k-1-i}u, D \text{trace}(Du)^i \rangle, \quad (1.7)$$

for all $k \geq 3$.

2 Edgeworth type expansions

The duality (1.2) and the commutation relation (1.3) show that

$$E[f'(\delta(u)) \langle u, u \rangle - \delta(u) f(\delta(u))] = -E[f'(\delta(u)) \langle u, \delta(Du) \rangle], \quad (2.1)$$

for $u \in \mathbb{D}_{1,2}(H)$, $F \in \mathbb{D}_{2,1}$ and $f \in \mathcal{C}_b^1(\mathbb{R})$. Applying the above relation (2.1) with $d = 1$ to the solution f_x of the Stein equation

$$\mathbf{1}_{(-\infty, x]}(z) - \Phi(z) = f'_x(z) - z f_x(z), \quad z \in \mathbb{R}, \quad (2.2)$$

satisfying the bounds $\|f_x\|_\infty \leq \sqrt{2\pi}/4$ and $\|f'_x\|_\infty \leq 1$, cf. Lemma 2.2-(v) of [4], yields the expansion

$$P(\delta(u) \leq x) - \Phi(x) = E[(1 - \langle u, u \rangle) f'_x(\delta(u))] + E[\langle u, \delta(Du) \rangle f'_x(\delta(u))], \quad x \in \mathbb{R},$$

around the Gaussian cumulative distribution function $\Phi(x)$, with $u \in \mathbb{D}_{1,2}(H)$. In the next proposition we extend (2.1) into an expansion of all orders that will be applied to Stein approximation in the next section. By comparison with Proposition 3.11 of [2], the last term in the expansion (2.3) below is not given by a cumulant operator.

Proposition 2.1. *Let $n \geq 1$ and assume that $u \in \mathbb{D}_{k,2}(H)$ for all $k = 1, \dots, n + 2$. Then for all $f \in \mathcal{C}_b^{n+1}(\mathbb{R})$ and $F \in \mathbb{D}_{2,1}$ we have*

$$E[F\delta(u)f(\delta(u))] = \sum_{k=0}^n E\left[f^{(k)}(\delta(u))\Gamma_{k+1}^u F\right] \quad (2.3)$$

$$+ \frac{1}{2}E\left[Ff^{(n+1)}(\delta(u))\langle(Du)^{n-1}u, D\langle u, u \rangle\rangle\right] + E\left[Ff^{(n+1)}(\delta(u))\langle(Du)^n u, \delta(Du)\rangle\right].$$

Proof. By the duality (1.2) between D and δ , the chain rule of derivation for D and the commutation relation (1.3), for $F \in \mathbb{D}_{2,1}$, $u \in \mathbb{D}_{n+1,2}(H)$, and all $k \in \mathbb{N}$ we have

$$\begin{aligned} & E[Ff(\delta(u))\langle(Du)^k u, \delta(D^*u)\rangle] - E[Ff'(\delta(u))\langle(D^*u)^{k+1}u, \delta(D^*u)\rangle] \\ &= E[\langle D^*u, D(Ff(\delta(u))(Du)^k u) \rangle] - E[Ff'(\delta(u))\langle(D^*u)^{k+1}u, \delta(D^*u)\rangle] \\ &= E[Ff'(\delta(u))\langle(Du)^{k+1}u, D\delta(u)\rangle] \\ &\quad - E[Ff'(\delta(u))\langle(D^*u)^{k+1}u, \delta(D^*u)\rangle] + E[f(\delta(u))\langle D^*u, D(F(Du)^k u) \rangle] \\ &= E[Ff'(\delta(u))\langle(Du)^{k+1}u, u \rangle] + E[Ff'(\delta(u))\langle(Du)^{k+1}u, \delta(D^*u)\rangle] \\ &\quad - E[Ff'(\delta(u))\langle(D^*u)^{k+1}u, \delta(D^*u)\rangle] + E[f(\delta(u))\langle D^*u, D(F(Du)^k u) \rangle] \\ &= E[Ff'(\delta(u))\langle(Du)^{k+1}u, u \rangle] + E[f(\delta(u))\langle D^*u, D(F(Du)^k u) \rangle] \\ &= E[Ff'(\delta(u))\langle(Du)^{k+1}u, u \rangle] + E[f(\delta(u))\langle(Du)^{k+1}u, DF \rangle] \\ &\quad + E[Ff(\delta(u))\langle D^*u, D((Du)^k u) \rangle], \end{aligned}$$

which shows that

$$\begin{aligned} & E[Ff(\delta(u))\langle(Du)^k u, \delta(D^*u)\rangle] - E[Ff'(\delta(u))\langle(D^*u)^{k+1}u, \delta(D^*u)\rangle] \quad (2.4) \\ &= E[Ff'(\delta(u))\langle(Du)^{k+1}u, u \rangle] - E[Ff(\delta(u))\langle D^*u, D((Du)^k u) \rangle] \\ &\quad + E[f(\delta(u))\langle(Du)^{k+1}u, DF \rangle]. \end{aligned}$$

Consequently, since $(Du)^{k-1}u \in \mathbb{D}_{(n+1)/k,1}(H)$ we have $\delta(u) \in \mathbb{D}_{(n+1)/(n-k+1),1}$, and by (2.4) we get

$$\begin{aligned} & E[Ff(\delta(u))\langle(Du)^k u, D\delta(u)\rangle] - E[Ff'(\delta(u))\langle(Du)^{k+1}u, D\delta(u)\rangle] \\ &= E[Ff(\delta(u))\langle(Du)^k u, u \rangle] + E[Ff(\delta(u))\langle(Du)^k u, \delta(D^*u)\rangle] \\ &\quad - E[Ff'(\delta(u))\langle(Du)^{k+1}u, u \rangle] - E[Ff'(\delta(u))\langle(Du)^{k+1}u, \delta(D^*u)\rangle] \\ &= E[Ff(\delta(u))\langle(Du)^k u, u \rangle] + E[Ff(\delta(u))\langle D^*u, D((Du)^k u) \rangle] + E[f(\delta(u))\langle(Du)^{k+1}u, DF \rangle] \\ &= E[f(\delta(u))\Gamma_{k+2}^u F], \end{aligned}$$

and therefore

$$\begin{aligned} E[F\delta(u)f(\delta(u))] &= E[Ff'(\delta(u))\langle u, D\delta(u) \rangle] + E[f(\delta(u))\langle u, DF \rangle] \\ &= E[f(\delta(u))\langle u, DF \rangle] + E\left[Ff^{(n+1)}(\delta(u))\langle(Du)^n u, D\delta(u)\rangle\right] \\ &\quad + \sum_{k=0}^{n-1} \left(E\left[Ff^{(k+1)}(\delta(u))\langle(Du)^k u, D\delta(u)\rangle\right] - E\left[Ff^{(k+2)}(\delta(u))\langle(Du)^{k+1}u, D\delta(u)\rangle\right] \right) \\ &= E[f(\delta(u))\langle u, DF \rangle] + \sum_{k=1}^n E\left[f^{(k)}(\delta(u))\Gamma_{k+1}^u F\right] + E\left[Ff^{(n+1)}(\delta(u))\langle(Du)^n u, D\delta(u)\rangle\right] \\ &= \sum_{k=0}^n E\left[f^{(k)}(\delta(u))\Gamma_{k+1}^u F\right] + E\left[Ff^{(n+1)}(\delta(u))\langle(Du)^n u, D\delta(u)\rangle\right] \\ &= E[f(\delta(u))\langle u, DF \rangle] + \sum_{k=1}^n E\left[f^{(k)}(\delta(u))\Gamma_{k+1}^u F\right] + E\left[Ff^{(n+1)}(\delta(u))\langle(Du)^n u, D\delta(u)\rangle\right] \end{aligned}$$

$$\begin{aligned}
 &= \sum_{k=0}^n E \left[f^{(k)}(\delta(u)) \Gamma_{k+1}^u F \right] + E \left[F f^{(n+1)}(\delta(u)) \langle (Du)^n u, u \rangle \right] \\
 &\quad + E \left[F f^{(n+1)}(\delta(u)) \langle (Du)^n u, \delta(Du) \rangle \right] \\
 &= \sum_{k=0}^n E \left[f^{(k)}(\delta(u)) \Gamma_{k+1}^u F \right] + \frac{1}{2} E \left[F f^{(n+1)}(\delta(u)) \langle (Du)^{n-1} u, D \langle u, u \rangle \rangle \right] \\
 &\quad + E \left[F f^{(n+1)}(\delta(u)) \langle (Du)^n u, \delta(Du) \rangle \right],
 \end{aligned}$$

where we used the relation (1.6). □

Based on Proposition 2.1 we make the following remarks for random isometries and quasi-nilpotent processes satisfying $\text{trace}(Du)^k = 0$, $k \geq 2$. Recall that the setting of quasi-nilpotent processes includes the particular case where $(u_t)_{t \in \mathbb{R}_+}$ is an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted process, cf. e.g. Lemma 3.5 of [15] and references therein, in which case $\delta(u)$ coincides with the Itô integral of u , cf. Proposition 1.3.11 of [13].

(i) Quasi-nilpotent processes. When $\text{trace}(Du)^k = 0$ for all $k = 2, \dots, n+1$ we have

$$\begin{aligned}
 E[\delta(u)f(\delta(u))] &= E[\langle u, u \rangle f'(\delta(u))] + \frac{1}{2} \sum_{k=2}^{n+1} E \left[\langle (Du)^{k-2} u, D \langle u, u \rangle \rangle f^{(k)}(\delta(u)) \right] \\
 &\quad + E \left[f^{(n+1)}(\delta(u)) \langle (Du)^n u, \delta(Du) \rangle \right], \quad n \geq 0.
 \end{aligned}$$

(ii) Random isometries. When $\langle u, u \rangle$ is deterministic we find

$$\begin{aligned}
 E[\delta(u)f(\delta(u))] &= \langle u, u \rangle E[f'(\delta(u))] + \sum_{k=1}^n E \left[\langle D^* u, D((Du)^{k-1} u) \rangle_{H \otimes H} f^{(k)}(\delta(u)) \right] \\
 &\quad + E \left[f^{(n+1)}(\delta(u)) \langle (Du)^n u, \delta(Du) \rangle \right], \quad n \geq 0.
 \end{aligned}$$

(iii) Multiple stochastic integral processes. Taking $u_t = I_n(f_{n+1}(*, t))$ where $n \in \mathbb{N}$ and f_{n+1} is a symmetric square-integrable function on \mathbb{R}_+^{n+1} , we have $\delta(u) = I_{n+1}(f_{n+1})$ and

$$\delta(D_t u) = n I_n(f_{n+1}(*, t)) = n u_t, \quad t \in \mathbb{R}_+. \tag{2.5}$$

Hence, applying again Proposition 2.1 and (1.6) to $u_t = I_{n-1}(f_n(*, t))$, $n \geq 1$, we get

$$\begin{aligned}
 &E[F I_n(f_n) f(I_n(f_n))] \\
 &= \sum_{k=0}^n E \left[f^{(k)}(I_n(f_n)) \Gamma_{k+1}^u F \right] + \frac{n}{2} E \left[F f^{(n+1)}(I_n(f_n)) \langle (Du)^{n-1} u, D \langle u, u \rangle \rangle \right].
 \end{aligned}$$

In the case of random and quasi-nilpotent isometries we get

$$E[\delta(u)f(\delta(u))] = \langle u, u \rangle E[f'(\delta(u))] + E \left[f^{(n+1)}(\delta(u)) \langle (Du)^n u, \delta(Du) \rangle \right],$$

which shows that $E[f^{(n+1)}(\delta(u)) \langle (Du)^n u, \delta(Du) \rangle] = 0$, $n \in \mathbb{N}$, and recovers the standard Gaussian integration by parts $E[\delta(u)f(\delta(u))] = \langle u, u \rangle E[f'(\delta(u))]$, cf. [18].

3 Stein approximation

From now on we work with $d = 1$ and a one-dimensional Brownian motion $(B_t)_{t \in \mathbb{R}_+}$, and we let $X \simeq \mathcal{N}(0, 1)$ denote a standard Gaussian random variable. In comparison with the results of [2], our bounds apply to a different stochastic integral representation.

Given $h : \mathbb{R} \rightarrow \mathbb{R}$ an absolutely continuous function with bounded derivative, the functional equation

$$h(z) - \mathbb{E}[h(X)] = f'(z) - zf(z), \quad z \in \mathbb{R}, \quad (3.1)$$

has a solution $f_h \in \mathcal{C}_b^1(\mathbb{R})$ which is twice differentiable and satisfies the bounds

$$\|f_h'\|_\infty \leq \|h'\|_\infty \quad \text{and} \quad \|f_h''\|_\infty \leq 2\|h'\|_\infty, \quad x \in \mathbb{R},$$

cf. Lemma 1.2-(v) of [9] and references therein. Let

$$d(F, G) = \sup_{h \in \mathcal{L}} |\mathbb{E}[h(F)] - \mathbb{E}[h(G)]|$$

denote the Wasserstein distance between the laws of F and G , where \mathcal{L} denotes the class of 1-Lipschitz functions. In the sequel we let $\|u\|_2 = \|u\|_{L^2(\Omega \times \mathbb{R}_+)}$.

Proposition 3.1. *Let $u \in \bigcap_{k=1}^3 \mathbb{D}_{k,2}(H)$. We have*

$$d(\delta(u), X) \leq E [1 - \langle u, u \rangle - \text{trace}(Du)^2] + \|u\|_2 \|D\langle u, u \rangle\|_2 + 2E [|\langle (Du)u, \delta(Du) \rangle|]. \quad (3.2)$$

Proof. For $n = 1$ and $F = 1$, Proposition 2.1 shows that

$$\begin{aligned} E[\delta(u)f(\delta(u))] &= E[f'(\delta(u))\Gamma_2^u \mathbf{1}] \\ &\quad + \frac{1}{2}E[f''(\delta(u))\langle u, D\langle u, u \rangle \rangle] + E[f''(\delta(u))\langle (Du)u, \delta(Du) \rangle], \end{aligned}$$

hence for any continuous function $h : \mathbb{R} \rightarrow [0, 1]$, denoting by f_h the solution to (3.1) we have

$$\begin{aligned} E[h(\delta(u))] - E[h(X)] &= E[\delta(u)f_h(\delta(u)) - f_h'(\delta(u))] \\ &= E[f_h'(\delta(u))(\Gamma_2^u \mathbf{1} - 1)] + \frac{1}{2}E[f_h''(\delta(u))\langle u, D\langle u, u \rangle \rangle] + 2E[f_h''(\delta(u))\langle (Du)u, \delta(Du) \rangle], \end{aligned}$$

hence

$$\begin{aligned} |E[\delta(u)h(\delta(u))] - E[h(X)]| \\ \leq \|h'\|_\infty E [1 - \Gamma_2^u \mathbf{1}] + \|h'\|_\infty E [|\langle u, D\langle u, u \rangle \rangle|] + 2\|h'\|_\infty E [|\langle (Du)u, \delta(Du) \rangle|], \end{aligned}$$

which yields (3.2) by the relation

$$\Gamma_2^u \mathbf{1} = \langle u, u \rangle + \langle D^*u, Du \rangle_{H \otimes H} = \langle u, u \rangle + \text{trace}(Du)^2. \quad (3.3)$$

□

By (3.2) and (3.3) we find

$$d(\delta(u), X) \leq \|1 - \langle u, u \rangle\|_2 + \|\text{trace}(Du)^2\|_2 + \|u\|_2 \|D\langle u, u \rangle\|_2 + 2E [|\langle (Du)u, \delta(Du) \rangle|],$$

which, as in Section 2, yields the following remarks.

- (i) Quasi-nilpotent processes. When $\text{trace}(Du)^2 = 0$, and in particular when $(u_t)_{t \in \mathbb{R}_+}$ is an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted process, we have

$$d(\delta(u), X) \leq E[|1 - \langle u, u \rangle|] + \|u\|_2 \|D\langle u, u \rangle\|_2 + 2E[|\langle (Du)u, \delta(Du) \rangle|]. \quad (3.4)$$

- (ii) Random isometries. When $\langle u, u \rangle$ is deterministic we find

$$d(\delta(u), X) \leq |1 - \langle u, u \rangle| + \|\text{trace}(Du)^2\|_2 + 2E[|\langle (Du)u, \delta(Du) \rangle|].$$

As another consequence of Proposition 3.1 and of the Skorohod isometry

$$\text{Var}[\delta(u)] = E[\delta(u)^2] = E[\langle u, u \rangle] + E[\text{trace}(Du)^2],$$

we also find the bound

$$\begin{aligned} d(\delta(u), X) \leq & |1 - \text{Var}[\delta(u)]| + \sqrt{\text{Var}[\|u\|_H^2 + \text{trace}(Du)^2]} \\ & + \|u\|_2 \|D\langle u, u \rangle\|_2 + 2E[|\langle (Du)u, \delta(Du) \rangle|], \end{aligned} \quad (3.5)$$

for $u \in \bigcap_{k=1}^3 \mathbb{D}_{k,2}(H)$, which will be applied below to multiple stochastic integrals. In particular we have the following.

- (i) Quasi-nilpotent processes. When $\text{trace}(Du)^2 = 0$ the bound (3.5) yields

$$d(\delta(u), X) \leq |1 - \text{Var}[\delta(u)]| + \sqrt{\text{Var}[\|u\|_H^2]} + \|u\|_2 \|D\langle u, u \rangle\|_2 + 2E[|\langle (Du)u, \delta(Du) \rangle|].$$

- (ii) Unit variance. In case $\text{Var}[\delta(u)] = 1$, (3.5) shows that

$$d(\delta(u), X) \leq \sqrt{E[(\|u\|_H^2 + \text{trace}(Du)^2)^2]} - 1 + \|u\|_2 \|D\langle u, u \rangle\|_2 + 2E[|\langle (Du)u, \delta(Du) \rangle|].$$

- (iii) Multiple stochastic integral processes. Taking $u_t = I_{n-1}(f_n(*, t))$ where f_n is a symmetric square-integrable function on \mathbb{R}_+^n , and applying again (1.6) and (2.5), by (3.5) we get

$$d(I_n(f_n), X) \leq |1 - n! \|f_n\|_2^2| + \sqrt{\text{Var}[\|u\|_H^2 + \text{trace}(Du)^2]} + (n-1)E[|\langle u, D\langle u, u \rangle \rangle|]. \quad (3.6)$$

The above bound (3.6) will be computed in terms of the kernel function f_n in the next section.

When $\delta(u)$ has unit variance and in addition $\text{trace}(Du)^2 = 0$ or $(u_t)_{t \in \mathbb{R}_+}$ is an adapted process, we find

$$d(\delta(u), X) \leq \sqrt{E[\|u\|_H^4]} - 1 + \|u\|_2 \|D\langle u, u \rangle\|_2 + 2E[|\langle (Du)u, \delta(Du) \rangle|].$$

4 Applications

a) Stochastic differential equations

Consider the stochastic differential equation

$$dX_t = \sigma(X_t) dW_t, \quad X_0 = x_0,$$

where $\sigma \in C_b^1(\mathbb{R})$. From Theorem 2.2.1 and Exercise 2.2.1 of [13], we have $X_t \in \text{Dom}(D)$, $t \in [0, T]$, and

$$D_s X_r = \mathbf{1}_{[0,r]}(s)\sigma(X_s)e^{\int_s^r \sigma'(X_u)dW_u - \int_s^r |\sigma'(X_u)|^2 du/2}, \quad 0 \leq s \leq r. \tag{4.1}$$

Since $X_T = \delta(\mathbf{1}_{[0,T]}\sigma(X))$, and taking $H = L^2([0, T])$, from (3.4) we get

$$d(X_T, X) \leq E[|1 - \langle \sigma(X), \sigma(X) \rangle|] + \|\sigma(X)\|_2 \|D\langle \sigma(X), \sigma(X) \rangle\|_2 + 2E[|\langle (D\sigma(X))\sigma(X), \delta(D\sigma(X)) \rangle|], \tag{4.2}$$

where $\langle (D\sigma(X))\sigma(X), \delta(D\sigma(X)) \rangle$ is given by

$$D_r \langle \sigma(X), \sigma(X) \rangle = 2\sigma(X_r) \int_r^T \sigma(X_s)\sigma'(X_s)e^{\int_r^s \sigma'(X_u)dW_u - \int_r^s |\sigma'(X_u)|^2 du/2} ds, \quad r \in \mathbb{R}_+.$$

In order to bound the last term in (4.2) we note that

$$\delta(D_r \sigma(X)) = \sigma(X_r) \int_r^T \sigma'(X_t)e^{\int_r^t \sigma'(X_u)dW_u - \int_r^t |\sigma'(X_u)|^2 du/2} dW_t, \quad 0 \leq r \leq T,$$

and by (4.1) we have

$$\begin{aligned} & \langle (D\sigma(X))\sigma(X), \delta(D\sigma(X)) \rangle \\ &= \int_0^T \int_0^T \sigma(X_s)D_s \sigma(X_r)ds\delta(D_r \sigma(X))dr \\ &= \int_0^T \sigma'(X_t) \int_0^t |\sigma(X_s)|^2 e^{\int_s^t \sigma'(X_u)dW_u - \int_s^t |\sigma'(X_u)|^2 du/2} \int_s^t \sigma(X_r)\sigma'(X_r)drdsdW_t, \end{aligned}$$

hence the last term in (4.2) can be bounded as

$$\begin{aligned} & E[|\langle (D\sigma(X))\sigma(X), \delta(D\sigma(X)) \rangle|] \\ & \leq \sqrt{E \left[\int_0^T |\sigma'(X_t)|^2 \left(\int_0^t |\sigma(X_s)|^2 e^{\int_s^t \sigma'(X_u)dW_u - \int_s^t |\sigma'(X_u)|^2 du/2} \int_s^t \sigma(X_r)\sigma'(X_r)drds \right)^2 dt \right]} \\ & \leq \sqrt{E \left[\int_0^T t |\sigma'(X_t)|^2 \int_0^t |\sigma(X_s)|^4 e^{2 \int_s^t \sigma'(X_u)dW_u - \int_s^t |\sigma'(X_u)|^2 du} (t-s) \int_s^t |\sigma(X_r)\sigma'(X_r)|^2 drdsdt \right]} \\ & \leq \frac{T^{5/2}}{\sqrt{15}} \|\sigma\|_\infty^3 \|\sigma'\|_\infty^2 e^{T\|\sigma'\|_\infty/2}, \end{aligned}$$

hence (4.2) provides an asymptotic bound on the distance $d(X_T, X)$ as $\|\sigma'\|_\infty$ tends to 0.

b) Multiple stochastic integrals

We now show that (3.5) can be used to recover the results [9] on multiple stochastic integrals. From (3.6) we get

$$d(I_n(f_n), X) \leq |1 - n! \|f_n\|_2^2| + \sqrt{\text{Var}[\|u\|_H^2 + \text{trace}(Du)^2]} + (n-1)\|u\|_2 \|D\langle u, u \rangle\|_2.$$

By the multiplication formula for multiple stochastic integrals, cf. e.g. Relation (2.29) in [9] we have

$$\langle u, u \rangle = \int_0^\infty (I_{n-1}(f_n(*, t)))^2 dt = \sum_{k=1}^n (k-1)! \binom{n-1}{k-1}^2 I_{2n-2k}(f_n \otimes_k f_n),$$

and, since $D_s u(t) = (n - 1)I_{n-2}(f_n(*, s, t))$,

$$\begin{aligned} \text{trace}(Du)^2 &= (n - 1)^2 \int_0^\infty \int_0^\infty I_{n-2}(f_n(*, s, t))I_{n-2}(f_n(*, t, s))dsdt \\ &= (n - 1)^2 \sum_{k=0}^{n-2} k! \binom{n-2}{k}^2 \int_0^\infty \int_0^\infty I_{2n-4-2k}(f_n(*, s, t) \otimes_k f_n(*, s, t))dsdt \\ &= (n - 1)^2 \sum_{k=2}^n (k - 2)! \binom{n-2}{k-2}^2 I_{2n-2k}(f_n \otimes_k f_n), \end{aligned}$$

hence

$$\begin{aligned} \Gamma_2^u \mathbf{1} &= I_{2n-2}(f_n \otimes_1 f_n) \\ &\quad + (n - 1)^2 \sum_{k=2}^n \left((k - 2)! \binom{n-2}{k-2}^2 + (k - 1)! \binom{n-1}{k-1}^2 \right) I_{2n-2k}(f_n \otimes_k f_n) \\ &= \sum_{k=1}^n k! \binom{n-1}{k-1}^2 I_{2n-2k}(f_n \otimes_k f_n), \end{aligned}$$

and

$$\text{Var}[\Gamma_2^u \mathbf{1}] = \sum_{k=1}^{n-1} k!^2 \binom{n-1}{k-1}^4 \|f_n \otimes_k f_n\|_2^2.$$

We also have

$$D_r \langle u, u \rangle = 2 \sum_{k=1}^{n-1} (n - k)(k - 1)! \binom{n-1}{k-1}^2 I_{2n-2k-1}((f_n \otimes_k f_n)(* , r)),$$

hence

$$\begin{aligned} E \left[\int_0^\infty |D_r \langle u, u \rangle|^2 dr \right] &= 4 \sum_{k=1}^{n-1} ((n - k)(k - 1)!)^2 \binom{n-1}{k-1}^4 \int_0^\infty \|(f_n \otimes_k f_n)(* , r)\|_2^2 dr \\ &= 4 \sum_{k=1}^{n-1} ((n - k)(k - 1)!)^2 \binom{n-1}{k-1}^4 \|f_n \otimes_k f_n\|_2^2. \end{aligned}$$

Finally we get

$$\begin{aligned} d(I_n(f_n), X) &\leq |1 - n! \|f_n\|_2^2| + \sqrt{\sum_{k=1}^{n-1} k!^2 \binom{n-1}{k-1}^4 \|f_n \otimes_k f_n\|_2^2} \\ &\quad + 2(n - 1)\sqrt{(n - 1)! \|f_n\|_2} \sqrt{\sum_{k=1}^{n-1} ((n - k)(k - 1)!)^2 \binom{n-1}{k-1}^4 \|f_n \otimes_k f_n\|_2^2}, \end{aligned}$$

which recovers Proposition 3.2 of [9], with different constants.

References

- [1] A.D. Barbour. Asymptotic expansions based on smooth functions in the central limit theorem. *Probab. Theory Relat. Fields*, 72(2):289–303, 1986.
- [2] H. Biermé, A. Bonami, I. Nourdin, and G. Peccati. Optimal Berry-Esseen rates on the Wiener space: the barrier of third and fourth cumulants. *ALEA Lat. Am. J. Probab. Math. Stat.*, 9(2):473–500, 2012.

- [3] S. Campese. Optimal convergence rates and one-term edge-worth expansions for multidimensional functionals of Gaussian fields. *ALEA Lat. Am. J. Probab. Math. Stat.*, 10(2):881–919, 2013.
- [4] L.H.Y. Chen, L. Goldstein, and Q.-M. Shao. *Normal approximation by Stein's method*. Probability and its Applications (New York). Springer, Heidelberg, 2011.
- [5] P. Hall. *The bootstrap and Edgeworth expansion*. Springer Series in Statistics. Springer-Verlag, New York, 1992.
- [6] P. McCullagh. *Tensor methods in statistics*. Monographs on Statistics and Applied Probability. Chapman & Hall, London, 1987.
- [7] I. Nourdin. Lectures on Gaussian approximations with Malliavin calculus. In *Séminaire de Probabilités XLV*, volume 2078 of *Lecture Notes in Math.*, pages 3–89. Springer, 2013.
- [8] I. Nourdin and G. Peccati. Stein's method and exact Berry-Esseen asymptotics for functionals of Gaussian fields. *Ann. Probab.*, 37(6):2231–2261, 2009.
- [9] I. Nourdin and G. Peccati. Stein's method on Wiener chaos. *Probab. Theory Related Fields*, 145(1-2):75–118, 2009.
- [10] I. Nourdin and G. Peccati. Cumulants on the Wiener space. *J. Funct. Anal.*, 258(11):3775–3791, 2010.
- [11] I. Nourdin and G. Peccati. *Normal approximations with Malliavin calculus*, volume 192 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2012.
- [12] I. Nourdin and G. Peccati. The optimal fourth moment theorem. *Proc. Amer. Math. Soc.*, 143:3123–3133, 2015.
- [13] D. Nualart. *The Malliavin calculus and related topics*. Probability and its Applications. Springer-Verlag, Berlin, second edition, 2006.
- [14] N. Privault. Moment identities for Skorohod integrals on the Wiener space and applications. *Electron. Commun. Probab.*, 14:116–121 (electronic), 2009.
- [15] N. Privault. Laplace transform identities and measure-preserving transformations on the Lie-Wiener-Poisson spaces. *J. Funct. Anal.*, 263:2993–3023, 2012.
- [16] N. Privault. Cumulant operators and moments of the Itô and Skorohod integrals. *C. R. Math. Acad. Sci. Paris*, 351(9-10):397–400, 2013.
- [17] N. Privault. Cumulant operators for Lie-Wiener-Itô-Poisson stochastic integrals. *J. Theoret. Probab.*, 28(1):269–298, 2015.
- [18] A. S. Üstünel and M. Zakai. Random rotations of the Wiener path. *Probab. Theory Relat. Fields*, 103(3):409–429, 1995.

Acknowledgments. This work was supported by NTU MOE Tier 2 Grant No. M4020140.