



Vol. 13 (2008), Paper no. 45, pages 1283–1306.

Journal URL

<http://www.math.washington.edu/~ejpecp/>

Semiclassical analysis and a new result for Poisson - Lévy excursion measures

Ian M Davies

Department of Mathematics, Swansea University,
Singleton Park, Swansea, SA2 8PP, Wales, UK

Email: I.M.Davies@Swansea.ac.uk

Abstract

The Poisson-Lévy excursion measure for the diffusion process with small noise satisfying the Itô equation $dX^\varepsilon = b(X^\varepsilon(t))dt + \sqrt{\varepsilon}dB(t)$ is studied and the asymptotic behaviour in ε is investigated. The leading order term is obtained exactly and it is shown that at an equilibrium point there are only two possible forms for this term - Lévy or Hawkes - Truman. We also compute the next to leading order.

Key words: excursion measures; asymptotic expansions.

AMS 2000 Subject Classification: Primary 60H10, 41A60.

Submitted to EJP on March 22, 2007, final version accepted June 5, 2008.

1 Introduction

Consider a one-dimensional diffusion process defined by

$$dX(t) = b(X(t)) dt + dB(t), \quad X(0) = a,$$

where b is a Lipschitz-continuous function and $B(t)$ is a standard Brownian motion. The generator, G , of the above diffusion is

$$G = \frac{1}{2} \frac{d^2}{dx^2} + b(x) \frac{d}{dx},$$

and the putative invariant density is

$$\rho_0(x) = \exp \left(2 \int^x b(u) du \right).$$

If $\rho_0 \in L^1(\mathbb{R}, dx)$ the boundary $\{-\infty, \infty\}$ is inaccessible. We assume this in what follows. The transition density

$$p_t(x, y) = \mathbb{P}(X(t) \in dy | X(0) = x) / dy,$$

satisfies

$$\begin{aligned} \frac{\partial p_t(x, y)}{\partial t} &= \frac{\partial}{\partial y} \left(\frac{1}{2} \frac{\partial p_t(x, y)}{\partial y} - b(y) p_t(x, y) \right), \\ &= (G_y^* p_t)(x, y), \end{aligned}$$

$$\lim_{t \downarrow 0} p_t(x, y) = \delta_x(y),$$

G_y^* being the L^2 adjoint of G_y , and δ being the Dirac delta function. The density of the diffusion

$$\rho^t(y) = \int \rho_0(x) p_t(x, y) dx$$

therefore satisfies

$$\frac{\partial \rho^t}{\partial t} = (G_y^* \rho^t)(y).$$

Evidently,

$$(G_y^* \rho_0)(y) = 0, \quad \frac{\partial \rho_0(y)}{\partial t} = 0,$$

so ρ_0 is the invariant density.

Crucial in what follows is the operator identity for any well-behaved f

$$Gf = - \left(\rho_0^{-1/2} H \rho_0^{1/2} \right) f,$$

or

$$G = - \left(\rho_0^{-1/2} H \rho_0^{1/2} \right)$$

and

$$G^* = - \left(\rho_0^{1/2} H \rho_0^{-1/2} \right),$$

where H is the one dimensional Schrödinger operator with potential $V = \frac{1}{2}(b^2 + b')$,

$$H = -\frac{1}{2} \frac{d^2}{dx^2} + V(x).$$

This follows because $(H\rho_0^{1/2}) \equiv 0$, i.e. $\rho_0^{1/2}$ is the ground state of H . For convenience, we will assume $V \in C^2(\mathbb{R})$, V bounded below together with V'' , V polynomially bounded with derivatives.

2 Excursion Theory

The map $s \mapsto X(s)$ is continuous and so $\{s > 0 : X(s) \geq a\}$ is an open subset of \mathbb{R} . Therefore, $\{s > 0 : X(s) \geq a\}$ can be decomposed into a countable union of open intervals – $\begin{matrix} \text{upward} \\ \text{downward} \end{matrix}$ excursion intervals. Define

$$L^\pm(t) = \text{Leb}\{s \in [0, t] : X(s) \geq a\},$$

and the local time at a

$$L^a(t) = \lim_{h \downarrow 0} h^{-1} \text{Leb}\{s \in [0, t] : X(s) \in (a - h/2, a + h/2)\}.$$

$L^a(t)$ has inverse $\gamma^a(t)$, the time required to wait until L^a equals t . It can be seen that $\gamma^a(t)$ is a stopping time with $X(\gamma^a(t)) = a$. Moreover, as is intuitively obvious,

$$\text{Jumps in } \gamma^a(t) = \text{Excursions of } X \text{ from } a \text{ up to } L^a \text{ equals } t.$$

Example 1 Lévy [1954]

Lévy proved that for $b \equiv 0$, for each $\lambda > 0$,

$$\mathbb{E}_a \exp(-\lambda \gamma^a(t)) = \exp \left\{ -t \int_0^\infty (1 - e^{-\lambda s}) d\nu_a(s) \right\},$$

with Poisson-Lévy excursion measure

$$\nu_a[s, \infty) = \left(\frac{2}{\pi}\right)^{1/2} s^{-1/2}, s > 0. \tag{2.1}$$

Equating powers of λ in the above, we conclude that

$$\sharp(s, t) = \text{Number of excursions of duration exceeding } s \text{ up to } L^a \text{ equals } t$$

is Poisson with

$$\mathbb{P}(\sharp(s, t) = N) = \exp(-t\nu_a[s, \infty)) (t\nu_a[s, \infty))^N / N!,$$

for $N = 0, 1, 2, \dots$, and so the expected number of excursions of duration exceeding s per unit local time at a is $\nu_a[s, \infty)$, the Poisson-Lévy excursion measure.

Example 2 Hawkes and Truman [1991]

For the Ornstein-Uhlenbeck process $b(x) = -kx$, where k is a positive constant, the Hamiltonian is just

$$H = \frac{1}{2} \left(-\frac{d^2}{dx^2} + k^2x^2 - k \right)$$

and $\rho_0(x) = C \exp(-kx^2)$. This leads to

$$\mathbb{E}_0 \exp(-\lambda\gamma^a(t)) = \exp \left\{ -t \int_0^\infty (1 - e^{-\lambda s}) d\nu_0(s) \right\}, \lambda > 0,$$

with

$$\nu_0[s, \infty) = \frac{2k^{1/2}}{\pi^{1/2}} \left(e^{2ks} - 1 \right)^{-1/2}. \tag{2.2}$$

We discuss generalisations of the above to upward and downward excursions. Note that ^{upward}_{downward} excursions can only be affected by values of $b(x)$ for $x \gtrless a$. Therefore it is natural to define the symmetrised potential

$$V_{\text{symm}}^+ = \begin{cases} V(x), & x > a, \\ V(2a - x), & x < a. \end{cases}$$

with V_{symm}^- being defined in a similar manner. In an analogous manner we also define

$$H^\pm = -\frac{1}{2} \frac{d^2}{dx^2} + V_{\text{symm}}^\pm(x).$$

We now have the result due to Truman and Williams [1991]

Proposition 1. *Modulo the above assumptions*

$$\mathbb{E}_a \exp(-\lambda L^\pm(\gamma^a(t))) = \exp \left\{ -t \int_0^\infty (1 - e^{-\lambda s}) d\nu_a^\pm(s) \right\}$$

with

$$\nu_a^\pm[t, \infty) = \int_{y \gtrless a} dy \frac{\rho_0^{1/2}(y)}{\rho_0^{1/2}(a)} \frac{\partial}{\partial x} \Big|_{x=a} \exp(-tH^\pm)(x, y).$$

Remarks

1. $L^\pm(\gamma^a(t))$ are independent with $L^+(\gamma^a(t)) + L^-(\gamma^a(t)) = \gamma^a(t)$.
2. Jumps in $L^\pm(\gamma^a(t)) = \begin{smallmatrix} \text{upward} \\ \text{downward} \end{smallmatrix}$ excursions from a up to L^a equals t .
3. $\nu_a^\pm[s, \infty)$ is the expected number of ^{upward}_{downward} excursions of duration exceeding s per unit local time at a .

Proof. (Outline) The proof uses the result of Lévy [1954]

$$\mathbb{E}_a \exp(-\lambda \gamma^a(t)) = \exp(-t/\tilde{p}_\lambda(a, a)),$$

where $\tilde{p}_\lambda(x, y) = \int_0^\infty e^{-\lambda s} p_s(x, y) ds$ and $p_s(\cdot, \cdot)$ is the transition density.

We can deduce that

$$\tilde{p}_\lambda^{-1}(a, a) = \lambda \int_{-\infty}^\infty \frac{\rho_0(x)}{\rho_0(a)} \mathbb{E} e^{-\lambda \tau_x(a)} dx,$$

where $\tau_x(a) = \inf\{s > 0 : X(s) = a | X(0) = x\}$. Here the point is that for any point a intermediate to x and y

$$p_t(x, y) = \int_0^\infty \mathbb{P}(\tau_x(a) \in du) p_{t-u}(a, y).$$

Since the right hand side is a convolutional product, taking Laplace transforms and letting $y \rightarrow a$ gives

$$\mathbb{E} e^{-\lambda \tau_x(a)} = \tilde{p}_\lambda(x, a) / \tilde{p}_\lambda(a, a).$$

Now multiply both sides by $\rho_0(x)$ and integrate with respect to x (using the fact that ρ_0 is the invariant density) to get the desired result for \tilde{p}_λ^{-1} . Some elementary computation then leads to the result in Proposition 1.

3 The Poisson-Lévy Excursion Measure for Small Noise

We will now consider the $\begin{smallmatrix} \text{upward} \\ \text{downward} \end{smallmatrix}$ excursions from the equilibrium point 0 for the one-dimensional time-homogeneous diffusion process with small noise, $X^\varepsilon(t)$, where

$$dX^\varepsilon(t) = b(X^\varepsilon(t)) dt + \sqrt{\varepsilon} dB(t).$$

Introducing the small noise term into the Truman-Williams Law seen in the previous section, we get:

Proposition 2. *The expected number of $\begin{smallmatrix} \text{upward} \\ \text{downward} \end{smallmatrix}$ excursions from 0 of duration exceeding s , per unit local time at 0 is given by*

$$\nu_0^\pm[s, \infty) = \pm \int_{y \gtrless 0} \frac{\rho_0^{\frac{1}{2}}(y)}{\rho_0^{\frac{1}{2}}(0)} \varepsilon \frac{\partial}{\partial x} \Big|_{x=0} \exp\left(-\frac{sH^\pm}{\varepsilon}\right)(x, y) dy,$$

where ρ_0 is the invariant density and H^\pm is the symmetrized Hamiltonian for $V = \frac{1}{2}(b^2 + \varepsilon b')$.

One should note the form of V , in particular the presence of ε as a multiplier of b' . This rather specific dependence originates from the Shrödinger operator mentioned earlier. Consequently, we are unable to resort to the usual methods for resolving such a dependence.

We now give a result due to Davies and Truman Davies and Truman [1982].

Proposition 3. Let $X_{min}(\cdot)$ be the minimising path for the classical action

$$A(z) = 2^{-1} \left(\int_0^t \dot{z}^2(s) ds + \int_0^t b^2(z(s)) ds \right) \text{ with } z(0) = x, z(t) = y.$$

Set $A(X_{min}) = A(x, y, t)$. Then for the self-adjoint quantum mechanical Hamiltonian $H(\varepsilon) = \left[-\frac{\varepsilon^2}{2} \Delta + V_\varepsilon \right]$, where $V_\varepsilon = \frac{1}{2}(b^2 + \varepsilon b') \in C^\infty(R)$ and is convex (where $V_\varepsilon = V_0 + \varepsilon V_1 \in C^4$, bounded below with $V_0'' \geq -|\beta|$), then for each finite time $t \geq 0$ (for $t \leq \pi/|\beta|^{\frac{1}{2}}$).

$$\begin{aligned} & \exp \left(-\frac{tH(\varepsilon)}{\varepsilon} \right) (x, y) \\ &= (2\pi\varepsilon)^{-\frac{1}{2}} \exp \left(-\frac{A(x, y, t)}{\varepsilon} \right) \left\{ \left| \frac{\partial^2 A(x, y, t)}{\partial x \partial y} \right|^{\frac{1}{2}} \left(1 + \varepsilon K + O(\varepsilon^2) \right) \right\}. \end{aligned}$$

K is a rather complicated expression with many terms involving sums and products of b (and its derivatives), V (and its derivatives) and the Feynman-Green function $G(\tau, \sigma)$ of the Sturm-Liouville differential operator $\frac{d^2}{d\sigma^2} - V''(X_{min}(\tau))$ with zero boundary conditions i.e. $G(0, \tau) = G(t, \tau) = 0$, and discontinuity of derivative across $\tau = \sigma$ of 1.

For a proof of this result see Davies and Truman [1982].

Henceforth, for simplicity we assume that $b^2(x)$ is an even function of x so that $\nu_0^+ = \nu_0^-$.

Theorem 1. Using the notation and assumptions of Proposition 3, the leading term of the Poisson-Lévy excursion measure, for excursions away from the position of stable equilibrium 0, where $b(0) = 0$, and $b'(0) \leq 0$ is given by

$$\begin{aligned} \nu_0^+[t, \infty) &\sim (2\pi\varepsilon)^{-\frac{1}{2}} \int_0^\infty dy \exp \left\{ -\frac{y^2}{2\varepsilon} \left(\frac{\partial^2 A(0, 0, t)}{\partial y^2} - b'(0) \right) - \frac{A(0, 0, t)}{\varepsilon} \right\} \\ &\times \left\{ \left| \frac{\partial^2 A}{\partial x \partial y} \right|_{x=0}^{\frac{1}{2}} \left(-\frac{\partial A}{\partial x} \right)_{x=0} \exp \left(\frac{1}{2} \int_0^t |b'(X_{min}(s))| ds \right) \right\}, \end{aligned}$$

with the action $A(x, y, t) = \frac{1}{2} \int_0^t \dot{z}^2(s) ds + \int_0^t V(z(s)) ds$, where $V = \frac{1}{2}b^2$.

Proof. As usual, the classical path $X_{min}(t) = X(x, y, t)$ satisfies, correct to first order in ε

$$\ddot{X}_{min} \sim V_0'(X_{min}) + \varepsilon V_1'(X_{min}).$$

For $V_0 = \frac{1}{2}b^2$ (assumed to be convex with $V_0(0) = 0$, $V_0'(0) = 0$, and $V_0''(0) > 0$, for example $V_0(x) = \frac{1}{2}x^2$, $b(x) = -x$), and $V_1 = \frac{1}{2}b'$, then to leading order $\ddot{X}_{min} = V_0'(X_{min})$.

The contribution to the action $A(x, y, t) = \frac{1}{2} \int_0^t \dot{z}^2 ds + \int_0^t V(z(s)) ds$ from V_1 is to leading order

$$\begin{aligned} \varepsilon \int_0^t V_1(X_{min}(s)) ds &= \frac{\varepsilon}{2} \int_0^t b'(X_{min}(s)) ds \\ &= -\frac{\varepsilon}{2} \int_0^t |b'(X_{min}(s))| ds, \end{aligned}$$

introducing $|\cdot|$ for convenience. Therefore, from Proposition 3,

$$\exp\left(-\frac{tH(\varepsilon)}{\varepsilon}\right)(x, y) \sim (2\pi\varepsilon)^{-\frac{1}{2}} \exp\left(-\frac{A(x, y, t)}{\varepsilon}\right) \left|\frac{\partial^2 A(x, y, t)}{\partial x \partial y}\right|^{\frac{1}{2}},$$

and so, the contribution to term $\exp\left(-\frac{A(x, y, t)}{\varepsilon}\right)$ from V_1 is

$$\exp\left(\frac{1}{2} \int_0^t |b'(X_{min}(s))| ds\right), \quad \text{the Zero Point Energy term.}$$

Therefore, we have using Proposition 2 the leading order term in the Poisson-Lévy excursion measure, for upward excursions from stable equilibrium point 0 given by

$$\begin{aligned} & \nu_0^+[t, \infty) \\ & \sim (2\pi\varepsilon)^{-\frac{1}{2}} \int_0^\infty dy \exp\left(\frac{1}{\varepsilon} \int_0^y b(u) du\right) \\ & \times \varepsilon \frac{\partial}{\partial x} \Big|_{x=0} \left[\exp\left(\frac{1}{2} \int_0^t |b'(X_{min}(s))| ds\right) \exp\left(-\frac{A(x, y, t)}{\varepsilon}\right) \left|\frac{\partial^2 A}{\partial x \partial y}\right|^{\frac{1}{2}} \right]. \end{aligned}$$

Hence, for small ε , the leading order term is

$$\begin{aligned} \nu_0^+[t, \infty) & \sim (2\pi\varepsilon)^{-\frac{1}{2}} \int_0^\infty dy \exp\left(\frac{1}{\varepsilon} \left(\int_0^y b(u) du - A(0, y, t)\right)\right) \\ & \times \left[\left|\frac{\partial^2 A}{\partial x \partial y}\right|^{\frac{1}{2}} \left(-\frac{\partial A}{\partial x}\right) \left(\exp\left(\frac{1}{2} \int_0^t |b'(X_{min}(s))| ds\right)\right) \right]_{x=0} + O(\varepsilon). \end{aligned} \tag{3.1}$$

Comparing this to the Laplace Integral

$$I(\varepsilon) = \int_a^b e^{-\frac{\phi(x)}{\varepsilon}} \theta(x) dx,$$

where the main contribution comes from the asymptotic behaviour at points $x_i \in [a, b]$ with $\phi'(x_i) = 0$, we can see that the main contribution to the integral in equation 3.1 comes from those $y(t, 0)$ satisfying

$$b(y) = \frac{\partial A(0, y, t)}{\partial y}.$$

If we expand $\phi(y) = \int_0^y b(u) du - A(0, y, t)$ in a Taylor series about $y(t, x) = 0$ we get

$$\begin{aligned} \phi(y) & = \phi(0) + (y - y(t, x))\phi'(0) + \frac{1}{2}(y - y(t, x))^2 \phi''(0) + \dots \\ & = \phi(0) + \frac{1}{2}(y - y(t, x))^2 \phi''(0) + \dots \end{aligned}$$

Hence,

$$\begin{aligned} \nu_0^+[s, \infty) &\sim (2\pi\varepsilon)^{-\frac{1}{2}} \int_0^\infty dy \exp \left\{ \frac{1}{\varepsilon} \left(-A(0, 0, s) \right. \right. \\ &\quad \left. \left. + \frac{1}{2}(y - y(s, x))^2 \frac{\partial^2}{\partial y^2} \left[\int_0^y b(u) du - A(0, y, s) \right] \right) \Big|_{y=y(s, x)=0} \right\} \\ &\times \left\{ \left(\exp \left(\frac{1}{2} \int_0^t |b'(X_{min}(s))| ds \right) \left| \frac{\partial^2 A}{\partial x \partial y} \right|_{x=0}^{\frac{1}{2}} \left(-\frac{\partial A}{\partial x} \right)_{x=0} \right) \right\}, \end{aligned}$$

giving,

$$\begin{aligned} \nu_0^+[s, \infty) &\sim (2\pi\varepsilon)^{-\frac{1}{2}} \int_0^\infty dy \exp \left\{ \frac{-A(0, 0, s)}{\varepsilon} - \frac{1}{2\varepsilon} \left[\frac{\partial^2 A(0, y, s)}{\partial y^2} - b'(y) \right]_{y=0} y^2 \right\} \\ &\times \left\{ \left(\exp \left(\frac{1}{2} \int_0^t |b'(X_{min}(s))| ds \right) \left(\frac{\partial^2 A}{\partial x \partial y} \right)_{x=0}^{\frac{1}{2}} \left(-\frac{\partial A}{\partial x} \right)_{x=0} \right) \right\}, \end{aligned}$$

and so the result follows. \square

4 Poisson-Lévy Excursion Measure – leading order behaviour

We have seen in the previous section that in order to calculate the Poisson-Lévy excursion measure $\nu_0^+[t, \infty)$ for a general process $X(t) = X[x, y, t]$, we require expressions for the following derivatives of the action $A(x, y, t)$.

$$\begin{aligned} \left. \frac{\partial A(x, y, t)}{\partial x} \right|_{x=0} &\quad \text{where} \quad p_0 = -\frac{\partial A}{\partial x}, \\ \left| \frac{\partial^2 A}{\partial x \partial y} \right|_{x=0}^{\frac{1}{2}} &\quad \text{where} \quad \frac{\partial X(t)}{\partial p_0} = \left(\frac{\partial^2 A}{\partial x \partial y} \right)^{-1} \quad (\text{the Van Vleck identity}), \\ \left. \frac{\partial^2 A}{\partial y^2} \right|_{x=0} &\quad \text{where} \quad b(y) = \frac{\partial A}{\partial y}(x, y, t). \end{aligned}$$

These expressions are evaluated in the following propositions :-

Proposition 4. For $p(0) = b(0) = 0$, $|b'(0)| \neq 0$,

$$-\frac{\partial A}{\partial x}(0, y, t) \sim \frac{|b'(0)| y}{\sinh |b'(0)| t}, \quad \text{as } y \rightarrow 0.$$

Proof. Observe that $p_x(y, t) = -\frac{\partial A(x, y, t)}{\partial x}$ = initial momentum at x needed to reach y in time t , satisfies

$$t = \int_x^y \frac{du}{\left(p_0^2(y, t) + b^2(u) - b^2(x) \right)^{\frac{1}{2}}}.$$

Therefore, $p_0(y, t)$ satisfies

$$t = \int_0^y \frac{du}{\left(p_0^2(y, t) + b^2(u)\right)^{\frac{1}{2}}}. \quad (4.1)$$

Changing integration variable $u = yv$,

$$t = \int_0^1 \frac{dv}{\left(\frac{p_0^2(y, t)}{y^2} + \frac{b^2(yv)}{y^2}\right)^{\frac{1}{2}}},$$

since to first order $V = \frac{1}{2}b^2$ and $V(0) = 0$ by assumption,

$$\frac{p_0(y) - p_0(0)}{y} \rightarrow p'_0(0), \quad \text{as } y \rightarrow 0.$$

Therefore,

$$t = \int_0^1 \frac{dv}{\left(p_0'^2(0, t) + v^2 b^2(0)\right)^{\frac{1}{2}}}, \quad \text{as } y \rightarrow 0.$$

Letting $v = \left|\frac{p'_0(0)}{b'(0)}\right| \sinh w$ we get

$$t = \frac{1}{|b'(0)|} \sinh^{-1} \left| \frac{b'(0)}{p'(0)} \right|, \quad (4.2)$$

giving, for $b'(0) \neq 0$,

$$|p'_0(0, t)| = \frac{|b'(0)|}{\sinh |b'(0)| t}. \quad (4.3)$$

□

Note that for $b'(0) = 0$, we get $p'_0(0, t) = \pm 1/t$ and so equation 4.2 has the correct limiting behaviour.

Proposition 5. For $|b'(0)| \neq 0$,

$$\frac{\partial^2 A}{\partial x \partial y}(0, y, t) \sim \left(\frac{\sinh |b'(0)| t}{|b'(0)|} \right)^{-1}, \quad \text{as } y \rightarrow 0.$$

Proof. Using the fact that $p_0(y) = -\partial A / \partial x$ and equation 4.1 we quickly get

$$\frac{-1}{p'_0(y)} = p_0(y) \left(p_0(y)^2 + b(y)^2\right)^{\frac{1}{2}} \int_0^y \frac{du}{\left(p_0(y)^2 + b(u)^2\right)^{\frac{3}{2}}}.$$

Again, changing the variable of integration $u = yv$, we get

$$\frac{-1}{p'_0(y)} = \frac{p_0(y)}{y} \left(\left(\frac{p_0(y)}{y}\right)^2 + \left(\frac{b(y)}{y}\right)^2 \right)^{\frac{1}{2}} \int_0^1 \frac{dv}{\left(\left(\frac{p_0(y)}{y}\right)^2 + \left(\frac{b(yv)}{y}\right)^2 \right)^{\frac{3}{2}}}.$$

Now, following the previous argument, as $y \rightarrow 0$,

$$\frac{-1}{p'_0(y)} \rightarrow p'_0(0, t) (p'_0(0, t)^2 + b'(0)^2)^{\frac{1}{2}} \int_0^1 \frac{dv}{(p'_0(0)^2 + b'(0)^2 v^2)^{\frac{3}{2}}}.$$

Letting $v = \left| \frac{p'_0(0)}{b'(0)} \right| \sinh w$ in the above equation gives

$$\begin{aligned} \frac{-1}{p'_0(y)} &\rightarrow \frac{(p'_0(0)^2 + b'(0)^2)^{\frac{1}{2}}}{|b'(0)| p'_0(0)^2} \int_0^{\sinh^{-1} \left| \frac{b'(0)}{p'_0(0)} \right|} \frac{1}{\cosh^2 w} dw \\ &= \frac{\sinh |b'(0)|t}{|b'(0)|}, \end{aligned}$$

using

$$|p'_0(0)| = \frac{|b'(0)|}{\sinh |b'(0)|t}.$$

□

Proposition 6. For $b'(0) \leq 0$, we have

$$b'(0) - \frac{\partial^2 A(0, 0, t)}{\partial y^2} = -|b'(0)|(1 + \coth |b'(0)|t).$$

($\frac{\partial A}{\partial y}$ is the momentum at y given that y is reached from x in time t .)

Proof. From

$$\frac{\partial^2 A}{\partial y^2} = \frac{\partial}{\partial y} p(y) = \frac{\partial \left(p_0^2(y, t) + b^2(y) \right)^{\frac{1}{2}}}{\partial y},$$

$$\begin{aligned} b'(y) - \frac{\partial^2 A}{\partial y^2} &= b'(y) - \frac{\frac{\partial p_0(y)}{\partial y} + \frac{b(y)}{p_0(y)} b'(y)}{\left(1 + \left(\frac{b(y)}{p_0(y)} \right)^2 \right)^{\frac{1}{2}}} \\ &\rightarrow b'(0) - \frac{p'_0(0) + \frac{b'(0)}{p'_0(0)} b'(0)}{\left(1 + \left(\frac{b'(0)}{p'_0(0)} \right)^2 \right)^{\frac{1}{2}}} \quad \text{as } y \rightarrow 0, \\ &= b'(0) - \frac{\frac{|b'(0)|}{\sinh |b'(0)|t} + b'(0)^2 \frac{\sinh |b'(0)|t}{|b'(0)|}}{\left(1 + \sinh^2 |b'(0)|t \right)^{\frac{1}{2}}} \\ &= b'(0) - |b'(0)| \frac{\cosh |b'(0)|t}{\sinh |b'(0)|t} \end{aligned}$$

Therefore

$$\left\{ b'(y) - \frac{\partial^2 A(0, y, t)}{\partial y^2} \right\} \Big|_{y=0} \longrightarrow -|b'(0)| (1 + \coth |b'(0)|t),$$

and result follows. \square

We now come to our main result for excursions from an equilibrium point 0.

Theorem 2. *For the diffusion X^ε with small noise satisfying*

$$dX^\varepsilon(t) = b(X^\varepsilon(t)) dt + \sqrt{\varepsilon} dB(t),$$

denote the Poisson-Lévy measure for excursions from 0 by ν_0^\pm .

Assuming b is continuous, b having right and left derivatives at 0, with $b(0^\pm) \leq 0$ and $b(0) = 0$, then if $V_0^\pm = \frac{1}{2} b^2$ satisfies

$$V_0^{\pm''} \geq -|\beta_\pm|, \quad \text{for } t \leq \frac{\pi}{|\beta_\pm|^{\frac{1}{2}}},$$

$$\nu_0^\pm[s, \infty) \sim \left(\frac{\varepsilon k_\pm}{\pi} \right)^{\frac{1}{2}} \left(e^{2k_\pm t} - 1 \right)^{-\frac{1}{2}}$$

with $k_\pm = |b'(0^\pm)|$. When $|b'(0^\pm)| = 0$ the limiting behaviour is correct and yields

$$\nu_0^\pm[t, \infty) \sim \left(\frac{\varepsilon}{2\pi} \right)^{\frac{1}{2}} t^{-\frac{1}{2}}.$$

Proof. Using Theorem 1, and the expressions obtained in Propositions 4, 5 and 6, for the derivatives of the action as $y \rightarrow 0$, we get as the leading order term for excursions from the stable equilibrium position 0, (dropping \pm again for convenience),

$$\begin{aligned} & \nu_0^+[s, \infty) \\ & \sim (2\pi\varepsilon)^{-\frac{1}{2}} e^{\frac{s|b'(0)|}{2}} \left| \frac{\sinh |b'(0)|t}{|b'(0)|} \right|^{-\frac{1}{2}} \left(\frac{|b'(0)|}{\sinh |b'(0)|t} \right) \\ & \times \int_0^\infty dy y \exp \left(-\frac{y^2}{2\varepsilon} |b'(0)|(1 + \coth |b'(0)|t) \right) \\ & = (2\pi\varepsilon)^{-\frac{1}{2}} e^{\frac{t|b'(0)|}{2}} |b'(0)|^{\frac{1}{2}} (\sinh |b'(0)|t)^{-\frac{1}{2}} \left(\frac{\varepsilon}{\cosh |b'(0)|t + \sinh |b'(0)|t} \right) \\ & = (2\pi)^{-\frac{1}{2}} \varepsilon^{\frac{1}{2}} |b'(0)|^{\frac{1}{2}} e^{-\frac{t|b'(0)|}{2}} \left(\frac{1}{2} (e^{t|b'(0)|} - e^{-t|b'(0)|}) \right)^{-\frac{1}{2}}. \end{aligned}$$

\square

These results correspond to the Poisson-Lévy excursion measures for the examples seen earlier.

5 Poisson-Lévy Excursion Measure – higher order behaviour

In calculating higher order terms in the Poisson-Lévy excursion measure $\nu_0^+[s, \infty)$, we obtain the surprising result that the next order term is identically zero. We now write the leading term as $\varepsilon^{\frac{1}{2}} \nu_{\frac{1}{2}}^+$.

Once again we must emphasise the particular dependence of V_ε on ε and how this requires us to follow a rather complicated route in determining the higher order dependencies on ε . This arises due to our study originating from stochastic mechanics where the Schrödinger equation and operator hold sway.

Theorem 3. *For the diffusion process with small noise, assuming $b(x) \leq 0$ for all x , the Poisson-Lévy excursion measure is given by*

$$\nu_\varepsilon^+ \cong \varepsilon^{\frac{1}{2}} \nu_{\frac{1}{2}}^+ + O(\varepsilon^{\frac{3}{2}})$$

i. e. the second order term is identically zero.

Proof - First part. For the derivation of the next order term of the Poisson- Lévy excursion measure $\nu_0^+[s, \infty)$ about $x = 0$, we must include the second order term in the expression for kernel $\exp\left(-\frac{tH(\varepsilon)}{\varepsilon}\right)(x, y)$ given in Proposition 3. Hence, from Propositions 2 and 3

$$\begin{aligned} \nu_0^+[s, \infty) &\sim (2\pi\varepsilon)^{-\frac{1}{2}} \int_{y<0} \frac{\rho_0^{\frac{1}{2}}(y)}{\rho_0^{\frac{1}{2}}(0)} \varepsilon \frac{\partial}{\partial x} \Big|_{x=0} \left[\exp\left(-\frac{A(x, y, t)}{\varepsilon}\right) \right. \\ &\quad \left. \times \exp\left(\frac{1}{2} \int_0^t |b'(X_{min}(s))| ds\right) \left| \frac{\partial^2 A}{\partial x \partial y} \right|^{\frac{1}{2}} [1 + \varepsilon K] \right] dy, \end{aligned}$$

where K , recall, is a very complicated expression involving the Feynman-Green function.

Therefore, up to order ε , we have assuming $\frac{\partial^2 A}{\partial x \partial y} \neq 0$

$$\begin{aligned}
& \nu_0^+[s, \infty) \\
&= (2\pi\varepsilon)^{-\frac{1}{2}} \int_0^\infty dy \exp\left(\frac{1}{\varepsilon} \int_0^y b(u) du\right) \\
&\times \varepsilon \left\{ \frac{1}{\varepsilon} \left[\left(-\frac{\partial A}{\partial x}\right)_{x=0} \exp\left(-\frac{A(0, y, t)}{\varepsilon}\right) \exp\left(\frac{1}{2} \int_0^t |b'(X_{min}(s))| ds\right) \left| \frac{\partial^2 A}{\partial x \partial y} \right|_{x=0}^{\frac{1}{2}} \right] \right. \\
&+ \varepsilon^0 \left[\frac{1}{2} \left| \frac{\partial^2 A}{\partial x \partial y} \right|_{x=0}^{-\frac{1}{2}} \frac{\partial}{\partial x} \right]_{x=0} \left(\frac{\partial^2 A}{\partial x \partial y} \right) \exp\left(-\frac{A(0, y, t)}{\varepsilon}\right) \exp\left(\frac{1}{2} \int_0^t |b'(X_{min}(s))| ds\right) ds \\
&- \frac{1}{2} \frac{\partial}{\partial x} \left[\int_0^t b'(X_{min}(s)) ds \right]_{x=0} \left| \frac{\partial^2 A}{\partial x \partial y} \right|_{x=0}^{\frac{1}{2}} \exp\left(-\frac{A(0, y, t)}{\varepsilon}\right) \exp\left(\frac{1}{2} \int_0^t |b'(X_{min}(s))| ds\right) \\
&+ \left. \left| \frac{\partial^2 A}{\partial x \partial y} \right|_{x=0}^{\frac{1}{2}} \exp\left(-\frac{A(0, y, t)}{\varepsilon}\right) \exp\left(\frac{1}{2} \int_0^t |b'(X_{min}(s))| ds\right) \left(-\frac{\partial A}{\partial x}\right)_{x=0} K \right\} \\
&+ O(\varepsilon^2). \tag{5.1}
\end{aligned}$$

We now use the result Olver [1974].

Proposition 7.

$$\begin{aligned}
& \int_0^\infty \exp\left(-\frac{f(y)}{\varepsilon}\right) g(y) dy \\
&= \int_0^\infty \exp\left(-\frac{f(y)}{\varepsilon}\right) [g_0(y) + \varepsilon g_1(y) + \frac{\varepsilon^2}{2!} g_2(y) + \dots] dy \tag{5.2}
\end{aligned}$$

$$\sim \Gamma(1) \varepsilon \frac{g'_0}{2} \frac{1}{\frac{f''}{2}} + \Gamma\left(\frac{3}{2}\right) \varepsilon^{\frac{3}{2}} \left(\frac{g''_0}{4} - 3 \frac{f'''}{4 \frac{f''}{2}} g'_0 \right) \frac{1}{\left(\frac{f''}{2}\right)^{\frac{3}{2}}} + \Gamma\left(\frac{1}{2}\right) \varepsilon^{\frac{3}{2}} \frac{g_1}{2} \frac{1}{\left(\frac{f''}{2}\right)^{\frac{1}{2}}} + \dots \tag{5.3}$$

Proof. For a proof of this standard result on asymptotic approximations see Olver [1974]. \square

Comparing equation 5.2 with our expression for $\nu_0^+[s, \infty)$ to second order in equation 5.1, we get

$$g_0(y) = (2\pi\varepsilon)^{-\frac{1}{2}} \left\{ \left| \frac{\partial^2 A(x, y, t)}{\partial x \partial y} \right|_{x=0}^{\frac{1}{2}} \exp\left(\frac{1}{2} \int_0^t |b'(X(s))| ds\right) \left(-\frac{\partial A(x, y, t)}{\partial x}\right) \right\}_{x=0},$$

and

$$\begin{aligned}
& g_1(y) = (2\pi\varepsilon)^{-\frac{1}{2}} \left| \frac{\partial^2 A(0, y, t)}{\partial x \partial y} \right|_{x=0}^{\frac{1}{2}} \exp\left(\frac{1}{2} \int_0^t |b'(X(s))| ds\right) \\
&\times \left\{ \frac{1}{2} \frac{\partial}{\partial x} \int_0^t |b'(X(s))| ds + \frac{1}{2} \left| \frac{\partial^2 A(x, y, t)}{\partial x \partial y} \right|_{x=0}^{-1} \frac{\partial}{\partial x} \left(\frac{\partial^2 A(x, y, t)}{\partial x \partial y} \right) - K \frac{\partial A(x, y, t)}{\partial x} \right\}_{x=0}.
\end{aligned}$$

Since we know for $x = 0$

$$\left. \frac{\partial A}{\partial x} \right|_{x=0} = -p_0(y), \text{ so } -\left. \frac{\partial A}{\partial x} \right|_{x=0} = p_0(y) (> 0), \text{ and } \left. \frac{\partial^2 A}{\partial x \partial y} \right|_{x=0} = -\frac{\partial p_0(y)}{\partial y},$$

giving

$$\left| \frac{\partial^2 A}{\partial x \partial y} \right| = \frac{\partial p_0(y)}{\partial y} (> 0), \quad \frac{\partial}{\partial x} \left| \frac{\partial^2 A}{\partial x \partial y} \right| = \frac{\partial^2 p_0(y)}{\partial x \partial y},$$

we can write the expressions for $g_0(y)$ and $g_1(y)$ as

$$g_0(y) = (2\pi\varepsilon)^{-\frac{1}{2}} \left\{ \left(\frac{\partial p_0(y)}{\partial y} \right)^{\frac{1}{2}} \exp \left(\frac{1}{2} \int_0^t |b'(X(s))| ds \right) (p_0(y)) \right\}_{x=0},$$

and

$$\begin{aligned} g_1(y) &= (2\pi\varepsilon)^{-\frac{1}{2}} \left(\frac{\partial p_0(y)}{\partial y} \right)^{\frac{1}{2}} \exp \left(\frac{1}{2} \int_0^t |b'(X(s))| ds \right) \\ &\times \left\{ \frac{1}{2} \frac{\partial}{\partial x} \int_0^t |b'(X(s))| ds + \frac{1}{2} \left(\frac{\partial p_0(y)}{\partial y} \right)^{-1} \left(\frac{\partial^2 p_0(y)}{\partial x \partial y} \right) - K p_0(y) \right\}_{x=0}. \end{aligned}$$

If we now expand each term in the expression for $g_0(y)$ in a Taylor series we get

$$\begin{aligned} g_0(y) &= (2\pi\varepsilon)^{-\frac{1}{2}} \left\{ (p'_0(0) + y p''_0(0) + \dots)^{\frac{1}{2}} \exp \left(\frac{1}{2} \int_0^t |b'(X(s))| ds \right) \right. \\ &\quad \left. + \frac{1}{2} y \frac{\partial}{\partial y} \int_0^t |b'(X(s))| ds \right|_{y=0} + \dots \left. \right\} (p_0(0) + y p'_0(0) + \dots). \end{aligned}$$

Therefore, we can now see that in order to obtain the first order expressions for $g_0(y)$ and $g_1(y)$, the following terms need to be evaluated :

$$\begin{array}{ccc} \left(\frac{\partial p_0(y)}{\partial y} \right)^{-1} & \frac{\partial^2 p_0(y)}{\partial y^2} & \frac{\partial^2 p_0(y)}{\partial x \partial y} \\ \int_0^t |b'(X(s))| ds & \frac{\partial}{\partial x} \int_0^t |b'(X(s))| ds & \frac{\partial}{\partial y} \int_0^t |b'(X(s))| ds. \end{array}$$

Let us be very thankful that an evaluation of K is not needed in this rather complicated computation.

Each of these terms is evaluated in the following Propositions. Recall that we have already seen in equation 4.3

$$|p'_0(0)| = \frac{\partial p_0(y)}{\partial y} = \frac{|b'(0)|}{\sinh |b'(0)|t}.$$

□

Proposition 8. For $p_0(0) = b(0) = 0$, as $y \rightarrow 0$,

$$\frac{\partial^2 p_0(y)}{\partial y^2} = \frac{b''(0) (\cosh |b'(0)|t - 1)^2}{\sinh^3 |b'(0)|t}. \quad (5.4)$$

Proof. In order to calculate $\frac{\partial^2 p_0(y)}{\partial y^2}$ we return to the identity,

$$t = \int_0^y \frac{du}{(p_0^2(y) + b^2(u))^{\frac{1}{2}}}.$$

Differentiating the equation above w.r.t. y , and then using the change of variable $u = yv$, gives dropping some inessential modulus signs for ease of presentation

$$\begin{aligned} \frac{1}{p_0'(y)} &= p_0(y) (p_0^2(y) + b^2(y))^{\frac{1}{2}} \int_0^y \frac{du}{(p_0^2(y) + b^2(u))^{\frac{3}{2}}} \\ &= \frac{p_0(y)}{y} \left(\left(\frac{p_0(y)}{y} \right)^2 + \left(\frac{b(y)}{y} \right)^2 \right)^{\frac{1}{2}} \int_0^1 \frac{dv}{\left(\left(\frac{p_0(y)}{y} \right)^2 + \left(\frac{b(yv)}{yv} \right)^2 v^2 \right)^{\frac{3}{2}}} \\ &\rightarrow p_0'(0) (p_0'(0)^2 + b'(0)^2)^{\frac{1}{2}} \int_0^1 \frac{dv}{(p_0'(0)^2 + b'(0)^2 v^2)^{\frac{3}{2}}} \quad \text{as } y \rightarrow 0, \end{aligned} \quad (5.5)$$

using the same argument as seen in Proposition 4

Expanding the r.h.s. of equation 5.5 in a Taylor Series, using for simplicity the notation $p = p_0(0)$ and $b = b(0)$, gives

$$\begin{aligned} &\left(p' + \frac{y}{2} p'' + \dots \right) \left(\left(p' + \frac{y}{2} p'' + \dots \right)^2 + \left(b' + \frac{y}{2} b'' + \dots \right)^2 \right)^{\frac{1}{2}} \\ &\quad \times \int_0^1 \frac{dv}{\left(\left(p' + \frac{y}{2} p'' + \dots \right)^2 + \left(b' + \frac{yv}{2} b'' + \dots \right)^2 v^2 \right)^{\frac{3}{2}}} \\ &\rightarrow \left(p' + \frac{y}{2} p'' + \dots \right) (p'^2 + b'^2)^{\frac{1}{2}} \left(1 + y \frac{p' p'' + b' b''}{p'^2 + b'^2} + \dots \right)^{\frac{1}{2}} \\ &\quad \times \int_0^1 dv (p'^2 + b'^2 v^2)^{-\frac{3}{2}} \left(1 + y \frac{p' p'' + b' b'' v^3}{p'^2 + b'^2 v^2} + \dots \right)^{-\frac{3}{2}} \\ &= \left(p' + \frac{y}{2} p'' + \dots \right) \left((p'^2 + b'^2)^{\frac{1}{2}} + \frac{y}{2} \left(\frac{p' p'' + b' b''}{(p'^2 + b'^2)^{\frac{1}{2}}} \right) + \dots \right) \\ &\quad \times \int_0^1 dv \left((p'^2 + b'^2 v^2)^{-\frac{3}{2}} - \frac{3}{2} y \left(\frac{p' p'' + b' b'' v^3}{(p'^2 + b'^2 v^2)^{\frac{5}{2}}} \right) + \dots \right) \\ &= \text{zero order term} + f.y + \text{higher order terms}(y^2 \dots). \end{aligned}$$

The zero order term is

$$p' (p'^2 + b'^2)^{\frac{1}{2}} \int_0^1 dv (p'^2 + b'^2 v^2)^{-\frac{3}{2}}.$$

Now the integral term in the equation above can be written as

$$\frac{1}{b'^3} \int_0^1 \frac{dv}{\left(\frac{p'^2}{b'^2} + v^2\right)^{\frac{3}{2}}}.$$

Using the change of variable $v = \left|\frac{p'}{b'}\right| \sinh w$ in the equation above gives

$$\begin{aligned} & \frac{1}{b'^3} \int_0^{\sinh^{-1}\left|\frac{b'}{p'}\right|} dw \frac{\left|\frac{p'}{b'}\right| \cosh w}{\left(\frac{p'^2}{b'^2} + \frac{p'^2}{b'^2} \sinh^2 w\right)^{\frac{3}{2}}} \\ &= \frac{1}{b' p'^2} \int_0^{\sinh^{-1}\left|\frac{b'}{p'}\right|} dw \frac{1}{\cosh^2 w} = \frac{1}{|p'|^3} \cdot \frac{1}{\cosh |b'(0)|t}. \end{aligned}$$

Therefore, the zero order term is (because of equation 4.3)

$$p' (p'^2 + b'^2)^{\frac{1}{2}} \cdot \frac{1}{p'^3} \cdot \frac{1}{\cosh |b'(0)|t} = \frac{1}{|p'|}.$$

The coefficient of y is given by

$$\begin{aligned} f &= \frac{p''}{2} (p'^2 + b'^2)^{\frac{1}{2}} \int_0^1 dv (p'^2 + b'^2 v^2)^{-\frac{3}{2}} + \frac{p' p' p'' + b' b''}{2 (p'^2 + b'^2)^{\frac{1}{2}}} \int_0^1 dv (p'^2 + b'^2 v^2)^{-\frac{3}{2}} \\ &\quad - \frac{3}{2} p' (p'^2 + b'^2)^{\frac{1}{2}} \int_0^1 dv \frac{p' p'' + b' b'' v^3}{(p'^2 + b'^2 v^2)^{\frac{5}{2}}}. \end{aligned}$$

Therefore, we have that

$$\frac{\partial p}{\partial y} = \frac{1}{p'^{-1} + yf + \dots} = \frac{p'}{1 + y p' f + \dots} = p'(1 - y p' f - \dots). \quad (5.6)$$

Now, by letting $|p'| = -\frac{|b'|}{a}$ giving $a = -\sinh |b'(0)|t$, we can write

$$\begin{aligned} p'^2 f &= \frac{p''}{2} + \frac{p'^3}{2} \frac{p' p'' + b' b''}{p'(1+a^2)^{\frac{1}{2}}} p'^{-3} (1+a^2)^{-\frac{1}{2}} \\ &\quad - \frac{3}{2} p'^3 p' (1+a^2)^{\frac{1}{2}} \left\{ \frac{p' p''}{p'^5} \left(\frac{2}{3(1+a^2)} + \frac{1}{3(1+a^2)^{\frac{3}{2}}} \right) \right. \\ &\quad \left. + \frac{b' b''}{p'^5} \left(\frac{2}{3a^2} - \frac{2+3a^2}{3a^4(1+a^2)^{\frac{3}{2}}} \right) \right\}, \end{aligned}$$

which simplifies to

$$p'^2 f = \frac{p''}{2} + \frac{1}{2} \frac{p'' - ab''}{(1+a^2)} - \frac{3}{2} p'' \left(\frac{2}{3} + \frac{1}{3(1+a^2)} \right) - \frac{3}{2} (-a) b'' \left(\frac{2(1+a^2)^{\frac{1}{2}}}{3a^4} - \frac{2+3a^2}{3a^4(1+a^2)} \right).$$

Hence, from equation 5.6

$$-p'' = \frac{p''}{2} + \frac{1}{2} \frac{p'' - ab''}{(1+a^2)} - p'' - \frac{p''}{2(1+a^2)} + \frac{3ab''}{2} \left(\frac{2(1+a^2)^{\frac{1}{2}}}{3a^4} - \frac{2+3a^2}{3a^4(1+a^2)} \right),$$

giving

$$0 = \frac{p''}{2} - \frac{ab''}{2(1+a^2)} + \frac{b''(1+a^2)^{\frac{1}{2}}}{a^3} - \frac{1+\frac{3}{2}a^2}{a^3(1+a^2)} b''.$$

Now since $a = -\frac{|b'|}{|p'|}$ and $|p'| = \frac{|b'(0)|}{\sinh|b'(0)|t}$, and again using the obvious notation

$$s = \sinh|b'(0)|t \quad \text{and} \quad c = \cosh|b'(0)|t,$$

we can write the equation above as

$$\begin{aligned} 0 &= p'' + b'' \left(-\frac{s}{c^2} + 2\frac{c}{s^3} - \frac{2+3s^2}{s^3c^2} \right) \\ &= p'' + \frac{b''}{s^3} \left(-\frac{s^4}{c^2} + 2c - \frac{2+3s^2}{c^2} \right) \\ &= p'' + \frac{b''}{s^3} \left(-\frac{(c^4-2c^2+1)}{c^2} + \frac{2c^3}{c^2} - \frac{2+3(c^2-1)}{c^2} \right) \\ &= p'' + \frac{b''}{s^3} \left(\frac{-c^4+2c^3-c^2}{c^2} \right). \end{aligned}$$

Hence, we get the result

$$p_0''(0) = \frac{b''(0) (\cosh|b'(0)|t - 1)^2}{\sinh^3|b'(0)|t}.$$

□

Proposition 9. As $y \rightarrow 0$,

$$\frac{\partial^2 p_0}{\partial x \partial y} \rightarrow \frac{b''(0) (\cosh|b'(0)|t - 1)^2}{\sinh^3|b'(0)|t} = p_0''(0).$$

Proof. We begin with

$$t = \int_x^y \frac{du}{(p^2(x, y) + b^2(u) - b^2(x))^{\frac{1}{2}}}.$$

Differentiating both sides w.r.t. x gives,

$$0 = -\frac{1}{|p(x, y)|} - \int_x^y du \frac{p \frac{\partial p}{\partial x} - b(x) \frac{\partial b(x)}{\partial x}}{(p^2(x, y) + b^2(u) - b^2(x))^{\frac{3}{2}}}.$$

Therefore, as $x \rightarrow 0$

$$0 = \frac{1}{|p_0(y)|} + p_0(y) \frac{\partial p}{\partial x} \Big|_{x=0} \int_0^y \frac{du}{(p_0^2(y) + b^2(u))^{\frac{3}{2}}}.$$

Hence,

$$\frac{\partial p}{\partial x} \Big|_{x=0} = \frac{-1}{p_0^2(y) \int_0^y (p_0^2(y) + b^2(u))^{-3/2} du}. \quad (5.7)$$

If we consider the quotient term on the r.h.s. of equation 5.7, with a change of variable $u = yv$ and again letting $y \rightarrow 0$, we get

$$r.h.s. = \frac{-1}{p_0^2(0) \int_0^1 (p_0^2(0) + b^2(0)v^2)^{-3/2} dv}.$$

Expanding the denominator of equation 5.7 in a Taylor series [$p(0) = 0$, $p'(0) \neq 0$], using the same notation as in the previous Proposition

$$\begin{aligned} & (p' + \frac{y}{2} p'' + \dots)^2 \int_0^1 \frac{dv}{((p' + \frac{y}{2} p'' + \dots)^2 + (b' + \frac{yv}{2} b'' + \dots)^2 v^2)^{\frac{3}{2}}} \\ &= (p'^2 + y p' p'' + \dots) \int_0^1 \frac{dv}{(p'^2 + b^2 v^2 + y(p' p'' + v^3 b' b'') + \dots)^{\frac{3}{2}}} \\ &= (p'^2 + y p' p'' + \dots) \int_0^1 \frac{1}{(p'^2 + b^2 v^2)^{\frac{3}{2}}} \left\{ 1 - \frac{3}{2} y \frac{p' p'' + v^3 b' b''}{(p' + b^2 v^2)} - \dots \right\} dv. \end{aligned}$$

Therefore,

$$\left(\frac{\partial p}{\partial x} \right)_{x=0}^{-1} = \text{zero order term} + fy + \dots,$$

where f is the coefficient of the y term as shown below

$$\begin{aligned} \left(\frac{\partial p}{\partial x} \right)_{x=0}^{-1} &= p'^2 \int_0^1 \frac{dv}{(p'^2 + b^2 v^2)^{\frac{3}{2}}} + y \left(p' p'' \int_0^1 \frac{dv}{(p'^2 + b^2 v^2)^{\frac{3}{2}}} \right. \\ &\quad \left. - \frac{3}{2} p'^2 \int_0^1 \frac{p' p'' + b' b'' v^3}{(p'^2 + b^2 v^2)^{\frac{5}{2}}} dv \right) + \dots \end{aligned}$$

Inverting this equation gives

$$\begin{aligned} \frac{\partial p}{\partial x} \Big|_{x=0} &= \left(p'^2 \int_0^1 \frac{dv}{(p'^2 + b^2 v^2)^{\frac{3}{2}}} \right)^{-1} \left\{ 1 - y \left(\frac{p' p'' \int_0^1 \frac{dv}{(p'^2 + b^2 v^2)^{\frac{3}{2}}} - \frac{3}{2} p'^2 \int_0^1 \frac{p' p'' + b' b'' v^3}{(p'^2 + b^2 v^2)^{\frac{5}{2}}} dv}{p'^2 \int_0^1 \frac{dv}{(p'^2 + b^2 v^2)^{\frac{3}{2}}} \right) - \dots \right\}. \end{aligned}$$

Now since

$$\frac{\partial p(x, y)}{\partial x} \Big|_{x=0} = \frac{\partial p(x, 0)}{\partial x} \Big|_{x=0} + y \frac{\partial^2 p(x, 0)}{\partial x \partial y} \Big|_{x=0} + \dots$$

Comparing y -terms yields

$$\frac{\partial^2 p(x, 0)}{\partial x \partial y} \Big|_{x=0} = \frac{p' p'' \int_0^1 \frac{dv}{(p'^2 + b'^2 v^2)^{\frac{3}{2}}} - \frac{3}{2} p'^2 \int_0^1 \frac{p' p'' + b' b'' v^3}{(p'^2 + b'^2 v^2)^{\frac{5}{2}}} dv}{\left(p'^2 \int_0^1 \frac{dv}{(p'^2 + b'^2 v^2)^{\frac{3}{2}}} \right)^2}.$$

Letting $a = -\left| \frac{b'}{p'} \right|$,

$$\begin{aligned} \frac{\partial^2 p}{\partial x \partial y} \Big|_{x=0} &= \frac{\frac{p''}{p'^2} \int_0^1 \frac{dv}{(1+a^2 v^2)^{\frac{3}{2}}} - \frac{3}{2} \frac{1}{p'^3} \int_0^1 \frac{p' p'' + b' b'' v^3}{(1+a^2 v^2)^{\frac{5}{2}}} dv}{\left(\frac{1}{p'} \int_0^1 \frac{dv}{(1+a^2 v^2)^{\frac{3}{2}}} \right)^2} \\ &= -\frac{1}{2} \frac{p''}{p'} \frac{1}{(1+a^2)} - \frac{1}{2} \frac{b' b''}{p'^2} \frac{(1+a^2)^{\frac{1}{2}}}{a^2} + \frac{1}{2} \frac{b' b''}{p'^2} \frac{1}{a^2(1+a^2)}. \end{aligned}$$

Now, substituting for p' and p'' gives the result. \square

Remark. A by product of the above is

$$\frac{\partial p_0(y)}{\partial x} \rightarrow -p'_0 \cosh |b'(0)| t.$$

Proposition 10. As $y \rightarrow 0$,

$$\int_0^t |b'(X_{\min}(s))| ds \rightarrow t |b'(0)|$$

with X_{\min} satisfying

$$\ddot{X}_{\min}(s) = b(X_{\min}(s)) b'(X_{\min}(s)) \quad \text{and} \quad X_{\min}(0) = x, X_{\min}(t) = y.$$

Proof. For $u = X_{\min}(s)$, $du = \dot{X}_{\min}(s) ds$, and considering

$$\begin{aligned} \int_0^t |b'(X_{\min}(s))| ds &= s |b'(X_{\min}(s))| \Big|_0^t + \int_0^t s b''(X_{\min}(s)) \dot{X}_{\min}(s) ds \\ &= t |b'(y)| + \int_x^y s(u) b''(u) du \\ &= t |b'(y)| + \int_x^y du \int_x^u dv \frac{b''(u)}{(p^2(x, y) + b^2(v) - b^2(x))^{\frac{1}{2}}} \end{aligned} \tag{5.8}$$

since,

$$t(u) = \int_x^y \frac{du}{(p^2(x, y) + b^2(u) - b^2(x))^{\frac{1}{2}}}.$$

Therefore, the integral term on the r.h.s. of equation 5.8 $\rightarrow 0$ as $y \rightarrow 0$, and so the result follows (recall $y > x > 0$). \square

Proposition 11.

$$\begin{aligned} \frac{\partial}{\partial x} \Big|_{x=0} \int_0^t |b'(X_{min}(s))| ds &\rightarrow -b''(0) p'_0(0) \frac{\partial p}{\partial x} \Big|_{x=0} \int_0^1 du \int_u^1 \frac{dv}{(p_0'^2(0) + b^2(0)v^2)^{\frac{3}{2}}} \\ &= \frac{b''(0)}{|b'(0)|} \frac{\cosh |b'(0)|t - 1}{\sinh |b'(0)|t}, \quad \text{as } y \rightarrow 0. \end{aligned} \quad (5.9)$$

Proof. Using equations 5.7 and 5.8, a calculation along the lines of the proof of Theorem 4, using integration by parts, yields

$$\begin{aligned} \frac{\partial}{\partial x} \Big|_{x=0} \int_0^t |b'(X_{min}(s))| ds &= \int_0^y du \left\{ p_0(y) \frac{\partial p}{\partial x} \Big|_{x=0} \int_0^u \frac{dv}{(p_0'^2(y) + b^2(v))^{\frac{3}{2}}} \right\} b''(u) \\ &\rightarrow -b''(0) p'_0(0) \frac{\partial p}{\partial x} \Big|_{y=x=0} \int_0^1 du \int_u^1 \frac{dv}{(p_0'^2(0) + b^2(0)v^2)^{\frac{3}{2}}} \quad \text{as } y \rightarrow 0 \end{aligned}$$

and result follows. \square

Proposition 12.

$$\begin{aligned} \frac{\partial}{\partial y} \int_0^t |b'(X(s))| ds &\rightarrow b''(0)(p'_0(0))^2 \int_0^1 du \int_0^u \frac{dv}{(p_0'^2(0) + b^2(0)v^2)^{\frac{3}{2}}} \\ &= \frac{b''(0)}{|b'(0)|} \frac{\cosh |b'(0)|t - 1}{\sinh |b'(0)|t}, \quad \text{as } y \rightarrow 0. \end{aligned} \quad (5.10)$$

Proof. The proof of this is similar to that of Proposition 11. \square

Proof - Second part. Therefore, by substituting equations 4.3, 5.4, 5.7, 5.9, 5.10 and Proposition 9 into the expressions obtained for $g_0(y)$ and $g_1(y)$, and using Proposition 7, we can complete the proof of Theorem 3.

$$\begin{aligned} g_0(y) &\sim (2\pi\varepsilon)^{-\frac{1}{2}} (p' + p''y + \dots)^{\frac{1}{2}} \exp\left(\frac{1}{2}|b'(X(0))|t\right. \\ &\quad \left. + \frac{1}{2}y \frac{b''(0)}{|b'(0)|} \frac{\cosh |b'(0)|t - 1}{\sinh |b'(0)|t} + \dots\right) \left(p(0) + yp'(0) + \frac{y^2}{2}p''(0) + \dots\right), \quad \text{as } y \rightarrow 0 \\ &= (2\pi\varepsilon)^{-\frac{1}{2}} (p'_0)^{\frac{3}{2}} \exp\left(\frac{t|b'(0)|}{2}\right) \left(1 + \frac{p''}{p'}y + \dots\right)^{\frac{1}{2}} \\ &\quad \times \left(y + \frac{p''}{p'} \frac{y^2}{2} - \frac{b''(0)}{2|b'(0)|} \frac{\cosh |b'(0)|t - 1}{\sinh |b'(0)|t} y^2 - \frac{b''(0)p''}{4|b'(0)|p'} \frac{\cosh |b'(0)|t - 1}{\sinh |b'(0)|t} y^3 + \dots\right) \\ &= (2\pi\varepsilon)^{-\frac{1}{2}} (p')^{\frac{3}{2}} \exp\left(\frac{t|b'(0)|}{2}\right) \left[y + \frac{1}{2}y^2 \left(2\frac{p''}{p'} - \frac{b''(0)}{|b'(0)|} \frac{\cosh |b'(0)|t - 1}{\sinh |b'(0)|t}\right) + \dots\right] \end{aligned}$$

Similarly,

$$\begin{aligned} g_1(y) &\sim (2\pi\varepsilon)^{-\frac{1}{2}} (p'_0)^{\frac{1}{2}} \exp\left(\frac{t|b'(0)|}{2}\right) \\ &\quad \times \left(-\frac{1}{2} \frac{b''(0)}{|b'(0)|} \frac{\cosh |b'(0)|t - 1}{\sinh |b'(0)|t} + \frac{1}{2} \frac{b''(0)}{|b'(0)|} \frac{(\cosh |b'(0)|t - 1)^2}{(\sinh |b'(0)|t)^2} + y(\dots) + \dots\right). \end{aligned}$$

Therefore,

$$g'_0(0) = (2\pi\varepsilon)^{-\frac{1}{2}} (p')^{\frac{3}{2}} \exp\left(\frac{t|b'(0)|}{2}\right),$$

and

$$g''_0(0) = (2\pi\varepsilon)^{-\frac{1}{2}} (p')^{\frac{3}{2}} \exp\left(\frac{t|b'(0)|}{2}\right) \left(2\frac{p''}{p'} - \frac{b''(0)}{|b'(0)|} \frac{\cosh |b'(0)|t - 1}{\sinh |b'(0)|t}\right).$$

Now, in our case, the expression for $f(y)$ in Proposition 7 is

$$f(y) = A(0, y, t) - \int_0^y b(u) du,$$

so

$$f(0) = A(0, 0, t) - \int_0^0 b(u) du = 0.$$

Also, since $p(t) = \sqrt{p_0^2(y) + b^2(y)}$,

$$f'(y) = \frac{\partial A(0, y, t)}{\partial y} - \frac{\partial}{\partial y} \int_0^y b(u) du = p_0(t) - b(y) = \sqrt{p_0^2(y) + b^2(y)} - b(y),$$

so

$$f'(0) = \sqrt{p_0^2(0) + b^2(0)} - b(0) = 0,$$

since $p_0(0) = 0$, (the momentum required to go from $x = 0$ to $y = 0$).

Now, $f''(y) = \frac{\partial}{\partial y} p(0, y, t) - b'(y)$, so differentiating $p(t) = \sqrt{p_0^2(y) + b^2(y)}$ twice with respect to y , and using the fact that $b(0) = 0$ and $p(t) \rightarrow 0$ as $y \rightarrow 0$, we get

$$\left\{ \left(\frac{\partial p(t)}{\partial y} \right)^2 + p(t) \frac{\partial^2 p(t)}{\partial y^2} \right\} \Bigg|_{x=0} = p'_0(y)^2 + p_0(y) p''_0(y) + b'(y)^2 + b(y) b''(y). \quad (5.11)$$

As $y \rightarrow 0$,

$$\left(\frac{\partial p_0(t)}{\partial y} \right)^2 \rightarrow p'_0(0)^2 + b'(0)^2,$$

since $p_0(0) = b(0) = 0$, and $p_0(t) \frac{\partial^2 p_0(t)}{\partial y^2} \rightarrow 0$, as $y \rightarrow 0$. Hence,

$$\begin{aligned} \left(\frac{\partial p_0(t)}{\partial y} \right)^2 \Bigg|_{x=0} &= p'_0(0)^2 + b'(0)^2 \\ &= \frac{(b'(0))^2}{(\sinh |b'(0)|t)^2} + b'(0)^2 \\ &= (b'(0))^2 \left(\frac{1 + \sinh^2 |b'(0)|t}{\sinh^2 |b'(0)|t} \right) \\ &= |b'(0)|^2 \coth^2 |b'(0)|t. \end{aligned}$$

Therefore,

$$f''(y) \rightarrow b'(0) \coth |b'(0)|t - b'(0) = |b'(0)| (\coth |b'(0)|t + 1),$$

since we are dealing with $b(x) < 0$.

Similarly,

$$f'''(y) = \frac{\partial^2 p}{\partial y^2}(0, y, t) - b''(y).$$

Differentiating equation 5.11 again w.r.t. y gives,

$$\left. \left\{ \frac{\partial^3 p(t)}{\partial y^3} p(t) + 3 \frac{\partial p(t)}{\partial y} \frac{\partial^2 p(t)}{\partial y^2} \right\} \right|_{x=0} = 3 p'_0(y) p''_0(y) + p_0(y) p'''(y) \\ + 3 b'(y) b'''(y) + b(y) b''''(y).$$

And again, since $p_0(0) = b(0) = 0$, and $p_0(t) \rightarrow 0$ as $y \rightarrow 0$, we get

$$\left. \left\{ \frac{\partial p_0(t)}{\partial y} \frac{\partial^2 p_0(t)}{\partial y^2} \right\} \right|_{y=0} = p'_0(0) p''_0(0) + b'(0) b''(0),$$

giving, after a little calculation

$$\left. \left\{ \frac{\partial^2 p_0(t)}{\partial y^2} \right\} \right|_{y=0} = \frac{p'_0(0) p''_0(0) + |b'(0)| b''(0)}{|b'(0)| \coth |b'(0)| t}.$$

Therefore, as $y \rightarrow 0$,

$$f'''(y) \rightarrow \frac{p'_0 p''_0 + |b'(0)| b''(0)}{|b'(0)| \coth |b'(0)| t} - b''(0) \\ = b''(0) \left(\frac{(\cosh |b'(0)| t - 1)^2}{\cosh |b'(0)| t (\sinh |b'(0)| t)^3} - \frac{\sinh |b'(0)| t}{\cosh |b'(0)| t} - 1 \right).$$

Finally, we conclude the proof of Theorem 3 by substituting into equation 5.3 to get

$$\nu_0^+[t, \infty) = \int_0^\infty dy \exp \left(\frac{1}{\varepsilon} \int_0^y b(u) du - A(0, y, t) \right) (g_0(y) + \varepsilon g_1(y) + \dots) dy \\ \sim \frac{\varepsilon}{2} (2\pi\varepsilon)^{-\frac{1}{2}} (p')^{\frac{3}{2}} \exp \left(\frac{t |b'(0)|}{2} \right) \frac{1}{f''} \\ + \sqrt{\frac{\pi}{2f''}} \varepsilon^{\frac{3}{2}} \left\{ \frac{(2\pi\varepsilon)^{-\frac{1}{2}}}{4} (p')^{\frac{3}{2}} \exp \left(\frac{t |b'(0)|}{2} \right) \left(\frac{2p''_0}{p'_0} - \frac{b''(0) \cosh |b'(0)| t - 1}{|b'(0)| \sinh |b'(0)| t} \right) \right. \\ \left. - \frac{3}{4} \frac{f'''}{f''} (2\pi\varepsilon)^{-\frac{1}{2}} (p')^{\frac{3}{2}} \exp \left(\frac{t |b'(0)|}{2} \right) \right\} \frac{1}{f''} \\ + \sqrt{\frac{\pi}{2f''}} \varepsilon^{\frac{3}{2}} (2\pi\varepsilon)^{-\frac{1}{2}} (p')^{\frac{1}{2}} \exp \left(-\frac{t |b'(0)|}{2} \right) \\ \times \left(-\frac{1}{2} \frac{b''(0) \cosh |b'(0)| t - 1}{|b'(0)| \sinh |b'(0)| t} + \frac{1}{2} \frac{b''(0) (\cosh |b'(0)| t - 1)^2}{|b'(0)| (\sinh |b'(0)| t)^2} \right).$$

Substituting for p' and p'' eventually gives, using an obvious shorthand notation

$$\begin{aligned}
\nu_0^+[t, \infty) &\sim \varepsilon(2\pi\varepsilon)^{-\frac{1}{2}} (p')^{\frac{3}{2}} \exp\left(\frac{t|b'(0)|}{2}\right) \frac{1}{f''} \\
&+ \varepsilon^{\frac{3}{2}}(2\pi\varepsilon)^{-\frac{1}{2}} (p')^{\frac{3}{2}} \exp\left(\frac{t|b'(0)|}{2}\right) \left(\frac{\pi}{2f''}\right)^{\frac{1}{2}} \left\{ \left[\frac{b''(0)(c-1)^2}{2p's^3} - \frac{b''(0)(c-1)}{4|b'(0)|s} \right. \right. \\
&- \left. \left. \frac{b''}{4} \left(\frac{(c-1)^2 - s^4 - cs^3}{b'cs^2(c+s)} \right) \right] \frac{2s}{b'(c+s)} - \frac{b''(c-1)}{2p'b's} + \frac{b''(c-1)^2}{2p'b's^2} \right\} \\
&= \varepsilon(2\pi\varepsilon)^{-\frac{1}{2}} (p')^{\frac{3}{2}} \exp\left(\frac{t|b'(0)|}{2}\right) \frac{1}{f''} + \varepsilon^{\frac{3}{2}}(2\pi\varepsilon)^{-\frac{1}{2}} (p')^{\frac{1}{2}} \exp\left(\frac{t|b'(0)|}{2}\right) \\
&\times \sqrt{\frac{\pi}{2f''}} \left(-\frac{b''(0)}{2|b'(0)|}\right) \left[\frac{c-1}{s} - \frac{(c-1)^2}{s^2} + \frac{1}{1+\frac{c}{s}} \left(\frac{c-1}{s^2} - \frac{2(c-1)^2}{s^3} \right) \right. \\
&+ \left. \frac{1}{s(1+\frac{c}{s})^2} \left(\frac{(c-1)^2}{cs^3} - \frac{s}{c} - 1 \right) \right] \\
&= \varepsilon(2\pi\varepsilon)^{-\frac{1}{2}} (p')^{\frac{3}{2}} \exp\left(\frac{t|b'(0)|}{2}\right) \frac{1}{f''} \\
&+ \frac{\varepsilon}{2} (p')^{\frac{1}{2}} \exp\left(\frac{t|b'(0)|}{2}\right) \frac{1}{(f'')^{\frac{1}{2}}} \left(-\frac{b''(0)}{2|b'(0)|}\right) [\dots].
\end{aligned}$$

A tedious calculation shows that the terms within the $[\dots]$ cancel leaving the result,

$$\begin{aligned}
\nu_0^+[t, \infty) &= \varepsilon(2\pi\varepsilon)^{-\frac{1}{2}} (p_0')^{\frac{3}{2}} \exp\left(\frac{t|b'(0)|}{2}\right) \frac{1}{f''} + O(\varepsilon^{\frac{3}{2}}), \\
&= \left(\frac{\varepsilon|b'(0)|}{\pi}\right)^{\frac{1}{2}} \left(e^{2|b'(0)|t} - 1\right)^{-\frac{1}{2}} + O(\varepsilon^{\frac{3}{2}}).
\end{aligned}$$

□

Acknowledgements

The author wishes to thank Aubrey Truman for his crucial input into an earlier joint collaboration concentrating on the leading order behaviour of the Poisson - Lévy excursion measure Davies and Truman [1998]. I am, as always, indebted to David Williams for his help and never-ending enthusiasm for probability.

References

P. Lévy, *Processus Stochastiques et Mouvement Brownien*, (Gauthier Villars, Paris, 1954) MR0190953

J. Hawkes and A. Truman, *Statistics of Local Time and Excursions for the Ornstein-Uhlenbeck Process*, Lecture Notes of the London Mathematical Society **167**, 91 (1991) MR1166408

A. Truman and D. Williams, *Excursions and Itô Calculus in Nelson's Stochastic Mechanics*, "Recent Developments in Quantum Mechanics", (Kluwer Academic Publishers, Netherlands, 1991)

I. M. Davies and A. Truman, *Laplace asymptotic expansions of conditional Wiener Integrals and generalized Mehler kernel formulas*, J. Math. Phys. **23**, 2059 (1982) MR0680002

I. M. Davies and A. Truman, *Semiclassical analysis and a new result in excursion theory*, in "Probability Theory and Mathematical Statistics, Proceedings of the Seventh Vilnius Conference (1998)", edited by B. Grigelionis et al, (Utrecht: VSP, Vilnius: TEV, 1999), 701 – 706

F. W. J. Olver, *Asymptotics and Special Functions*, (Academic Press, New York and London, 1974) MR0435697