

Convergence of mixing times for sequences of random walks on finite graphs

D.A. Croydon* B.M. Hambly† T. Kumagai‡

Abstract

We establish conditions on sequences of graphs which ensure that the mixing times of the random walks on the graphs in the sequence converge. The main assumption is that the graphs, associated measures and heat kernels converge in a suitable Gromov-Hausdorff sense. With this result we are able to establish the convergence of the mixing times on the largest component of the Erdős-Rényi random graph in the critical window, sharpening previous results for this random graph model. Our results also enable us to establish convergence in a number of other examples, such as finitely ramified fractal graphs, Galton-Watson trees and the range of a high-dimensional random walk.

Keywords: random walk ; mixing ; Gromov-Hausdorff convergence ; random graph ; Galton-Watson tree ; fractal graph.

AMS MSC 2010: Primary 60J10, Secondary 05C80.

Submitted to EJP on March 4, 2011, final version accepted on December 9, 2011.

1 Introduction

The geometric and analytic properties of random graphs have been the subject of much recent research. One strand of this development has been to examine sequences of random subgraphs of vertex transitive graphs that are, in some sense, at or near criticality. A key example is the percolation model and, for bond percolation above the upper critical dimension, we expect to see mean-field behavior in the sequence of finite graphs in the critical window. That is, the natural scaling exponents for the volume and diameter of the graph and for the mixing time are of the same order as those for the Erdős-Rényi random graph in the critical window, as given in [36].

This mean-field behavior is seen in other natural models of sequences of critical random graphs. For example [6] obtained general conditions for the geometric properties of percolation clusters on sequences of finite graphs and discussed examples such

*Department of Statistics, University of Warwick, Coventry, CV4 7AL, United Kingdom
E-mail: d.a.croydon@warwick.ac.uk

†Mathematical Institute, University of Oxford, 24-29 St Giles', Oxford, OX1 3LB, United Kingdom
E-mail: hambly@maths.ox.ac.uk

‡RIMS, Kyoto University, Kyoto 606-8502, Japan
E-mail: kumagai@kurims.kyoto-u.ac.jp

as the high dimensional torus and the n -cube, while the random walk on critical percolation clusters on the high-dimensional torus is treated in [24]. Motivated by these results we will focus on the asymptotic behavior of mixing times for random walks on sequences of finite graphs. We consider general sequences of graphs but under some strong conditions which will enable us to establish the convergence of the mixing time.

In order to demonstrate our main result we consider the Erdős-Rényi random graph. Let $G(N, p)$ be the random subgraph of the complete graph on N labeled vertices $\{1, \dots, N\}$ in which each edge is present with probability p independently of the other edges. It is a classical result that if we set $p = c/N$, then as $N \rightarrow \infty$, if $c > 1$ there is a giant component containing a positive fraction of the vertices, while for $c < 1$ the largest component is of size $\log N$. However, if $p = N^{-1} + \lambda N^{-4/3}$ for some $\lambda \in \mathbb{R}$, we are in the so-called critical window, and it is known that the largest connected component \mathcal{C}^N , is of order $N^{2/3}$. The recent work of [1] has shown that the scaling limit of the graph, \mathcal{M} , exists and can be constructed from the continuum random tree.

For the Erdős-Rényi random graph above criticality, [19] and [4] established mixing time bounds for the simple random walk on the giant component. The simple random walk on this graph is the discrete time Markov chain with transition probabilities determined by $p(x, y) = 1/\deg(x)$ for all y such that (x, y) is an edge in \mathcal{C}^N . For the random graph in the critical window, the following result on the mixing time $t_{\text{mix}}^1(\mathcal{C}^N)$ (a precise definition will be given later in (1.8), see also Remark 1.3) of the lazy random walk (a version of the simple random walk which remains at its current vertex with probability $1/2$, otherwise it moves as the simple random walk) was obtained by Nachmias and Peres ([36, Theorem 1.1]).

Theorem 1.1. *Let \mathcal{C}^N be the largest connected component of $G(N, (1 + \lambda N^{-1/3})/N)$ for some $\lambda \in \mathbb{R}$. Then, for any $\epsilon > 0$, there exists $A = A(\epsilon, \lambda) < \infty$ such that for all large N ,*

$$P(t_{\text{mix}}^1(\mathcal{C}^N) \notin [A^{-1}N, AN]) < \epsilon.$$

It is natural to ask for more refined results on the behavior of the family of mixing times. The purpose of this paper is to give a general criteria for the convergence of mixing times for a sequence of simple random walks on finite graphs in the setting where the graphs can be embedded nicely in a compact metric space. Due to the recent work of [1] and [9] we can apply our main result to the case of the Erdős-Rényi random graph, to obtain the following result.

Theorem 1.2. *Fix $p \in [1, \infty]$. If $t_{\text{mix}}^p(\rho^N)$ is the L^p -mixing time of the simple random walk on \mathcal{C}^N started from its root ρ^N , then*

$$N^{-1}t_{\text{mix}}^p(\rho^N) \rightarrow t_{\text{mix}}^p(\rho),$$

in distribution, where the random variable $t_{\text{mix}}^p(\rho) \in (0, \infty)$ is the L^p -mixing time of the Brownian motion on \mathcal{M} started from ρ .

We will later illustrate our main result with a number of other examples of random walks on sequences of finite graphs. In order to state it, though, we start by describing the general framework in which we work. Firstly, let (F, d_F) be a compact metric space and let π be a non-atomic Borel probability measure on F with full support. We will assume that balls $B_F(x, r) := \{y \in F : d_F(x, y) < r\}$ are π -continuity sets (i.e. $\pi(\partial B_F(x, r)) = 0$ for every $x \in F, r > 0$). Secondly, take $X^F = (X_t^F)_{t \geq 0}$ to be a π -symmetric Hunt process on F (for definition and properties see [20]), which will typically be the Brownian motion on the limit of the sequence of graphs. We suppose the

following:

- X^F is conservative, i.e. its semigroup $(P_t)_{t \geq 0}$ satisfies $P_t 1 = 1$, π -a.e., $\forall t > 0$, (1.1)

- there exists a jointly continuous transition density $(q_t(x, y))_{x, y \in F, t > 0}$ of X^F , (1.2)

- for every $x, y \in F$ and $t > 0$, $q_t(x, y) > 0$, (1.3)

- for every $x \in F$ and $t > 0$, $q_t(x, \cdot)$ is not identically equal to 1, (1.4)

where conditions (1.3) and (1.4) are assumed to exclude various trivial cases, and by transition density we mean the kernel $q_t(x, y)$ such that

$$\mathbf{E}_x[f(X_t^F)] = \int_F q_t(x, y)f(y)\pi(dy),$$

for all bounded continuous function f on F . Furthermore, we will say that the transition density $(q_t(x, y))_{x, y \in F, t > 0}$ converges to stationarity in an L^p sense for some $p \in [1, \infty]$ if it holds that

$$\lim_{t \rightarrow \infty} D_p(x, t) = 0, \tag{1.5}$$

for every $x \in F$, where $D_p(x, t) := \|q_t(x, \cdot) - 1\|_{L^p(\pi)}$. If this previous condition is satisfied, then it is possible to check that the L^p -mixing time of F ,

$$t_{\text{mix}}^p(F) := \inf \left\{ t > 0 : \sup_{x \in F} D_p(x, t) \leq 1/4 \right\}, \tag{1.6}$$

is a finite quantity (see Section 3). Finally, note that $t_{\text{mix}}^p(F) \leq t_{\text{mix}}^{p'}(F)$ for $p \leq p'$, which can easily be shown using the Hölder inequality.

We continue by introducing some general notation for graphs and their associated random walks. First, fix $G = (V(G), E(G))$ to be a finite connected graph with at least two vertices, where $V(G)$ denotes the vertex set and $E(G)$ the edge set of G , and suppose d_G is a metric on $V(G)$. In some examples, d_G will be a rescaled version of the usual shortest path graph distance, by which we mean that $d_G(x, y)$ is some multiple of the number of edges in the shortest path from x to y in G , but this is not always the most convenient choice. Define a symmetric weight function $\mu^G : V(G)^2 \rightarrow \mathbb{R}_+$ that satisfies $\mu_{xy}^G > 0$ if and only if $\{x, y\} \in E(G)$. The discrete time random walk on the weighted graph G is then the Markov chain $((X_m^G)_{m \geq 0}, \mathbf{P}_x^G, x \in V(G))$ with transition probabilities $(P_G(x, y))_{x, y \in V(G)}$ defined by $P_G(x, y) := \mu_{xy}^G / \mu_x^G$, where $\mu_x^G := \sum_{y \in V(G)} \mu_{xy}^G$. If we define a measure π^G on $V(G)$ by setting, for $A \subseteq V(G)$, $\pi^G(A) := \sum_{x \in A} \mu_x^G / \sum_{x \in V(G)} \mu_x^G$, then π^G is the invariant probability measure for X^G . The transition density of X^G , with respect to π^G , is given by $(p_m^G(x, y))_{x, y \in V(G), m \geq 0}$, where

$$p_m^G(x, y) := \frac{\mathbf{P}_x^G(X_m = y)}{\pi^G(\{y\})}.$$

Due to parity concerns for bipartite graphs, we will consider a smoothed version of this function $(q_m^G(x, y))_{x, y \in V(G), m \geq 0}$ obtained by setting

$$q_m^G(x, y) := \frac{p_m^G(x, y) + p_{m+1}^G(x, y)}{2}, \tag{1.7}$$

and define the L^p -mixing time of G by

$$t_{\text{mix}}^p(G) := \inf \left\{ m > 0 : \sup_{x \in V(G)} D_p^G(x, m) \leq 1/4 \right\}, \tag{1.8}$$

where $D_p^G(x, m) := \|q_m^G(x, \cdot) - 1\|_{L^p(\pi^G)}$. Finally, in the case that we are considering a sequence of graphs $(G^N)_{N \geq 1}$, we will usually abbreviate π^{G^N} to π^N and q^{G^N} to q^N , etc.

Remark 1.3. In [36], the mixing time of \mathcal{C}^N is defined in terms of the total variation distance, that is

$$T_{\text{mix}}(\mathcal{C}^N) = \min\{t : \|P_t(x, \cdot) - \pi(\cdot)\|_{\text{TV}} \leq 1/8, \forall x \in V(\mathcal{C}^N)\}, \quad (1.9)$$

where $P_t(x, B) = \sum_{y \in B} p_t^N(x, y)\pi(y)$ for $B \subset V(\mathcal{C}^N)$, $p_t^N(x, y)$ is the transition density for the random walk and $\|\mu - \nu\|_{\text{TV}} = \max_{B \subset V(\mathcal{C}^N)} |\mu(B) - \nu(B)|$ for probability measures μ, ν on $V(\mathcal{C}^N)$. (To be precise, 1/8 in (1.9) is 1/4 in [36], but this only affects the constants in the results.) However, noting that

$$\|\mu - \nu\|_{\text{TV}} = \frac{1}{2} \sum_{x \in V(\mathcal{C}^N)} |\mu(\{x\}) - \nu(\{x\})|,$$

(see, for example [35, Proposition 4.2]), one sees that $T_{\text{mix}}(\mathcal{C}^N) = t_{\text{mix}}^1(\mathcal{C}^N)$. Also note that [36] considers the lazy walk on the graph to avoid parity issues, but the same techniques will apply to the mixing time defined in terms of the smoothed heat kernel introduced at (1.7).

We are now ready to state the assumption under which we are able to prove the convergence of mixing times for the random walks on a sequence of graphs. This captures the idea that, when suitably rescaled, the discrete state spaces, invariant measures and transition densities of a sequence of graphs converge to (F, d_F) , π and $(q_t(x, y))_{x, y \in F, t > 0}$, respectively. Its formulation involves a spectral Gromov-Hausdorff topology, the definition of which is postponed until Section 2, and a useful sufficient condition for it will be given in Proposition 2.4 below. Note that we extend the definition of the discrete transition densities on graphs to all positive times by linear interpolation of $(q_m^G(x, y))_{m \geq 0}$ for each pair of vertices $x, y \in V(G)$. Note also that the extended transition densities are different from those of continuous time Markov chains.

Assumption 1. $(G^N)_{N \geq 1}$ is a sequence of finite connected graphs with at least two vertices for which there exists a sequence $(\gamma(N))_{N \geq 1}$ such that, for any compact interval $I \subset (0, \infty)$,

$$\left((V(G^N), d_{G^N}), \pi^{G^N}, (q_{\gamma(N)t}^N(x, y))_{x, y \in V(G^N), t \in I} \right) \rightarrow ((F, d_F), \pi, (q_t(x, y))_{x, y \in F, t \in I})$$

in a spectral Gromov-Hausdorff sense.

In the case where we have random graphs, we will typically assume that we have the above convergence holding in distribution. Our main conclusion is then the following.

Theorem 1.4. Suppose that Assumption 1 is satisfied. If $p \in [1, \infty]$ is such that the transition density $(q_t(x, y))_{x, y \in F, t > 0}$ converges to stationarity in an L^p sense, then $t_{\text{mix}}^p(F) \in (0, \infty)$ and

$$\gamma(N)^{-1} t_{\text{mix}}^p(G^N) \rightarrow t_{\text{mix}}^p(F). \quad (1.10)$$

In Section 3.2, we will explain how to derive a variation of Theorem 1.4 that concerns the convergence of mixing times of processes started at a distinguished point in the state space.

We emphasize that a key part of our paper is to verify Assumption 1 and apply Theorem 1.4 in various interesting examples (including the Erdős-Rényi random graphs in the critical window as mentioned above). Therefore, we devote considerable space to applying our results to such examples.

The organization of the paper is as follows. In Section 2, we give a precise definition of the spectral Gromov-Hausdorff convergence and give some of its basic properties. In Section 3, we prove Theorem 1.4 and derive a variation of the theorem for distinguished starting points. Some sufficient conditions for (1.1)-(1.5) are given in Section 4. A selection of examples where the assumptions of Theorem 1.4 can be verified, and hence we have convergence of the mixing time sequence, are given in Section 5. In Section 6 we introduce some geometric conditions on graphs for upper and lower bounds on the mixing times for the corresponding symmetric Markov chains. We use these ideas to derive tail estimates of mixing times on random graphs in the case of the continuum random tree and the Erdős-Rényi random graph. The proofs of these results can be found in the Appendix.

2 Spectral Gromov-Hausdorff convergence

The aim of this section is to define a spectral Gromov-Hausdorff distance on triples consisting of a metric space, a measure and a heat kernel-type function that will allow us to make Assumption 1 precise. We will also derive an equivalent characterization of this assumption that will be applied in the subsequent section when proving our mixing time convergence result, and present a sufficient condition for Assumption 1 that will be useful when it comes to checking it in examples. Note that we do not need to assume (1.3), (1.4) in this section, and only use (1.1) to deduce Proposition 2.4 from a result of [14].

First, for a compact interval $I \subset (0, \infty)$, let $\tilde{\mathcal{M}}_I$ be the collection of triples of the form (F, π, q) , where $F = (F, d_F)$ is a non-empty compact metric space, π is a Borel probability measure on F and $q = (q_t(x, y))_{x, y \in F, t \in I}$ is a jointly continuous real-valued function of (t, x, y) . We say two elements, (F, π, q) and (F', π', q') , of $\tilde{\mathcal{M}}_I$ are equivalent if there exists an isometry $f : F \rightarrow F'$ such that $\pi \circ f^{-1} = \pi'$ and $q'_t \circ f = q_t$ for every $t \in I$, by which we mean $q'_t(f(x), f(y)) = q_t(x, y)$ for every $x, y \in F, t \in I$. Define \mathcal{M}_I to be the set of equivalence classes of $\tilde{\mathcal{M}}_I$ under this relation. We will often abuse notation and identify an equivalence class in \mathcal{M}_I with a particular element of it. Now, set

$$\begin{aligned} \Delta_I((F, \pi, q), (F', \pi', q')) \\ := \inf_{Z, \phi, \phi', \mathcal{C}} \left\{ d_H^Z(\phi(F), \phi'(F')) + d_P^Z(\pi \circ \phi^{-1}, \pi' \circ \phi'^{-1}) \right. \\ \left. + \sup_{(x, x'), (y, y') \in \mathcal{C}} \left(d_Z(\phi(x), \phi'(x')) + d_Z(\phi(y), \phi'(y')) + \sup_{t \in I} |q_t(x, y) - q'_t(x', y')| \right) \right\}, \end{aligned}$$

where the infimum is taken over all metric spaces $Z = (Z, d_Z)$, isometric embeddings $\phi : F \rightarrow Z, \phi' : F' \rightarrow Z$, and correspondences \mathcal{C} between F and F' , d_H^Z is the Hausdorff distance between compact subsets of Z , and d_P^Z is the Prohorov distance between Borel probability measures on Z . Note that, by a correspondence \mathcal{C} between F and F' , we mean a subset of $F \times F'$ such that for every $x \in F$ there exists at least one $x' \in F'$ such that $(x, x') \in \mathcal{C}$ and conversely for every $x' \in F'$ there exists at least one $x \in F$ such that $(x, x') \in \mathcal{C}$.

In the following lemma, we check that the above definition gives us a metric and that the corresponding space is separable. (The latter fact will be useful when it comes to making convergence in distribution statements regarding the mixing times of sequences of random graphs, as is done in Sections 5.2 and 5.3, for example). Before this, however, let us make a few remarks about the inspiration for the distance in question. In the infimum characterizing Δ_I , the first term is simply that used in the standard Gromov-Hausdorff distance (see [7, Definition 7.3.10], for example). In fact, as far as

the topology is considered, this term could have been omitted since it is absorbed by the other terms in the expression, but we find that it is technically convenient and somewhat instructive to maintain it. The second term is that considered by the authors of [23] in defining their ‘Gromov-Prohorov’ distance between metric measure spaces. The final term is closely related to one used in [17, Section 6] when defining a distance between spatial trees – real trees equipped with a continuous function. Indeed, the notion of a correspondence is quite standard in the Gromov-Hausdorff setting as a way to relate two compact metric spaces. One can, for example, alternatively define the Gromov-Hausdorff distance between compact metric spaces as half the infimum of the distortion of the correspondences between them (see [7, Theorem 7.3.25]).

Lemma 2.1. *For any compact interval $I \subset (0, \infty)$, $(\mathcal{M}_I, \Delta_I)$ is a separable metric space.*

Proof. Fix a compact interval $I \subset (0, \infty)$. That Δ_I is a non-negative function and is symmetric is obvious. To prove that it is also the case that $\Delta_I((F, \pi, q), (F', \pi', q')) < \infty$ for any choice of $(F, \pi, q), (F', \pi', q') \in \mathcal{M}_I$, simply consider Z to be the disjoint union of F and F' , setting $d_Z(x, x') := \text{diam}(F, d_F) + \text{diam}(F', d_{F'})$ for any $x \in F, x' \in F'$, and suppose that $\mathcal{C} = F \times F'$.

We next show that Δ_I is positive definite. Suppose $(F, \pi, q), (F', \pi', q') \in \mathcal{M}_I$ are such that $\Delta_I((F, \pi, q), (F', \pi', q')) = 0$. For every $\varepsilon > 0$, we can thus choose $Z, \phi, \phi', \mathcal{C}$ such that the sum of quantities in the defining infimum of Δ_I is bounded above by ε . Moreover, there exists a $\delta \in (0, \varepsilon]$ such that

$$\sup_{\substack{x_1, x_2, y_1, y_2 \in F: \\ d_F(x_1, x_2), d_{F'}(y_1, y_2) \leq \delta}} \sup_{t \in I} |q_t(x_1, y_1) - q_t(x_2, y_2)| \leq \varepsilon. \tag{2.1}$$

Now, let $(x_i)_{i=1}^\infty$ be a dense sequence of disjoint elements of F (in the case F is finite, we suppose that the sequence terminates after having listed all of the elements of F). By the compactness of F , there exists an integer N_ε such that $(B_F(x_i, \delta))_{i=1}^{N_\varepsilon}$ is a cover for F . Define $A_1 := B_F(x_1, \delta)$, and $A_i := B_F(x_i, \delta) \setminus \cup_{j=1}^{i-1} B_F(x_j, \delta)$ for $i = 2, \dots, N_\varepsilon$, so that $(A_i)_{i=1}^{N_\varepsilon}$ is a disjoint cover of F , and then consider a function $f_\varepsilon : F \rightarrow F'$ obtained by setting

$$f_\varepsilon(x) := x'_i$$

on A_i , where x'_i is chosen such that $(x_i, x'_i) \in \mathcal{C}$ for each $i = 1, \dots, N_\varepsilon$. Clearly, by definition, f_ε is a measurable function. It is further the case that it satisfies, for any $x \in F$,

$$d_Z(\phi(x), \phi'(f_\varepsilon(x))) \leq d_Z(\phi(x), \phi(x_i)) + d_Z(\phi(x_i), \phi'(x'_i)) \leq 2\varepsilon,$$

where, in the above, we assume that $i \in \{1, \dots, N_\varepsilon\}$ is such that $x \in A_i$. From this, it readily follows that:

$$\sup_{x, y \in F} |d_F(x, y) - d_{F'}(f_\varepsilon(x), f_\varepsilon(y))| \leq 4\varepsilon \tag{2.2}$$

and

$$d_P^{F'}(\pi \circ f_\varepsilon^{-1}, \pi') \leq 3\varepsilon, \tag{2.3}$$

where $d_P^{F'}$ is the Prohorov distance on F' . By applying (2.1), we also have that

$$\sup_{x, y \in F, t \in I} |q_t(x, y) - q'_t(f_\varepsilon(x), f_\varepsilon(y))| \leq 2\varepsilon. \tag{2.4}$$

To continue, we use a diagonalization argument to deduce the existence of a sequence $(\varepsilon_n)_{n \geq 1}$ such that $f_{\varepsilon_n}(x_i)$ converges to some limit $f(x_i) \in F'$ for every $i \geq 1$. From (2.2), we obtain that $d_{F'}(f(x_i), f(x_j)) = d_F(x_i, x_j)$ for every $i, j \geq 1$, and so we can extend the map f continuously to the whole of F ([7, Proposition 1.5.9]). This construction

immediately implies that f is distance preserving. Moreover, reversing the roles of F and F' , we are able to find a distance preserving map from F' to F . Hence f must be an isometry. To check that (F, π, q) and (F', π', q') are equivalent, it therefore remains to check that $\pi \circ f^{-1} = \pi'$ and $q'_t \circ f = q_t$ for every $t \in I$. Fix $\varepsilon > 0$ and recall that the definition of $(x_i)_{i=1}^{N_\varepsilon}$ means that it is an ε -net for F . Let $\varepsilon' \in (0, \varepsilon]$ be such that $d_{F'}(f_{\varepsilon'}(x_i), f(x_i)) \leq \varepsilon$ for every $i = 1, \dots, N_\varepsilon$. Then,

$$d_{F'}(f_{\varepsilon'}(x), f(x)) \leq d_{F'}(f_{\varepsilon'}(x), f_{\varepsilon'}(x_i)) + d_{F'}(f_{\varepsilon'}(x_i), f(x_i)) + d_{F'}(f(x_i), f(x)) \leq 7\varepsilon, \quad (2.5)$$

where we are again assuming that $i \in \{1, \dots, N_\varepsilon\}$ is such that $x \in A_i$, and have applied (2.2) and the distance-preserving property of f . In particular, this implies that

$$d_P^{F'}(\pi \circ f^{-1}, \pi') \leq d_P^{F'}(\pi \circ f^{-1}, \pi \circ f_{\varepsilon'}^{-1}) + d_P^{F'}(\pi \circ f_{\varepsilon'}^{-1}, \pi') \leq 10\varepsilon,$$

where we use (2.3) to deduce the second inequality. Since $\varepsilon > 0$ was arbitrary, this yields that $\pi \circ f^{-1} = \pi'$. Finally, (2.4) and (2.5) imply that

$$\sup_{x,y \in F, t \in I} |q_t(x, y) - q'_t(f(x), f(y))| \leq 2\varepsilon + \sup_{\substack{x'_1, x'_2, y'_1, y'_2 \in F': \\ d_{F'}(x'_1, x'_2), d_{F'}(y'_1, y'_2) \leq 7\varepsilon}} \sup_{t \in I} |q'_t(x'_1, y'_1) - q_t(x'_2, y'_2)|,$$

and so $q'_t \circ f = q_t$ for every $t \in I$ follows from the continuity properties of q' . This completes the proof of the fact that: if $\Delta_I((F, \pi, q), (F', \pi', q')) = 0$, then the triples (F, π, q) and (F', π', q') are equivalent in the sense described at the start of the section. Consequently, Δ_I is indeed positive definite on the set of equivalence classes \mathcal{M}_I .

For the triangle inequality, we closely follow the proof of [23, Lemma 5.2]. Let $(F^{(i)}, \pi^{(i)}, q^{(i)})$ be an element of \mathcal{M}_I , $i = 1, 2, 3$. Suppose that

$$\Delta_I((F^{(1)}, \pi^{(1)}, q^{(1)}), (F^{(2)}, \pi^{(2)}, q^{(2)})) < \delta_1,$$

so that we can find a metric space Z_1 , isometric embeddings $\phi_{1,1} : F^{(1)} \rightarrow Z_1$ and $\phi_{2,1} : F^{(2)} \rightarrow Z_1$ and correspondence \mathcal{C}_1 between $F^{(1)}$ and $F^{(2)}$ such that the sum of quantities in the defining infimum of Δ_I is bounded above by δ_1 . If

$$\Delta_I((F^{(2)}, \pi^{(2)}, q^{(2)}), (F^{(3)}, \pi^{(3)}, q^{(3)})) < \delta_2,$$

we define $Z_2, \phi_{2,2}, \phi_{3,2}, \mathcal{C}_2$ in an analogous way. Now, set Z to be the disjoint union of Z_1 and Z_2 , and define a distance on it by setting $d_Z|_{Z_i \times Z_i} = d_{Z_i}$ for $i = 1, 2$, and for $x \in Z_1, y \in Z_2$,

$$d_Z(x, y) := \inf_{z \in F^{(2)}} (d_{Z_1}(x, \phi_{2,1}(z)) + d_{Z_2}(\phi_{2,2}(z), y)).$$

Abusing notation slightly, it is then the case that, after points separated by a 0 distance have been identified, (Z, d_Z) is a metric space into which there is a natural isometric embedding ϕ_i of Z_i , $i = 1, 2$. In this space, we have that

$$\begin{aligned} d_H^Z(\phi_1(\phi_{1,1}(F^{(1)})), \phi_2(\phi_{3,2}(F^{(3)}))) \\ \leq d_H^{Z_1}(\phi_{1,1}(F^{(1)}), \phi_{2,1}(F^{(2)})) + d_H^{Z_2}(\phi_{2,2}(F^{(2)}), \phi_{3,2}(F^{(3)})), \end{aligned}$$

where we have applied the fact that $\phi_1(\phi_{2,1}(y)) = \phi_2(\phi_{2,2}(y))$ for every $y \in F^{(2)}$, and so $\phi_1(\phi_{2,1}(F^{(2)})) = \phi_2(\phi_{2,2}(F^{(2)}))$ as subsets of Z . A similar bound applies to the embedded measures. Now, let

$$\mathcal{C} := \{(x, z) \in F^{(1)} \times F^{(3)} : \exists y \in F^{(2)} \text{ such that } (x, y) \in \mathcal{C}_1, (y, z) \in \mathcal{C}_2\},$$

then if $(x, z) \in \mathcal{C}$,

$$d_Z(\phi_1(\phi_{1,1}(x)), \phi_2(\phi_{3,2}(z))) \leq d_{Z_1}(\phi_{1,1}(x), \phi_{2,1}(y)) + d_{Z_2}(\phi_{2,2}(y), \phi_{3,2}(z)),$$

where $y \in F^{(2)}$ is chosen such that $(x, y) \in \mathcal{C}_1$ and $(y, z) \in \mathcal{C}_2$, and we again note $\phi_1(\phi_{2,1}(y)) = \phi_2(\phi_{2,2}(y))$. Proceeding in the same fashion, one can deduce a corresponding bound involving $q^{(i)}$, $i = 1, 2, 3$. Putting these pieces together, it is elementary to deduce that

$$\Delta_I((F^{(1)}, \pi^{(1)}, q^{(1)}), (F^{(3)}, \pi^{(3)}, q^{(3)})) \leq \delta_1 + \delta_2,$$

and the triangle inequality follows. Thus we have proved that $(\mathcal{M}_I, \Delta_I)$ is a metric space.

To complete the proof, we only need to show separability. This is straightforward, however, as for any element of \mathcal{M}_I , one can construct an approximating sequence that incorporates only: metric spaces with a finite number of points and rational distances between them, probability measures on these with a rational mass at each point, and functions that are defined (at each coordinate pair) to be equal to rational values at a finite collection of rational time points and are linear between these. To be more explicit, let (F, π, q) be an element of \mathcal{M}_I , and then define a sequence $(F^N, \pi^N, q^N)_{N \geq 1}$ as follows. First, let F^N be a finite N^{-1} -net of F , which exists because F is compact. By perturbing d_F , it is possible to define a metric d_{F^N} on F^N such that $|d_{F^N}(x, y) - d_F(x, y)| \leq N^{-1}$ and moreover $d_{F^N}(x, y) \in \mathbb{Q}$ for all $x, y \in F^N$. Now, since F^N is an N^{-1} -net of F , it is possible to choose a partition $(A_x)_{x \in F^N}$ of F such that $x \in A_x$ and the diameter of A_x (with respect to d_F) is no greater than $2N^{-1}$. Moreover, it is possible to choose the partition in such a way that A_x is measurable for each $x \in F^N$. We construct a probability measure on F^N by choosing $\pi^N(\{x\}) \in \mathbb{Q}$ such that $|\pi^N(\{x\}) - \pi(A_x)| \leq N^{-1}$ (subject to the constraint that $\sum_{x \in F^N} \pi^N(\{x\}) = 1$). Finally, define ε_N by setting

$$\varepsilon_N := \sup_{\substack{s, t \in I: \\ |s-t| \leq N^{-1}}} \sup_{\substack{x, x', y, y' \in F^N: \\ d_F(x, x'), d_F(y, y') \leq N^{-1}}} |q_s(x, y) - q_t(x', y')|,$$

so that, by the joint continuity of q , $\varepsilon_N \rightarrow 0$ as $N \rightarrow \infty$. Let $\inf I \leq t_0 \leq t_1 \leq \dots \leq t_K \leq \sup I$ be a set of rational times such that $|t_0 - \inf I|, |\sup I - t_K|, |t_{i+1} - t_i| \leq N^{-1}$, choose $q_{t_i}^N(x, y) \in \mathbb{Q}$ such that $|q_{t_i}^N(x, y) - q_{t_i}(x, y)| \leq N^{-1}$ for each $x, y \in F^N$, and then extend q^N to have domain $F^N \times F^N \times I$ by linear interpolation in t at each pair of vertices. This construction readily yields that $\Delta_I((F, \pi, q), (F^N, \pi^N, q^N)) \leq 6N^{-1} + 3\varepsilon_N \rightarrow 0$. Since the class of triples from which the approximating sequence is chosen is clearly countable, this completes the proof of separability. \square

We will say that a sequence in \mathcal{M}_I converges in a spectral Gromov-Hausdorff sense if it converges to a limit in this space with respect to the metric Δ_I . We note that in the framework of compact Riemannian manifolds, different but related notions of spectral distances were introduced by Bérard, Besson and Gallot ([5]) and by Kasue and Kumura ([25]). Moreover, by applying our characterization of spectral Gromov-Hausdorff convergence, we are able to deduce that if Assumption 1 holds, then we can isometrically embed all the rescaled graphs, measures and transition densities upon them into a common metric space (E, d_E) so that they converge to the relevant limit objects in a more standard way, as the following lemma makes precise. Note that in the proof of the result and henceforth we define balls in the space (E, d_E) by setting $B_E(x, r) := \{x \in E : d_E(x, y) < r\}$.

Lemma 2.2. *Suppose that Assumption 1 is satisfied. For any compact interval $I \subset (0, \infty)$, there exist isometric embeddings of $(V(G^N), d_{G^N})$, $N \geq 1$, and (F, d_F) into a common metric space (E, d_E) such that*

$$\lim_{N \rightarrow \infty} d_H^E(V(G^N), F) = 0, \tag{2.6}$$

$$\lim_{N \rightarrow \infty} d_P^E(\pi^N, \pi) = 0, \tag{2.7}$$

and also,

$$\lim_{N \rightarrow \infty} \sup_{x, y \in F} \sup_{t \in I} \left| q_{\gamma(N)t}^N(g_N(x), g_N(y)) - q_t(x, y) \right| = 0, \tag{2.8}$$

where, for brevity, we have identified the spaces $(V(G^N), d_{G^N})$, $N \geq 1$, and (F, d_F) , and the measures upon them with their isometric embeddings in (E, d_E) . For each $x \in F$, we define $g_N(x)$ to be a vertex in $V(G^N)$ minimizing $d_E(x, y)$ over $y \in V(G^N)$.

Proof. Fix a compact interval $I \subset (0, \infty)$. By Assumption 1, for each $N \geq 1$ it is possible to find metric spaces (E_N, d_N) , isometric embeddings $\phi_N : (V(G^N), d_{G^N}) \rightarrow (E_N, d_N)$, $\phi'_N : (F, d_F) \rightarrow (E_N, d_N)$ and correspondences \mathcal{C}_N between $V(G^N)$ and F such that, identifying the original objects and their embeddings,

$$d_H^{E^N}(V(G^N), F) + d_P^{E^N}(\pi^N, \pi) + \sup_{(x, x'), (y, y') \in \mathcal{C}_N} \left(d_N(x, x') + d_N(y, y') + \sup_{t \in I} \left| q_{\gamma(N)t}^N(x, y) - q_t(x', y') \right| \right) \leq \varepsilon_N, \tag{2.9}$$

where $\varepsilon_N \rightarrow 0$. Now, proceeding similarly to the proof of the triangle inequality in Lemma 2.1, set E to be the disjoint union of E^N , $N \geq 1$, and define a distance on it by setting $d_E|_{E^N \times E^N} = d_N$ for $N \geq 1$, and for $x \in E^N$, $x' \in E^{N'}$, $N \neq N'$, set

$$d_E(x, x') := \inf_{y \in F} (d_N(x, y) + d_{N'}(y, x')).$$

Quotienting out points that are separated by distance 0 results in a metric space (E, d_E) (again, this is a slight abuse of notation), into which we have natural isometric embeddings of the metric spaces $(V(G^N), d_{G^N})$, $N \geq 1$, and (F, d_F) . Moreover, in the metric space (E, d_E) , it readily follows from (2.9) that the relevant isometrically embedded objects satisfy (2.6) and (2.7). To prove (2.8), first note that for every $x \in V(G^N)$, $N \geq 1$, there exists an $x' \in F$ such that $(x, x') \in \mathcal{C}_N$. This implies that $d_E(x, x') \leq \varepsilon_N$, and so, for any $\delta > 0$,

$$\begin{aligned} & \sup_{\substack{x, y, z \in V(G^N): \\ d_{G^N}(y, z) \leq \delta}} \sup_{t \in I} \left| q_{\gamma(N)t}^N(x, y) - q_{\gamma(N)t}^N(x, z) \right| \\ & \leq 2\varepsilon_N + \sup_{\substack{x, y, z \in F: \\ d_F(y, z) \leq \delta + 2\varepsilon_N}} \sup_{t \in I} |q_t(x, y) - q_t(x, z)|. \end{aligned} \tag{2.10}$$

Now, for every $x \in F$ and $N \geq 1$, there exists an $x' \in V(G^N)$ such that $(x', x) \in \mathcal{C}_N$, and so $d_E(x', x) \leq \varepsilon_N$. Therefore, since $g_N(x)$ is the closest vertex of $V(G^N)$ to x ,

$$g_N(x) \in B_E(x, 2\varepsilon_N) \cap V(G^N) \subseteq B_E(x', 3\varepsilon_N) \cap V(G^N) = B_{V(G^N)}(x', 3\varepsilon_N).$$

Consequently,

$$\begin{aligned} & \sup_{x, y \in F} \sup_{t \in I} \left| q_{\gamma(N)t}^N(g_N(x), g_N(y)) - q_t(x, y) \right| \\ & \leq \varepsilon_N + 2 \sup_{\substack{x, y, z \in V(G^N): \\ d_{G^N}(y, z) \leq 3\varepsilon_N}} \sup_{t \in I} \left| q_{\gamma(N)t}^N(x, y) - q_{\gamma(N)t}^N(x, z) \right| \\ & \leq 5\varepsilon_N + 2 \sup_{\substack{x, y, z \in F: \\ d_F(y, z) \leq 5\varepsilon_N}} \sup_{t \in I} |q_t(x, y) - q_t(x, z)|, \end{aligned}$$

where the second inequality is an application of (2.10). Letting $N \rightarrow \infty$ and applying the joint continuity of $(q_t(x, y))_{x, y \in F, t > 0}$, we obtain the desired result. \square

For our later convenience, let us note a useful tightness condition for the rescaled transition densities that was essentially established in the proof of the previous result.

Lemma 2.3. *Suppose that Assumption 1 holds. For any compact interval $I \subset (0, \infty)$,*

$$\lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_{\substack{x, y, z \in V(G^N): \\ d_{G^N}(y, z) \leq \delta}} \sup_{t \in I} \left| q_{\gamma(N)t}^N(x, y) - q_{\gamma(N)t}^N(x, z) \right| = 0. \quad (2.11)$$

Proof. Recalling the continuity property of q , taking the limit as $N \rightarrow \infty$ in (2.10) yields

$$\limsup_{N \rightarrow \infty} \sup_{\substack{x, y, z \in V(G^N): \\ d_{G^N}(y, z) \leq \delta}} \sup_{t \in I} \left| q_{\gamma(N)t}^N(x, y) - q_{\gamma(N)t}^N(x, z) \right| \leq \sup_{\substack{x, y, z \in F: \\ d_F(y, z) \leq \delta}} \sup_{t \in I} |q_t(x, y) - q_t(x, z)|.$$

Again appealing to the continuity of q , the right-hand side here converges to 0 as $\delta \rightarrow 0$, which completes the proof. \square

It is straightforward to reverse the conclusions of the previous two lemmas to check that if (2.6), (2.7), (2.8) and (2.11) hold, then so does Assumption 1. Indeed, under these assumptions, we have isometric embeddings of $(V(G^N), d_{G^N})$, $N \geq 1$, and (F, d_F) into a common metric space (E, d_E) for which: (2.6) gives the Hausdorff convergence of sets; (2.7) gives the Prohorov convergence of measures; and moreover, it is elementary to check from (2.8) and (2.11) that, with respect to the correspondences

$$\mathcal{C}_N := \{(x, x') \in F \times V(G^N) : d_E(x, x') \leq N^{-1}\},$$

the relevant transition densities converge uniformly, as described in the definition of the metric Δ_I . Thus, in examples, it will suffice to check these equivalent conditions when seeking to verify Assumption 1. In fact, it is further possible to weaken these assumptions slightly by appealing to a local limit theorem from [14]. To be precise, because we are assuming that the transition densities of the graph satisfy the tightness condition of (2.11), we can apply [14, Theorem 15], to replace the local convergence statement of (2.8) with a central limit-type convergence statement. Note that, although in [14] it was assumed that the metric on G^N was a shortest path graph distance, exactly the same argument yields the corresponding conclusion in our setting, and so we simply state the result.

Proposition 2.4 (cf. [14, Theorem 15]). *Suppose that $(V(G^N), d_{G^N})$, $N \geq 1$, and (F, d_F) can be isometrically embedded into a common metric space (E, d_E) in such a way that (2.6) and (2.7) are both satisfied. Moreover, assume that there exists a dense subset F^* of F such that, for any compact interval $I \subset (0, \infty)$, $x \in F^*$, $y \in F$, $r > 0$,*

$$\lim_{N \rightarrow \infty} \mathbf{P}_{g_N(x)}^{G^N} \left(X_{\lfloor \gamma(N)t \rfloor}^{G^N} \in B_E(y, r) \right) = \mathbf{P}_x^F \left(X_t^F \in B_E(y, r) \right) \quad (2.12)$$

uniformly for $t \in I$, and also (2.11) holds. Then Assumption 1 holds.

To complete this section, let us observe that [14] also provides two ways to check (2.11): one involving a resistance estimate on the graphs in the sequence ([14, Proposition 17]), and one involving the parabolic Harnack inequality ([14, Proposition 16]). Since the first of these two methods will be applied in several of our examples later, let us recall the result here. To allow us to state the result, we define $R_{G^N}(x, y)$ to be the resistance between x and y in $V(G^N)$ (see (6.1)), when we suppose that G^N is an electrical network with conductances of edges being given by the weight function μ^{G^N} . This

defines a metric on $V(G^N)$, for which the following result is proved as [14, Proposition 17]. As above, note that although it was a shortest path graph distance considered in [14], the same proof applies for a general distance on the graph in question. Moreover, the statement of the lemma is slightly different from that of the corresponding result in [14], because there the scaling $\alpha(n)$ was absorbed into the definition of the metric.

Lemma 2.5 (cf. [14, Proposition 17]). *Suppose that there exists a sequence $(\alpha(N))_{N \geq 1}$ and constants $\kappa, c_1, c_2, c_3 \in (0, \infty)$ such that*

$$R_{G^N}(x, y) \leq c_1 (\alpha(N) d_{G^N}(x, y))^\kappa, \quad \forall x, y \in V(G^N),$$

and also

$$c_2 \gamma(N) \leq \alpha(N)^\kappa \beta(N) \leq c_3 \gamma(N),$$

where $\beta(N) := \sum_{x, y \in V(G^N)} \mu_{xy}^{G^N}$, then (2.11) holds.

3 Convergence of L^p -mixing times

3.1 Proof of Theorem 1.4

In this subsection we prove the mixing time convergence result of Theorem 1.4. Throughout, we will suppose that Assumption 1 holds and that the graphs G^N and limiting metric space F have been embedded into a common metric space (E, d_E) in the way described by Lemma 2.2.

Recall from the introduction the definition of $D_p(x, t) = \|q_t(x, \cdot) - 1\|_{L^p(\pi)}$, the L^p -distance from stationarity of the process X^F started from x at time t . By applying the continuity of $(q_t(x, y))_{x, y \in F, t > 0}$, compactness of F and finiteness of π , it is easy to check that this quantity is finite for every $x \in F$ and $t > 0$. The next lemma collects together a number of other basic properties of $D_p(x, t)$ that we will apply later (the first part is a minor extension of [8, Proposition 3.1], in our setting).

Lemma 3.1. *Let $p \in [1, \infty]$. For every $x \in F$, the function $t \mapsto D_p(x, t)$ is continuous and strictly decreasing. Furthermore, we have*

$$\lim_{t \rightarrow 0} D_p(x, t) \geq 2. \tag{3.1}$$

Proof. That the function $t \mapsto D_p(x, t)$ is continuous is clear from (1.2), and so we turn to checking that it is strictly decreasing. First, a standard argument involving an application of Jensen's inequality and the invariance of π allows one to deduce that $\|P_t f\|_{L^p(\pi)} \leq \|f\|_{L^p(\pi)}$ for any $f \in L^p(F, \pi)$, where $(P_t)_{t \geq 0}$ is the semigroup naturally associated with the transition density $(q_t(x, y))_{x, y \in F, t > 0}$. Now, suppose $f \in L^p(F, \pi)$ is such that $\|P_t f\|_{L^p(\pi)} = \|f\|_{L^p(\pi)}$, and define $f_1(y) := |P_t f(y)|^p$ and $f_2(y) := P_t(|f|^p)(y)$. By the assumption on f and the fact that X^F is conservative and π -symmetric, we have

that

$$\begin{aligned}
 \int_F f_1 d\pi &= \int_F |P_t f(y)|^p \pi(dy) \\
 &= \int_F |f(y)|^p \pi(dy) \\
 &= \int_F |f(y)|^p \int_F q_t(y, z) \pi(dz) \pi(dy) \\
 &= \int_F \int_F |f(y)|^p q_t(z, y) \pi(dy) \pi(dz) \\
 &= \int_F P_t(|f|^p)(z) \pi(dz) \\
 &= \int_F f_2 d\pi.
 \end{aligned}$$

Furthermore, Jensen's inequality implies $f_1(y) \leq f_2(y)$. Thus, it must be the case that $f_1(y) = f_2(y)$, π -a.e. In particular, because π is a probability measure, there exists a $y \in F$ such that $f_1(y) = f_2(y)$.

In the case $p > 1$, the conclusion of the previous paragraph readily implies that f is constant $q_t(y, z)\pi(dz)$ -a.e. Recalling the assumption that $q_t(y, z) > 0$ everywhere, namely (1.3), it must therefore hold that f is constant π -a.e. Observing that for $s, t > 0$ we can write $D_p(x, s + t) = \|P_s(q_t(x, \cdot) - 1)\|_{L^p(\pi)}$, it follows that $D_p(x, s + t) < D_p(x, t)$ if and only if $q_t(x, \cdot) = 1$, π -a.e. However, condition (1.4) and the assumption that the transition density is continuous imply that there exists a non-empty open set on which $q_t(x, \cdot) \neq 1$. Thus, because π has full support, it is not the case that $q_t(x, \cdot) = 1$, π -a.e., and we must have $D_p(x, s + t) < D_p(x, t)$, as desired.

For $p = 1$, the result $f_1(y) = f_2(y)$ implies that f is either non-negative or non-positive, π -a.e. Consequently, if we suppose that $D_p(x, s + t) = \|P_s(q_t(x, \cdot) - 1)\|_{L^p(\pi)} = D_p(x, t)$ for some $s > 0$, then it must be the case that $q_t(x, \cdot) - 1$ is either non-negative or non-positive. However, since $\int_F (q_t(x, y) - 1)\pi(dy) = 0$ (due to (1.1)) and (1.4) holds, we arrive at a contradiction. In particular, it must be the case that $D_p(x, s + t) < D_p(x, t)$, and this completes the proof of strict monotonicity.

To establish the limit in (3.1), it will suffice to prove the result in the case $p = 1$ (obtaining the result for other values of p is then simply Jensen's inequality). Let $x \in F$ and $r > 0$, then

$$\begin{aligned}
 D_1(x, t) &\geq \int_{B_E(x, r)} (q_t(x, y) - 1)\pi(dy) + \int_{B_E(x, r)^c} (1 - q_t(x, y))\pi(dy) \\
 &= 2\mathbf{P}_x(X_t^F \in B_E(x, r)) - 2\pi(B_E(x, r)),
 \end{aligned}$$

where (1.1) is used in the last equality. Since X^F is a Hunt process, the first term here converges to 2 as $t \rightarrow 0$. Furthermore, because π is non-atomic, the second term can be made arbitrarily small by suitable choice of r . The result follows. \square

We continue by defining the L^p -mixing time at $x \in F$ by setting

$$t_{\text{mix}}^p(x) := \inf\{t > 0 : D_p(x, t) \leq 1/4\}.$$

In fact, the previous lemma yields that $t_{\text{mix}}^p(x)$ is the unique value of $t \in (0, \infty)$ such that $D_p(x, t) = 1/4$ (when (1.5) holds at x). Similarly, define the L^p -mixing time of $x \in V(G^N)$ by setting

$$t_{\text{mix}}^{N,p}(x) := \inf\{t > 0 : D_p^N(x, t) \leq 1/4\},$$

where $D_p^N(x, m) = \|q_m^N(x, \cdot) - 1\|_{L^p(\pi^N)}$. That the discrete mixing times at a point converge when suitably rescaled to the continuous mixing time there is the conclusion of the following proposition.

Proposition 3.2. *Suppose that Assumption 1 is satisfied. If $p \in [1, \infty]$ is such that (1.5) holds for $x \in F$, then*

$$\lim_{N \rightarrow \infty} \gamma(N)^{-1} t_{\text{mix}}^{N,p}(g_N(x)) = t_{\text{mix}}^p(x),$$

where, as in the statement of Lemma 2.2, $g_N(x)$ is a vertex in $V(G^N)$ that minimizes the distance $d_E(x, y)$ over $V(G^N)$.

Proof. Suppose $p \in [1, \infty]$ is such that (1.5) holds for $x \in F$, set $t_0 := t_{\text{mix}}^p(x) \in (0, \infty)$, and fix $\varepsilon > 0$. By (1.2) and the tightness of Lemma 2.3, there exists a $\delta > 0$ such that

$$\sup_{t \in I} \sup_{\substack{y, z \in F: \\ d_E(y, z) < 2\delta}} |q_t(x, y) - 1|^p - |q_t(x, z) - 1|^p < \varepsilon, \tag{3.2}$$

$$\limsup_{N \rightarrow \infty} \sup_{t \in I} \sup_{\substack{y, z \in V(G^N): \\ d_{G^N}(y, z) < 3\delta}} \left| |q_{\gamma(N)t}^N(g_N(x), y) - 1|^p - |q_{\gamma(N)t}^N(g_N(x), z) - 1|^p \right| < \varepsilon, \tag{3.3}$$

where $I := [t_0/2, 2t_0]$. Moreover, by the compactness of F , there exists a finite collection of balls $(B_E(x_i, \delta))_{i=1}^k$ covering F . Define $A_1 := B(x_1, 2\delta)$, and $A_i := B_E(x_i, 2\delta) \setminus \bigcup_{j=1}^{i-1} B_E(x_j, 2\delta)$ for $i = 2, \dots, k$, so that $(A_i)_{i=1}^k$ is a disjoint cover of the δ -enlargement of F .

We observe

$$|D_p(x, t)^p - D_p^N(g_N(x), \gamma(N)t)^p| \leq T_1 + T_2 + T_3 + T_4,$$

where

$$\begin{aligned} T_1 &:= \left| \int_F |q_t(x, y) - 1|^p \pi(dy) - \sum_{i=1}^k |q_t(x, x_i) - 1|^p \pi(A_i) \right|, \\ T_2 &:= \left| \sum_{i=1}^k |q_t(x, x_i) - 1|^p \pi(A_i) - \sum_{i=1}^k |q_t(x, x_i) - 1|^p \pi^N(A_i) \right|, \\ T_3 &:= \left| \sum_{i=1}^k |q_t(x, x_i) - 1|^p \pi^N(A_i) - \sum_{i=1}^k |q_{\gamma(N)t}^N(g_N(x), g_N(x_i)) - 1|^p \pi^N(A_i) \right|, \\ T_4 &:= \left| \sum_{i=1}^k |q_{\gamma(N)t}^N(g_N(x), g_N(x_i)) - 1|^p \pi^N(A_i) - \int_{V(G^N)} |q_{\gamma(N)t}^N(g_N(x), y) - 1|^p \pi^N(dy) \right|. \end{aligned}$$

Now, suppose $t \in I$. From (3.2), we immediately deduce that $T_1 \leq \varepsilon$. For T_2 , we first observe that the fact balls are π -continuity sets implies that A_1, \dots, A_k are also π -continuity sets. Hence $\pi^N(A_i) \rightarrow \pi(A_i)$ for each $i = 1, \dots, k$, and so $T_2 \leq \varepsilon$ for large N . That $T_3 \leq \varepsilon$ for large N is a straightforward consequence of Lemma 2.2. Finally, applying the fact that $d_H^E(F, V(G^N)) \rightarrow 0$, we deduce that, for large N , $(A_i)_{i=1}^k$ is a disjoint cover for $V(G^N)$. Since $g_N(x_i) \in B_E(x_i, \delta)$ for large N , we also have that $d_{G^N}(y, g_N(x_i)) \leq 3\delta$, uniformly over $y \in A_i$, $i = 1, \dots, k$. Thus we can appeal to (3.3) to deduce that it is also the case that $T_4 \leq \varepsilon$ for large N . In fact, each of these bounds can be assumed to hold uniformly over $t \in I$, thereby demonstrating that

$$\lim_{N \rightarrow \infty} \sup_{t \in I} |D_p(x, t) - D_p^N(g_N(x), \gamma(N)t)| = 0. \tag{3.4}$$

Since $t \mapsto D_p^N(g_N(x), \gamma(N)t)$ is a decreasing function in t for every N (cf. [8, Proposition 3.1]) and $t \mapsto D_p(x, t)$ is strictly decreasing, the proposition follows. \square

Remark 3.3. In the case $p = 2$, the proof of the previous result greatly simplifies. In particular, we note that

$$D_2(x, t)^2 = \|q_t(x, \cdot) - 1\|_2^2 = q_{2t}(x, x) - 1, \tag{3.5}$$

and similarly

$$D_2^N(x, \gamma(N)t)^2 = \|q_{\gamma(N)t}^N(x, \cdot) - 1\|_2^2 = q_{2\gamma(N)t}^N(x, x) - 1.$$

Hence the limit at (3.4) is an immediate consequence of the local limit result of (2.8), and we do not have to concern ourselves with estimating the relevant integrals directly.

To extend the above proposition to the corresponding result for the mixing times of the entire spaces, we will appeal to the following lemma, which establishes a continuity property for the L^p -mixing times from fixed starting points in the limiting space, and a related tightness property for the discrete approximations.

Lemma 3.4. *Suppose $p \in [1, \infty]$ is such that (1.5) holds for $x \in F$, then the following statements are true.*

- (a) *The function $y \mapsto t_{\text{mix}}^p(y)$ is continuous at x .*
- (b) *Under Assumption 1, it is the case that*

$$\lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_{\substack{y \in V(G^N): \\ d_{G^N}(g_N(x), y) < \delta}} \gamma(N)^{-1} \left| t_{\text{mix}}^{N,p}(y) - t_{\text{mix}}^{N,p}(g_N(x)) \right| = 0.$$

Proof. Consider $p \in [1, \infty]$ such that (1.5) holds for $x \in F$, so that $t_0 := t_{\text{mix}}^p(x)$ is finite, and let $\varepsilon \in (0, t_0/2)$. Since the function $t \mapsto D_p(x, t)$ is strictly decreasing (by Lemma 3.1), there exists an $\eta > 0$ such that $D_p(x, t_0 - \varepsilon) > D_p(x, t_0) + \eta = 1/4 + \eta$ and also $D_p(x, t_0 + \varepsilon) < 1/4 - \eta$. By the continuity of $(q_t(x, y))_{x,y \in F, t > 0}$, there also exists a $\delta > 0$ such that

$$\sup_{t \in [t_0 - \varepsilon, t_0 + \varepsilon]} \sup_{\substack{y \in F: \\ d_F(x, y) < \delta}} |D_p(x, t) - D_p(y, t)| < \eta.$$

Hence if $y \in B_F(x, \delta)$, then

$$D_p(y, t_0 - \varepsilon) > D_p(x, t_0 - \varepsilon) - \eta > \frac{1}{4},$$

$$D_p(y, t_0 + \varepsilon) < D_p(x, t_0 + \varepsilon) + \eta < \frac{1}{4}.$$

This implies that $t_{\text{mix}}^p(y) \in [t_0 - \varepsilon, t_0 + \varepsilon]$, and (a) follows.

The proof of part (b) is similar. In particular, choose η as above and note that (3.4) implies that $D_p^N(g_N(x), \gamma(N)(t_0 - \varepsilon)) > 1/4 + \eta/2$ and $D_p^N(g_N(x), \gamma(N)(t_0 + \varepsilon)) < 1/4 - \eta/2$ for large N . Furthermore, by the transition density tightness of Lemma 2.3, there exists a $\delta > 0$ such that

$$\sup_{t \in [t_0 - \varepsilon, t_0 + \varepsilon]} \sup_{\substack{y \in V(G^N): \\ d_{G^N}(g_N(x), y) < \delta}} |D_p^N(g_N(x), \gamma(N)t) - D_p^N(y, \gamma(N)t)| < \frac{\eta}{2},$$

for large N . Hence if N is large and $y \in V(G^N)$ is such that $d_{G^N}(g_N(x), y) < \delta$, then $D_p^N(y, \gamma(N)(t_0 - \varepsilon)) > 1/4$, and $D_p^N(y, \gamma(N)(t_0 + \varepsilon)) < 1/4$. This implies that $\gamma(N)^{-1} t_{\text{mix}}^{N,p}(y) \in [t_0 - \varepsilon, t_0 + \varepsilon]$. Since it is trivially true that, once N is large enough, this result can be applied with $y = g_N(x)$, the result follows. \square

We are now ready to give the proof of our main result.

Proof of Theorem 1.4. Observe that, under the assumptions of the theorem, Lemma 3.4(a) implies that the function $(t_{\text{mix}}^p(x))_{x \in F}$ is continuous. Since F is compact, the supremum of $(t_{\text{mix}}^p(x))_{x \in F}$ is therefore finite. Now, it is an elementary exercise to check that we can write the L^p -mixing time of F , as defined at (1.6), in the following way:

$$t_{\text{mix}}^p(F) = \sup_{x \in F} t_{\text{mix}}^p(x). \tag{3.6}$$

Consequently $t_{\text{mix}}^p(F) \in (0, \infty)$, as desired.

To complete the proof, we are required to demonstrate the convergence statement of (1.10). Fix $\varepsilon > 0$. For every $x \in F$, Proposition 3.2 and Lemma 3.4(b) allow us to choose $\delta(x) > 0$ and $N(x) < \infty$ such that

$$\begin{aligned} \sup_{N \geq N(x)} \left| \gamma(N)^{-1} t_{\text{mix}}^{N,p}(g_N(x)) - t_{\text{mix}}^p(x) \right| &\leq \varepsilon, \\ \sup_{N \geq N(x)} \sup_{\substack{y \in V(G^N): \\ d_{G^N}(g_N(x), y) < 4\delta(x)}} \gamma(N)^{-1} \left| t_{\text{mix}}^{N,p}(g_N(x)) - t_{\text{mix}}^{N,p}(y) \right| &\leq \varepsilon. \end{aligned}$$

Since $(B_E(x, \delta(x)))_{x \in F}$ is an open cover for F , by compactness it admits a finite subcover $(B_E(x, \delta(x)))_{x \in \mathcal{X}}$. Moreover, because $d_H^E(F, V(G^N)) \rightarrow 0$, there exists an $N_0 > 0$ such that if $N \geq N_0$, then $(B_E(x, 2\delta(x)))_{x \in \mathcal{X}}$ is a cover for $V(G^N)$. Applying this choice of \mathcal{X} , we have for $N \geq N_0 \vee \max_{x \in \mathcal{X}} N(x)$ that

$$\gamma(N)^{-1} t_{\text{mix}}^p(G^N) \leq \sup_{x \in \mathcal{X}} \gamma(N)^{-1} t_{\text{mix}}^{N,p}(g_N(x)) + \varepsilon \leq \sup_{x \in \mathcal{X}} t_{\text{mix}}^p(x) + 2\varepsilon \leq t_{\text{mix}}^p(F) + 2\varepsilon,$$

where we note that, similarly to (3.6), the L^p -mixing time of the graph G^N can be written as

$$t_{\text{mix}}^p(G^N) = \sup_{x \in V(G^N)} t_{\text{mix}}^{N,p}(x).$$

Furthermore, if $x_0 \in F$ is chosen such that $t_{\text{mix}}^p(x_0) \geq t_{\text{mix}}^p(F) - \varepsilon$, then, for large N ,

$$\gamma(N)^{-1} t_{\text{mix}}^{N,p}(G^N) \geq \gamma(N)^{-1} t_{\text{mix}}^{N,p}(g_N(x_0)) \geq t_{\text{mix}}^p(x_0) - \varepsilon \geq t_{\text{mix}}^p(F) - 2\varepsilon,$$

where we have again made use of Proposition 3.2. Since $\varepsilon > 0$ was arbitrary, we are done. \square

3.2 Distinguished starting points

In certain situations, convergence of transition densities might only be known with respect to a single distinguished starting point. This is the case, for instance, in two of the most important examples we present in Section 5 – critical Galton-Watson trees and the critical Erdős-Rényi random graph. In such settings, it is only possible to prove a convergence result for the mixing time from the distinguished point. It is the purpose of this subsection to present a precise conclusion of this kind.

Consider, for a compact interval $I \subset (0, \infty)$, the space of triples of the form (F, π, q) , where $F = (F, d_F, \rho)$ is a non-empty compact metric space with distinguished vertex ρ , π is a Borel probability measure on F and $q = (q_t(x, y))_{x, y \in F, t \in I}$ is a jointly continuous real-valued function of (t, x, y) ; this is the same as the collection $\tilde{\mathcal{M}}_I$ defined in Section 2, though we have added the supposition that the metric spaces are pointed. We say two such elements, (F, π, q) and (F', π', q') , are equivalent if there exists an isometry $f : F \rightarrow F'$ such that $f(\rho) = \rho'$, $\pi \circ f^{-1} = \pi'$ and $q'_t \circ f = q_t$ for every $t \in I$. By

following the proof of Lemma 2.1, one can check that it is possible to define a metric on the equivalence classes of this relation by simply including in the definition of Δ_I the condition that the correspondence \mathcal{C} must contain (ρ, ρ') . We define convergence in a spectral pointed Gromov-Hausdorff sense to be with respect to this metric. The distinguished starting point version of Assumption 1 is then as follows.

Assumption 2. Let $(G^N)_{N \geq 1}$ be a sequence of finite connected graphs with at least two vertices and one, ρ^N say, distinguished, for which there exists a sequence $(\gamma(N))_{N \geq 1}$ such that, for any compact interval $I \subset (0, \infty)$,

$$\left((V(G^N), d_{G^N}, \rho^N), \pi^N, \left(q_{\gamma(N)t}^N(\rho^N, x) \right)_{x \in V(G^N), t \in I} \right)$$

converges to $((F, d_F, \rho), \pi, (q_t(\rho, x))_{x \in F, t \in I})$ in a spectral pointed Gromov-Hausdorff sense, where ρ is a distinguished point in F .

The following result can then be proved in an almost identical fashion to Proposition 3.2, simply replacing $g_N(x)$ by ρ^N and x by ρ . In doing this, it is useful to note that if Assumption 1 is replaced by Assumption 2, then we are able to include in the conclusions of Lemma 2.2 that ρ^N converges to ρ in E .

Theorem 3.5. Suppose that Assumption 2 is satisfied. If $p \in [1, \infty]$ is such that (1.5) holds for $x = \rho$, then

$$\gamma(N)^{-1} t_{\text{mix}}^{N,p}(\rho^N) \rightarrow t_{\text{mix}}^p(\rho).$$

4 Convergence to stationarity of the transition density

Before continuing to present example applications of the mixing time convergence results proved so far, we describe how to check the L^p convergence to stationarity of the transition density of X^F in the case when we have a spectral decomposition for it and a spectral gap. In the same setting, we will also explain how to check the non-triviality conditions on the transition density that were made in the introduction.

Write the generator of the conservative Hunt process X^F as $-\Delta$, and suppose that Δ has a compact resolvent. Then there exists a complete orthonormal basis of $L^2(F, \pi)$, $(\varphi_k)_{k \geq 1}$ say, such that $\Delta \varphi_k = \lambda_k \varphi_k$ for all $k \geq 0$, $0 \leq \lambda_0 \leq \lambda_1 \leq \dots$ and $\lim_{k \rightarrow \infty} \lambda_k = \infty$. By expanding as a Fourier series, we can consequently write the transition density of X^F as

$$\begin{aligned} q_t(x, y) &= \sum_{k \geq 0} \left(\int_F q_t(x, z) \varphi_k(z) \pi(dz) \right) \varphi_k(y) \\ &= \sum_{k \geq 0} P_t^F \varphi_k(x) \varphi_k(y) \\ &= \sum_{k \geq 0} e^{-\lambda_k t} \varphi_k(x) \varphi_k(y), \end{aligned}$$

where $(P_t^F)_{t \geq 0}$ is the associated semigroup, and the final equality holds as a simple consequence of the fact that $\frac{d}{dt}(P_t^F \varphi_k) = -P_t^F \Delta \varphi_k = -\lambda_k P_t^F \varphi_k$. Now by (1.1), it holds that $1 = P_t^F 1$ is in the domain of Δ . A standard argument thus yields $\Delta 1 = \Delta P_t^F 1 = -\frac{d}{dt}(P_t^F 1) = 0$, and so there is no loss of generality in presupposing that $\lambda_0 = 0$ and $\varphi_0 \equiv 1$ in this setting. The only additional assumption we make on the transition density $(q_t(x, y))_{x, y \in F, t > 0}$ is that it is jointly continuous in (t, x, y) (i.e. (1.2) holds).

Lemma 4.1. *Suppose that the operator Δ has a compact resolvent, so that the above spectral decomposition holds. If there is a spectral gap, i.e. $\lambda_1 > 0$, then $(q_t(x, y))_{x, y \in F, t > 0}$ converges to stationarity in an L^p sense (namely (1.5) holds) for any $p \in [1, \infty]$.*

Proof. Recall from (3.5) that $D_2(x, t)^2 = q_{2t}(x, x) - 1$. Under the assumptions of the lemma, it follows that

$$D_2(x, t)^2 = \sum_{k \geq 1} e^{-2\lambda_k t} \varphi_k(x)^2 \rightarrow 0, \tag{4.1}$$

as $t \rightarrow \infty$, which completes the proof of the result for $p = 2$. To extend this to any p , we first use Cauchy-Schwarz to deduce

$$\begin{aligned} (q_t(x, y) - 1)^2 &= \left(\sum_{k \geq 1} e^{-\lambda_k t} \varphi_k(x) \varphi_k(y) \right)^2 \\ &\leq \sum_{k \geq 1} e^{-\lambda_k t} \varphi_k(x)^2 \sum_{k \geq 1} e^{-\lambda_k t} \varphi_k(y)^2 \\ &= (q_t(x, x) - 1)(q_t(y, y) - 1). \end{aligned}$$

Consequently, we have that

$$\begin{aligned} D_\infty(x, t)^2 &= \sup_{y \in F} (q_t(x, y) - 1)^2 \\ &\leq (q_t(x, x) - 1) \sup_{y \in F} (q_t(y, y) - 1) \\ &\leq D_2(x, t/2)^2 \sup_{y \in F} D_\infty(y, 1) \end{aligned}$$

for any $t \geq 1$, where the second inequality involves an application of the monotonicity property proved as part of Lemma 3.1. Now, by (1.2), the term $\sup_{y \in F} D_\infty(y, 1)$ is a finite constant, and so combining the above bound with (4.1) implies that $D_\infty(x, t) \leq CD_2(x, t/2) \rightarrow 0$ as $t \rightarrow \infty$. The result for general $p \in [1, \infty]$ is an immediate consequence of this. \square

We now give a lemma that explains how to check conditions (1.3) and (1.4).

Lemma 4.2. *Suppose that the operator Δ has a compact resolvent and there is a spectral gap, then the conditions (1.3) and (1.4) are automatically satisfied.*

Proof. Firstly, assume that $q_t(x, y) = 0$ for some $x, y \in F$, $t > 0$. If $s \in (0, t)$, then the Chapman-Kolmogorov equations yield $0 = q_t(x, y) = \int_F q_s(x, z) q_{t-s}(z, y) \pi(dz)$. Since π has full support, using (1.2), it follows that $q_s(x, z) q_{t-s}(z, y) = 0$ for every $z \in F$. In particular, $q_s(x, y) q_{t-s}(y, y) = 0$. Noting that $q_{t-s}(y, y) = D_2^2(y, t/2) + 1 \geq 1$, we deduce that $q_s(x, y) = 0$. Now, define a function $f : (0, \infty) \rightarrow \mathbb{R}_+$ by setting $f(s) := q_s(x, y)$. Letting $(\lambda'_i)_{i \geq 0}$ represent the distinct eigenvalues of Δ , we can write

$$f(s) = \sum_{i \geq 0} a_i e^{-\lambda'_i s},$$

where $a_i := \sum_{j: \lambda_j = \lambda'_i} \varphi_j(x) \varphi_j(y)$. In fact, since Cauchy-Schwarz implies $\sum_{i \geq 0} |a_i e^{-\lambda'_i s}| \leq (q_s(x, x) q_s(y, y))^{1/2} < \infty$, this series converges absolutely whenever $s \in (0, \infty)$. Thus $f(z) := \sum_{i \geq 0} a_i e^{-\lambda'_i z}$ defines an analytic function on the whole half-plane $\Re(z) > 0$. By our previous observation regarding $q_s(x, y)$, this analytic function is equal to 0 on the set $(0, t]$, and therefore it must be 0 everywhere on $\Re(z) > 0$. However, this contradicts

the fact that $f(t) = q_t(x, y) \rightarrow 1$ as $t \rightarrow \infty$, which was proved in Lemma 4.1. Hence, $q_t(x, y) > 0$ for every $x, y \in F, t > 0$.

Secondly, suppose that $q_t(x, \cdot) \equiv 1$ for some $x \in F$ and $t > 0$. Then $1 = q_t(x, x) = 1 + \sum_{i \geq 1} \varphi_i(x)^2 e^{-\lambda_i t}$, and so $\varphi_i(x) = 0$ for every $i \geq 1$. This implies that $q_t(x, x) = 1$ for every $t > 0$. However, by following the proof of (3.1), one can deduce that

$$\lim_{t \rightarrow 0} (q_t(x, x) - 1) = \lim_{t \rightarrow 0} D_2^2(x, t/2) \geq \lim_{t \rightarrow 0} D_1^2(x, t/2) \geq 2,$$

and so the previous conclusion can not hold. Consequently, we have shown that $q_t(x, \cdot) \not\equiv 1$ for any $x \in F, t > 0$, as desired. \square

To summarize, the above results demonstrate that to verify all the conditions on the transition density that are required to apply our mixing time convergence results, it will suffice to check that the conservative Hunt process X^F has a jointly continuous transition density and the corresponding non-negative self-adjoint operator, Δ , has a compact resolvent and exhibits a spectral gap. As the following corollary explains, this is a particularly useful observation in the case that the Dirichlet form $(\mathcal{E}, \mathcal{F})$ associated with X^F is a resistance form. A precise definition of such an object appears in [27, Definition 3.1], for example, but the key property is the finiteness of the corresponding resistance, i.e.

$$R(x, y) := \sup \left\{ \frac{|f(x) - f(y)|^2}{\mathcal{E}(f, f)} : f \in \mathcal{F}, \mathcal{E}(f, f) > 0 \right\}$$

is finite for any $x, y \in F$.

Corollary 4.3. *Suppose that X^F is a π -symmetric Hunt process on F such that the associated Dirichlet form $(\mathcal{E}, \mathcal{F})$ is a resistance form, then (1.1)-(1.5) are automatically satisfied.*

Proof. The fact that X^F is conservative is clear since for a resistance form $1 \in \mathcal{F}$ and $\mathcal{E}(1, 1) = 0$. That (1.2) holds is proved in [27, Lemma 10.7]. Moreover, we can check that the non-negative operator corresponding to $(\mathcal{E}, \mathcal{F})$ has a compact resolvent (see [27, Lemma 9.7] and [30, Theorem B.1.13]) and exhibits a spectral gap (this is an easy consequence of the fact that, for a resistance form, $\mathcal{E}(f, f) = 0$ if and only if f is constant). Thus, by Lemma 4.1 and Lemma 4.2, the transition density of X^F also satisfies (1.3)-(1.5). \square

5 Examples

The mixing time results of the previous sections have many applications. To begin with a particularly simple one, consider G^N to be a discrete d -dimensional box of side-length N , i.e. vertex set $\{1, 2, \dots, N\}^d$ and nearest neighbor connections. By applying classical results about the convergence of the simple random walk on this graph to Brownian motion on $[0, 1]^d$ reflected at the boundary, Theorem 1.4 readily implies that the L^p -mixing time of the simple random walk on $\{1, 2, \dots, N\}^d$, when rescaled by N^{-2} , converges to the L^p -mixing time of the limit process for any $p \in [1, \infty]$. A similar result could be proved for the random walk on the discrete torus $(\mathbb{Z}/N\mathbb{Z})^d$. More interestingly, however, as we will now demonstrate, it is possible to apply our main results in a number of examples where the graphs, and sometimes limiting spaces, are random: self-similar fractal graphs with random weights, critical Galton-Watson trees, the critical Erdős-Rényi random graph, and the range of the random walk in high dimensions. For the second and third of these, we will in the next section go on to describe how the convergence in distribution of mixing times we establish can be applied to relate tail asymptotics for mixing time distributions of the discrete and continuous models.

5.1 Self-similar fractal graphs with random weights

Although the results we have proved apply more generally to self-similar fractal graphs (see below for some further comments on this point), to keep the presentation concise we restrict our attention here to graphs based on the classical Sierpinski gasket, the definition of which we now recall. Suppose p_1, p_2, p_3 are the vertices of an equilateral triangle in \mathbb{R}^2 . Define the similitudes

$$\psi_i(x) := p_i + \frac{z - p_i}{2}, \quad i = 1, 2, 3.$$

Since $(\psi_i)_{i=1}^3$ is a family of contraction maps, there exists a unique non-empty compact set F such that $F = \cup_{i=1}^3 \psi_i(F)$ – this is the Sierpinski gasket. We will suppose d_F is the intrinsic shortest path metric on F defined in [28], and note that this induces the same topology as the Euclidean metric. Moreover, we suppose π is the $(\ln 3)/(\ln 2)$ -Hausdorff measure on F with respect to the Euclidean metric, normalized to be a probability measure. This measure is non-atomic, has full support and satisfies $\pi(\partial B(x, r)) = 0$ for every $x \in F, r > 0$ (see [14, Lemma 25]).

We now define a sequence of graphs $(G^N)_{N \geq 0}$ by setting

$$V(G^N) := \bigcup_{i_1, \dots, i_N=1}^3 \psi_{i_1 \dots i_N}(V_0),$$

where $V_0 := \{p_1, p_2, p_3\}$ and $\psi_{i_1 \dots i_n} := \psi_{i_1} \circ \dots \circ \psi_{i_n}$, and

$$E(G^N) := \{\{\psi_{i_1 \dots i_N}(x), \psi_{i_1 \dots i_N}(y)\} : x, y \in V_0, x \neq y, i_1, \dots, i_N \in \{1, 2, 3\}\}.$$

We set $d_{G^N} := d_F|_{V(G^N) \times V(G^N)}$, so that $(V(G^N), d_{G^N})$ converges to (F, d_F) with respect to the Hausdorff distance between compact subsets of F . Weights $(\mu_e^N)_{e \in E(G^N), N \geq 0}$ will be selected independently at random from a common distribution, which we assume is supported on an interval $[c_1, c_2]$, where $0 < c_1 \leq c_2 < \infty$. By the procedure described in the introduction, we define from these weights a sequence of random measures $(\pi^N)_{N \geq 0}$ on the vertex sets of our graphs in the sequence $(G^N)_{N \geq 0}$. That π^N weakly converges to π as Borel probability measures on F , almost-surely, can be checked by applying [14, Lemma 26].

To describe the scaling limit of the random walks associated with the random weights μ^N , we appeal to the homogenization result of [31]. To describe this, we first introduce the Dirichlet form associated with the walk on the level N graph by setting, for $f \in \mathbb{R}^{V(G^N)}$,

$$\mathcal{E}^N(f, f) := \sum_{i_1, \dots, i_N=1}^3 \sum_{x, y \in V_0, x \neq y} \mu_{\psi_{i_1 \dots i_N}(x)\psi_{i_1 \dots i_N}(y)}^N (f(\psi_{i_1 \dots i_N}(x)) - f(\psi_{i_1 \dots i_N}(y)))^2. \tag{5.1}$$

Let $\Lambda^N = (\Lambda_{xy}^N)_{x, y \in V_0, x \neq y}$ be the collection of weights such that the associated random walk on G^0 is the trace of X^{G^N} onto V_0 . It then follows from [31, Theorem 3.4] that there exists a deterministic constant $C \in (0, \infty)$ such that

$$\lim_{n \rightarrow \infty} \mathbf{E} \left| \left(\frac{5}{3} \right)^N \Lambda_{xy}^N - C \right| = 0,$$

for any $x, y \in V_0, x \neq y$. Now, suppose \mathcal{E}_C^N is a quadratic form on $\mathbb{R}^{V(G^N)}$ which satisfies (5.1) with $\mu_{\psi_{i_1 \dots i_N}(x)\psi_{i_1 \dots i_N}(y)}^N$ replaced by C in each summand, then define

$$\mathcal{E}(f, f) = \lim_{N \rightarrow \infty} \left(\frac{5}{3} \right)^N \mathcal{E}_C^N(f|_{V(G^N)}, f|_{V(G^N)})$$

for $f \in \mathcal{F}$, where \mathcal{F} is the subset of $C(F, \mathbb{R})$ such that the right-hand side above exists and is finite. It is known that $(\mathcal{E}, \mathcal{F})$ is a local, regular Dirichlet form on $L^2(F, \pi)$, which is also a resistance form (see [30], for example). Thus, by Corollary 4.3, the associated π -symmetric diffusion X^F , which (modulo the scaling constant C) is known as Brownian motion on the Sierpinski gasket, satisfies (1.1)-(1.5).

For the case of unbounded fractal graphs, a probabilistic version of (2.12) was proved as [14, Proposition 30(i)] by applying the homogenization result for processes of [32] (cf. [31]). Since the Sierpinski gasket is a finitely ramified fractal, it is a relatively straightforward technical exercise to adapt this result to the compact case by considering a decomposition of the sample paths of the relevant processes into segments started at one of the outer corners of the gasket and stopped upon hitting another (a full proof of this can be found in [15]).

Finally, a probabilistic version of the tightness condition of (2.11) is easily checked by applying (a probabilistic version of) Lemma 2.5, using known resistance estimates for nested fractals (cf. [14, Proposition 30(ii)]), and so Assumption 1 holds in probability due to Proposition 2.4. Thus we are able to apply Theorem 1.4 to deduce the following.

Theorem 5.1. *If $t_{\text{mix}}(G^N)$ is the mixing time of the random walk on the level N approximation to the Sierpinski gasket equipped with uniformly bounded, independently and identically distributed random weights, then*

$$5^{-N} t_{\text{mix}}(G^N) \rightarrow t_{\text{mix}}(F)$$

in probability, where $t_{\text{mix}}(F)$ is the mixing time of the diffusion X^F .

Let us remark that the same argument will yield at least two generalizations of this theorem. Firstly, it is not necessary for the weights to be independent and identically distributed, but rather it will be sufficient for them only to be ‘cell independent’, i.e. each collection $(\mu_{\psi_{i_1 \dots i_N}(x)\psi_{i_1 \dots i_N}(y)})_{x,y \in V_0, x \neq y}$ is independent and identically distributed as $(\mu_{xy})_{x,y \in V_0, x \neq y}$. (We note that without a symmetry condition, though, the limiting diffusion will no longer be guaranteed to be the Brownian motion on the Sierpinski gasket.) Secondly, the Sierpinski gasket is just one example of a nested fractal. Identical arguments could be applied to obtain corresponding mixing time results for sequences of graphs based on any of the highly-symmetric fractals that come from this class (since the key references [14], [31] and [32] all incorporate nested fractals already).

Finally, variations on the above mixing time convergence result can also be established for examples along the lines of those appearing in [14, Sections 7.4 and 7.5]. These include: an almost-sure statement for Vicsek set-type graphs (which complements the mixing time bounds for deterministic versions of these graphs proved in [22]); a convergence of mixing times for deterministic Sierpinski carpet graphs; and a subsequential limit for Sierpinski carpets with random weights. Since many of the ideas needed for these applications are similar to those discussed above, we omit the details.

5.2 Critical Galton-Watson trees

The connection between critical Galton-Watson processes and α -stable trees is now well-known, and so we will be brief in introducing it. Let ξ be a mean 1 random variable whose distribution is aperiodic (not supported on a sub-lattice of \mathbb{Z}). Furthermore, suppose that ξ is in the domain of attraction of a stable law with index $\alpha \in (1, 2)$, by which we mean that there exists a sequence $a_N \rightarrow \infty$ such that

$$\frac{\xi[N] - N}{a_N} \rightarrow \Xi, \tag{5.2}$$

in distribution, where $\xi^{[N]}$ is the sum of N independent copies of ξ and the limit random variable satisfies $\mathbf{E}(e^{-\lambda \bar{\xi}}) = e^{-\lambda^\alpha}$ for $\lambda > 0$. If \mathcal{T}_N is a Galton-Watson tree with offspring distribution ξ conditioned to have total progeny N , then it is the case that

$$N^{-1} a_N \mathcal{T}_N \rightarrow \mathcal{T}^{(\alpha)}, \tag{5.3}$$

in distribution with respect to the Gromov-Hausdorff distance between compact metric spaces, where $\mathcal{T}^{(\alpha)}$ is an α -stable tree normalized to have total mass equal to 1 (see [34, Theorem 4.3], which is a corollary of a result originally proved in [16]). Note that the left-hand side here is shorthand for the metric space $(V(\mathcal{T}_N), N^{-1} a_N d_{\mathcal{T}_N})$, where $V(\mathcal{T}_N)$ is the vertex set of \mathcal{T}_N and $d_{\mathcal{T}_N}$ is the shortest path graph distance on this set.

The α -stable tree $\mathcal{T}^{(\alpha)}$ is almost-surely a compact metric space. Moreover, there is a natural non-atomic probability measure upon it, $\pi^{(\alpha)}$ say, which has full support, and appears as the limit of the uniform measure on the approximating graph trees. Usefully, we can decompose this measure in terms of a collection of measures of level sets of the tree. More specifically, in the construction of the α -stable tree from an excursion we can naturally choose a root $\rho \in \mathcal{T}^{(\alpha)}$. We define $\mathcal{T}^{(\alpha)}(r) := \{x \in \mathcal{T}^{(\alpha)} : d_{\mathcal{T}^{(\alpha)}}(\rho, x) = r\}$ to be the collection of vertices at height r above this vertex. For almost-every realization of $\mathcal{T}^{(\alpha)}$, there then exists a càdlàg sequence of finite measures on $\mathcal{T}^{(\alpha)}$, $(\ell^r)_{r>0}$, such that ℓ^r is supported on $\mathcal{T}^{(\alpha)}(r)$ for each r and $\pi^{(\alpha)} = \int_0^\infty \ell^r dr$ (see [17, Section 4.2]). Clearly this implies that $\pi^{(\alpha)}(\partial B_{\mathcal{T}^{(\alpha)}}(\rho, r)) = 0$ for every $r > 0$, for almost-every realization of $\mathcal{T}^{(\alpha)}$. Since α -stable trees satisfy a root-invariance property (see [17, Theorem 4.8]), one can easily extend this result to hold for $\pi^{(\alpha)}$ -a.e. $x \in \mathcal{T}^{(\alpha)}$. Although this is not quite the assumption of the introduction that $\pi^{(\alpha)}(\partial B_{\mathcal{T}^{(\alpha)}}(x, r)) = 0$ for every $x \in \mathcal{T}^{(\alpha)}$, $r > 0$, by a minor tweak of the proof of Proposition 3.2, we are still able to apply our mixing time convergence results in the same way.

Upon almost-every realization of the metric measure space $(\mathcal{T}^{(\alpha)}, \pi^{(\alpha)})$, it is possible to define a corresponding Brownian motion $X^{(\alpha)}$ (to do this, apply [29, Theorem 5.4], in the way described in [10, Section 2.2]). This is a conservative $\pi^{(\alpha)}$ -symmetric Hunt process, and the associated Dirichlet form $(\mathcal{E}^{(\alpha)}, \mathcal{F}^{(\alpha)})$ is actually a resistance form. Thus we can again apply Corollary 4.3 to confirm that (1.1)-(1.5) hold for some corresponding transition density, $q^{(\alpha)}$ say. Now, in [13], it was demonstrated that if $\mathbf{P}_{\rho_N}^{\mathcal{T}_N}$ is the law of the random walk on \mathcal{T}_N started from its root (original ancestor) ρ_N and π^N is its stationary probability measure, then, after embedding all the objects into an underlying Banach space in a suitably nice way, the conclusion of (5.3) can be extended to the distributional convergence of

$$\left(N^{-1} a_N \mathcal{T}_N, \pi^N(N a_N^{-1} \cdot), \mathbf{P}_{\rho_N}^{\mathcal{T}_N} \left(\left(N^{-1} a_N X_{[N^2 a_N^{-1} t]}^{\mathcal{T}_N} \right)_{t \in [0,1]} \in \cdot \right) \right)$$

to $(\mathcal{T}^{(\alpha)}, \pi^{(\alpha)}, \mathbf{P}_\rho^{(\alpha)})$, where $\mathbf{P}_\rho^{(\alpha)}$ is the law of $X^{(\alpha)}$ started from ρ . By applying the fixed starting point version of the local limit result of Proposition 2.4 (cf. [14, Theorem 1]), similarly to the argument of [14, Section 7.2], for the Brownian continuum random tree, which corresponds to the case $\alpha = 2$, one can obtain from this a distributional version of Assumption 2. (The tightness condition of (2.11) is easily checked by applying Lemma 2.5.)

Lemma 5.2. *For any compact interval $I \subset (0, \infty)$,*

$$\left((V(\mathcal{T}_N), N^{-1} a_N d_{\mathcal{T}_N}, \rho^N), \pi^N, \left(q_{N^2 a_N^{-1} t}^N(\rho^N, x) \right)_{x \in V(\mathcal{T}_N), t \in I} \right)$$

converges in distribution to $((\mathcal{T}^{(\alpha)}, d_{\mathcal{T}^{(\alpha)}}, \rho), \pi^{(\alpha)}, (q_t^{(\alpha)}(\rho, x))_{x \in \mathcal{T}^{(\alpha)}, t \in I})$ in a spectral pointed Gromov-Hausdorff sense.

Consequently, since the space in which the above convergence in distribution occurs is separable, we can use a Skorohod coupling argument to deduce from this and Theorem 3.5 the following mixing time convergence result. We remark that the $\sqrt{2}$ that appears in the finite variance result is simply an artefact of the particular scaling we have described here, and could alternatively have been absorbed in the scaling of metrics.

Theorem 5.3. *Fix $p \in [1, \infty]$. If $t_{\text{mix}}^p(\rho^N)$ is the L^p -mixing time of the random walk on \mathcal{T}_N started from its root ρ^N , then*

$$N^{-2} a_N t_{\text{mix}}^p(\rho^N) \rightarrow t_{\text{mix}}^p(\rho),$$

in distribution, where $t_{\text{mix}}^p(\rho) \in (0, \infty)$ is the L^p -mixing time of the Brownian motion on $\mathcal{T}^{(\alpha)}$ started from ρ . In particular, in the case when the offspring distribution has finite variance σ , it is the case that

$$\frac{\sigma}{\sqrt{2}} N^{-3/2} t_{\text{mix}}^p(\rho^N) \rightarrow t_{\text{mix}}^p(\rho),$$

in distribution.

Remark 5.4. We note that it was only for convenience that the convergence of the random walks on the trees \mathcal{T}_N , $N \geq 1$, to the Brownian motion on $\mathcal{T}^{(\alpha)}$ was proved from a single starting point in [13]. We do not anticipate any significant problems in extending this result to hold for arbitrary starting points. Indeed, the first step would be to make the obvious adaptations to the proof of [13, Lemma 4.2] to extend the result, which demonstrates convergence of simple random walks (and related additive functionals) on subtrees of \mathcal{T}_N consisting of a finite number of branch segments to the corresponding continuous objects, from the case when all the random walks start from the root to an arbitrary starting point version. An argument identical to the remainder of [13, Section 4] could then be used to obtain the convergence of simple random walks on the whole trees, at least in the case when the starting point of the diffusion is in one of the finite subtrees considered. Since the union of the finite subtrees is dense in the limiting space, we could subsequently use the heat kernel continuity properties to obtain the non-pointed spectral Gromov-Hausdorff version of Lemma 5.2. However, we do not pursue this approach here as it would require a substantial amount of space and new notation that is not relevant to the main ideas of this article. Were it to be checked, Theorem 1.4 would imply, for any $p \in [1, \infty]$, the distributional convergence of $t_{\text{mix}}^p(\mathcal{T}_N)$, the L^p -mixing time of the random walk on \mathcal{T}_N , when rescaled appropriately, to $t_{\text{mix}}^p(\mathcal{T}^{(\alpha)}) \in (0, \infty)$, the L^p -mixing time of the Brownian motion on $\mathcal{T}^{(\alpha)}$.

5.3 Critical Erdős-Rényi random graph

Closely related to the random trees of the previous section is the Erdős-Rényi random graph at criticality. In particular, let $G(N, p)$ be the random graph in which every edge of the complete graph on N labeled vertices $\{1, \dots, N\}$ is present with probability p independently of the other edges. Supposing $p = N^{-1} + \lambda N^{-4/3}$ for some $\lambda \in \mathbb{R}$, so that we are in the so-called critical window, it is known that the largest connected component \mathcal{C}^N , equipped with its shortest path graph metric $d_{\mathcal{C}^N}$, satisfies

$$\left(V(\mathcal{C}^N), N^{-1/3} d_{\mathcal{C}^N} \right) \rightarrow (\mathcal{M}, d_{\mathcal{M}})$$

in distribution, again with respect to the Gromov-Hausdorff distance between compact metric spaces, where $(\mathcal{M}, d_{\mathcal{M}})$ is a random compact metric space [1]. (In fact, this

and all the results given in this subsection hold for a family of i -th largest connected components for all $i \in \mathbb{N}$. For notational simplicity, we only discuss the largest connected component \mathcal{C}^N .) Moreover, in [9], it was shown that the associated random walks started from a root vertex ρ^N satisfy a distributional convergence result of the form

$$\left(N^{-1/3} X_{[Nt]}^{\mathcal{C}^N} \right)_{t \geq 0} \rightarrow (X_t^{\mathcal{M}})_{t \geq 0},$$

where $X^{\mathcal{M}}$ is a diffusion on the space \mathcal{M} started from a distinguished vertex $\rho \in \mathcal{M}$. Although the invariant probability measures of the random walks, π^N say, were not considered in [9], it is not difficult to extend this result to include them since the hard work regarding their convergence has already been completed (see [9, Lemma 6.3], in particular). Hence, by again applying the fixed starting point version of the local limit result of Proposition 2.4 (using Lemma 2.5 again to deduce the relevant tightness condition), we are able to obtain the analogue of Lemma 5.2 in this setting.

Lemma 5.5. *For any compact interval $I \subset (0, \infty)$,*

$$\left(\left(V(\mathcal{C}^N), N^{-1/3} d_{\mathcal{C}^N}, \rho^N \right), \pi^N, (q_{Nt}^N(\rho^N, x))_{x \in V(\mathcal{T}_N), t \in I} \right),$$

converges in distribution to $((\mathcal{M}, d_{\mathcal{M}}, \rho), \pi^{\mathcal{M}}, (q_t^{\mathcal{M}}(\rho, x))_{x \in \mathcal{M}, t \in I})$, where $\pi^{\mathcal{M}}$ is the invariant probability measure of $X^{\mathcal{M}}$ and $(q_t^{\mathcal{M}}(x, y))_{x, y \in \mathcal{M}, t > 0}$ is its transition density with respect to this measure, in a spectral pointed Gromov-Hausdorff sense.

In order to proceed as above, we must of course check that $\pi^{\mathcal{M}}$ and $q^{\mathcal{M}}$ satisfy a number of technical conditions. To do this, first observe that a typical realization of \mathcal{M} looks like a (rescaled) typical realization of the Brownian continuum random tree $\mathcal{T}^{(2)}$ glued together at a finite number of pairs of points [1]. Since $\pi^{\mathcal{M}}$ can be considered as the image of the canonical measure $\pi^{(2)}$ on $\mathcal{T}^{(2)}$ under this gluing map, it is elementary to obtain from the statements of the previous section regarding $\pi^{(2)}$ that $\pi^{\mathcal{M}}$ is almost-surely non-atomic, has full support and satisfies $\pi^{\mathcal{M}}(\partial B_{\mathcal{M}}(x, r)) = 0$ for $\pi^{\mathcal{M}}$ -a.e. $x \in \mathcal{M}$ and every $r > 0$, as desired. For $q^{\mathcal{M}}$, we simply observe that because the Dirichlet form corresponding to $X^{\mathcal{M}}$ is a resistance form ([9, Proposition 2.1]), we can once again apply Corollary 4.3 to establish conditions (1.1)-(1.5).

Given these results, pointwise mixing time convergence follows from Theorem 3.5.

Theorem 5.6. *Fix $p \in [1, \infty]$. If $t_{\text{mix}}^p(\rho^N)$ is the L^p -mixing time of the random walk on \mathcal{C}^N started from its root ρ^N , then*

$$N^{-1} t_{\text{mix}}^p(\rho^N) \rightarrow t_{\text{mix}}^p(\rho),$$

in distribution, where $t_{\text{mix}}^p(\rho) \in (0, \infty)$ is the L^p -mixing time of the Brownian motion on \mathcal{M} started from ρ .

Remark 5.7. As discussed in Remark 5.4, we do not expect any major barriers in extending the above result to arbitrary starting points. The first task in doing this would be to adapt the convergence result proved in [9] regarding the convergence of simple random walks on subgraphs of \mathcal{C}_1^n formed of a finite number of line segments ([9, Lemma 6.4]) to arbitrary starting points. One could then extend this to obtain the desired convergence result for simple random walks on the entire space using ideas from [9, Section 7] and heat kernel continuity. It would also be necessary to introduce a new Gromov-Hausdorff-type topology to state the result, as the one used in [9] is only suitable for the pointed case. Again, we suspect taking these steps will simply be a

lengthy technical exercise, and choose not to follow them through here. We do though reasonably expect that $t_{\text{mix}}^p(\mathcal{C}^N)$, the L^p -mixing time of the random walk on \mathcal{C}^N , when rescaled appropriately, converges in distribution to $t_{\text{mix}}^p(\mathcal{M}) \in (0, \infty)$, the L^p -mixing time of the Brownian motion on \mathcal{M} , for any $p \in [1, \infty]$.

5.4 Random walk on range of random walk in high dimensions

Let $S = (S_n)_{n \geq 0}$ be the simple random walk on \mathbb{Z}^d started from 0, built on an underlying probability space with probability measure \mathbf{P} , and define the range of S up to time N to be the graph G^N with vertex set

$$V(G^N) := \{S_n : 0 \leq n \leq N\}, \tag{5.4}$$

and edge set

$$E(G^N) := \{\{S_{n-1}, S_n\} : 1 \leq n \leq N\}. \tag{5.5}$$

In this section, we will explain how to prove that if $d \geq 5$, which is an assumption henceforth, then the mixing times of the sequence of graphs $(G^N)_{N \geq 1}$ grows asymptotically as cN^2 , \mathbf{P} -a.s., where c is a deterministic constant. Since doing this primarily depends on making relatively simple adaptations of the high-dimensional scaling limit result of [12] for the random walk on the entire range of S (i.e. the $N = \infty$ case) to the finite length setting, we will be brief with the details.

First, suppose that $S = (S_n)_{n \in \mathbb{Z}}$ is a two-sided extension of $(S_n)_{n \geq 0}$ such that $(S_{-n})_{n \geq 0}$ is an independent copy of $(S_n)_{n \geq 0}$. The set of cut-times for this process,

$$\mathcal{T} := \{n : S_{(-\infty, n]} \cap S_{[n+1, \infty)} = \emptyset\},$$

is known to be infinite \mathbf{P} -a.s. ([18]). Thus we can write $\mathcal{T} = \{T_n : n \in \mathbb{Z}\}$, where $\dots T_{-1} < T_0 \leq 0 < T_1 < T_2 < \dots$. The corresponding set of cut-points is given by $\mathcal{C} := \{C_n : n \in \mathbb{Z}\}$, where $C_n := S_{T_n}$. For these objects, an ergodicity argument can be applied to obtain that, \mathbf{P} -a.s., as $|n| \rightarrow \infty$,

$$\frac{T_n}{n} \rightarrow \tau(d) := \mathbf{E}(T_1 | 0 \in \mathcal{T}) \in [1, \infty), \tag{5.6}$$

$$\frac{d_G(0, C_n)}{|n|} \rightarrow \delta(d) := \mathbf{E}(d_G(0, C_1) | 0 \in \mathcal{T}) \in [1, \infty),$$

where d_G is the shortest path graph distance on the range G of the entire two-sided walk S , which is defined analogously to (5.4) and (5.5). In particular, see [12, Lemma 2.2], for a proof of the same convergence statements under the measure $\mathbf{P}(\cdot | 0 \in \mathcal{T})$, and note that the conditioning can be removed by using the relationship between \mathbf{P} and $\mathbf{P}(\cdot | 0 \in \mathcal{T})$ described in [12, Lemma 2.1]. Given these results, it is an elementary exercise to check that the metric space $(V(G^N), \tau(d)\delta(d)^{-1}N^{-1}d_{G^N})$, where d_{G^N} is the shortest path graph distance on G^N , converges \mathbf{P} -a.s. with respect to the Gromov-Hausdorff distance to the interval $[0, 1]$ equipped with the Euclidean metric. Moreover, the same ideas readily yield an extension of this result to a spectral Gromov-Hausdorff one including that π^N , the invariant measure of the associated simple random walk, converges to Lebesgue measure on $[0, 1]$.

Now, for a fixed realization of G , let $X = (X_n)_{n \geq 0}$ be the simple random walk on G started from 0. Define the hitting times by X of the set of cut-points \mathcal{C} by $H_0 := \min\{m \geq 0 : X_m \in \mathcal{C}\}$, and, for $n \geq 1$, $H_n := \min\{m > H_{n-1} : X_m \in \mathcal{C}\}$. We use these times to define a useful indexing process $Z = (Z_n)_{n \geq 0}$ taking values in \mathbb{Z} . In particular, if $n < H_0$, define Z_n to be the unique $k \in \mathbb{Z}$ such that $X_{H_0} = C_k$. Similarly, if $n \in [H_{m-1}, H_m)$ for some $m \geq 1$, then define Z_n to be the unique $k \in \mathbb{Z}$ such that $X_{H_m} = C_k$. Noting that

this definition precisely coincides with the definition of Z in [12], from Lemma 3.5 of that article we have that: for \mathbf{P} - a.e. realization of G ,

$$(N^{-1}\tau(d)Z_{\lfloor tN^2 \rfloor})_{t \geq 0} \rightarrow (B_{t\kappa_2(d)})_{t \geq 0}, \tag{5.7}$$

in distribution, where $(B_t)_{t \geq 0}$ is a standard Brownian motion on \mathbb{R} started from 0, and $\kappa_2(d) \in (0, \infty)$ is the deterministic constant defined in [12]. To deduce from (5.7) the following scaling limit for X^N , the simple random walk on G^N , we proceed via a time-change argument that is essentially a reworking of parts of [12, Section 3].

Lemma 5.8. *For \mathbf{P} -a.e. realization of S , if X^N is started from 0, then*

$$\left(\tau(d)\delta(d)^{-1}N^{-1}d_{G^N}\left(0, X_{\lfloor \kappa_2(d)^{-1}N^2t \rfloor}^N\right)\right)_{t \geq 0} \rightarrow \left(B_t^{[0,1]}\right)_{t \geq 0},$$

in distribution, where $B^{[0,1]} = (B_t^{[0,1]})_{t \geq 0}$ is Brownian motion on $[0, 1]$ started at 0 and reflected at the boundary.

Proof. The following proof can be applied to any typical realization of S . To begin with, define a process $(A_n^{Z,N})_{n \geq 0}$ by setting

$$A_n^{Z,N} := \sum_{m=0}^{n-1} \mathbf{1}_{\{Z_m \in [0, T_N^{-1}]\}},$$

where $T_N^{-1} := \max\{n : T_n \leq N\}$. From (5.6), we have that $T_N^{-1} \sim \tau(d)^{-1}N$. Combining this observation with (5.7), one can check that, simultaneously with (5.7), $(N^{-2}A_{\lfloor tN^2 \rfloor}^N)_{t \geq 0}$ converges in distribution to $(\kappa_2(d)^{-1}A_{t\kappa_2(d)}^B)_{t \geq 0}$, where

$$A_t^B := \int_0^t \mathbf{1}_{\{B_s \in [0,1]\}} ds$$

(cf. [12, Lemma 3.5]).

We now apply the above result to establish a scaling limit for the process X observed on the vertex set $V(\tilde{G}^N) := \{S_n : T_1 \leq n \leq T_N^{-1}\}$. Specifically, set

$$A_n^N := \sum_{m=0}^{n-1} \mathbf{1}_{\{X_m, X_{m+1} \in V(\tilde{G}^N)\}}.$$

Similarly to the proof of [12, Lemma 3.6], one can check that

$$\sup_{0 \leq m \leq n} |A_m^N - A_m^{Z,N}| \leq \sum_{m=0}^n \mathbf{1}_{\{Z_m \in [0,1,2] \cup [T_N^{-1}-2, T_N^{-1}-1, T_N^{-1}]\}}.$$

It is therefore a simple consequence of (5.7) that $N^{-2} \sup_{0 \leq m \leq TN^2} |A_m^N - A_m^{Z,N}|$ converges to 0 in probability as $N \rightarrow \infty$ for any $T \in (0, \infty)$. Since we know from equation (16) of [12] that

$$N^{-1} \sup_{0 \leq m \leq TN^2} |d_G(0, X_m) - \delta(d)Z_m|$$

also converges to 0 in probability, we readily obtain

$$\left(\tau(d)\delta(d)^{-1}N^{-1}d_G\left(0, \tilde{X}_{\lfloor N^2t \rfloor}^N\right)\right)_{t \geq 0} \rightarrow \left(B_{\kappa_2(d)t}^{[0,1]}\right)_{t \geq 0}, \tag{5.8}$$

in distribution, where $\tilde{X}^N = (\tilde{X}_n^N)_{n \geq 0}$ is the random walk X observed on $V(\tilde{G}^N)$ – this is defined precisely by setting $\tilde{X}_n^N := X_{\alpha^N(n)}$, where $\alpha^N(n) := \max\{A_m^N \leq n\}$. We

remark that the particular limit process $B^{[0,1]}$ arises as a consequence of the fact that $(B_{\alpha^B}(t))_{t \geq 0}$, where α^B is the right-continuous inverse of A^B , has exactly the distribution of $B^{[0,1]}$.

Finally, since the process \tilde{X}^N is identical in law to the simple random walk X^N observed on $V(\tilde{G}^N)$, to replace \tilde{X}^N by X^N in (5.8) it will suffice to check that X^N spends only an asymptotically negligible amount of time in $V(G^N) \setminus V(\tilde{G}^N)$. Since doing this requires only a simple adaptation of the proof of [12, Lemma 3.8], we omit the details. To complete the proof, one then needs to replace d_G by d_{G^N} , but this is straightforward since

$$N^{-1} \sup_{0 \leq n \leq N} |d_G(0, S_n) - d_{G^N}(0, S_n)| \leq N^{-1} (T_1 + T_{T_N^{-1}+1} - T_{T_N^{-1}}) \rightarrow 0,$$

as $N \rightarrow \infty$. □

Although the previous lemma only contains a convergence statement for the random walks started from the particular vertex 0, there is no difficulty in extending this to the case when X^N is started from a point $x_0^N \in V(G^N)$ such that $d_{G^N}(0, x_0^N) \sim \tau(d)^{-1} \delta(d) N x_0$, and $B^{[0,1]}$ is started from $x_0 \in [0, 1]$. Applying the local limit result of Proposition 2.4 (to establish (2.11), we once again appeal to Lemma 2.5), we are able to deduce from this that Assumption 1 holds for \mathbf{P} -a.e. realization of the original random walk.

Lemma 5.9. *For \mathbf{P} -a.e. realization of S , if $I \subset (0, \infty)$ is a compact interval, then*

$$\left((V(G^N), \tau(d) \delta(d)^{-1} N^{-1} d_{G^N}), \pi^N, \left(q_{\kappa_2(d)^{-1} N^2 t}^N(x, y) \right)_{x, y \in V(G^N), t \in I} \right),$$

converges in $(\mathcal{M}_I, \Delta_I)$ to the triple consisting of: $[0, 1]$ equipped with the Euclidean metric, Lebesgue measure on this set and the transition density of Brownian motion on $[0, 1]$ reflected at the boundary.

Since it is clear that (1.1)-(1.5) hold in this case, we can therefore apply Theorem 1.4 to obtain the desired convergence of mixing times.

Theorem 5.10. *Fix $p \in [1, \infty]$. If $t_{\text{mix}}^p(S_{[0, N]})$ is the L^p -mixing time of the simple random walk on the range of S up to time N , then \mathbf{P} -a.s.,*

$$\kappa_2(d) N^{-2} t_{\text{mix}}^p(S_{[0, N]}) \rightarrow t_{\text{mix}}^p([0, 1]),$$

where $t_{\text{mix}}^p([0, 1])$ is the L^p -mixing time of the Brownian motion on $[0, 1]$ reflected at the boundary.

6 Mixing time tail estimates

In this section, we give some sufficient conditions for deriving upper and lower estimates for mixing times of random walks on finite graphs, primarily using techniques adapted from [36]. We will also discuss how to apply these general estimates to concrete random graphs (see Section 6.3). In order to shorten the paper, we skip most of the proofs, which can be found in the extended version of the paper that appears on arXiv [15].

As will be illustrated by our examples, the results in this section are robust and convenient for obtaining mixing time tail estimates. Moreover, when the convergence of mixing times (as in Theorem 1.4) is available for a sequence of graphs, we highlight how, by first deriving estimates for the relevant continuous mixing time distribution

(where similar techniques are sometimes applicable, see Remark 6.3), it can be possible to deduce results regarding the asymptotic tail behavior of random graph mixing times that are difficult to obtain directly (see the proof of Proposition 6.6 or Remark 6.9, for example).

We start by fixing our notation. Let $G = (V(G), E(G))$ be a finite connected graph and μ^G be a weight function, as in the introduction. Suppose here that d_G is the shortest path metric on the graph G , and denote, for a distinguished vertex $\rho \in V(G)$,

$$B(R) = \{y : d_G(\rho, y) < R\}, \quad V(R) := \sum_{x \in B(R)} \sum_{y: y \sim x} \mu_{xy}^G = \pi^G(B(R))\mu(G), \quad R \in (0, \infty),$$

where we write $x \sim y$ if $\mu_{xy}^G > 0$ and set $\mu(G) := \sum_{x, y \in V(G)} \mu_{xy}^G$. For the Markov chain X^G , let

$$\tau_R = \tau_{B(\rho, R)} = \min\{n \geq 0 : X_n^G \notin B(R)\}.$$

We define a quadratic form \mathcal{E} by

$$\mathcal{E}(f, g) = \frac{1}{2} \sum_{\substack{x, y \in V(G) \\ x \sim y}} \mu_{xy}^G (f(x) - f(y))(g(x) - g(y)),$$

and let $H^2 = \{f \in \mathbb{R}^{V(G)} : \mathcal{E}(f, f) < \infty\}$. For disjoint subsets A, B of G , the effective resistance between them is then given by:

$$R_{\text{eff}}(A, B)^{-1} = \inf\{\mathcal{E}(f, f) : f \in H^2, f|_A = 1, f|_B = 0\}. \quad (6.1)$$

If we further define $R_{\text{eff}}(x, y) = R_{\text{eff}}(\{x\}, \{y\})$, and $R_{\text{eff}}(x, x) = 0$, then one can check that $R_{\text{eff}}(\cdot, \cdot)$ is a metric on $V(G)$ (see [30, Section 2.3]). We will call this the resistance metric. The resistance metric enjoys the following important (but easy to deduce) estimate,

$$|f(x) - f(y)|^2 \leq R_{\text{eff}}(x, y)\mathcal{E}(f, f), \quad \forall f \in L^2(G, \mu^G).$$

Moreover, it is easy to verify that if $c_1^{-1} := \inf_{x, y \in G: x \sim y} \mu_{xy}^G > 0$, then

$$R_{\text{eff}}(x, y) \leq c_1 d_G(x, y) \quad \forall x, y \in G. \quad (6.2)$$

Let $v, r : \{0, 1, \dots, \text{diam}_{d_G}(G) + 1\} \rightarrow [0, \infty)$ be strictly increasing functions with $v(0) = r(0) = 0$, $v(1) = r(1) = 1$, which satisfy

$$C_1^{-1} \left(\frac{R}{R'}\right)^{d_1} \leq \frac{v(R)}{v(R')} \leq C_1 \left(\frac{R}{R'}\right)^{d_2}, \quad C_2^{-1} \left(\frac{R}{R'}\right)^{\alpha_1} \leq \frac{r(R)}{r(R')} \leq C_2 \left(\frac{R}{R'}\right)^{\alpha_2} \quad (6.3)$$

for all $0 < R' \leq R \leq \text{diam}_{d_G}(G) + 1$, where $C_1, C_2 \geq 1$, $1 \leq d_1 \leq d_2$ and $0 < \alpha_1 \leq \alpha_2 \leq 1$. In what follows, $v(\cdot)$ will give the volume growth order and $r(\cdot)$ the resistance growth order. For convenience, we extend them to functions on $[0, \text{diam}_{d_G}(G) + 1]$ by linear interpolation. For the rest of the paper, C_1, C_2, d_1, d_2 and α_1, α_2 stand for the constants given in (6.3).

6.1 General upper and lower bounds

In this subsection, we give general upper and lower bounds for mixing times. Note that, since $t_{\text{mix}}^p(\rho) \leq t_{\text{mix}}^p(G)$ and $t_{\text{mix}}^p(G) \leq t_{\text{mix}}^{p'}(G)$ for $p \leq p'$, it will be enough to estimate $t_{\text{mix}}^\infty(G)$ for the upper bound¹ and $t_{\text{mix}}^1(\rho)$ for the lower bound.

Upper bound: We first give an upper bound of the mixing times that is a reworking of [36, Corollary 4.2], in our setting.

¹In fact, for the upper bound it is enough to estimate $t_{\text{mix}}^2(G)$. Indeed, the Cauchy-Schwarz inequality and (3.5) implies the following known fact for mixing times of symmetric Markov chains; $t_{\text{mix}}^\infty(G) \leq 2 t_{\text{mix}}^2(G; 1/2)$, where $t_{\text{mix}}^2(G; 1/2)$ is the L^2 -mixing time of G with $1/2$ instead of $1/4$ in the definition (1.8).

Lemma 6.1. For any weighted graph (G, μ^G) ,

$$t_{\text{mix}}^\infty(G) \leq 4 \text{diam}_R(G) \mu(G),$$

where $\text{diam}_R(G)$ is the diameter of G with respect to the resistance metric R_{eff} .

Lower bound: We next give the mixing time lower bound. Let $\lambda \geq 1, H_0, \dots, H_3 > 0$, and let $C_3 := 2^{-2/\alpha_1} C_2^{-1/\alpha_2}$ where C_2 is the constant in (6.3). We give the following two conditions concerning the volume and resistance growth.

$$R_{\text{eff}}(\rho, y) \leq \lambda^{H_0} r(d_G(\rho, y)), \quad \forall y \in B(R), \quad \text{and} \quad V(R) \leq \lambda^{H_1} v(R), \quad (6.4)$$

$$R_{\text{eff}}(\rho, B(R)^c) \geq \lambda^{-H_2} r(R) \quad \text{and} \quad V(C_3 \lambda^{-(H_0+H_2)/\alpha_1} R) \geq \lambda^{-H_3} v(C_3 \lambda^{-(H_0+H_2)/\alpha_1} R). \quad (6.5)$$

Proposition 6.2. *i) For $\lambda, R > 1$, assume that $\mu(G) \geq 4V(R)$, and that (6.4), (6.5) hold for R , then*

$$t_{\text{mix}}^1(G) > C_4 \lambda^{-H'_2 - H_3} v(R) r(R), \quad (6.6)$$

where $H'_2 = H_2 + (H_0 + H_2)d_2/\alpha_1$.

ii) For $\lambda, R > 1$, assume that $\mu(G) \geq 4V(R)$, and (6.4), (6.5) hold for R and $\varepsilon_0(\lambda)R$, where $\varepsilon_0(\lambda) := c_1 \lambda^{-(H_0 + \sum_{i=0}^3 H_i + H'_2)/\alpha_1}$ for some $c_1 > 0$ small enough. Then

$$t_{\text{mix}}^1(\rho) > C_4 \lambda^{-H'_2 - H_3} v(\varepsilon_0(\lambda)R) r(\varepsilon_0(\lambda)R).$$

Remark 6.3. Essentially the same argument can be applied to deduce the corresponding mixing time upper and lower bounds in the continuous setting when we suppose that we have a process whose Dirichlet form is a resistance form (see [15], Remark A.1, for the upper bound in particular).

6.2 Random graph case

We now consider a probability space $(\Omega, \mathcal{F}, \mathbf{P})$ carrying a family of random weighted graphs $G^N(\omega) = (V(G^N(\omega)), E(G^N(\omega)), \mu^{N(\omega)}; \omega \in \Omega)$. We assume that, for each $N \in \mathbb{N}$ and $\omega \in \Omega$, $G^N(\omega)$ is a finite, connected graph containing a marked vertex ρ^N , and $\#V(G^N(\omega)) \leq M_N$ for some non-random constant $M_N < \infty$. (Here, for a set A , $\#A$ is the number of elements in A .) Let $d_{G^N(\omega)}(\cdot, \cdot)$ be a graph distance, $B(R) := B_\omega(\rho^N, R)$, and $V(R) := V_\omega(\rho^N, R)$. We write $X = (X_n, n \geq 0, P_\omega^x, x \in G^N(\omega))$ for the random walk on $G^N(\omega)$, and denote by $p_n^\omega(x, y)$ its transition density with respect to π^ω . Furthermore, we introduce a strictly increasing function $h : \mathbb{N} \cup \{0\} \rightarrow [0, \infty)$ with $h(0) = 0$, which will roughly describe the diameter of G^N with respect to the graph distance. We then set $\gamma(\cdot) = v(h(\cdot)) \cdot r(h(\cdot))$. Finally, for $i = 1, 2$, we suppose $p_i : [1, \infty) \rightarrow [0, 1]$ are functions such that $\lim_{\lambda \rightarrow \infty} p_i(\lambda) = 0$. We then have the following. (Note that C_2, d_2 in the statement are the constant in (6.3).)

Proposition 6.4. (1) Suppose that the following holds:

$$\mathbf{P}(\text{diam}_R(G^N) \geq \lambda r(h(N))) \leq p_1(\lambda), \quad \mathbf{P}(\mu^N(G^N) \geq \lambda v(h(N))) \leq p_2(\lambda), \quad (6.7)$$

then

$$\mathbf{P}(t_{\text{mix}}^\infty(G^N) \geq \lambda \gamma(N)) \leq \inf_{\theta \in [0, 1]} (p_1(\lambda^\theta/8) + p_2(\lambda^{1-\theta})).$$

(2) Suppose there exist $c_1 \leq 1$ and $J \geq (1 + H_1)/d_2$ such that the following holds:

$$\mathbf{P}((6.4) \wedge (6.5) \text{ for } R = c_1 \lambda^{-J} h(N)) \geq 1 - p_1(\lambda), \quad \mathbf{P}(\mu^N(G^N) < \lambda^{-1} v(h(N))) \leq p_2(\lambda),$$

then there exist $c_2, p_0 > 0$ such that

$$\mathbf{P}(t_{\text{mix}}^1(G^N) \leq c_2 \lambda^{-p_0} \gamma(N)) \leq 2p_1(\lambda) + p_2(\lambda/(4C_1 c_1^{d_2})).$$

(3) Suppose there exist $c_1 \leq 1$ and $J \geq (1 + H_1)/d_2$ such that the following holds:

$$\begin{aligned} &\mathbf{P}((6.4) \wedge (6.5) \text{ for } R = c_1 \lambda^{-J} h(N) \text{ and for } \varepsilon_0(\lambda)R \geq 1 - p_1(\lambda), \\ &\mathbf{P}(\mu^N(G^N) < \lambda^{-1} v(h(N))) \leq p_2(\lambda), \end{aligned} \tag{6.8}$$

where $\varepsilon_0(\lambda)$ is as in Proposition 6.2 ii), then there exist $c_2, p_0 > 0$ such that

$$\mathbf{P}(t_{\text{mix}}^1(\rho^N) \leq c_2 \lambda^{-p_0} \gamma(N)) \leq 2p_1(\lambda) + p_2(\lambda/(4C_1 c_1^{d_2})).$$

To illustrate this result, we consider the case when the random graphs $G^N(\omega)$ are obtained as components of percolation processes on finite graphs, thereby recovering [36, Theorem 1.2(c)]. (In [36], it was actually the lazy random walk was considered to avoid parity concerns, but the same techniques apply when we consider $q_m^G(\cdot, \cdot)$ as in (1.7) instead.)

Proposition 6.5. *Let \hat{G}^N be a graph with N vertices and with the maximum degree $d \in [3, N - 1]$. Let \mathcal{C}^N be the largest component of the percolation subgraph of \hat{G}^N for $0 < p < 1$. Let $p \leq \frac{1+\lambda n^{-1/3}}{d-1}$ for some fixed $\lambda \in \mathbb{R}$, and assume that there exist $c_1, \theta_1 \in (0, \infty)$ and $K_1 \in \mathbb{N}$ such that*

$$\mathbf{P}(\#\mathcal{C}^N \leq A^{-1} N^{2/3}) \leq c_1 A^{-\theta_1}, \quad \forall A, N \geq K_1, \tag{6.9}$$

then there exist $c_2, \theta_2 \in (0, \infty)$ and $K_2 \in \mathbb{N}$ such that, for all $p \in [1, \infty]$,

$$\mathbf{P}(A^{-1} N \leq t_{\text{mix}}^p(\mathcal{C}^N) \leq AN) \geq 1 - c_2 A^{-\theta_2}, \quad \forall A, N \geq K_2. \tag{6.10}$$

Finally, below is a list of exponents for each example in Section 5.

Section	$v(R)$	$r(R)$	$h(N)$	$\gamma(N)$
5.1	$R^{\log K / \log L}$	$R^{\log \lambda / \log L}$	L^N	$(K\lambda)^N$
5.2 with $a_N = N^{1/\alpha}, \alpha \in (1, 2]$	$R^{\alpha/(\alpha-1)}$	R	$N^{1-1/\alpha}$	$N^{2-1/\alpha}$
5.3	R^2	R	$N^{1/3}$	N
5.4	R	R	N	N^2

Here the Euclidean distance is used instead of the intrinsic shortest path metric for the examples in Section 5.1. Note that when $\alpha = 2$ in Section 5.2 (the finite variance case), the growth of $v(R)$ and $r(R)$ is of the same order as in Section 5.3. The difference of scaling exponents of mixing times (namely $\gamma(N)$) is due to the difference of scaling exponents for graph distances (namely $h(N)$). We also observe that the convergence to a stable law at (5.2) forces the scaling constants to be of the form $a_N = N^{1/\alpha} L(N)$ for some slowly varying function L (see [21, Section 35]), and hence the above table captures all the most important first order behavior for the examples in Section 5.2.

6.3 Examples

Critical Galton-Watson trees of Section 5.2

By combining the results in this section with our mixing time convergence result, we can establish asymptotic bounds for the distributions of mixing times of graphs in the sequence $(\mathcal{T}_N)_{N \geq 1}$ in the case when we have a finite variance offspring distribution.

Proposition 6.6. *In the case when the offspring distribution has finite variance, there exist constants $c_1, c_2, c_3, c_4 \in (0, \infty)$ such that*

$$\limsup_{N \rightarrow \infty} \mathbf{P} \left(N^{-3/2} t_{\text{mix}}^\infty(\mathcal{T}_N) \geq \lambda \right) \leq c_1 e^{-c_2 \lambda^2}, \quad \forall \lambda \geq 0, \quad (6.11)$$

and also

$$\limsup_{N \rightarrow \infty} \mathbf{P} \left(N^{-3/2} t_{\text{mix}}^1(\rho_N) \leq \lambda^{-1} \right) \leq c_3 e^{-c_4 \lambda^{1/25}}, \quad \forall \lambda \geq 0. \quad (6.12)$$

Proof. To prove (6.11), we apply the general mixing time upper bound of Lemma 6.1 to deduce that

$$\mathbf{P} \left(N^{-3/2} t_{\text{mix}}^\infty(\mathcal{T}_N) \geq \lambda \right) \leq \mathbf{P} \left(8N^{-1/2} \text{diam}_{d_{\mathcal{T}_N}}(\mathcal{T}_N) \geq \lambda \right),$$

where $\text{diam}_{d_{\mathcal{T}_N}}(\mathcal{T}_N)$ is the diameter of \mathcal{T}_N with respect to $d_{\mathcal{T}_N}$, and we note that $\#E(\mathcal{T}_N)$ is equal to $2(N-1)$. By (5.3), the right-hand side here converges to $\mathbf{P}(8 \text{diam}_{d_{\mathcal{T}^{(2)}}}(\mathcal{T}^{(2)}) \geq \lambda)$. By construction, the diameter of the continuum random tree $\mathcal{T}^{(2)}$ is bounded above by twice the supremum of the Brownian excursion of length 1. We can thus use the known distribution of the latter random variable (see [26], for example) to deduce the relevant bound.

For (6.12), we first apply the convergence in distribution of Theorem 5.3 to deduce that

$$\limsup_{N \rightarrow \infty} \mathbf{P} \left(N^{-3/2} t_{\text{mix}}^1(\rho_N) \leq \lambda^{-1} \right) \leq \mathbf{P} \left(t_{\text{mix}}^1(\rho) \leq \lambda^{-1} \right).$$

Now, for the continuum random tree, define

$$J(\lambda) = \{r > 0 : \lambda^{-1} r^2 \leq \pi^{(2)}(B_{\mathcal{T}^{(2)}}(\rho, r)) \leq \lambda r^2, R_{\mathcal{T}^{(2)}}^{(2)}(\rho, B_{\mathcal{T}^{(2)}}(\rho, r)^c) \geq \lambda^{-1} r\},$$

where $R_{\mathcal{T}^{(2)}}$ is the resistance on the continuum random tree (see [11, (20)]). Then

$$\mathbf{P}(r \in J(\lambda)) \geq 1 - e^{-c\lambda}, \quad \forall r \in (0, \frac{1}{2}], \lambda \geq 1,$$

(see [11, Lemmas 4.1 and 7.1]). As a consequence of this, we can apply the continuous version of the mixing time lower bound discussed in Remark 6.3 (with $H_0 = 0, H_1 = H_2 = H_3 = 1, H'_2 = 3, \alpha_i = 1$ and $d_i = 2$) to deduce the desired result. \square

Remark 6.7. The above proof already gives an estimate for the lower tail of $t_{\text{mix}}^1(\rho)$. That the bound corresponding to (6.11) holds for the limiting tree, i.e.

$$\mathbf{P} \left(t_{\text{mix}}^\infty(\mathcal{T}^{(2)}) \geq \lambda \right) \leq c_1 e^{-c_2 \lambda^2},$$

can be proved similarly to the discrete case (see Remark 6.3).

Critical Erdős-Rényi random graph of Section 5.3

Let \mathcal{C}^N be the largest component of the Erdős-Rényi random graph in the critical window. Then the following holds.

Proposition 6.8. *There exist constants $c_1, c_2, c_3, N_0, \theta \in (0, \infty)$ such that*

$$\sup_{N \geq 1} \mathbf{P} \left(N^{-1} t_{\text{mix}}^\infty(\mathcal{C}^N) \geq \lambda \right) \leq c_1 e^{-c_2 \lambda}, \quad \forall \lambda \geq 0, \quad (6.13)$$

$$\sup_{N \geq N_0} \mathbf{P} \left(N^{-1} t_{\text{mix}}^1(\mathcal{C}^N) \leq \lambda^{-1} \right) \leq c_3 \lambda^{-\theta}, \quad \forall \lambda \geq 0. \quad (6.14)$$

Proof. By [36, Proposition 1.4] and [37, Theorem 1], (6.13) is an application of Proposition 6.4 with $p_1(\lambda) = c_4 e^{-c_5 \lambda^{3/2}}$ and $p_2(\lambda) = c_6 e^{-c_7 \lambda^3}$. (6.14) is a consequence of Proposition 6.5. \square

Remark 6.9. (1) The tail estimates for $t_{\text{mix}}^1(\mathcal{C}^N)$ are given in [36, Theorem 1.1] without quantitative bounds. (In fact, reading the paper very carefully, it can be checked that the bounds similar to Proposition 6.8 are available in the paper.)

(2) It does not seem possible to apply current estimates for the graphs $(\mathcal{C}^N)_{N \geq 1}$ and techniques for bounding mixing times to replace $t_{\text{mix}}^1(\mathcal{C}^N)$ by $t_{\text{mix}}^1(\rho^N)$ in the latter estimate (see [15], Remark A.4), or even prove that the sequence $(N/t_{\text{mix}}^1(\rho^N))_{N \geq 1}$ is tight, i.e.

$$\lim_{\lambda \rightarrow \infty} \limsup_{N \rightarrow \infty} \mathbf{P} \left(N^{-1} t_{\text{mix}}^1(\rho^N) \leq \lambda^{-1} \right) = 0.$$

That this final statement is nonetheless true is a simple consequence of Theorem 5.6.

References

- [1] L. Addario-Berry, N. Broutin, and C. Goldschmidt, *The continuum limit of critical random graphs*, Probab. Theory Related Fields, to appear.
- [2] D. Aldous and J. Fill, *Reversible Markov chains and random walks on graphs*, Preprint <http://www.stat.berkeley.edu/~aldous/RWG/book.html>
- [3] M.T. Barlow, A.A. Járai, T. Kumagai and G. Slade, *Random walk on the incipient infinite cluster for oriented percolation in high dimensions*, Comm. Math. Phys. **278** (2008), 385–431.
- [4] I. Benjamini, G. Kozma and N. Wormald, *The mixing time of the giant component of a random graph*, preprint.
- [5] P. Bérard, G. Besson and S. Gallot, *Embedding Riemannian manifolds by their heat kernel*, Geom. Funct. Anal. **4** (1994), 373–398.
- [6] C. Borgs, J.T. Chayes, R. van der Hofstad, G. Slade and J. Spencer, *Random subgraphs of finite graphs: I. The scaling window under the triangle condition*, Random Structures Algorithms, **27** (2005), 137–184.
- [7] D. Burago, Y. Burago, and S. Ivanov, *A course in metric geometry*, Graduate Studies in Mathematics, vol. 33, American Mathematical Society, Providence, RI, 2001.
- [8] G.-Y. Chen and L. Saloff-Coste, *The cutoff phenomenon for ergodic Markov processes*, Electron. J. Probab. **13** (2008), 26–78.
- [9] D. A. Croydon, *Scaling limit for the random walk on the largest connected component of the critical random graph*, Publ. RIMS. Kyoto Univ., to appear.
- [10] ———, *Convergence of simple random walks on random discrete trees to Brownian motion on the continuum random tree*, Ann. Inst. Henri Poincaré Probab. Stat. **44** (2008), 987–1019.
- [11] ———, *Volume growth and heat kernel estimates for the continuum random tree*, Probab. Theory Related Fields **140** (2008), 207–238.
- [12] ———, *Random walk on the range of random walk*, J. Stat. Phys. **136** (2009), 349–372.
- [13] ———, *Scaling limits for simple random walks on random ordered graph trees*, Adv. in Appl. Probab. **42** (2010), 528–558.
- [14] D. A. Croydon and B. M. Hambly, *Local limit theorems for sequences of simple random walks on graphs*, Potential Anal. **29** (2008), 351–389.
- [15] D. A. Croydon, B. M. Hambly and T. Kumagai, *Convergence of mixing times for sequences of random walks on finite graphs* (extended version), arXiv:1111.0108.
- [16] T. Duquesne, *A limit theorem for the contour process of conditioned Galton-Watson trees*, Ann. Probab. **31** (2003), 996–1027.
- [17] T. Duquesne and J.-F. Le Gall, *Probabilistic and fractal aspects of Lévy trees*, Probab. Theory Related Fields **131** (2005), 553–603.

- [18] P. Erdős and S. J. Taylor, *Some intersection properties of random walk paths*, Acta Math. Acad. Sci. Hungar. **11** (1960), 231–248.
- [19] N. Fountoulakis and B.A. Reed, *The evolution of the mixing rate of a simple random walk on the giant component of a random graph*, Random Structures Algorithms, **33** (2008), 68–86.
- [20] M. Fukushima, Y. Oshima and M. Takeda, *Dirichlet forms and symmetric Markov processes*, de Gruyter Studies in Mathematics, 19. Walter de Gruyter & Co., Berlin, 2011.
- [21] B. V. Gnedenko and A. N. Kolmogorov, *Limit distributions for sums of independent random variables*, Addison-Wesley Publishing Company, Inc., Cambridge, Mass., 1954, Translated and annotated by K. L. Chung. With an Appendix by J. L. Doob.
- [22] S. Goel, R. Montenegro, and P. Tetali, *Mixing time bounds via the spectral profile*, Electron. J. Probab. **11** (2006), 1–26.
- [23] A. Greven, P. Pfaffelhuber, and A. Winter, *Convergence in distribution of random metric measure spaces (Λ -coalescent measure trees)*, Probab. Theory Related Fields **145** (2009), 285–322.
- [24] M. Heydenreich and R. van der Hofstad, *Random graph asymptotics on high-dimensional tori II. Volume, diameter and mixing time*, Probab. Theory Related Fields, **149** (2011), 397–415.
- [25] A. Kasue and H. Kumura, *Spectral convergence of Riemannian manifolds*, Tôhoku Math. J. **46** (1994), 147–179.
- [26] D. P. Kennedy, *The distribution of the maximum Brownian excursion*, J. Appl. Probab. **13** (1976), 371–376.
- [27] J. Kigami, *Resistance forms, quasisymmetric maps and heat kernel estimates*, Memoirs Amer. Math. Soc. to appear.
- [28] J. Kigami, *Hausdorff dimensions of self-similar sets and shortest path metrics*, J. Math. Soc. Japan **47** (1995), 381–404.
- [29] J. Kigami, *Harmonic calculus on limits of networks and its application to dendrites*, J. Funct. Anal. **128** (1995), 48–86.
- [30] J. Kigami, *Analysis on fractals*, Cambridge Tracts in Mathematics, vol. 143, Cambridge University Press, Cambridge, 2001.
- [31] T. Kumagai, *Homogenization on finitely ramified fractals*, Stochastic analysis and related topics in Kyoto, Adv. Stud. Pure Math., vol. 41, Math. Soc. Japan, Tokyo, 2004, pp. 189–207.
- [32] T. Kumagai and S. Kusuoka, *Homogenization on nested fractals*, Probab. Theory Related Fields **104** (1996), 375–398.
- [33] T. Kumagai and J. Misumi, *Heat kernel estimates for strongly recurrent random walk on random media*, J. Theoret. Probab. **21** (2008), 910–935.
- [34] J.-F. Le Gall, *Random real trees*, Ann. Fac. Sci. Toulouse Math. (6) **15** (2006), 35–62.
- [35] D. Levin, Y. Peres and E. Wilmer, *Markov chains and mixing times*, Amer. Math. Soc., Providence, RI, 2009.
- [36] A. Nachmias and Y. Peres, *Critical random graphs: diameter and mixing time*, Ann. Probab. **36** (2008), 1267–1286.
- [37] A. Nachmias and Y. Peres, *The critical random graph, with martingales*, Israel J. Math. **176** (2010), 29–41.