

J o u r n a l
o f
E l e c t r o n i c
P r o b a b i l i t y

Vol. 6 (2001) Paper no. 8, pages 1–26

Journal URL

<http://www.math.washington.edu/~ejpecp/>

Paper URL

<http://www.math.washington.edu/~ejpecp/EjpVol6/paper8.abs.html>

**THE PRINCIPLE OF LARGE DEVIATIONS
FOR MARTINGALE ADDITIVE FUNCTIONALS
OF RECURRENT MARKOV PROCESSES ¹**

Matthias K. Heck

HypoVereinsbank, 150 East 42nd New York, NY 100117, USA
matthias_heck@america.hypovereinsbank.com

Faïza Maaouia

Faculté des Sciences de Tunis, Département de Mathématiques,
Campus Universitaire, 1060 - Tunis, Tunisia
Faiza.Maaouia@fst.rnu.tn

Abstract We give a principle of large deviations for a generalized version of the strong central limit theorem. This generalized version deals with martingale additive functionals of a recurrent Markov process.

Keywords Central Limit Theorem (CLT), Large Deviations Principle (LDP), Markov Processes, Autoregressive Model (AR1), Positive Recurrent Processes, Martingale Additive Functional (MAF)

AMS subject classification Primary 60F05, 60F10, 60F15; Secondary 60F17, 60J25.

Submitted to EJP on April 26, 2000. Final version accepted on March 2, 2001.

¹This work was done while the first named author was supported by a research fellowship of the Deutsche Forschungsgemeinschaft and the second named author was supported by a fellowship of the Tunisian University and the Courant Institute of Mathematical Sciences (NYU).

1 Introduction

This paper presents a natural extension of the (ASCLT) due to Brosamler [1], [2] and Schatte [27]. In the last few years the Almost Sure Central Limit Theorem (ASCLT) has emerged as an area of probability theory in which an intensive research activity has taken place. In this context we should in particular mention the work of Lacey & Philipp [17], Berkes & Dehling [3], Csörgö & Horváth [6], Rodzik & Rychlik [26] and Touati [29].

The aim of this paper is to establish the Large Deviations Principle (LDP) for a generalized version of the (ASCLT) for Martingale Additive Functionals (MAF's). This result can be regarded as an extension of the (ASCLT) for (MAF's), proved by the second named author (see Maâouia [21]) as well as an extension of the (LDP) for the (ASCLT) for i.i.d. random variables, proved by the first named author (see Heck [14]). For a slightly weaker version of the (LDP) for the (ASCLT) for i.i.d. random variables see also March and Seppäläinen [22].

1.1 Notation, terminology and data

$X = \{\Omega, \mathcal{F}, (\mathbb{P}_x)_{x \in E}, \mathbb{F} = (\mathcal{F}_k)_{k \in \mathbb{N}}, (X_k)_{k \in \mathbb{N}}\}$ denotes the canonical version of a homogeneous Markov process indexed by \mathbb{N} (non negative integers) with values in a measurable space (E, \mathcal{E}) ; \mathbb{F} being its natural filtration and \mathbb{P}_x its law starting from x .

We denote by Π the transition probability of the Markov chain X and by $(R_p)_{p \in]0,1[}$ its resolvent:

$$(1-1) \quad R_p(x, A) = \mathbb{E}_x \left(\sum_{k=1}^{\infty} p^{k-1} \mathbf{1}_{\{X_k \in A\}} \right) = \sum_{k=1}^{\infty} p^{k-1} \Pi_k(x, A) \text{ for } x \in E, A \in \mathcal{E}.$$

Using Duflo's [10] and Meyn & Tweedie's [23] terminology, we call a set $C \in \mathcal{E}$ a **small set** for the Markov chain X , if there exists a probability measure ν on \mathcal{E} (with $\nu(C) = 1$), $p_0 \in]0,1[$ and $b \in]0,1[$, such that:

$$(1-2) \quad \forall (x, A) \in E \times \mathcal{E}, p_0 R_{p_0}(x, A) \geq b \mathbf{1}_C(x) \nu(A).$$

From now on, the expression "X is a **positive recurrent Markov chain**" means that X has a small set C , with the following properties (1-3) and (1-4)

$$(1-3) \quad \mathbb{E}_x \left(\limsup_{n \rightarrow \infty} (\mathbf{1}_{\{X_n \in C\}}) \right) = \mathbb{E}_x \left(\sum_{k=0}^{\infty} \mathbf{1}_{\{X_k \in C\}} = +\infty \right) = 1, \forall x \in E,$$

$$(1-4) \quad \sup_{x \in C} \mathbb{E}_x (T_C) < \infty \text{ with } T_C = \inf \{k \geq 1, X_k \in C\}.$$

In this case there is a probability measure μ , invariant under Π , such that X is **Harris recurrent**:

$$(1-5) \quad \forall A \in \mathcal{E} \text{ with } \mu(A) > 0, \mathbb{E}_x \left(\overline{\lim}_k (\mathbf{1}_{\{X_k \in A\}}) \right) = 1.$$

We shall say that X is **Riemannian recurrent of order k** , for each $k \in \mathbb{N}$, if (1-3) and

$$(1-6) \quad \sup_{x \in C} \mathbb{E}_x (\mathbf{T}_C^k) < \infty \text{ for each } k \in \mathbb{N}$$

hold.

Conversely, the existence of an invariant probability measure μ for X satisfying (1-5) implies the existence of a small set satisfying (1-3) and (1-4), if we assume that the σ -algebra \mathcal{E} is countably generated (cf. Dufflo [10], Meyn & Tweedie [23]).

We remind that an **additive functional** (**AF**) $A = (A_k)_{k \in \mathbb{N}}$ of X is an \mathbb{F} -adapted process, vanishing at 0, such that:

$$(1-7) \quad A_{k+l} = A_k + A_l \circ \theta_k \quad (\mathbb{P}_\nu\text{- a.s.}) \quad \forall k, l \in \mathbb{N},$$

for any initial law ν . Here $(\theta_k)_{k \in \mathbb{N}}$ are the standard translation operators on (Ω, \mathcal{F}) .

A **martingale additive functional** (**MAF**), $M = (M_k)_{k \in \mathbb{N}}$ of X is an (**AF**) which is also an $(\mathbb{F}, \mathbb{P}_\nu)$ martingale, for any initial law ν or equivalently

$$(1-8) \quad \mathbb{E}_x(M_k) = 0, \quad \forall k \in \mathbb{N}, \quad \forall x \in E.$$

Next, we will use the following notation and terminology.

(1-9) $\mathcal{C}_0([0,1])$ is the space of continuous functions from $[0,1]$ to \mathbb{R} vanishing in 0.

(1-10) $\mathcal{M}_1(\mathcal{C}_0([0,1]))$ is the space of probability measures on the Borel sets of $\mathcal{C}_0([0,1])$, endowed with the topology of weak convergence.

(1-11) $\mathcal{N} = \mathcal{N}(0,1)$ denotes the Gaussian law with mean 0 and variance 1 on \mathbb{R} .

For $a \in]0, 1]$ we introduce the function

$$(1-12) \quad \vartheta_a: \mathcal{C}_0([0,1]) \longrightarrow \mathcal{C}_0([0,1]) \text{ with } \vartheta_a(\omega)(t) = \frac{\omega(at)}{\sqrt{a}}.$$

Using these functions ϑ_a we call a measure $Q \in \mathcal{M}_1(\mathcal{C}_0([0,1]))$ ϑ -invariant if $Q = Q \circ \vartheta_a^{-1}$ for all $a \in]0, 1]$. Furthermore, for two probability measures η, ρ on a measurable space, we denote by $H(\eta|\rho)$ the relative entropy of η relative to ρ , i.e.

$$(1-13) \quad H(\eta|\rho) = \begin{cases} \int \ln \left(\frac{d\eta}{d\rho} \right) d\rho & : \text{if } \eta \ll \rho, \text{ i.e. if } \eta \text{ is abs. continuous w.r.t. } \rho \\ \infty & : \text{else.} \end{cases}$$

Now we define the rate function $\mathcal{H} : \mathcal{M}_1(\mathcal{C}_0([0, 1])) \longrightarrow [0, \infty]$ as follows:

$$(1-14) \quad \mathcal{H}(Q) = \begin{cases} \lim_{a \downarrow 0} \frac{1}{\ln(a^{-1})} H(Q \circ |_{[a,1]}^{-1} | W \circ |_{[a,1]}^{-1}) & : \text{if } Q \text{ is } \vartheta\text{-invariant} \\ \infty & : \text{else} \end{cases}$$

where W is the Wiener measure on $\mathcal{C}_0([0,1])$ and $|_{[a,1]}$ denotes the restriction operator. That \mathcal{H} is well defined has already been shown in Heck [14, 15].

1.2 ASCLT for MAF of a recurrent Markov process

The second named author proved the following general version of the (ASCLT) (see Maâouia [21]).

Theorem A *Let $X = (\Omega, \mathcal{F}, (\mathbb{P}_x)_{x \in E}, \mathbb{F} = (\mathcal{F}_k)_{k \in \mathbb{N}}, (X_k)_{k \in \mathbb{N}})$ be a positive recurrent Markov chain. Then every (MAF) M of X satisfying the assumption:*

$$(1-15) \quad \mathbb{E}_x(M_k^2) < \infty, \quad \forall k \in \mathbb{N}, \forall x \in E \quad \text{and} \quad \mathbb{E}_\mu(M_1^2) = \sigma_M^2 \in]0, \infty[$$

satisfies a functional ASCLT (FASCLT) under \mathbb{P}_x for all initial states x . More precisely, \mathbb{P}_x -almost-surely for every x , we have the following properties:

(FASCLT) *The random measures $(W_n)_{n \in \mathbb{N}^*}$:*

$$(1-16) \quad W_n(\cdot) = L(n)^{-1} \sum_{k=1}^n k^{-1} \delta_{\{\Psi_k \in \cdot\}}$$

converge weakly to W , the Wiener measure on $\mathcal{C}_0([0,1])$, where $L(n) = \sum_{k=1}^n k^{-1}$ and $(\Psi_n)_{n \in \mathbb{N}^}$ is defined by:*

$$(1-17) \quad \Psi_n(t) = \sigma_M^{-1} n^{-1/2} \{M_{[nt]} + (nt - [nt])(M_{[nt]+1} - M_{[nt]})\}. \quad \diamond$$

2 Main results

Our results are stated for (MAF), $M = (M_k)_{k \in \mathbb{N}}$ of the Markov process X which satisfies the assumption (2-1) below.

$$(2-1) \quad \begin{cases} \mathbb{E}_\mu(|M_1|^\beta) < \infty, \quad \forall \beta > 0; \\ \mathbb{E}_\mu(M_1^2) = \sigma_M^2 \in]0, \infty[. \end{cases}$$

For every (MAF) M satisfying the assumptions (2-1) we consider the processes $(\Psi_n)_n$ and the measures $(W_n)_n$, defined as in Theorem A.

Theorem 2.1 *Let $X = (\Omega, \mathcal{F}, (\mathbb{P}_x)_{x \in E}, \mathbb{F} = (\mathcal{F}_k)_{k \in \mathbb{N}}, (X_k)_{k \in \mathbb{N}})$ be a Riemannian recurrent Markov chain of order k , for every $k \in \mathbb{N}$, with a small set C . Then every (MAF) M of X satisfying the assumption (2-1) satisfies the (LDP) with constants $(\ln(n))_{n \in \mathbb{N}^*}$ and rate function \mathcal{H} w.r.t. \mathbb{P}_x for μ -a.a. initial states $x \in E$; i.e. for every Borel set $A \subset \mathcal{M}_1(\mathcal{C}_0([0,1]))$:*

$$(2-2) \quad -\inf_{\overset{\circ}{A}} \mathcal{H} \leq \liminf_{n \rightarrow \infty} \frac{1}{\ln(n)} \ln(\mathbb{P}_x \{W_n \in A\})$$

$$\leq \limsup_{n \rightarrow \infty} \frac{1}{\ln(n)} \ln (\mathbb{P}_x \{W_n \in A\}) \leq -\frac{\inf \mathcal{H}}{A}$$

for μ - a.a. initial states $x \in E$. \diamond

The results we present could easily be generalized to the continuous time parameter case. However for the proof of the continuous parameter case we would need rather technical oscillation estimates very similar to those used in Heck [15] in order to reduce the continuous case to the discrete time case. These lengthy technical estimates would increase the size of the paper considerably without presenting any new ideas. Therefore we decided to restrict ourselves to the discrete time parameter case.

3 The identification of an autoregressive process

In this section we shall apply our result Theorem 2.1 to autoregressive models. The latter models have a great interest in mathematical finance (for example: risk management, derivative securities like options, stochastic volatility,..., see e.g. Hull [16], section 19.6) .

On a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ we consider a sequence $\beta = (\beta_n)_{n \in \mathbb{N}^*}$ of i.i.d. real random variables with mean 0 and variance $\sigma^2 > 0$; called **white noise**. To this sequence β and a given random variable X_0 we associate the first order autoregressive process (**AR1**):

$$(3-1) \quad X_{n+1} = \theta X_n + \beta_{n+1}$$

or

$$(3-2) \quad X_{n+1} = \alpha + \theta X_n + \beta_{n+1}$$

where α and θ are unknown real parameters. These parameters α and β are to be estimated.

In the following we shall assume that the random variables β satisfies the moment condition

$$(3-3) \quad \mathbb{E} \left(|\beta_1|^{2\delta} \right) < \infty$$

for some $\delta > 1$.

For the (**AR1**), defined by (3-1), the least squares estimator of θ :

$$(3-4 \text{ a}) \quad \hat{\theta}_n = \left(\sum_{k=1}^n X_{k-1}^2 \right)^{-1} \left(\sum_{k=1}^n X_{k-1} X_k \right) \text{ for each } n \in \mathbb{N}^*$$

satisfies

$$(3-4 \text{ b}) \quad \widehat{\theta}_n - \theta = \left(\sum_{k=1}^n X_{k-1}^2 \right)^{-1} \left(\sum_{k=1}^n X_{k-1} \beta_k \right).$$

Under the hypothesis $\mathbb{E}(\beta_1^2) < \infty$, $(\widehat{\theta}_n)$ has the following asymptotic properties (see [10] for more details).

(3-5) $(\widehat{\theta}_n)_{n \in \mathbb{N}^*}$ is a strongly consistent estimator of the arbitrary unknown real parameter θ .

In the **stable case** ($|\theta| < 1$), $(\widehat{\theta}_n)$ satisfies:

$$(3-6) \quad \sqrt{n} (\widehat{\theta}_n - \theta) \xrightarrow[n \rightarrow \infty]{\Longrightarrow} \mathcal{N}(0, 1 - \theta^2)$$

$$(3-7) \quad \limsup_{n \rightarrow \infty} \sqrt{\frac{n}{\ln n}} \left| \widehat{\theta}_n - \theta \right| = \sqrt{1 - \theta^2} \quad \text{a.s. .}$$

Under the hypothesis (3-3) and in the stable case, the following result hold under \mathbb{P}_x for all starting state x :

$$(3-8) \quad W_n^\theta = L(n)^{-1} \sum_{k=1}^n k^{-1} \delta_{\Psi_k^\theta} \xrightarrow[n \rightarrow \infty]{\Longrightarrow} W$$

where $\Psi_n^\theta \in \mathcal{C}_0([0, 1])$ is linear on $[\frac{k}{n}, \frac{k+1}{n}]$ and

$$(3-9) \quad \Psi_n^\theta \left(\frac{k}{n} \right) = \sqrt{n (1 - \theta^2)^{-1} \frac{k}{n}} (\widehat{\theta}_k - \theta) \quad \text{for each } k \in \{1, \dots, n\}$$

and " \Longrightarrow " denotes weak convergence.

The property (3.8) is a consequence of the FASCLT for the martingales obtained by Chaâbane [5]. It is also consequence of Theorem A above, if we assume that the noise β satisfy (3.3) and the distribution of β_1 has a non vanishing density part. In fact, under these hypotheses, we prove the existence of a small set for the AR(1) Markov chain X (see Lemma 4.8).

The next Proposition gives the LDP associated with the property (3-8).

Proposition 3.1 *Let $X = (X_n)_{n \in \mathbb{N}}$ be the first order stable autoregressive process (AR1) satisfying (3-1) constructed from a white noise $\beta = (\beta_n)_{n \in \mathbb{N}^*}$ with variance σ^2 satisfying the hypothesis (3-3) for all $\delta > 1$, such that the distribution of β_1 has a non vanishing density part and an unknown real parameter $\theta \in]-1, 1[$. Then for all $X_0 \equiv x \neq 0$ the following result holds for the least squares estimator.*

(3-10) $(W_n^\theta)_n$ satisfies the (PLD) with constants $(\ln n)_n$ and rate function \mathcal{H} w.r.t. \mathbb{P}_x ; i.e. for every Borel set $A \subset \mathcal{M}_1(\mathcal{C}_0([0, 1]))$:

$$\begin{aligned}
-\inf_{\overset{\circ}{A}} \mathcal{H} &\leq \liminf_{n \rightarrow \infty} \frac{1}{\ln n} \ln (\mathbb{P}_x \{W_n^\theta \in A\}) \\
&\leq \limsup_{n \rightarrow \infty} \frac{1}{\ln n} \ln (\mathbb{P}_x \{W_n^\theta \in A\}) \leq -\inf_A \mathcal{H}. \diamond
\end{aligned}$$

For the (AR1) model, defined by (3-2), we can estimate α and θ by:

$$(3-11) \quad \begin{cases} \hat{\theta}_n &= D_n^{-1} \left\{ \left(\frac{1}{n} \sum_{k=1}^n X_{k-1} X_k \right) - \left(\frac{1}{n} \sum_{k=1}^n X_k \right) \left(\frac{1}{n} \sum_{k=1}^n X_{k-1} \right) \right\}, \\ \hat{\alpha}_n &= D_n^{-1} \left\{ \left(\frac{1}{n} \sum_{k=1}^n X_k \right) \left(\frac{1}{n} \sum_{k=1}^n X_{k-1}^2 \right) - \left(\frac{1}{n} \sum_{k=1}^n X_{k-1} \right) \left(\frac{1}{n} \sum_{k=1}^n X_{k-1} X_k \right) \right\}; \end{cases}$$

for each $n \in \mathbb{N}^*$, with $D_n = \left(\frac{1}{n} \sum_{k=1}^n X_{k-1}^2 \right) - \left(\frac{1}{n} \sum_{k=1}^n X_{k-1} \right)^2$.

These estimators satisfy

$$(3-12) \quad \begin{cases} D_n (\hat{\theta}_n - \theta) &= \left(\frac{1}{n} \sum_{k=1}^n X_{k-1} \right) \left(\frac{1}{n} \sum_{k=1}^n \beta_k \right) - \left(\frac{1}{n} \sum_{k=1}^n X_{k-1} \beta_k \right), \\ D_n (\hat{\alpha}_n - \alpha) &= \left(\frac{1}{n} \sum_{k=1}^n X_{k-1}^2 \right) \left(\frac{1}{n} \sum_{k=1}^n \beta_k \right) - \left(\frac{1}{n} \sum_{k=1}^n X_{k-1} \right) \left(\frac{1}{n} \sum_{k=1}^n X_{k-1} \beta_k \right), \\ (\hat{\alpha}_n - \alpha) &= \left(\frac{1}{n} \sum_{k=1}^n X_{k-1} \right) (\hat{\theta}_n - \theta) + \left(\frac{1}{n} \sum_{k=1}^n \beta_k \right); \end{cases}$$

and they have the following asymptotic properties:

(3-13) $(\hat{\theta}_n)_{n \geq 1}$ and $(\hat{\alpha}_n)_{n \geq 1}$ are strongly consistent estimators of the arbitrary unknown parameters θ and α .

In the **stable case** ($|\theta| < 1$), $(\hat{\theta}_n)_{n \geq 1}$ and $(\hat{\alpha}_n)_{n \geq 1}$ satisfy:

$$(3-14) \quad \begin{cases} \sqrt{n} (\hat{\theta}_n - \theta) \xrightarrow[n \rightarrow \infty]{} \mathcal{N}(0, 1 - \theta^2), \\ \sqrt{n} (\hat{\alpha}_n - \alpha) \xrightarrow[n \rightarrow \infty]{} \mathcal{N}\left(0, (1 - \theta^2) \left(\frac{\sigma^2}{1 - \theta^2} + \frac{\alpha^2}{(1 - \theta)^2} \right)\right); \end{cases}$$

$$(3-15) \quad \begin{cases} \limsup_{n \rightarrow \infty} \sqrt{\frac{n}{\ln \ln n}} |\hat{\theta}_n - \theta| = \sqrt{1 - \theta^2} \quad a.s., \\ \limsup_{n \rightarrow \infty} \sqrt{\frac{n}{\ln \ln n}} |\hat{\alpha}_n - \alpha| = \sqrt{(1 - \theta^2) \left(\frac{\sigma^2}{1 - \theta^2} + \frac{\alpha^2}{(1 - \theta)^2} \right)} \quad a.s.; \end{cases}$$

$$(3-16) \quad \begin{cases} W_n^\theta = L(n)^{-1} \sum_{k=1}^n k^{-1} \delta_{\Psi_k^\theta} \xrightarrow[n \rightarrow \infty]{} W, \\ W_n^\alpha = L(n)^{-1} \sum_{k=1}^n k^{-1} \delta_{\Psi_k^\alpha} \xrightarrow[n \rightarrow \infty]{} W; \end{cases}$$

where $\Psi_n^\theta, \Psi_n^\alpha \in \mathcal{C}_0([0, 1])$ are linear on $[\frac{k}{n}, \frac{k+1}{n}]$ and

$$(3-17) \quad \begin{cases} \Psi_n^\theta(\frac{k}{n}) = \sqrt{\frac{n}{(1-\theta^2)}} \frac{k}{n} (\hat{\theta}_k - \theta), \\ \Psi_n^\alpha(\frac{k}{n}) = \sqrt{\frac{n}{(1-\theta^2)(\frac{\sigma^2}{1-\theta^2} + \frac{\alpha^2}{(1-\theta)^2})}} \frac{k}{n} (\hat{\alpha}_k - \alpha); \forall k \in \{1, \dots, n\}. \end{cases}$$

Proposition 3.2 *Let $X = (X_n)_{n \in \mathbb{N}}$ be the (AR1) model satisfying (3-2) constructed from a white noise $\beta = (\beta_n)_{n \in \mathbb{N}^*}$ with variance σ^2 satisfying the hypothesis (3-3) for all $\delta > 1$, such that the distribution of β_1 has a non vanishing density part and an unknown real parameters $\theta \in]-1, 1[$ and α . Then for all $X_0 \equiv x \neq 0$ the following result holds for the least squares estimator:*

$$(3-18) \quad (W_n^\theta)_n \text{ and } (W_n^\alpha)_n \text{ satisfy the (LDP) with constants } (\ln n)_n \text{ and rate function } \mathcal{H} \text{ w.r.t. } \mathbb{P}_x. \diamond$$

4 Proofs

4.1 An ASCLT for i.i.d. random variables

The proof of Theorem 2.1 is essentially based on a reduction to a version of the (ASCLT) for i.i.d. random variables. In order to formulate this version we introduce some notations.

For random variables $(\xi_n, \tau_n)_{n \in \mathbb{N}^*}$ as in Proposition 4.1 below we denote by S_n and T_n the corresponding partial sums, i.e. $S_n = \sum_{k=1}^n \xi_k$ and $T_n = \sum_{k=1}^n \tau_k$ and let for $t \geq 0$ $N_t = \inf\{k \geq 0 : T_{k+1} > t\}$. Further let $S_n^* = S_{N_n}$.

As in the introduction we define random functions $\tilde{\Psi}_n \in \mathcal{C}_0([0, 1])$ by

$$\tilde{\Psi}_n(t) = \frac{\sqrt{m}}{\sigma\sqrt{n}} \left\{ S_{[nt]}^* + (nt - [nt]) (S_{[nt]+1}^* - S_{[nt]}^*) \right\} \forall t \in [0, 1].$$

Finally we define random measures $\tilde{W}_n \in \mathcal{M}_1(\mathcal{C}_0([0, 1]))$ by $\tilde{W}_n = \frac{1}{L(n)} \sum_{k=1}^n \frac{1}{k} \delta_{\tilde{\Psi}_k}$.

Proposition 4.1 *Let $(\xi_n, \tau_n)_{n \in \mathbb{N}^*}$ be independent random variables on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, such that for some $n_0 \in \mathbb{N}^*$*

$$(4-1 \text{ a}) \quad \mathbb{E}(\xi_n) = 0 \text{ for all } n \geq 1, \sup_{n \geq 1} \mathbb{E}(|\xi_n|^\beta) < \infty \text{ for all } \beta > 0, \text{ and } \mathbb{E}(\xi_n^2) = \sigma^2 > 0 \text{ for all } n \geq n_0.$$

$$(4-1 \text{ b}) \quad \tau_n \geq 0 \text{ for all } n \geq 1, \sup_{n \geq 1} \mathbb{E}(\tau_n^\beta) < \infty \text{ for all } \beta > 0 \text{ and } \mathbb{E}(\tau_n) = m > 0 \text{ for all } n \geq n_0.$$

Then $(\tilde{W}_n)_{n \geq 1}$ satisfies the (LDP) with constants $(\ln n)_{n \geq 1}$ and rate function \mathcal{H} . \diamond

Remark 4.2 We shall remark that in the special case $(\xi_k)_{k \in \mathbb{N}^*}$ i.i.d. and $\tau_k \equiv 1$, i.e. $N_n \equiv n$, the above proposition states the (LDP) for the (ASCLT) for i.i.d. random variables. This result is exactly the contents of Theorem 1.2 in Heck [14].

In order to prove Proposition 4.1 we shall recall for the readers convenience some simple facts:

Lemma 4.3 *Let $Y_n, Z_n, n \in \mathbb{N}^*$ be random variables with values in a separable metric space (E, d) such that for all $\varepsilon > 0$*

$$(4-2) \quad \lim_{n \rightarrow \infty} \frac{1}{\ln n} \ln (\mathbb{P} \{d(Y_n, Z_n) > \varepsilon\}) = -\infty .$$

Then $\left(\frac{1}{L(n)} \sum_{k=1}^n \frac{1}{k} \delta_{Y_k} \right)_{n \in \mathbb{N}^*}$ and $\left(\frac{1}{L(n)} \sum_{k=1}^n \frac{1}{k} \delta_{Z_k} \right)_{n \in \mathbb{N}^*}$ are equivalent w.r.t. the (LDP) i.e. $\left(\frac{1}{L(n)} \sum_{k=1}^n \frac{1}{k} \delta_{Y_k} \right)_{n \in \mathbb{N}^*}$ satisfies the (LDP) with constants $(\ln n)_{n \in \mathbb{N}^*}$ if and only if $\left(\frac{1}{L(n)} \sum_{k=1}^n \frac{1}{k} \delta_{Z_k} \right)_{n \in \mathbb{N}^*}$ satisfies the (LDP) with constants $(\ln n)_{n \in \mathbb{N}^*}$ and the same rate function \mathcal{H} . \diamond

Lemma 4.3 is a minor modification of Lemma 2.7 in Heck [14]. Details shall be omitted.

Lemma 4.4 *Let $(M_n)_{n \in \mathbb{N}}$ be random variables with $M_0 \equiv 0$ and let η be a random variable with values in \mathbb{N} .*

a) *For all $\beta \geq 1$ and all $p, q > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$ ($\beta q > 1$) there exists $C_1 > 0$ such that*

$$\mathbb{E} \left(|M_\eta|^\beta \right) \leq C_1 \sup_{k \geq 1} \left\{ \mathbb{E} \left(|M_k - M_{k-1}|^{q\beta} \right)^{1/q} \right\} \mathbb{E} \left(\eta^{p(\beta+2)} \right)^{1/p} .$$

b) *If in addition $(M_n)_{n \in \mathbb{N}}$ is a martingale then there exists $C_2 > 0$ such that for all $\beta \geq 2$*

$$\mathbb{E} \left(|M_\eta|^\beta \right) \leq C_2 \sup_{k \geq 1} \left\{ \mathbb{E} \left(|M_k - M_{k-1}|^{q\beta} \right)^{1/q} \right\} \mathbb{E} \left(\eta^{p(\beta/2+2)} \right)^{1/p} . \quad \diamond$$

Proof. Let $p, q > 1$ such that $\frac{1}{p} + \frac{1}{q} = 1$. By Hölder's inequality and Chebychev's inequality

$$(4-3) \quad \begin{aligned} \mathbb{E} \left(|M_\eta|^\beta \right) &= \sum_{k=1}^{\infty} \mathbb{E} \left(|M_k|^\beta 1_{\{\eta=k\}} \right) \leq \sum_{k=1}^{\infty} \mathbb{E} \left(|M_k|^{\beta q} \right)^{\frac{1}{q}} \mathbb{P} (\{\eta = k\})^{\frac{1}{p}} \\ &\leq \sum_{k=1}^{\infty} \mathbb{E} \left(|M_k|^{\beta q} \right)^{\frac{1}{q}} \mathbb{E} \left(\eta^{p(\beta+2)} \right)^{\frac{1}{p}} k^{-(\beta+2)} \end{aligned}$$

In order to prove Part a) we shall use the following inequalities and $r = 1$ in eq. (4-3),

$$\mathbb{E}\left(|M_k|^{\beta q}\right) \leq k^{\beta q} \sup_{k \geq 1} \mathbb{E}\left(|M_n - M_{n-1}|^{\beta q}\right).$$

In order to prove part b) we shall take $r = \frac{1}{2}$ in eq. (4-3), proceed as in a) and use Burkholder-Davis-Gundy inequality to estimate $\mathbb{E}\left(|M_k|^{\beta q}\right)$ for $\beta q > 1$,

$$\begin{aligned} \mathbb{E}\left(|M_k|^{\beta q}\right) &\leq \mathbb{E}\left(\sup_{j \leq k} |M_j|^{\beta q}\right) \leq \text{const.} \mathbb{E}\left(\left\{\sum_{j=1}^k (M_j - M_{j-1})^2\right\}^{\frac{\beta q}{2}}\right) \\ &\leq \text{const.} k^{\frac{\beta q}{2}} \sup_{n \geq 1} \mathbb{E}\left(|M_n - M_{n-1}|^{\beta q}\right). \quad \square \end{aligned}$$

We shall remark that one can in particular choose for $(M_n)_{n \in \mathbb{N}}$ the partial sums of independent random variables with expectation 0.

Lemma 4.5 *Let $(\tau_n)_{n \in \mathbb{N}^*}$ be independent random variables satisfying (4-1 b).*

a) *For every $\beta \geq 1$ there exists $C_4 > 0$ such that for all $n \in \mathbb{N}^*$, $\mathbb{E}(|N_n|^\beta) \leq C_4 n^{\beta+1}$.*

b) *For all $\alpha \in]\frac{1}{2}, 1[$ and $\gamma > 0$ there exists $C_5 > 0$ such that for all sufficiently large $n \in \mathbb{N}^*$, $\mathbb{P}\{|N_n - \frac{n}{m}| > n^\alpha\} \leq C_5 n^{-\gamma}$. \diamond*

Proof. In order to prove part a) we observe that by Lemma 4.4 and Chebychev's inequality for $k \in \mathbb{N}^*$ with $k > \frac{2}{m}n + 2n_0$

$$\begin{aligned} \mathbb{P}\{N_n = k\} &\leq \mathbb{P}\{T_k \leq n\} \leq \mathbb{P}\{|T_k - \mathbb{E}(T_k)| \geq \frac{m}{2}k\} \\ &\leq \left(\frac{2}{m}\right)^{2\beta+8} k^{-2\beta-8} \mathbb{E}\left\{|T_k - \mathbb{E}(T_k)|^{2\beta+8}\right\} \\ &\leq \left(\frac{2}{m}\right)^{2\beta+8} C_6 k^{\beta+6} k^{-2\beta-8} = C_7 k^{-\beta-2}. \end{aligned}$$

This inequality implies $\mathbb{E}(N_n^\beta) \leq \sum_{k \leq \frac{2}{m}n + 2n_0} k^\beta + \sum_{k > \frac{2}{m}n + 2n_0} C_7 k^{-2}$, which in turn implies part a).

For the proof of part b) we note that again Lemma 4.4 and Chebychev's inequality imply for sufficiently large n

$$\begin{aligned}
\mathbb{P} \left\{ \left| N_n - \frac{n}{m} \right| > n^\alpha \right\} &= \sum_{\left| k - \frac{n}{m} \right| > n^\alpha} \mathbb{P} \{ T_k \leq n < T_{k+1} \} \\
&\leq \sum_{k=0}^{\lfloor \frac{n}{m} - n^\alpha \rfloor} \mathbb{P} \{ T_{k+1} \geq n \} + \sum_{k=\lfloor \frac{n}{m} + n^\alpha \rfloor + 1}^{\infty} \mathbb{P} \{ T_k \leq n \} \\
&\leq \sum_{k=0}^{\lfloor \frac{n}{m} - n^\alpha \rfloor} \mathbb{P} \left\{ |T_{k+1} - \mathbb{E}(T_{k+1})| \geq \frac{m}{2} n^\alpha \right\} + \sum_{k=\lfloor \frac{n}{m} + n^\alpha \rfloor + 1}^{\infty} \mathbb{P} \left\{ |T_k - \mathbb{E}(T_k)| \geq \mathbb{E}(T_k) - n \right\} \\
&\leq \sum_{k=0}^{\lfloor \frac{n}{m} - n^\alpha \rfloor} C_8 n^{\beta+2} n^{-2\alpha\beta} + \sum_{k=\lfloor \frac{n}{m} + n^\alpha \rfloor + 1}^{\infty} C_9 k^{\beta+2} (mk - mn_0 - n)^{-2\beta} \leq C_{10} n^{(1-2\alpha)\beta+3}
\end{aligned}$$

where we used that for $n \in \mathbb{N}^*$ sufficiently large and $k \leq \lfloor \frac{n}{m} - n^\alpha \rfloor$

$$n - \mathbb{E}(T_{k+1}) \geq n - \mathbb{E}\left(T_{\lfloor \frac{n}{m} - n^\alpha \rfloor + 1}\right) \geq n - m\left(\frac{n}{m} - n^\alpha - n_0 + 1\right) - \mathbb{E}(T_{n_0}) \geq \frac{m}{2} n^\alpha.$$

Hence we conclude the proof of part b) by choosing β sufficiently large. \square

Proof of Proposition 4.1 For the special case $\tau \equiv 1$ Proposition 4.1 has already been proved in Heck [14] (see Remark 4.2). Therefore, by Lemma 4.3 the proof of Proposition 4.1 is complete if we show that there exists a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with random variables $\mathcal{X} = (X_i, x_i)_{i \in \mathbb{N}^*}$ and $\mathcal{Y} = (Y_i, y_i)_{i \in \mathbb{N}^*}$ such that \mathcal{X} has the same distribution as $(\xi_i, \tau_i)_{i \in \mathbb{N}^*}$, Y_i , $i \in \mathbb{N}^*$ are independent, $\mathcal{N}(0,1)$ -distributed, $y_i \equiv 1$ and

$$(4-4) \quad \left(\widetilde{W}_n^{\mathcal{X}} \right)_{n \in \mathbb{N}^*} \text{ and } \left(\widetilde{W}_n^{\mathcal{Y}} \right)_{n \in \mathbb{N}^*} \text{ are equivalent w.r.t. the LDP.}$$

Here $\widetilde{W}_n^{\mathcal{X}}$ denotes the random measure \widetilde{W}_n constructed from the sequence \mathcal{X} . Similar we shall use the notations $S_n^{\mathcal{X}}$ and $\widetilde{\Psi}_n^{\mathcal{X}}$, to indicate that the functions are constructed from the sequence \mathcal{X} .

By Skorokhod's representation theorem there exists a probability space $(\widetilde{\Omega}, \widetilde{\mathcal{F}}, \widetilde{\mathbb{P}})$, a random variable $\widetilde{B}: \widetilde{\Omega} \rightarrow \mathcal{C}_0([0, \infty[)$ and $\widetilde{\mathbb{P}}$ -a.s. finite stopping times $0 = \widetilde{R}_0 \leq \widetilde{R}_1 \leq \widetilde{R}_2 \leq \dots$ such that

(4-5 a) \widetilde{B} is a Brownian motion

(4-5 b) $\left(\widetilde{B}_{\widetilde{R}_i} - \widetilde{B}_{\widetilde{R}_{i-1}} \right)_{i \in \mathbb{N}^*}$ have the same distribution as $(\xi_i)_{i \in \mathbb{N}^*}$

(4-5 c) $\widetilde{R}_i - \widetilde{R}_{i-1}$ are independent with $\sup_{n \geq 1} \mathbb{E} \left(\left(\widetilde{R}_n - \widetilde{R}_{n-1} \right)^\beta \right) < \infty$ for all $\beta > 0$ and $\mathbb{E} \left(\widetilde{R}_n - \widetilde{R}_{n-1} \right) = \sigma^2$ for $n \geq n_0$.

(See e.g. Chapter 1, Theorem 117 in Freedman [11] and Brosamler [1], p. 570 regarding the moments for the stopping times.)

Now let $\Omega = \tilde{\Omega} \times \mathbb{R}^{\mathbb{N}^*}$, \mathcal{F} the corresponding product- σ -field and

$$\mathbb{P}(A \times A_1 \times \dots \times A_n \times \mathbb{R} \times \dots) = \int_A \prod_{i=1}^n \eta_i \left(\tilde{B}_{\tilde{R}_i} - \tilde{B}_{\tilde{R}_{i-1}}, A_i \right) d\tilde{\mathbb{P}}.$$

Here η_i denotes the conditional distribution $\eta_i(t, \cdot) = \mathbb{P}(\tau_i \in \cdot \mid \xi_i = t)$.

If we let $B(\tilde{\omega}, (t_n)_{n \in \mathbb{N}^*}) = \tilde{B}(\tilde{\omega})$, $R(\tilde{\omega}, (t_n)_{n \in \mathbb{N}^*}) = \tilde{R}(\tilde{\omega})$, then (4.5) still hold for \tilde{B} and \tilde{R}_i replaced by B and R_i .

Hence by scaling properties of Brownian motion, if we let $X_n = B_{R_n} - B_{R_{n-1}}$,

$x_n(\tilde{\omega}, (t_k)_{k \in \mathbb{N}^*}) = t_n$, $Y_n = \frac{\sqrt{m}}{\sigma} \left(B_{\frac{n\sigma^2}{m}} - B_{\frac{(n-1)\sigma^2}{m}} \right)$ and finally $y_n \equiv 1$ then obviously it remains to prove (4-4) for this special choice for \mathcal{X} and \mathcal{Y} .

By Lemma 4.3 part b) the proof of (4-4) is complete if we show that for all $\varepsilon > 0$

$$(4-6) \quad \lim_{n \rightarrow \infty} \frac{1}{\ln n} \ln \left(\mathbb{P} \left\{ \left\| \tilde{\Psi}_n^{\mathcal{X}} - \tilde{\Psi}_n^{\mathcal{Y}} \right\|_{\infty} > \varepsilon \right\} \right) = -\infty.$$

Note that $\tilde{\Psi}_n^{\mathcal{X}} \left(\frac{k}{n} \right) = \frac{\sqrt{m}}{\sigma\sqrt{n}} B_{R_{N_k}}$ and $\tilde{\Psi}_n^{\mathcal{Y}} \left(\frac{k}{n} \right) = \frac{\sqrt{m}}{\sigma\sqrt{n}} B_{\frac{n\sigma^2}{m}}$. Hence the definition of $\tilde{\Psi}_n$ via interpolation implies for $k \in \{0, 1, \dots, n-1\}$ and $t \in \left[\frac{k}{n}, \frac{k+1}{n} \right]$

$$(4-7) \quad \left| \tilde{\Psi}_n^{\mathcal{X}}(t) - \tilde{\Psi}_n^{\mathcal{Y}}(t) \right| \leq \frac{\sqrt{m}}{\sigma\sqrt{n}} \max \left\{ \left| B_{R_{N_k}} - B_{\frac{k\sigma^2}{m}} \right|, \left| B_{R_{N_{k+1}}} - B_{\frac{(k+1)\sigma^2}{m}} \right| \right\}.$$

Hence the proof of (4-6) is complete if we show that for all $\varepsilon > 0$

$$(4-8) \quad \lim_{n \rightarrow \infty} \frac{1}{\ln(n)} \ln \left(\mathbb{P} \left\{ \max_{k=1, \dots, n} \left\{ \left| B_{R_{N_k}} - B_{\frac{k\sigma^2}{m}} \right| \right\} > \varepsilon\sqrt{n} \right\} \right) = -\infty.$$

Let $\varepsilon, \gamma > 0$. Fix $p > 1$, $\beta > 0$ such that $\beta > 9 + 4\gamma$. By Lemma 4.4 part b), (4-5 b) and Lemma 4.5 part a) we conclude that for sufficiently large $n \in \mathbb{N}$ and $k \in \{1, \dots, [n^{3/4}]\}$

$$(4-9) \quad \mathbb{P} \left\{ \left| B_{R_{N_k}} \right| \geq \frac{\varepsilon}{2}\sqrt{n} \right\} \leq \left(\frac{2}{\varepsilon} \right)^{2\beta} C_{12} n^{-\beta} \mathbb{E} \left(N_k^{p(\beta+2)} \right)^{\frac{1}{p}} \leq \left(\frac{2}{\varepsilon} \right)^{2\beta} C_{13} n^{-\beta} k^{\beta+2+\frac{1}{p}}$$

$$(4-10) \quad \leq \left(\frac{2}{\varepsilon} \right)^{2\beta} C_{13} n^{-\frac{1}{4}\beta + \frac{9}{4}} \leq \left(\frac{2}{\varepsilon} \right)^{2\beta} C_{13} n^{-\gamma}.$$

Using Lemma 4.4 part b) we conclude that for sufficiently large $n \in \mathbb{N}^*$ and $k \in \{1, \dots, [n^{3/4}]\}$

$$(4-11) \quad \mathbb{P} \left\{ \left| B_{\frac{k\sigma^2}{m}} \right| > \frac{\varepsilon}{2} \sqrt{n} \right\} \leq C_{14} \varepsilon^{-2\beta} n^{-\gamma}.$$

For the following we assume that $n \in \mathbb{N}^*$ is sufficiently large and $k \in \{[n^{1/4}], \dots, n\}$. Observing that

$$(4-12) \quad \mathbb{P} \left\{ \left| B_{R_{N_k}} - B_{\frac{k\sigma^2}{m}} \right| > \varepsilon \sqrt{n} \right\} \leq \mathbb{P} \left\{ \left| B_{R_{N_k}} - B_{\frac{k\sigma^2}{m}} \right| > \varepsilon \sqrt{n}, \left| R_{N_k} - \frac{k\sigma^2}{m} \right| \leq \sigma^2 k^{\frac{4}{5}} \right\} \\ + \mathbb{P} \left\{ \left| R_{N_k} - \frac{k\sigma^2}{m} \right| > \sigma^2 k^{\frac{4}{5}}, \left| N_k - \frac{k}{m} \right| \leq k^{\frac{3}{4}} \right\} + \mathbb{P} \left\{ \left| N_k - \frac{k}{m} \right| > k^{\frac{3}{4}} \right\}.$$

Obviously Lemma 4.5 part b) implies that

$$(4-13) \quad \mathbb{P} \left\{ \left| N_k - \frac{k}{m} \right| > k^{\frac{3}{4}} \right\} \leq C_{15} k^{-\frac{16}{3}\gamma} \leq C_{15} n^{-\gamma}.$$

Keeping in mind that B is a Brownian motion, the symmetry properties of Brownian motion and

$$(4-14) \quad \mathbb{P} \left\{ \sup_{t \in [a, b]} \{|B_t - B_a|\} > c \right\} \leq 2 \exp \left\{ -\frac{c^2}{2(b-a)} \right\}$$

for $c > 0$, $0 \leq a < b$ show that

$$(4-15 \text{ a}) \quad \mathbb{P} \left\{ \left| B_{R_{N_k}} - B_{\frac{k\sigma^2}{m}} \right| > \varepsilon \sqrt{n}, \left| R_{N_k} - \frac{k\sigma^2}{m} \right| \leq \sigma^2 k^{\frac{4}{5}} \right\} \\ (4-15 \text{ b}) \quad \leq \mathbb{P} \left\{ \sup_{t \in \left[\frac{k\sigma^2}{m} - \sigma^2 k^{\frac{4}{5}}, \frac{k\sigma^2}{m} + \sigma^2 k^{\frac{4}{5}} \right]} \left| B_t - B_{\frac{k\sigma^2}{m}} \right| > \varepsilon \sqrt{n} \right\} \\ (4-15 \text{ c}) \quad \leq 2 \mathbb{P} \left\{ \sup_{t \in [0, \sigma^2 k^{\frac{4}{5}}]} |B_t| > \varepsilon \sqrt{n} \right\} \\ (4-15 \text{ d}) \quad \leq 4 \exp \left\{ -\frac{\varepsilon^2}{2\sigma^2} n k^{-\frac{4}{5}} \right\} \leq 4 \exp \left\{ -\frac{\varepsilon^2}{2\sigma^2} n^{\frac{1}{5}} \right\}.$$

Applying Lemma 4.4 part b) we obtain for β sufficiently large

$$(4-16 \text{ a}) \quad \mathbb{P} \left\{ \left| R_{N_k} - \frac{k\sigma^2}{m} \right| > \sigma^2 k^{\frac{4}{5}}, \left| N_k - \frac{k}{m} \right| \leq k^{\frac{3}{4}} \right\} \\ (4-16 \text{ b}) \quad \leq \sum_{l=-[k^{3/4}]-1}^{[k^{3/4}]+2} \mathbb{P} \left\{ \left| R_{l+\lceil \frac{k}{m} \rceil} - \frac{k\sigma^2}{m} \right| > \sigma^2 k^{\frac{4}{5}} \right\} \\ (4-16 \text{ c}) \quad \leq \sum_{l=-[k^{3/4}]-1}^{[k^{3/4}]+2} \mathbb{P} \left\{ \left| R_{l+\lceil \frac{k}{m} \rceil} - \mathbb{E} \left(R_{l+\lceil \frac{k}{m} \rceil} \right) \right| > \frac{1}{2} \sigma^2 k^{\frac{4}{5}} \right\}$$

$$(4-16 \text{ d}) \quad \leq \sum_{l=-[k^{3/4}]-1}^{[k^{3/4}]+2} C_{16} \frac{(l + [\frac{k}{m}])^{\beta+2}}{k^{\frac{8}{5}\beta}}$$

$$(4-16 \text{ e}) \quad \leq \sum_{l=-[k^{3/4}]-1}^{[k^{3/4}]+2} C'_{16} k^{-\frac{3}{5}\beta+2} \leq C_{17} k^{-\frac{3}{5}\beta+3} \leq C_{17} n^{-\gamma}.$$

Now we conclude by the inequalities (4-9) to (4-16) that

$$(4-17) \quad \limsup_{n \rightarrow \infty} \frac{1}{\ln n} \ln \left(\mathbb{P} \left\{ \max_{k=1, \dots, n} \left| \mathbb{B}_{\mathbb{R}^N}_k - \mathbb{B}_{\frac{k\sigma^2}{m}} \right| > \varepsilon \sqrt{n} \right\} \right) \leq -\gamma + 1$$

and hence (4-8) by letting $\gamma \rightarrow \infty$ in (4-17). \square

4.2 Proof of Theorem 2.1

The proof of Theorem 2.1 is divided into three steps. First we shall consider the case where the small set is also an atom for the Markov chain, second we shall prove the theorem under the additional assumption that (1-2) already holds for the transition function Π it self instead of pR_p . And finally in the third and last step we shall prove the general case.

We shall remark that this technique has been already used by several authors (see e.g. Dufflo [10], Touati [28]). In connection with (ASCLT's) this technique was introduced by Maâouia [19-21].

First of all we shall prove

Lemma 4-6 *Let $X = (\Omega, \mathcal{F}, (\mathbb{P}_x)_{x \in E}, \mathbb{F} = (\mathcal{F}_k)_{k \in \mathbb{N}}, (X_k)_{k \in \mathbb{N}})$ be a Riemannian recurrent Markov chain of order k , for each $k \in \mathbb{N}$ with invariant measure μ . We have:*

a) *Each measurable function $g: E \rightarrow [0, \infty[$ such that $\mu(g) < \infty$ satisfies*

$$\sup_n \mathbb{E}_x (g(X_n)) < \infty, \mu - a.e.$$

b) *Each positive random variable Y on (Ω, \mathcal{F}) such that $\mathbb{E}_\mu (Y) < \infty$ satisfies*

$$\sup_n \mathbb{E}_x (Y \circ \theta_n) < \infty, \mu - a.e.. \quad \diamond$$

Proof. To prove this lemma we shall use Theorem 2 of [24]. Indeed in order to apply Theorem 2 we have to verify that under our assumptions X is a positive Harris recurrent chain with an irreducible kernel Π , a maximal irreducible measure μ and convergence parameter 1 (see e.g. [24]). By Theorem 2.1 of [25] we know that there exist an integer $m_0 \geq 1$, a positive function s (called small function) satisfying $\mu(s) \in]0, \infty[$ and a bounded positive measure η on (Ω, \mathcal{F}) (called small measure) such that the following minoration condition holds

$$\Pi_{m_0}(x, A) \geq s(x)\eta(A), \forall x \in E, \forall A \in \mathcal{E}.$$

Then by the Theorem 2 of [25] we can see that

$$\sup_{1 \leq h \leq g} |\mathbb{E}_x(h(X_n)) - \mu(h)| \xrightarrow{n \rightarrow \infty} 0, \mu - a.e.,$$

letting $h = g$ and using the fact that $\mu(g) < \infty$ and μ is Π -invariant, we have

$$\mathbb{E}_x(g(X_n)) < \infty \quad \mu - a.e., \quad \mathbb{E}_x(g(X_n)) \leq |\mathbb{E}_x(g(X_n)) - \mu(g)| + \mu(g)$$

and then

$$\sup_n \mathbb{E}_x(g(X_n)) < \infty, \mu - a.e. .$$

So the first part of the Lemma is proved.

In order to prove the second part we simply apply part a) to the function

$$g : x \in E \mapsto \mathbb{E}_x(Y). \quad \square$$

4.2.1 Case I: Atomic chains

For the following we shall assume that X not only has a small set A but also that A is an atom for the Markov chain X .

Let T_A denote the first entry time into A , i.e. $T_A = \inf \{k > 0, X_k \in A\}$ and let $T_0 \equiv 0$, and $T_{k+1} = T_A \circ \theta_{T_k} + T_k$. Further for $k \in \mathbb{N}^*$ let $\xi_k = M_{T_k} - M_{T_{k-1}}$ and $\tau_k = T_k - T_{k-1}$. Since the chain is positive recurrent, it is well known that the invariant distribution is given by

$$(4-18) \quad \mu(\cdot) = \frac{1}{\mathbb{E}_a(T_A)} \mathbb{E}_a \left(\sum_{k=0}^{T_A-1} 1_{\{X_k \in \cdot\}} \right) \text{ where } a \in A \text{ is arbitrary.}$$

Further, since A is an atom, the Markov property implies that $\Xi = (\xi_k, \tau_k)_{k \in \mathbb{N}^*}$ is a sequence of independent random variables and $(\xi_k, \tau_k)_{k \geq 2}$ are identically distributed w.r.t. \mathbb{P}_x for all $x \in E$. Keeping in mind that the chain is Riemannian recurrent of order k for all $k \in \mathbb{N}^*$ the Markov property shows that for $x \in E$ and $\beta > 0$

$$\mathbb{E}_x \left(|T_2 - T_1|^\beta \right) = \mathbb{E}_a \left(|T_A|^\beta \right) < \infty.$$

By Proposition 8.3.23 in Dufflo [10] we conclude that $\mathbb{E}_\mu \left(T_1^\beta \right) < \infty$ and hence $\mathbb{E}_x \left(T_1^\beta \right) < \infty$ for $\mu - a.a. x \in E$. This together with the identical distribution for $k \geq 2$ implies

$$(4-19) \quad \sup_{n \in \mathbb{N}^*} \mathbb{E}_x \left(\tau_n^\beta \right) < \infty \text{ for } \mu - a.a. x \in E.$$

Since $T_A \geq 1$,

$$(4-20) \quad \mathbb{E}_x(\tau_k) = m > 0.$$

Using Lemma 4.4 part a) we conclude for $\mu - a.a. x \in E$,

$$\begin{aligned} \mathbb{E}_x(|\xi_1|^\beta) &= \mathbb{E}_x(|M_{T_1}|^\beta) < \infty \text{ and} \\ \mathbb{E}_x(|\xi_2|^\beta) &= \mathbb{E}_x(|M_{T_2} - M_{T_1}|^\beta) \leq 2^\beta \left\{ \mathbb{E}_x(|M_{T_2}|^\beta) + \mathbb{E}_x(|M_{T_1}|^\beta) \right\} < \infty. \end{aligned}$$

This together with the identical distribution of the ξ_i , $i \geq 2$ implies for $\mu - a.a. x \in E$

$$(4-21) \quad \sup_{n \in \mathbb{N}^*} \mathbb{E}_x(|\xi_n|^\beta) < \infty.$$

We have that for $\mu - a.a. x \in E$

$$(4-22) \quad \mathbb{E}_x(\xi_1) = \mathbb{E}_x(M_{T_1}) = 0 \text{ and } \mathbb{E}_x(\xi_2) = \mathbb{E}_a(M_{T_A}) = 0.$$

Moreover by the Martingale property of M_n , the Markov property of X and (4-18)

$$\begin{aligned} (4-23) \quad \mathbb{E}_x(\xi_2^2) &= \mathbb{E}_a(M_{T_A}^2) = \mathbb{E}_a \left(\sum_{k=0}^{T_A-1} \mathbb{E}_{X_k}((M_1 - M_0)^2) \right) \\ &= \mathbb{E}_\mu(M_1^2) \mathbb{E}_a(T_A) = \sigma_M^2 m > 0. \end{aligned}$$

(4-19) to (4-23) imply that $\Xi = (\xi_n, \tau_n)_{n \in \mathbb{N}}$ satisfies the assumptions of Proposition 4.1. Hence Proposition 4.1 implies that $(\widetilde{W}_n^\Xi)_{n \in \mathbb{N}^*}$ satisfies the (LDP) with the constants $(\ln n)_{n \in \mathbb{N}^*}$ and rate function \mathcal{H} . Therefore the proof of Theorem 2.1 is in this case complete if we show that $(\widetilde{W}_n^\Xi)_{n \in \mathbb{N}^*}$ and $(\widetilde{W}_n)_{n \in \mathbb{N}^*}$ are equivalent w.r.t. \mathbb{P}_x for $\mu - a.a. x \in E$.

Following the same steps as in the proof of Proposition 4.1 we see that in order to verify this equivalence it suffices to prove that for $\mu - a.a. x \in E$

$$(4-24) \quad \lim_{n \rightarrow \infty} \frac{1}{\ln n} \ln \left(\mathbb{P} \left\{ \max_{k=1, \dots, n} \{|M_k - S_k^{\Xi}| \} > \varepsilon \sqrt{n} \right\} \right) = -\infty.$$

If we let $Y_n = \sup_{T_n \leq t < T_{n+1}} |M_t - M_{T_n}|$, and as in the prior section $N_n = \inf \{k \geq 0, T_{k+1} > n\}$, then it is easy to see that

$$(4-25) \quad |M_n - S_n^{\Xi}| \leq Y_{N_n}.$$

By Doob's inequality and (4-21) for $\mu - a.a. x \in E$ and $\beta > 1$

$$(4-26) \sup_{k \geq 0} \mathbb{E}_x \left(Y_k^\beta \right) \leq C_{18} \sup_{k \geq 0} \mathbb{E}_x \left(|M_{T_{k+1}} - M_{T_k}|^\beta \right) = C_{18} \sup_{k \geq 1} \mathbb{E}_x \left(|\xi_k|^\beta \right).$$

So by Lemma 4.5 part a)

$$(4-27 \text{ a}) \mathbb{E}_x \left(Y_{N_k}^\beta \right) = \sum_{l=0}^{\infty} \mathbb{E}_x \left(Y_l^\beta 1_{\{N_k=l\}} \right) \leq \sum_{l=0}^{\infty} \mathbb{E}_x \left(Y_l^{2\beta} \right)^{1/2} \mathbb{P}_x (N_k = l)^{1/2}$$

$$(4-27 \text{ b}) \leq C_{19} \sum_{l=0}^{\infty} \frac{\mathbb{E}_x(N_k^4)^{1/2}}{l^2} \leq C_{20} k^3.$$

This together with Chebychev's inequality and (4-25) implies

$$(4-28) \mathbb{P} \left\{ \max_{k=1, \dots, n} \{ |M_k - S_k^*| \} > \varepsilon \sqrt{n} \right\} \\ \leq \sum_{k=0}^n \mathbb{P}_x (|Y_{N_k}| > \varepsilon \sqrt{n}) \leq \sum_{k=0}^n C_{21} \frac{k^3}{n^{\beta+6}} \leq C_{22} n^{-\beta}.$$

This concludes the proof of Theorem 2.1 for the special case of atoms. \square

4.2.2 Case II: Chains with minoration property

We shall proof in this section Theorem 2.1 under the additional assumption, that there exist a set $C \in \mathcal{E}$, $b \in]0, 1[$ and a probability measure $\nu \in \mathcal{M}_1(\mathbb{E})$ with $\nu(C) = 1$ such that

$$(4-29) \quad \Pi(x, \cdot) \geq b 1_C(x) \nu(\cdot).$$

We shall remark that in particular C is a small set (see e.g. Duflo [10] p. 286). Using this small set we construct (as in [21] for example) a new chain called split chain, i.e. a canonical version of a homogeneous Markov process

$$\bar{X} = (\bar{\Omega}, \bar{\mathcal{F}}, (\bar{\mathbb{P}}_x)_{x \in \bar{\mathbb{E}}}, \bar{\mathbb{F}} = (\bar{\mathcal{F}}_k)_{k \in \mathbb{N}}, (\bar{X}_k)_{k \in \mathbb{N}})$$

with values in $\bar{\mathbb{E}} = \mathbb{E} \times \{0, 1\}$ and transition probability

$$(4-30) \bar{\Pi}((x, i), A \times B) = \begin{cases} \nu(A) \{ (1 - b 1_C(x)) \delta_1(B) + b 1_C(x) \delta_0(B) \} & \text{if } i = 0, \\ Q(x, A) \{ (1 - b 1_C(x)) \delta_1(B) + b 1_C(x) \delta_0(B) \} & \text{if } i = 1; \end{cases}$$

where $Q(x, A) = (1 - b 1_C(x))^{-1} (\Pi(x, A) - b 1_C(x) \nu(A))$. It is well known that

Remarks 4.7

a) $\mathbb{E} \times \{0\}$ is an atom for \bar{X} and

$$(4-31) \quad \sup_{\bar{x} \in \mathbb{E} \times \{0\}} \sup_{\bar{x} \in \mathbb{E} \times \{0\}} \bar{\mathbb{E}}_{\bar{x}} \left(T_{\mathbb{E} \times \{0\}}^k \right) \leq C_{22} \sup_{\bar{x} \in C} \mathbb{E}_x \left(T_C^k \right) < \infty \text{ for all } k \geq 1.$$

b) If we denote the invariant distribution (which obviously exists by part a) of this remark) by $\bar{\mu}$, then $\bar{\mu}$ is related to the invariant distribution μ of the original Markov chain through $\bar{\mu}(A \times \{0\}) = b\mu(A \cap C)$ and $\bar{\mu}(A \times \{1\}) = (1-b)\mu(A \cap C) + \mu(A \cap C^c)$.

c) Part b) of the remark implies easily that for all $Z \in L^1(\mu)$, $\bar{\mathbb{E}}_{\bar{\mu}}(\bar{Z}) = \mathbb{E}_{\mu}(Z)$ and $\mathbb{E}_x(Z) = b 1_{\{x \in C\}} \bar{\mathbb{E}}_{(x,0)}(\bar{Z}) + (1-b) 1_{\{x \in C^c\}} \bar{\mathbb{E}}_{(x,1)}(\bar{Z})$ for all $x \in \mathbb{E}$. Here \bar{Z} denotes the natural lift of Z from Ω to $\bar{\Omega} = \Omega \times \{0, 1\}^{\mathbb{N}}$, i.e. $\bar{Z}(\omega, (x_n)_{n \geq 0}) = Z(\omega)$.

For details on the above construction and the remark we refer to Duflo [10], section 8.2.4.

By Remark 4.3 b) we conclude

$$(4-32) \quad \bar{\mathbb{E}}_{\bar{\mu}} \left(\bar{M}_1^2 \right) = \mathbb{E}_{\mu} \left(M_1^2 \right) = \sigma_M^2 \text{ and } \bar{\mathbb{E}}_{\bar{\mu}} \left(|\bar{M}_1|^\beta \right) = \mathbb{E}_{\mu} \left(|M_1|^\beta \right) < \infty.$$

Since Theorem 2.1 has already been proved for chains with atoms, we conclude by (4-31) and (4-32) that $\left(\bar{W}_n^{\bar{M}} \right)_{n \geq 0}$ satisfies the (LDP) with constants $(\ln n)_{n > 0}$ and rate function \mathcal{H} w.r.t. $\bar{\mathbb{P}}_{\bar{x}}$ for $\bar{\mu}$ - a.a. $\bar{x} \in \bar{\mathbb{E}}$.

Here $\bar{W}_n^{\bar{M}}$ denote the empirical measure defined as in (1-16) with $(M_k)_{k > 0}$ replaced by $(\bar{M}_k)_{k > 0}$. It is not hard to see that $\bar{W}_n^{\bar{M}}$ is the lift of W_n . We therefore conclude by Remark 4.7 part c) for $x \in \mathbb{E} \setminus C$ $\mathbb{P}_x \{W_n \in \cdot\} = \bar{\mathbb{P}}_{(x,1)} \left\{ \bar{W}_n^{\bar{M}} \in \cdot \right\}$ and for $x \in C$ $\mathbb{P}_x \{W_n \in \cdot\} = b \bar{\mathbb{P}}_{(x,0)} \left\{ \bar{W}_n^{\bar{M}} \in \cdot \right\} + (1-b) \bar{\mathbb{P}}_{(x,1)} \left\{ \bar{W}_n^{\bar{M}} \in \cdot \right\}$ and hence $(W_n)_{n > 0}$ satisfies the (LDP) with constants $(\ln n)_{n > 0}$ and rate function \mathcal{H} w.r.t. \mathbb{P}_x for μ - a.a. $x \in \mathbb{E}$. \square

4.2.3 Case III: General case

In this section will shall finish the proof of Theorem 2.1.

By enlarging the space if necessary, we may assume without loss of generality that there exists a sequence of i.i.d. random variables $(\rho_k)_{k > 0}$ with $\mathbb{P}_x(\rho_1 = 0) = p_0$ and $\mathbb{P}_x(\rho_1 = k) = (1 - p_0)^2 p_0^{k-1}$ for $k \in \mathbb{N}^*$ and $x \in \mathbb{E}$ which in addition are independent of the Markov chain. Then

$$\mathbb{E}_x(\rho_1) = 1. \text{ Now let } \mathcal{R}_0 \equiv 0, \mathcal{R}_n = \sum_{k=1}^n \rho_k \text{ and } \hat{X}_n = X_{\mathcal{R}_n}.$$

Then $\hat{X} = \left(\hat{\Omega}, \hat{\mathcal{F}}, (\hat{\mathbb{P}}_x)_{x \in \mathbb{E}}, \hat{\mathbb{F}} = (\hat{\mathcal{F}}_k)_{k \in \mathbb{N}}, (\hat{X}_k)_{k \in \mathbb{N}} \right)$ is a Markov chain with transition probability $\hat{\Pi}(x, dy) = p_0 \delta_x(dy) + (1 - p_0)^2 R_{p_0}(x, dy)$.

Hence, since C is a small set for X (i.e.(1-2) holds) C is a small set for \widehat{X} which satisfies in addition (4-29). Further μ is also the invariant distribution for \widehat{X} .

Moreover, if we let $\widehat{M}_n = M_{\mathcal{R}_n}$ then $\widehat{M} = \left(\widehat{M}_n, \sigma \left(\widehat{X}_k, k \leq n \right) \right)_n$ is a Martingale additive functional.

We shall show now that \widehat{X} and \widehat{M} satisfy the assumption of Theorem 2.1. By Lemma 4.4 part a) and the fact that $\mathbb{E}_\mu \left(|M_1|^\beta \right) = \mathbb{E}_\mu \left(|M_n - M_{n-1}|^\beta \right) < \infty$ we obtain

$$\mathbb{E}_\mu \left(\left| \widehat{M}_1 \right|^\beta \right) = \mathbb{E}_\mu \left(|M_{\mathcal{R}_1}|^\beta \right) < \infty, \mathbb{E}_\mu \left(\widehat{M}_1^2 \right) = \mathbb{E}_\mu \left(M_1^2 \right) \mathbb{E} \left(\mathcal{R}_1 \right) = \sigma_M^2.$$

Furthermore if we let $\widehat{T}_C = \inf \left\{ k \geq 1, \widehat{X}_k \in C \right\}$ then

$$(4-33) \quad \sup_{x \in C} \mathbb{E}_x \left(\widehat{T}_C^\beta \right) \leq C_{23} \sup_{x \in C} \mathbb{E}_x \left(T_C^\beta \right).$$

For $\beta = 1$ this is exactly part 4) of Proposition 8.2.13 in Duflo [10]. The general case is proved by a straight forward modification of the proof for $\beta = 1$ given in Duflo. Details shall be omitted.

We therefore obtain from the previous part of the proof of Theorem 2.1 (case II) that $\left(W_n^{\widehat{M}} \right)_{n \geq 1}$ satisfies the (LPD) with constants $(\ln(n))_{n \geq 1}$ and rate function \mathcal{H} w.r.t. $\widehat{\mathbb{P}}_x$ for $\mu - a.a. x \in E$.

Therefore the proof of Theorem 2.1 is complete, if we show that $\left(W_n^{\widehat{M}} \right)_{n \geq 1}$ and $\left(W_n^M \right)_{n \geq 1}$ are equivalent w.r.t. the (LDP). The proof of the equivalence however is a straight forward modification of the proof of (4-4). For the readers convenience we shall sketch the proof below.

As for (4-4) the proof can be reduced to

$$(4-34) \quad \lim_{n \rightarrow \infty} \frac{1}{\ln n} \ln \left(\mathbb{P}_x \left\{ \max_{k=1, \dots, n} \left| \widehat{M}_k - M_k \right| > \varepsilon \sqrt{n} \right\} \right) = -\infty$$

for $\mu - a.a. x \in E$ (compare (4-8)). Observe that for $\mu - a.a. x \in E$

$$(4-35) \quad C_{24} = \sup_n \mathbb{E}_x \left(|M_n - M_{n-1}|^\beta \right) < \infty$$

and hence by Lemma 4.4

$$(4-36) \quad \sup_n \mathbb{E}_x \left(\left| \widehat{M}_n - \widehat{M}_{n-1} \right|^\beta \right) = \sup_n \mathbb{E}_x \left(|M_{\mathcal{R}_n} - M_{\mathcal{R}_{n-1}}|^\beta \right) < \infty.$$

Fix such an $x \in E$ and $\gamma > 0$, Using (4-35) we obtain as in (4-11) and (4-12) for sufficiently large $n \geq 1$ and $k \in \{1, \dots, [n^{3/4}]\}$

$$(4-37) \quad \mathbb{P}_x \{ |M_k| > \varepsilon \sqrt{n} \} \leq C_{25} n^{-\gamma} \text{ and } \mathbb{P}_x \left\{ \left| \widehat{M}_k \right| > \varepsilon \sqrt{n} \right\} \leq C_{26} n^{-\gamma}.$$

By Lemma 4.4 and Chebychev's inequality we conclude for $k \in \{[n^{3/4}], \dots, n\}$

$$(4-38) \quad \mathbb{P}_x \{ |\mathcal{R}_k - k| > k^{3/4} \} \leq C_{27} k^{-\frac{4}{3}\gamma} \leq C_{28} n^{-\gamma}.$$

Now by Doob's maximal inequality, the Burkholder-Davis-Gundy inequality and Lemma 4.4 for sufficiently large $n \in \mathbb{N}^*$

$$(4-39 \text{ a}) \quad \mathbb{P}_x \left\{ \left| \widehat{M}_k - M_k \right| > \varepsilon \sqrt{n}, |\mathcal{R}_k - k| \leq k^{3/4} \right\}$$

$$(4-39 \text{ b}) \quad \leq \mathbb{P}_x \left\{ \max_{l \in \{1, \dots, 2[k^{3/4}] + 1\}} \left| M_{k-[k^{3/4}]} - M_{l+k-[k^{3/4}]} \right| > \frac{\varepsilon}{2} \sqrt{n} \right\}$$

$$(4-39 \text{ c}) \quad \leq C_{29} \mathbb{E}_x \left(\left| M_{k+[k^{3/4}]} - M_{k-[k^{3/4}]} \right|^{2\beta} \right) n^{-\beta}$$

$$(4-39 \text{ d}) \quad \leq C_{30} k^{\frac{3}{4}(\beta+2)} n^{-\beta} \leq C_{30} n^{-\gamma}.$$

Using these estimates we conclude the proof of (4-34) in the same way as the proof of (4-8).

This completes the proof of Theorem 2.1. \square

4.3 Proof of Proposition 3.1 and Proposition 3.2

We shall prove only Proposition 3.1 because the proof of Proposition 3.2 is a straight forward modification of the proof of Proposition 3.1 and contains no new ideas.

We shall denote by $X^x = (X_k^x)_{k \in \mathbb{N}}$ the (AR1) given through (3-1) with $X_0 \equiv x$.

We observe first that if $X = (\Omega, \mathcal{F}, (\mathbb{P}_x)_{x \in E}, \mathbb{F} = (\mathcal{F}_k)_{k \in \mathbb{N}}, (X_k)_{k \in \mathbb{N}})$ is a standard Markov chain on \mathbb{R} with transition probability $\Pi(x, \cdot) = \mathbb{P}(\theta x + \beta_1 \in \cdot)$, then:

$$(4-40) \quad \text{The distribution of } X^x \text{ under } \mathbb{P} \text{ is equal to that of } X \text{ under } \mathbb{P}_x.$$

It is well known that in the stable case the Markov chain has an invariant measure μ , which is equal to the distribution of $\sum_{k=1}^{\infty} \theta^{k-1} \beta_k$.

We shall prove next

Lemma 4.8

$$\text{a) For every } x \in \mathbb{R} \text{ and } \delta > 0, \sup_{n \in \mathbb{N}^*} \mathbb{E}_x \left(|X_n|^\delta \right) < \infty.$$

$$\text{b) Let } C =]a, b[\text{ with } a < b, \mu(C) > 0 \text{ and } \varepsilon > 0. \text{ Then } \sup_{x \in C_\varepsilon} \mathbb{E}_x \left((T_{C_\varepsilon})^\delta \right) < \infty \text{ for every } \delta > 0, \text{ where } C_\varepsilon =]a - \varepsilon, b + \varepsilon[. \diamond$$

Proof. For the proof of part a) we may assume without loss of generality that $\delta \in 2\mathbb{N}^*$. Using Hölder's inequality and the identical distribution of the random variables $(\beta_n)_{n \in \mathbb{N}^*}$ and letting $\beta_0 = x$ we obtain

$$\mathbb{E}_x \left(|X_n|^\delta \right) \leq \sum_{l_1, \dots, l_\delta=0}^n |\theta|^{l_1} \dots |\theta|^{l_\delta} \mathbb{E}_x (|\beta_{l_1}| \dots |\beta_{l_\delta}|) \leq (1 - |\theta|)^{-\delta} \mathbb{E}_x \left((|\beta_0| + \dots + |\beta_\delta|)^\delta \right).$$

In order to prove part b) it obviously suffices to show that there exist $m \in \mathbb{N}^*$, $q \in]0, 1[$ and $C_{31} > 0$ such that for all $n \in \mathbb{N}^*$

$$(4-41) \quad \sup_{x \in C_\varepsilon} \mathbb{P}_x (T_{C_\varepsilon} > m n) \leq C_{31} q^n.$$

Using (3-1) we obtain inductively for $k, n \in \mathbb{N}^*$ with $k < n$

$$(4-42) \quad X_n = \theta^{n-k} X_k + \sum_{l=1}^{n-k} \theta^{n-k-l} \beta_{l+k} = \theta^{n-k} X_k + Z_{k,n}$$

For $m \in \mathbb{N}^*$ with $4|\theta|^m (|a| + |b| + 1) < \varepsilon$ we conclude for $x \in C_\varepsilon$

$$\begin{aligned} \mathbb{P}_x (T_{C_\varepsilon} > m n) &\leq \mathbb{P}_x \left(\bigcap_{k=1}^n \{X_{k m} \notin C_\varepsilon\} \right) = \mathbb{P}_x \left(\bigcap_{k=1}^n \{\theta^{k m} x + Z_{0, k m} \notin C_\varepsilon\} \right) \\ &\leq \mathbb{P}_x \left(\bigcap_{k=1}^n \{Z_{0, k m} \notin C_{\frac{\varepsilon}{2}}\} \right) = \alpha_n(0). \end{aligned}$$

We dropped the parameter x in $\alpha_n(0)$, since the distribution of $Z_{0, i m}$, $i \in \mathbb{N}^*$ under \mathbb{P}_x is independent of x . In the following we fix $x \in \mathbb{R}$. Analogously to (4-42) we obtain

$$(4-43) \quad Z_{0, i m} = \theta^m Z_{0, (i-1) m} + Z_{(i-1) m, i m}.$$

We observe next that $Z_{(i-1) m, i m}$, $i \in \mathbb{N}^*$ are i.i.d. and that the distribution of $Z_{0, m}$ converges (for $m \rightarrow \infty$) weakly to the invariant measure μ . Hence by the Portmanteau Lemma $\limsup_{m \rightarrow \infty} \mathbb{P}_x (Z_{0, m} \notin C) \leq \mu (\mathbb{R} \setminus C) \leq 1 - \mu (C) < 1$.

For the following fix $r \in]\mu (\mathbb{R} \setminus C), 1[$ and $m \in \mathbb{N}^*$ such that $\mathbb{P}_x (Z_{0, m} \notin C) \leq r$,

$4|\theta|^m (|a| + |b| + 1) < \varepsilon$ and $q = r + 2|\theta|^m < 1$.

Letting $\alpha_n(i) = \mathbb{P}_x \left(\bigcap_{k=1}^n \left\{ Z_{0, k m} \notin \left[- \left(\frac{|\theta|^{-m}}{2} \right)^i \varepsilon, \left(\frac{|\theta|^{-m}}{2} \right)^i \varepsilon \right] \right\} \right)$ for $i \in \mathbb{N}^*$. We shall see that for $i \in \mathbb{N}$ and $n \geq 2$

$$(4-44) \quad \alpha_n(i) \leq r \alpha_{n-1}(i) + \alpha_{n-1}(i+1)$$

Indeed, using (4-43) and the independence of the $Z_{(i-1) m, i m}$, $i \in \mathbb{N}^*$, we conclude

$$\begin{aligned}
\alpha_n(0) &= \mathbb{P}_x \left(\bigcap_{k=1}^n \{Z_{0,km} \notin C_{\frac{\varepsilon}{2}}\} \right) \\
&\leq \mathbb{P}_x \left(\bigcap_{k=1}^{n-1} \{Z_{0,km} \notin C_{\frac{\varepsilon}{2}}\} \cap \{Z_{0,(n-1)m} \in [-(2|\theta|^m)^{-1}\varepsilon, (2|\theta|^m)^{-1}\varepsilon]\} \right. \\
&\quad \left. \cap \{|\theta|^m Z_{0,(n-1)m} + Z_{(n-1)m, nm} \notin C_{\frac{\varepsilon}{2}}\} \right) \\
&\quad + \mathbb{P}_x \left(\bigcap_{k=1}^{n-1} \{Z_{0,km} \notin C_{\frac{\varepsilon}{2}}\} \cap \{Z_{0,(n-1)m} \notin [-(2|\theta|^m)^{-1}\varepsilon, (2|\theta|^m)^{-1}\varepsilon]\} \right) \\
&\leq \mathbb{P}_x \left(\bigcap_{k=1}^{n-1} \{Z_{0,km} \notin C_{\frac{\varepsilon}{2}}\} \cap \{Z_{(n-1)m, nm} \notin C\} \right) + \alpha_{n-1}(1) \\
&\leq \alpha_{n-1}(0) \mathbb{P}_x (Z_{(n-1)m, nm} \notin C) + \alpha_{n-1}(1) = r \alpha_{n-1}(0) + \alpha_{n-1}(1).
\end{aligned}$$

We used also the fact that (by the choice of m) $C_{\frac{\varepsilon}{2}} \subseteq [-(2|\theta|^m)^{-1}\varepsilon, (2|\theta|^m)^{-1}\varepsilon]$.

The case $i \geq 1$ is proved analogously.

Using (4-44) an easy induction argument shows that

$$\alpha_n(0) \leq C_{32} \sum_{i=0}^{n-1} \binom{n-1}{i} r^{n-1-i} \alpha_1(i)$$

Observing that by part a) and Chebychev's inequality $\alpha_1(i) \leq C_{32} (2|\theta|^m)^i$ for some $C_{32} > 0$, we conclude

$$\alpha_n(0) \leq C_{32} \sum_{i=0}^{n-1} \binom{n-1}{i} r^{n-1-i} (2|\theta|^m)^i = C_{32} (r + 2|\theta|^m)^{n-1} = \frac{C_{32}}{q} q^n. \quad \square$$

Next we shall show that

Lemma 4.9 *There exists a small set C such that the Markov chain is Riemannian of any order k . \diamond*

Proof. Since the distribution of β_i has a non vanishing density part, the distribution of $\beta_i + \theta\beta_{i-1}$ has a non vanishing density part with a continuous density say h . Hence there exists $a < b$ such that $\inf_{]a,b[} h > 0$.

A simple application of Borel-Cantelli Lemma shows that there exists a $n_0 \in \mathbb{N}^*$ such that for all $n \geq n_0$ and $x \in \mathbb{R}$

$$(4-45) \quad \mathbb{P}_x \left(\sum_{k=n+1}^{\infty} |\beta_k \theta^k| < \varepsilon \right) \geq \frac{1}{2}.$$

Let $\varepsilon \in]0,1[$ such that $\frac{a}{2(1-\theta^2)} + 2\varepsilon < \frac{b}{2(1-\theta^2)} - 2\varepsilon$ and let $n \geq n_0$ such that $|\theta|^n \left(2 - \frac{|a|+|b|}{1-\theta^2}\right) < \varepsilon$. Now observe that for $n \in \mathbb{N}^*$

$$(4-46) \quad X_{2n} = \theta^{2n} X_0 + \sum_{i=1}^n \theta^{2n-2i} (\beta_{2i} + \theta\beta_{2i-1}) = \theta^{2n} X_0 + Z_{0,2n}$$

and hence the distribution of X_{2n} w.r.t. \mathbb{P}_x has a non vanishing density part with a continuous density, say f_{2n}^x , such that

$$(4-47) \quad \inf_{x \in]\frac{a}{1-\theta^2}, \frac{b}{1-\theta^2}[} f_{2n}^x > 0.$$

Moreover since the invariant measure μ is equal to the distribution of $Z_{0,2n} + \sum_{k=2n+1}^{\infty} \theta^{k-1} \beta_k$, it is easy to see that by (4-45) and (4-47) μ has a non vanishing density part say g with

$$]a \frac{1-\theta^{2n}}{1-\theta^2} + 2\varepsilon, b \frac{1-\theta^{2n}}{1-\theta^2} - 2\varepsilon[\quad g > 0.$$

By (4-47) we see that $]a \frac{1-\theta^{2n}}{1-\theta^2} + 2\varepsilon, b \frac{1-\theta^{2n}}{1-\theta^2} - 2\varepsilon[$ is a small set, and by Lemma 4.8 the Markov chain is Riemannian recurrent of any order k . \square

Lemma 4.10 *For every $x \in \mathbb{R}$ and $\delta > 0$ there exists $C_{33} > 0$ such that*

$$(4-48) \quad \mathbb{E}_x \left(\left| \sum_{k=1}^n X_{k-1}^2 - n \frac{\sigma^2}{1-\theta^2} \right|^\delta \right) \leq C_{33} n^{\delta/2}. \quad \diamond$$

Proof. We remark that

$$(4-49) \quad \sum_{k=1}^n X_{k-1}^2 - n \frac{\sigma^2}{1-\theta^2} = \frac{1}{1-\theta^2} \left(\sum_{k=1}^n \{ X_k^2 - \mathbb{E}_x (X_k^2 / \mathcal{F}_{k-1}) \} - X_n^2 + X_0^2 \right)$$

hence we conclude the proof by applying the Burkholder-Davis-Gundy inequality to the martingale $\left(\sum_{k=1}^n \{ X_k^2 - \mathbb{E}_x (X_k^2 / \mathcal{F}_{k-1}) \} \right)_n$ and the part a) of Lemma 4.8. \square

Proof of Proposition 3.1. We observe first that $M_n = \sum_{k=1}^n X_{k-1} \beta_k$, $n \in \mathbb{N}^*$ is a (MFA) with $\sigma_M^2 = \mathbb{E}_\mu (M_1^2) = \frac{\sigma^4}{1-\theta^2}$. Hence, by Lemma 4.9 we can apply Theorem 2.1 to M and the Markov chain X .

Letting $M_n^x = \sum_{k=1}^n X_{k-1}^x \beta_k$ we conclude by Theorem 2.1 and (4-40) that $(\Psi_n^{M^x})_{n \in \mathbb{N}^*}$ satisfies the (LDP) with constants $(\ln(n))_{n \in \mathbb{N}^*}$ and rate function \mathcal{H} w.r.t. \mathbb{P}_x for $\mu - a.a.$ $x \in \mathbb{R}$.

Furthermore by (4-42) $M_n^x - M_n^y = (x - y) \sum_{k=1}^n \theta^{k-1} \beta_k$. Since with M_n^x and M_n^y also $M_n^x - M_n^y$ is a martingale, Chebychev's inequality and Doob's maximal inequality imply easily for $\gamma \in \mathbb{N}^*$

$$\begin{aligned} \mathbb{P} \left\{ \max_{k=1, \dots, n} |M_n^x - M_n^y| > \varepsilon \sqrt{n} \right\} &\leq C_{34} \varepsilon^{-2\gamma} n^{-\gamma} |x - y|^{2\gamma} \mathbb{E}_x \left(\left| \sum_{k=1}^n \theta^{k-1} \beta_k \right|^{2\gamma} \right) \\ &\leq C_{35} n^{-\gamma}. \end{aligned}$$

This in turn implies that $(W_n^{M^x})_{n \in \mathbb{N}^*}$ and $(W_n^{M^y})_{n \in \mathbb{N}^*}$ are equivalent w.r.t. the (LDP) (see the proof of (4-4) and in particular (4-6) and (4-8)).

Therefore it remains to prove that for all initial states $x \neq 0$ $(W_n^\theta)_{n \in \mathbb{N}^*}$ and $(W_n^{M^x})_{n \in \mathbb{N}^*}$ are equivalent w.r.t. (LDP).

The proof of the equivalence is again very similar to that of (4-42), so that it suffices to give only a sketch of the proof.

Fix $x \neq 0$. A simple calculation show that

$$\begin{aligned} \left| \Psi_n^\theta \left(\frac{k}{n} \right) - \Psi_n^{M^x} \left(\frac{k}{n} \right) \right| &= \left| \sqrt{\frac{n}{1-\theta^2}} \binom{k}{n} \left(\sum_{k=1}^k X_{k-1}^2 \right)^{-1} M_k^x - \sqrt{\frac{1-\theta^2}{n\sigma^4}} M_k^x \right| \\ &= \sqrt{\frac{1-\theta^2}{n\sigma^4}} |V_k^x| |M_k^x| |U_k^x|^{-1} \end{aligned}$$

where $U_n^x = \sum_{k=1}^n (X_{k-1}^x)^2$ and $V_n^x = n \frac{\sigma^2}{1-\theta^2} - U_n^x$.

Using again the same arguments as in the proof of (4-4) it remains to verify

$$(4-50) \quad \limsup_{n \rightarrow \infty} \mathbb{P} \left\{ \max_{k=1, \dots, n} |V_k^x| |M_k^x| |U_k^x|^{-1} > \varepsilon \sqrt{n} \right\} = -\infty$$

Observing that $U_n^x \geq (X_0^x)^2 = x^2 > 0$ we obtain by Chebychev's inequality, Hölder's inequality and Lemma 4.10 and Lemma 4.4 analogously to (4-9) that for $\gamma > 0$, n sufficiently large and $k \leq n^{1/8}$

$$(4-51) \quad \mathbb{P} \left\{ |V_k^x| |M_k^x| |U_k^x|^{-1} > \varepsilon \sqrt{n} \right\} \leq C_{36} (x^2 \varepsilon)^{-2\gamma} n^{\frac{\gamma}{4}} n^{-\frac{\gamma}{2}} \leq C_{37} n^{-\frac{\gamma}{4}}$$

and for $k \in \{[n^{1/8}], \dots, n\}$

$$(4-52) \quad \mathbb{P} \left\{ |V_k| > k^{\frac{3}{4}} \right\} \leq C_{38} n^{-\frac{\gamma}{4}}.$$

Finally again by Chebychev's inequality and Lemma 4.4 we conclude for n sufficiently large and $k \in \{[n^{1/8}], \dots, n\}$

$$\begin{aligned}
(4-53 \text{ a}) \quad & \mathbb{P} \left\{ |V_k^x| |M_k^x| |U_k^x|^{-1} > \varepsilon \sqrt{n}, \quad |V_k^x| \leq k^{\frac{3}{4}} \right\} \\
(4-53 \text{ b}) \quad & \leq \mathbb{P} \left\{ k^{\frac{3}{4}} |M_k^x| 2 (\sigma^2 (1 - \theta^2)^{-1} k)^{-1} > \varepsilon \sqrt{n} \right\} \\
(4-53 \text{ c}) \quad & \leq \mathbb{P} \left\{ |M_k^x| > \varepsilon n^{\frac{17}{32}} \right\} \leq C_{39} n^{-\gamma}.
\end{aligned}$$

Now (4-50) is immediate consequence of (4-51) to (4-53), since $\gamma > 0$ was arbitrary. This concludes the proof of Proposition 3.1. \square

5 Remarks

We shall conclude this paper with some remarks.

a) Theorem 2.1 implies (LDP) for further a.s. limit theorems, like the (ASCLT) on the real line, a.s. versions of arcsine law (Compare Corollary 2.10 and Examples 2.11 in Heck [14]).

b) The (LDP) for (ASCLT) implies in particular easily the (ASCLT) itself. Therefore, for random variables satisfying the assumption of Theorem 2.1, Theorem 2.1 can be regarded as an generalization of Theorem A (see Corollary 2.12 in Heck [14]).

c) The moment assumptions like (1-6) or (2-1) are necessary compare to the corresponding assumptions in Heck [14] and March and Seppäläinen [22]. Lifshits & Stankevich [18] prove the necessity of these hypotheses.

Acknowledgment: This work was done during a visit of the authors to the Courant Institute of Mathematical Sciences, New York. The authors would like to thank you the Courant Institute for his hospitality. The authors would like to thank you S.R.S. Varadhan for helpful comments and useful discussions.

References

- [1] G. A. BROSAMLER *An almost everywhere central limit theorem*, Math. Proc. Cambridge Philos., 104, (1988), p: 561-574.
- [2] G. A. BROSAMLER *A simultaneous almost everywhere central limit theorem, for diffusions and its application to path energy and eigenvalues of the Laplacian*, Illinois J. Math., 34 (1990), p: 526-556.
- [3] I. BERKES & H. DEHLING *Some limit theorems in log density*, Ann. Probab. Vol. 23, N. 3, (1993), p: 1640-1670.
- [4] F. CHAABANE, F. MAAOUIA & A. TOUATI *Généralisation du théorème de la limite centrale presque-sûr pour les martingales vectorielles*, C. R. Acad. Sci. Paris, T 326, Série I, (1998), p: 229-232.
- [5] F. CHAABANE *Invariance principles with logarithmic averaging for martingales*, To appear in Studia Math. Sci. Hungar., (2001).

- [6] M. CSÖRGÖ, L. HORVÁTH *Invariance principles for logarithmic averages*, Proc. Camb. Phil. Soc., (1992), p: 112-195.
- [7] D. DACUNHA-CASTELLE & M. DUFLO *Probabilités et statistiques (2. Problèmes à temps mobiles)*, Masson, (1983).
- [8] J.D. DEUSCHEL & D.W. STROOCK *Large deviations*, Academic Press, (1989).
- [9] M. D. DONSKER *An invariance principle for certain probability limit theorems* Mem. Amer. Math. Soc. 6, (1951).
- [10] M. DUFLO *Random iterative models*, Masson, (1997).
- [11] D. FREEDMAN *Brownian motion and diffusion*, Springer, New York, (1983).
- [12] P. HALL & C.C. HEYDE *Martingale limit theory and its application. Probability and Mathematical Statistics*, Academic Press, (1980).
- [13] M. K. HECK *Das Prinzip der großen Abweichungen für fast sicheren Zentralen Grenzwertsatz und für seine funktionale Verallgemeinerung*, Dissertation, Saarbrücken, Germany, (1995).
- [14] M. K. HECK *The principle of large deviations for the almost everywhere central limit theorem*. Stoch. Process. Appl. 76, N°1, (1998), p: 61-75.
- [15] M. K. HECK *Principle of large deviations for the empirical processes of Ornstein-Uhlenbeck Process*, J. Theoret. Probab. 12, N°1, (1999), p: 147-179.
- [16] J. T. HULL *Options, futures, and other derivatives* Prentice-Hall, Upper Saddle River, (1997).
- [17] M. T. LACEY & W. PHILLIP *A note on the almost sure central limit theorem*, Statist. probab. Let. 9, (1990), p: 201-205.
- [18] M. A. LIFSHITS & E. S. STANKEVICH *On the Large Deviation Principle for the Almost Sure CLT*, Preprint Université de Lille, France.
- [19] F. MAAOUIA *Comportements asymptotiques des fonctionnelles additives des processus de Markov récurrents au sens de Harris*, Thèse de 3^{ème} cycle, Université de Paris VII, (1987).
- [20] F. MAAOUIA *Versions fortes du théorème de la limite centrale pour les processus de Markov*, C. R. Acad. Sci. Paris, T 323, Série I, (1996), p: 293-296.
- [21] F. MAAOUIA *Principes d'invariance par moyennisation logarithmique pour les processus de Markov*, To appear in The Annals of Probability, (2001).
- [22] P. MARCH & T. SEPPÄLÄINEN *Large deviations from the almost everywhere central limit theorem*, J. Theor. Probab. 10, (1997), p: 935-965.
- [23] S. P. MEYN & R. L. TWEEDIE *Markov chains and stochastic stability*, Springer, (1993).
- [24] E. NUMMELIN *General irreducible Markov chains and negative operators*, Cambridge University Press, (1983).
- [25] E. NUMMELIN & R. L. TWEEDIE *Geometric ergodicity and R-positivity for general Markov chains*, Ann. of Probab. Vol. 6, (1978), p: 404-420.
- [26] B. RODZIK & Z. RYCHLIK *An almost sure central limit theorem for independent random variables*, Ann. IHP Vol. 30, N. 1, (1994), p: 1-11.
- [27] P. SCHATTE *On strong versions of the central limit theorem*, Math. Nachr. 137, (1988), p: 249-256.
- [28] A. TOUATI *Loi fonctionnelle du logarithme itéré pour des processus de Markov récurrents*, Ann. Prob. Vol. 18, n° 1, (1990), p: 140-159.
- [29] A. TOUATI *Sur les versions fortes du théorème de la limite centrale*, Preprint Université de Marne-La-Vallée, n° 23, (1995).
- [30] S. R. S. VARADHAN *Large deviations and applications*, SIAM, Philadelphia, (1984).