

AN EXTENSION OF KAZAMAKI'S RESULTS ON BMO DIFFERENTIALS¹

BY PHILIP PROTTER

Purdue University

Kazamaki has shown that if $(M^n)_{n \geq 1}$, M are BMO martingales with continuous paths and $\lim M^n = M$ in BMO, then $\mathcal{E}(M^n)$ converges in \mathcal{H}^1 to $\mathcal{E}(M)$, where $\mathcal{E}(M)$ denotes the stochastic exponential of M . While Kazamaki's result does not extend to the right continuous case, it does extend "locally." It is shown here that if M^n , M are semimartingales and M^n converges locally in \mathcal{H}^∞ (a semimartingale BMO-type norm) to M then X^n converges locally in \mathcal{H}^p ($1 \leq p < \infty$) to X , where X^n , X are respectively solutions of stochastic integral equations with Lipschitz-type coefficients and differentials dM^n , dM . (The coefficients are also allowed to vary.) This is a stronger stability than usually holds for solutions of stochastic integral equations, reflecting the strength of the \mathcal{H}^∞ norm.

1. Introduction. Recently Kazamaki [5] and Kazamaki and Sekiguchi [6] showed that if M is a *continuous* martingale, in BMO then the stochastic exponential $\mathcal{E}(M)$ is in \mathcal{H}^1 , and if M^n converges in the BMO martingale norm to a martingale M in BMO then $\mathcal{E}(M^n)$ converges to $\mathcal{E}(M)$ in the \mathcal{H}^1 martingale norm: Simple examples show that these results do not extend to the right continuous case: $M \in \text{BMO}$ does not necessarily imply that $\mathcal{E}(M) \in \mathcal{H}^1$ (cf. Remarks (3.7), #3). One does have, of course, that $M \in \text{BMO}$ implies that $\mathcal{E}(M)$ is locally in \mathcal{H}^1 , but this is not surprising since every local martingale is locally in \mathcal{H}^1 !

A consequence of the results presented here is that if $M \in \text{BMO}$ then $\mathcal{E}(M)$ is locally in \mathcal{H}^p for all p , $1 \leq p < \infty$. Moreover we show that if M^n converges locally in BMO to M then $\mathcal{E}(M^n)$ converges locally in \mathcal{H}^p to $\mathcal{E}(M)$ for all p ($1 \leq p < \infty$). We extend Kazamaki's results further, however, by working with semimartingales and general stochastic differential equations. Suppose X , X^n are respectively solutions of

$$(1.1) \quad X_t = J_t + \int_0^t (FX)_s - dM_s$$

$$(1.2) \quad X_t^n = J_t^n + \int_0^t (F^n X^n)_s - dM_s^n$$

where M , M^n , J , J^n are semimartingales and where F , $F^n \in \text{Lip}(K)$. (Precise definitions are given in Section 2.) Meyer [9] has extended the notion of BMO martingales to semimartingales. Such a semimartingale is said to be in \mathcal{H}^∞ . We show that if $M \in \mathcal{H}^\infty$ and $J \in \mathcal{H}^p$ for some p ($1 \leq p < \infty$), then X is locally in \mathcal{H}^p for the same p . Moreover we show that if M^n converges locally in \mathcal{H}^∞ to M , and if F^n converges to F and J^n converges to J in appropriate ways, then X^n converges locally in \mathcal{H}^p to X ($1 \leq p < \infty$ with p depending on the convergence of F^n , J^n to F and J). This principal result is the content of Theorem (3.4).

By insisting that the differentials converge locally in \mathcal{H}^∞ we obtain local convergence of the solutions: that is, we get $(X^n)^{T_k}$ converging to X^{T_k} in \mathcal{H}^p for stopping times T_k increasing to ∞ a.s. (X and X^n are the solutions respectively of (1.1) and (1.2)). The previous best results (Emery [3] and Protter [12]) obtained only the *weak-local* convergence of *subsequences*.

Section 2 consists of preliminaries including some recent developments not contained in

Received August 3, 1978; revised February 1979.

¹ Supported in part by NSF grant no. MCS77-00095 and the Institute for Advanced Study.

AMS 1970 subject classifications. Primary 60H10; secondary 60H20, 60G45.

Key words and phrases. Stochastic differential equations; BMO, martingales, semimartingales.

Meyer [7]. One innovation is the generalization of Emery’s idea of “carving” a semimartingale into small \mathcal{H}^∞ slices; by requiring only that the slices be in \mathcal{H}^p , we show in Lemma (2.15) that if M^n converges to M in \mathcal{H}^p then there exists an N such that for all $n > N$ the same stopping times that carve M are also carving times for M^n . The convergence theorem described above (Theorem (3.4)) is the content of Section 3. In Section 4 we extend a result of Garcia, Maillard, and Peltraut [4] by constructing a local martingale with a given *random* “multiplicative jump” at a given totally inaccessible stopping time. We then apply Theorem (3.4) to obtain a continuity theorem for martingales with multiplicative jumps.

2. Preliminaries. We use the notation of and assume the reader is familiar with the theory of the semimartingale calculus as given in Meyer [7]. Let (Ω, \mathcal{F}, P) be a complete probability space and let $(\mathcal{F}_t)_{t \geq 0}$ be a right continuous filtration with $\mathcal{F}_\infty = \mathcal{F}$ and where \mathcal{F}_0 contains all P -null sets.

Let \mathcal{C} denote the adapted processes whose paths are right continuous with left limits (cadlag). For $J \in \mathcal{C}$ let

$$\|J\|_{\mathcal{S}^p} = \|\sup_{s < \infty} |J_s|\|_{L^p} \tag{1 \leq p \leq \infty}.$$

For a stopping time T we say a process $X \in \mathcal{C}$ is *stopped at T^-* if

$$X_t = X_t^{T^-} = X_t 1_{[0, T[} + X_{T-} 1_{[T, \infty[}$$

where $X_{t-} = \lim_{s \rightarrow t, s < t} X_s$. We let

$$\delta X_T = X_T - X_{T-},$$

the jump at T , and we make the notational conventions that

$$\|X\|_{\mathcal{S}^p(T)} = \|X^T\|_{\mathcal{S}^p}; \quad \|X\|_{\mathcal{S}^p(T-)} = \|X^{T-}\|_{\mathcal{S}^p}.$$

Emery [1] and Meyer [9] have proposed \mathcal{H}^p norms ($1 \leq p \leq \infty$ and $p = \omega$) for semimartingales. (Here ω represents the first limit ordinal and is not a point in Ω .) Meyer has shown that the \mathcal{H}^p norms for semimartingales are equivalent to the martingale \mathcal{H}^p norms when the process in question is a martingale ($1 \leq p \leq \infty$ or $p = \omega$; $p = \omega$ corresponds to the BMO martingales).

For a semimartingale M with a decomposition $M = N + A$ where N is a local martingale and A is a VF process, we define ($1 \leq p \leq \infty$)

$$j_p(N, A) = \|[N, N]_\infty^{1/2} + \int_{0-}^\infty |dA_s|\|_{L^p}.$$

For $p = \omega$, $j_\omega(N, A)$ is the smallest constant c such that for any stopping time T

$$E\{([N, N]_\infty - [N, N]_{T-})^{1/2} + \int_{T-}^\infty |dA_s|\} \leq c \text{ a.s.}$$

We let

$$\|M\|_{\mathcal{H}^p} = \inf_{M=N+A} j_p(N, A), \tag{1 \leq p \leq \infty \text{ or } p = \omega},$$

where the infimum is taken over all possible decompositions of M . For notational convenience we write

$$\|X\|_{\mathcal{H}^p(T)} = \|X^T\|_{\mathcal{H}^p}; \quad \|X\|_{\mathcal{H}^p(T-)} = \|X^{T-}\|_{\mathcal{H}^p}.$$

We caution the reader, however, that Emery in [2] uses the notation $\|\cdot\|_{\mathcal{H}^p(T-)}$ differently: $\|X\|_{\mathcal{H}^p(T-)}$ denotes (in [2]) the infimum of the \mathcal{H}^p norms of all semimartingales L such that $L^{T-} = X^{T-}$.

The next proposition is elementary but it may give the reader some feeling for the $\|\cdot\|_{\mathcal{H}^\omega}$ norm, which can be thought of as an extension to semimartingales of the BMO norm for martingales.

(2.1). PROPOSITION. *Let X and Y be semimartingales with $\|X\|_{\mathcal{H}^\omega} < \infty$ and $\|Y\|_{\mathcal{H}^\omega} < \infty$, and let T be a stopping time. Then the following hold:*

$$\begin{aligned} (2.2) \quad & \|\delta X_T 1_{[T, \infty[}\|_{\mathcal{H}^\omega} \leq \|X\|_{\mathcal{H}^\omega(T)} \\ (2.3) \quad & \|X\|_{\mathcal{H}^\omega(T-)} \leq 2 \|X\|_{\mathcal{H}^\omega(T)} \\ (2.4) \quad & \|X\|_{\mathcal{H}^\omega(T)} \leq \|X\|_{\mathcal{H}^\omega} \\ (2.5) \quad & \|X + Y\|_{\mathcal{H}^\omega(T-)} \leq \|X\|_{\mathcal{H}^\omega(T-)} + \|Y\|_{\mathcal{H}^\omega(T-)}. \end{aligned}$$

PROOF. Let $X = N + A$ be any decomposition of X . Then $|\delta X_T| \leq |\delta N_T| + |\delta A_T|$. For any stopping time S we have

$$|\delta X_T| 1_{\{S \leq T\}} \leq ([N, N]_T - ([N, N]_{S-})^{1/2} + \int_{S-}^T |dA_s|,$$

and (2.2) follows upon conditioning with respect to \mathcal{F}_S .

To establish (2.3) note that $\|X\|_{\mathcal{H}^\omega(T-)} = \|X - \delta X_T 1_{[T, \infty[}\|_{\mathcal{H}^\omega(T)} \leq 2 \|X\|_{\mathcal{H}^\omega(T)}$. Inequality (2.4) is clear and (2.5) is merely the triangle inequality applied to the semimartingales X^{T-} and Y^{T-} . \square

We make repeated use of the following inequalities. A proof can be found in Meyer [9].

(2.6) EMERY-MEYER INEQUALITIES. *Let $1 \leq p \leq \infty$, $1 \leq q \leq \infty$, or $1 \leq p \leq \infty$ and $q = \omega$, and $(1/p) + (1/q) = (1/r)$. Let X be predictable and M be a semimartingale such that the stochastic integral $X \cdot M$ exists. Then*

$$\begin{aligned} (2.7) \quad & \|X \cdot M\|_{\mathcal{H}^r} \leq \|X\|_{\mathcal{H}^p} \|M\|_{\mathcal{H}^q} \\ (2.8) \quad & \|X \cdot M\|_{\mathcal{H}^p} \leq h_p \|X\|_{\mathcal{H}^p} \|M\|_{\mathcal{H}^\omega}. \end{aligned}$$

If $1 \leq p \leq \infty$, $1 \leq q \leq \infty$, $(1/p) + (1/q) = (1/r)$ but $r < \infty$, then

$$(2.9) \quad \|X \cdot M\|_{\mathcal{H}^r} \leq c_r \|X\|_{\mathcal{H}^p} \|M\|_{\mathcal{H}^\omega}$$

and if $1 \leq p < \infty$, then

$$(2.10) \quad \|X \cdot M\|_{\mathcal{H}^p} \leq s_p \|X\|_{\mathcal{H}^p} \|M\|_{\mathcal{H}^\omega}.$$

where h_p, c_r, s_p are universal constants.

We record here a trivial but useful observation.

(2.11). PROPOSITION. *Let X be predictable, let M be a semimartingale and suppose the stochastic integral $X \cdot M$ exists. For any stopping times S, T with $S < T$ a.s. we have for $1 \leq p \leq \infty$, $1 \leq q \leq \infty$ and $(1/p) + (1/q) = (1/r)$*

$$\|X \cdot M - X \cdot M^S\|_{\mathcal{H}^r(T-)} \leq \|X\|_{\mathcal{H}^p(T)} \|M - M^S\|_{\mathcal{H}^q(T-)}.$$

If $1 \leq p < \infty$,

$$\|X \cdot M - X \cdot M^S\|_{\mathcal{H}^p(T-)} \leq h_p \|X\|_{\mathcal{H}^p(T)} \|M - M^S\|_{\mathcal{H}^\omega(T-)}.$$

If $1 \leq p \leq \infty$, $1 \leq q \leq \infty$, $(1/p) + (1/q) = (1/r)$ but $r < \infty$, then

$$\|X \cdot M - X \cdot M^S\|_{\mathcal{H}^r(T-)} \leq c_r \|X\|_{\mathcal{H}^p(T)} \|M - M^S\|_{\mathcal{H}^q(T-)}$$

and if $1 \leq p < \infty$, then

$$\|X \cdot M - X \cdot M^S\|_{\mathcal{S}^p(T^-)} \leq s_p \|X\|_{\mathcal{S}^p(T)} \|M - M^S\|_{\mathcal{H}^\omega(T^-)}.$$

PROOF. Let $N = (M - M^S)^{T^-}$. The proposition then follows by an application of the Emery-Meyer inequalities (2.6). \square

We shall also use a technique developed by Doléans-Dade, Meyer, and Emery, the idea of which is contained in the next definition, which was first given in Emery [1].

(2.11 $\frac{1}{2}$). DEFINITION. Let $\epsilon > 0$ and M be a semimartingale. M is said to be *carved in slices smaller than ϵ* if there exists a finite sequence of stopping times (carving times) $0 = T_0 < T_1 < \dots < T_k$ such that $M = M^{T_k^-}$, $M \in \mathcal{H}^\infty$ and for $1 \leq i \leq k$,

$$(2.12) \quad \|M - M^{T_{i-1}^-}\|_{\mathcal{H}^\infty(T_i^-)} < \epsilon.$$

We write $M \in D^\infty(\epsilon)$ if M is cut in slices smaller than ϵ . We also write $M \in D^\infty(\epsilon, k)$ to signify the number k of nonzero carving times needed to cut M into slices smaller than ϵ .

(2.13). DEFINITION. For $\epsilon > 0$ we say that M is in $D^p(\epsilon, k)$, $1 \leq p \leq \infty$ or $p = \omega$ if there exists a finite sequence of stopping times $0 = T_0 < T_1 < \dots < T_k$ such that $M = M^{T_k^-}$, $M \in \mathcal{H}^p$, and for $1 \leq i \leq k$,

$$(2.14) \quad \|M - M^{T_{i-1}^-}\|_{\mathcal{H}^p(T_i^-)} < \epsilon.$$

Note that $M \in D^q(\epsilon, k)$ implies that $M \in D^p(\epsilon, k)$ for $q \geq p$, or $q = \infty$ and $p = \omega$.

The next lemma is due to Emery ([1] or [3]).

(2.15). LEMMA. Let M be a semimartingale. For each $\epsilon > 0$ there exists an arbitrarily large stopping time T and a constant k depending on T and ϵ such that $M^{T^-} \in D^p(\epsilon, k)$, $1 \leq p \leq \infty$ and $p = \omega$.

PROOF. It suffices to prove the result for the \mathcal{H}^∞ norm since it is stronger than the \mathcal{H}^p norm, $1 \leq p \leq \infty$ or $p = \omega$. Let $M = N + A$ be a decomposition of M . By letting $C_t = \sum_{s \leq t} \delta N_s 1_{\{|\delta N_s| \geq \gamma/2\}}$ and \tilde{C}_t be its dual predictable projection (also called its ‘‘compensator’’) we have that $\tilde{N} = N_t + (C_t - \tilde{C}_t)$ is a martingale. It is a simple matter to check that C_t has locally integrable variation and that γ is a bound for the jumps of \tilde{N} . (See Meyer [8] for the details). So we assume that $M = N + A$ with γ bounding the jumps of N . Let $R_0 = 0$ and inductively define:

$$R_{k+1} = \inf\{t \geq R_k : \int_{]R_k, t]} |dA_s| \geq \gamma \text{ or } \int_0^t |dA_s| \geq k\}.$$

For each k , $A^{R_k^-} \in D^\infty(\gamma)$. Let $S_0 = 0$ and inductively define:

$$S_{k+1} = \inf\{t \geq S_k : [N, N]_t - [N, N]_{T_k} \geq \gamma^2 \text{ or } [N, N]_t \geq k\}.$$

For each k , $N^{S_k^-}$ is in \mathcal{H}^∞ and since

$$(N - N^{S_k})_{S_{k+1}^-} = (N_{S_{k+1}}^{S_{k+1}^-} - N_{S_k}^{S_k^-}) - \delta N_{S_k} 1_{]S_k, S_{k+1}^-]}$$

we have

$$\begin{aligned} \|N - N^{S_k}\|_{\mathcal{H}^\infty(S_{k+1}^-)} &\leq \|([N, N]_{S_{k+1}} - [N, N]_{S_k})^{1/2} + |\delta N_{S_k}|\|_{L^\infty} \\ &\leq \|((\delta N_{S_k})^2 + [N, N]_{S_{k+1}} - [N, N]_{S_k})^{1/2} + |\delta N_{S_k}|\|_{L^\infty} \\ &= (\gamma^2 + \gamma^2)^{1/2} + \gamma = (1 + 2)^{1/2} \gamma. \end{aligned}$$

Thus taking $\gamma = \epsilon/2(1 + 2)^{1/2}$, we have $N \in D^\infty(\epsilon/2)$ and $A \in D^\infty(\epsilon/2)$. But it is a simple matter to check that the sum of two elements of $D^\infty(\epsilon/2)$ is in $D^\infty(\epsilon)$. \square

The next lemma will be used in the proof of Lemma (3.25), which in turn is crucial to the proof of Theorem (3.4). It is important because we can have $M^n \in D^p(\alpha, k)$ for $n \geq N$ where the constant k does not depend on n . The case of interest for us is $p = \omega$.

(2.16). PROPOSITION. Fix a p , $1 \leq p \leq \infty$, or $p = \omega$ and let M be a semimartingale such that $\|M\|_{\mathcal{H}^p} < \infty$. Suppose M^n is a sequence of semimartingales such that $\lim_{n \rightarrow \infty} \|M - M^n\|_{\mathcal{H}^p} = 0$. Then for any $\epsilon > 0$ there exists an arbitrarily large stopping time T and constants N and k such that $M \in D^p(\epsilon, k)$ and $M^n \in D^p(\epsilon, k)$ for all $n > N$.

PROOF. Lemma (2.15) assures us of the existence of an arbitrarily large T such that $M \in D^p(\epsilon/2, k)$. Choose $\gamma = \epsilon/8$ and choose N so that $n > N$ implies $\|M - M^n\|_{\mathcal{H}^p} < \gamma$. Let $0 = T_0 < T_1 < \dots < T_k = T$ be the stopping times that carve M into slices. Then

$$\begin{aligned} \|M^n - (M^n)^{T_{i-1}}\|_{\mathcal{H}^p(T_{i-1})} &= \|(M^n - M) - (M^n - M)^{T_{i-1}} + (M - M^{T_{i-1}})\|_{\mathcal{H}^p(T_{i-1})} \\ &\leq 2 \|M^n - M\|_{\mathcal{H}^p(T_{i-1})} = \|(M - M^{T_{i-1}})\|_{\mathcal{H}^p(T_{i-1})} \\ &\leq 4 \|M^n - M\|_{\mathcal{H}^p} + \epsilon/2 < \epsilon. \end{aligned}$$

Thus the same times T_0, \dots, T_k carve each M^n into k slices, each smaller than ϵ , for $n > N$. \square

The most general coefficients used in scalar stochastic differential equations for which unique solutions exist are those which satisfy the conditions described in the next definition.

(2.17). DEFINITION. Let $K > 0$ and let F be an operator mapping \mathcal{C} into itself. F is said to be in $\text{Lip}(K)$ if the following two conditions are satisfied:

(2.18) For X, Y in \mathcal{C} and each stopping time T , $X^{T-} = Y^{T-}$ implies $(FX)^{T-} = (FY)^{T-}$;

(2.19) $(FX - FY)^* \leq K(X - Y)^*$ as processes, where $X_t^* = \sup_{s \leq t} |X_s|$.

We state our results in terms of local convergences. Processes X^n converge locally in a norm $\|\cdot\|$ to X if there exists a sequence of stopping time $(T_k)_{k \geq 1}$ increasing to ∞ a.s. such that for each fixed k , $\lim_{n \rightarrow \infty} \|(X^n - X)^{T_k}\| = 0$.

We remark that in [12] we used a convergence designated "weak-local", and Emery in [2] uses simply the term "local" to denote weak-local convergence. The processes X^n converge to X weak-locally if $\lim_{n \rightarrow \infty} \|(X^n - X)^{T_k}\| = 0$ for each k . We will not need this type of convergence, but see Remark (3.7), (4), in Section 3.

3. Stochastic differential equations. We consider the following type of equation:

$$(3.1) \quad X_t = J_t + \sum_{i=1}^k \int_0^t (F_i X)_s - dM_s^i$$

where M^i are semimartingales, F_i are in $\text{Lip}(K)$, and J is either a semimartingale or $J \in \mathcal{C}(1 \leq i \leq k)$. For simplicity of notation, we assume $k = 1$. We are interested only in the case $\|M\|_{\mathcal{H}^p} < \infty$. We refer the reader to Emery [2, 3] for an account of recent results concerning this equation. In particular, unique solutions exist.

Whether $J \in \mathcal{C}$ or J is a semimartingale governs what norm we use for the solution X . The semimartingale-norm $\|\cdot\|_{\mathcal{H}^p}$ is a stronger norm than the \mathcal{C} -norm $\|\cdot\|_{\mathcal{C}^p}$; however, the following lemma allows us to work exclusively with $\|\cdot\|_{\mathcal{C}^p}$ and still deduce the results for $\|\cdot\|_{\mathcal{H}^p}$, should J be a semimartingale.

(3.2). LEMMA. Suppose M, J, M^n , and J^n are all semimartingales for $n \geq 1$. Let X and X^n be

solutions respectively of

$$X_t = J_t + \int_0^t (FX)_{s-} dM_s$$

$$X_t^n = J_t^n + \int_0^t (F^n X^n)_{s-} dM_s^n$$

where F and F^n are in $\text{Lip}(K)$, some K independent of n . Suppose that (1) J^n tends to J locally in \mathcal{H}^p , (2) $F^n X$ tends to FX locally in \mathcal{S}^p , and (3) $\|M\|_{\mathcal{H}^\omega} < \infty$ and M^n tends to M locally in \mathcal{H}^ω . Then X^n tends to X locally in \mathcal{H}^p if and only if X^n tends to X locally in \mathcal{S}^p ($1 \leq p < \infty$).

PROOF. Necessity is simply a consequence of the fact that the \mathcal{H}^p norm is stronger than the \mathcal{S}^p norm. Without loss of generality we may assume that the convergence hypothesized in conditions (1), (2), and (3) above is global, not local. Further, by stopping at an arbitrarily large time T if necessary, we may assume that $\|FX_-\|_{\mathcal{S}^p} < \infty$. Then by hypothesis (2), the assumption that $\|X^n - X\|_{\mathcal{S}^p}$ tends to 0, and the inequality

$$\begin{aligned} \|F^n X^n\|_{\mathcal{S}^p} &\leq \|F^n X^n - F^n X_-\|_{\mathcal{S}^p} + \|F^n X_- - FX_-\|_{\mathcal{S}^p} + \|FX_-\|_{\mathcal{S}^p} \\ &\leq K \|X^n - X_-\|_{\mathcal{S}^p} + \|F^n X - FX\|_{\mathcal{S}^p} + \|FX_-\|_{\mathcal{S}^p}, \end{aligned}$$

we may assume $\|F^n X^n\|_{\mathcal{S}^p}$ is bounded uniformly in n . Using the Emery-Meyer inequalities we have:

$$\begin{aligned} (3.3) \quad \|X^n - X\|_{\mathcal{H}^p} &\leq \|J^n - J\|_{\mathcal{H}^p} + \|FX_- - F^n X_-\|_{\mathcal{S}^p} \|M\|_{\mathcal{H}^\omega} \\ &\quad + \|F^n X_- - F^n X^n\|_{\mathcal{S}^p} \|M\|_{\mathcal{H}^\omega} + \|F^n X^n\|_{\mathcal{S}^p} \|M - M^n\|_{\mathcal{H}^\omega} \end{aligned}$$

and since $\|F^n X_- - F^n X^n\|_{\mathcal{S}^p} \leq K \|X_- - X^n\|_{\mathcal{S}^p}$ and $\|F^n X^n\|_{\mathcal{S}^p}$ is bounded uniformly in n , the right side of (3.3) tends to 0 as n tends to ∞ . \square

In view of Lemma (3.2) we need state the next theorem, our chief result, only for the case where $J \in \mathcal{C}$.

(3.4). THEOREM. Let $M, (M^n)_{n \geq 1}$ be semimartingales. Let $F, (F^n)_{n \geq 1}$ be in $\text{Lip}(K)$ for some K and all n . Suppose, $J, (J^n)_{n \geq 1}$ are in \mathcal{C} and that $X, (X^n)_{n \geq 1}$ are solutions respectively of

$$(3.5) \quad X_t = J_t + \int_0^t (FX)_{s-} dM_s$$

$$(3.6) \quad X_t^n = J_t^n + \int_0^t (F^n X^n)_{s-} dM_s^n.$$

Fix a p ($1 \leq p < \infty$) and suppose (1) J^n converges locally in \mathcal{S}^p to J , (2) $F^n X$ converges locally in \mathcal{S}^p to FX_- and (3) $\|M\|_{\mathcal{H}^\omega} < \infty$ and M^n converges locally in \mathcal{H}^ω to M . Then X^n converges locally in \mathcal{S}^p to X .

(3.7). REMARKS. (1) Because of the local nature of stochastic integrals (cf. Meyer [7], page 307), one can replace J_t and J_t^n in (3.5) and (3.6) with $\tilde{J}_t = J_t 1_{\wedge}$ and $\tilde{J}_t^n = J_t^n 1_{\wedge}$, for $\wedge \in \mathcal{F}_0$. Then it is elementary that $t \rightarrow X_t$ and $t \rightarrow \tilde{X}_t$ (\tilde{X} is the solution of (3.5) with J replaced by \tilde{J}) agree on \wedge a.s. (cf., e.g., Protter [11], page 48). This gives us a way to handle the situation if the initial conditions J_0 and J_0^n are not in L^p : if there exist sets $\wedge_k \in \mathcal{F}_0$ increasing to Ω such that $\|(J^n - J)1_{\wedge_k}\|_{\mathcal{S}^p}$ tends to 0 for each k , one can define $T_k = k 1_{\wedge_k}$ and one concludes that the modified stopped processes $X_{t \wedge T_k}^n 1_{\{T_k > 0\}}$ converge to $X_{t \wedge T_k} 1_{\{T_k > 0\}}$ in \mathcal{S}^p for each k . Of course, the T_k increase to ∞ a.s.

(2) If we know that $J, (J^n)_{n \geq 1}$ are also semimartingales and that J^n tends locally in \mathcal{H}^p to J , then Lemma (3.2) allows us to conclude that X^n tends to X locally in \mathcal{H}^p . We also remark that local \mathcal{H}^p convergence does not in general imply local \mathcal{H}^q convergence for $q > p$. See [12], page 344 for a simple example of processes which converge in \mathcal{H}^1 but do not converge locally in \mathcal{H}^p for any $p > 1$.

(3) Neither the \mathcal{H}^ω nor the stronger \mathcal{H}^∞ norm can be used to extend Kazamaki's result: if N_t is the compensated Poisson process of intensity one and jump size α , and if T is the first jump time, then $M_t = N_{t \wedge T}$ is in \mathcal{H}^∞ and \mathcal{H}^ω , but $\mathcal{E}(M)$ is in \mathcal{H}^1 if and only if $\alpha < 1$.

(4) As will be clear from the proof, the hypothesis that $F^n X$ converges locally in \mathcal{S}^p to FX can be weakened to $F^n X$ converges weak-locally in \mathcal{S}^p to FX (or in Emery's terminology, $F^n X$ converges "locally" to FX in \mathcal{S}^p .)

(5) One need not state the convergence of the coefficients in terms of X . For example, if as in [10] we assume F^n is of the form $F^n(\omega, t, x)$ with $F^n(\cdot, 0, x) = 0$ a.s., and if we require

$$\lim_{n \rightarrow \infty} p \{ \sup_{|x| \leq m} |F^n(\cdot, t, x) - F(\cdot, t, x)| > \epsilon \} = 0$$

for each $\epsilon > 0$ and each $m \in N$, then the conclusion of Theorem (3.12) still holds.

PROOF OF THEOREM (3.4). By stopping at a large stopping time if necessary, we may assume without loss of generality that:

- (a) $\lim_{n \rightarrow \infty} \|J^n - J\|_{\mathcal{S}^p} = 0$
- (b) $\lim_{n \rightarrow \infty} \|F^n X - FX\|_{\mathcal{S}^p} = 0$
- (c) $\lim_{n \rightarrow \infty} \|M^n - M\|_{\mathcal{H}^\omega} = 0.$

We choose and fix an α such that $0 < \alpha < 1/s_p K$. We know (Lemma (2.15)) that there exists an arbitrarily large stopping time T such that $M^{T-} \in D^\omega(\alpha/2, k)$ for some $k < \infty$. By Proposition (2.16) and reduction (c) above we know there exists an N such that for all $n > N$ we have $(M^n)^{T-} \in D^\omega(\alpha, k)$, where k does not depend on n . Let T be such a stopping time, and let $0 = T_0 < T_1 < \dots < T_k = T$ be the times that carve M (and also M^n for $n > N$) into slices less than α . We then have

- (d) $M^{T-} \in D(\alpha/2, k)$ and $(M^n)^{T-} \in D(\alpha, k)$
for all $n > N, k < \infty$, and $0 < \alpha < 1/s_p K$.

Observe that for any stopping time R we have that:

$$\begin{aligned} (3.8) \quad \|F^n X^n\|_{\mathcal{S}^p(R-)} &\leq \|F^n X^n - F^n O_-\|_{\mathcal{S}^p(R)} + \|F^n O_-\|_{\mathcal{S}^p(R)} \\ &+ \|F^n X_- - FX_-\|_{\mathcal{S}^p(R)} + \|FX_- - FO_-\|_{\mathcal{S}^p(R)} + \|FO_-\|_{\mathcal{S}^p(R)} \\ &\leq K \|X^n\|_{\mathcal{S}^p(R)} + 2K \|X_-\|_{\mathcal{S}^p(R)} + \|F^n X_- - FX_-\|_{\mathcal{S}^p(R)} + \|FO_-\|_{\mathcal{S}^p(R)}. \end{aligned}$$

Since FO_- is left continuous we let $R_n = \inf\{t > 0: |FO| \geq n\}$ and $\lim_{n \rightarrow \infty} R_n = \infty$ a.s. and $\|FO_-\|_{\mathcal{S}^p(R_n)} \leq n$. Thus without loss of generality we can assume

- (e) $\|FO_-\|_{\mathcal{S}^p} < \infty.$

Since $\|F^n X_- - FX_-\|_{\mathcal{S}^p}$ tends to 0 by reduction (b), it is bounded, and hence (3.8) implies that $\|F^n X^n\|_{\mathcal{S}^p(R-)}$ is bounded if $\|X\|_{\mathcal{S}^p(R-)} < \infty$ and $\sup_n \|X^n\|_{\mathcal{S}^p(R-)} < \infty$. That there exists an arbitrarily large stopping time R such that $\|X\|_{\mathcal{S}^p(R)} < \infty$ and $\sup_n \|X^n\|_{\mathcal{S}^p(R)} < \infty$ is perhaps the crux of proof. Given (d) this fact is the content of Lemma (3.25), the statement and proof of which follow this proof. Thus by (e), inequality (3.8), and Lemma (3.25) we may assume without loss of generality that

- (f) $\sup_n \|F^n X^n\|_{\mathcal{S}^p} < \infty.$

From equations (3.4) and (3.5) we have that

$$(3.9) \quad X_t^n - X_t = J_t^n - J_t + \int_0^t (F^n X_{s-} - F X_{s-}) dM_s + \int_0^t (F^n X_{s-}^n - F^n X_{s-}) dM_s + \int_0^t F^n X_{s-}^n d(M^n - M)_s.$$

By the Emery-Meyer inequalities and Proposition (2.11) we have for any stopping time $R > 0$ a.s. that equation (3.9) yields:

$$(3.10) \quad \|X^n - X\|_{\mathcal{S}^p(R-)} \leq \|J^n - J\|_{\mathcal{S}^p} + s_p \|F^n X_- - F X_-\|_{\mathcal{S}^p} \|M\|_{\mathcal{H}^\omega} + s_p K \|X^n - X\|_{\mathcal{S}^p(R-)} \|M\|_{\mathcal{H}^\omega(R-)} + s_p \|F^n X^n\|_{\mathcal{S}^p} \|M^n - M\|_{\mathcal{H}^\omega} = \gamma_n + s_p K \|M\|_{\mathcal{H}^\omega(R-)} \|X^n - X\|_{\mathcal{S}^p(R-)}$$

where s_p is given in (2.11). Note that by reductions (a), (b), (c) and (f) we know that $\lim_{n \rightarrow \infty} \gamma_n = 0$.

Let $0 = T_0 < T_1 \dots T_k = T$ be the ‘‘carving times’’ whose existence is assured in (d). Then for $R = T_1$ and letting $\beta(1, n) = \gamma_n$, inequality (3.10) yields:

$$\|X^n - X\|_{\mathcal{S}^p(T_1-)} \leq \beta(1, n) + r \|X^n - X\|_{\mathcal{S}^p(T_1-)}$$

where $r = s_p K \alpha < 1$. Subtraction yields

$$(3.11) \quad \|X^n - X\|_{\mathcal{S}^p(T_1-)} \leq \beta(1, n)/(1 - r).$$

Recall that $\delta X_t = X_t - X_{t-}$, the jump at t . One easily sees that:

$$(3.12) \quad \begin{aligned} \|\delta X_{T_1}^n - \delta X_{T_1}\|_{L^p} &\leq \|\delta J_{T_1}^n - \delta J_{T_1}\|_{L^p} \\ &+ \|F^n X - F X\|_{\mathcal{S}^p(T_1-)} \|\delta M_{T_1} 1_{[T_1, \infty[}\|_{\mathcal{H}^\omega} \\ &+ K \|X^n - X\|_{\mathcal{S}^p(T_1-)} \|\delta M_{T_1} 1_{[T_1, \infty[}\|_{\mathcal{H}^\omega} \\ &+ \|F^n X_{T_1-}^n\|_{\mathcal{S}^p(T_1-)} \|\delta(M^n - M)_{T_1} 1_{[T_1, \infty[}\|_{\mathcal{H}^\omega} \end{aligned}$$

and since $\|\delta M_{T_1} 1_{[T_1, \infty[}\|_{\mathcal{H}^\omega} \leq \|M\|_{\mathcal{H}^\omega}$ (Proposition (2.1)), combining (3.11) and (3.12) gives us that

$$\lim_{n \rightarrow \infty} \|X^n - X\|_{\mathcal{S}^p(T_1)} = 0.$$

Now suppose we have established that $\lim_{n \rightarrow \infty} \|X^n - X\|_{\mathcal{S}^p(T_i)} = 0$, and consider T_{i+1} . We have that X_t and X_t^n are solutions respectively of (3.4) and (3.5) with J_t and J_t^n replaced respectively with $X_{t \wedge T_i} + J_t - J_{t \wedge T_i}$ and $X_{t \wedge T_i}^n + J_t^n - J_{t \wedge T_i}^n$ and with M and M^n replaced respectively with $M - M^{T_i}$ and $M^n - (M^n)^{T_i}$. Inequality (3.10) then gives us, taking $R = T_{i+1}$,

$$\begin{aligned} \|X^n - X\|_{\mathcal{S}^p(T_{i+1}-)} &\leq \beta(i, n) + s_p K \|M - M^{T_i}\|_{\mathcal{H}^\omega(T_{i+1}-)} \|X^n - X\|_{\mathcal{S}^p(T_{i+1}-)} \\ &\leq \beta(i, n) + r \|X^n - X\|_{\mathcal{S}^p(T_{i+1}-)} \end{aligned}$$

where $r = s_p K \alpha < 1$. Thus

$$(3.13) \quad \|X^n - X\|_{\mathcal{S}^p(T_{i+1}-)} \leq \beta(i, n)/(1 - r)$$

where $\lim_{n \rightarrow \infty} \beta(i, n) = 0$. A calculation analogous to (3.12) shows that

$$\lim_{n \rightarrow \infty} \|X^n - X\|_{\mathcal{S}^p(T_{i+1})} = 0,$$

and hence by a (finite) induction we conclude that $\lim_{n \rightarrow \infty} \|X^n - X\|_{\mathcal{S}^p(T)} = 0$ and the proof is complete. \square

The next two lemmas were used in the proof of Theorem (3.4). Lemma (3.14) has some

interest in its own right, since it gives a bound on the \mathcal{S}^p norm of the solution of a stochastic differential equation in terms of other norms which are (at least theoretically) known.

(3.14). LEMMA. *Let M be a semimartingale, let $F \in \text{Lip}(K)$ and let $J \in \mathcal{C}$. Let X be the (unique) solution of*

$$(3.15) \quad Z_t = J_t + \int_0^t (FZ)_{s-} dM_s.$$

Suppose that (1) $\|M\|_{\mathcal{H}^\omega} = m < \infty$, (2) $\|J\|_{\mathcal{S}^p} = j < \infty$, and (3) $\|FO_-\|_{\mathcal{S}^p} = \tau < \infty$. Let α be a constant such that $0 < \alpha < 1/s_p K$, where s_p is given in (2.10). Let T be a stopping time such that $M^{T^-} \in D^\omega(\alpha, k)$. Then

$$\|X\|_{\mathcal{S}^p(T)} \leq C < \infty$$

where $C = C(p, j, K, \tau, m, \alpha, k)$ is a constant depending on the seven parameters in its argument.

PROOF. First suppose that $\|M\|_{\mathcal{H}^\omega} \leq \alpha$. Then by the Emery-Meyer inequalities (2.6) we have

$$(3.16) \quad \begin{aligned} \|X\|_{\mathcal{S}^p} &\leq \|J\|_{\mathcal{S}^p} + s_p \|FX_-\|_{\mathcal{S}^p} \|M\|_{\mathcal{H}^\omega} \\ &\leq j + s_p K \alpha (\|X\|_{\mathcal{S}^p} + \|FO_-\|_{\mathcal{S}^p}) \\ &\leq j + r\tau + r \|X\|_{\mathcal{S}^p} \end{aligned}$$

where $0 < r = s_p K \alpha < 1$. Then by subtraction we have

$$(3.17) \quad \|X\|_{\mathcal{S}^p} \leq (j + r\tau)/(1 - r)$$

provided that $\|X\|_{\mathcal{S}^p} < \infty$. We show this by successive approximation. Let $X_t^0 = J_t$ and recursively define

$$X_t^{n+1} = J_t + \int_0^t (FX^n)_{s-} dM_s.$$

Then clearly $\|X^n\|_{\mathcal{S}^p} < \infty$ for each n using induction and the Emery-Meyer inequalities. Moreover, one easily checks that for $l \geq 1$:

$$(3.18) \quad \|X^l - X^0\|_{\mathcal{S}^p} \leq (j + \tau) \sum_{i=1}^l r^i$$

where $0 < r = s_p K \alpha < 1$. Fix a large N . Then for $m > n > N$ we have:

$$(3.19) \quad \|X^m - X^n\|_{\mathcal{S}^p} \leq r^N \|X^{m-N} - X^{n-N}\|_{\mathcal{S}^p}$$

and combining (3.18) and (3.19) yields

$$(3.20) \quad \|X^m - X^n\|_{\mathcal{S}^p} \leq r^N [2(j + \tau)/(1 - r)]$$

for $m > n > N$, and the right side of (3.20) tends to 0 as $N \rightarrow \infty$. Thus $(X^n)_{n \geq 1}$ is Cauchy in the Banach space $(\mathcal{C}, \|\cdot\|_{\mathcal{S}^p})$. It is a simple matter to check that X^n converges in $\|\cdot\|_{\mathcal{S}^p}$ to a solution Z of (3.15), and then $Z = X$ by the uniqueness of the solution.

Now remove the assumption that $\|M\|_{\mathcal{H}^\omega} \leq \alpha$. We know that $M^{T^-} \in D^\omega(\alpha, k)$. Let $0 = T_0 < T_1 < \dots < T_k = T$ be the "carving times" of Definition (2.13). Let $N^1 = M^{T_1^-}$. By the unicity of solutions we know $X_t^{T_1^-}$ is the unique solution of

$$Z_t = J_t + \int_0^t (FX)_{s-} dN_s^1.$$

The preceding reduction gives

$$(3.21) \quad \|X\|_{\mathcal{S}^p(T_1^-)} \leq C(p, j, K, \tau, \alpha) < \infty.$$

Since $X_{T_1} = X_{T_1-} + \delta J_{T_1} + FX_{T_1-} - \delta M_{T_1}$, we have that

$$(3.22) \quad \|X\|_{\mathcal{S}^p(T_1)} \leq \|X\|_{\mathcal{S}^p(T_1-)} + 2j + s_p m K \|X\|_{\mathcal{S}^p(T_1-)} + s_p m K \tau.$$

Inequalities (3.21) and (3.22) imply

$$(3.23) \quad \|X\|_{\mathcal{S}^p(T_1)} \leq C(p, j, K, \tau, m, \alpha) < \infty.$$

Now set $N_t^{i+1} = M_t^{T_{i+1}-} - M_t^{T_i}$ for $1 \leq i \leq k - 1$, and note that $X_t^{T_{i+1}-}$ is the solution of the following equation:

$$Z_t = X_{t \wedge T_i} + (J_t^{T_{i+1}-} - J_t^{T_i}) + \int_0^t (FZ)_s - dN_s^{i+1}.$$

Since $\|N^{i+1}\|_{\mathcal{H}^\omega} \leq \alpha$ we get as in (3.17) that

$$(3.24) \quad \|X^{T_{i+1}-}\|_{\mathcal{S}^p} \leq (\|X\|_{\mathcal{S}^p(T_i)} + 2j + r\tau)/(1 - r)$$

and since

$$X_{T_{i+1}} = X_{T_{i+1}-} + \delta J_{T_{i+1}} + FX_{T_{i+1}-} - \delta M_{T_{i+1}}$$

it follows as in (3.21) through (3.23) that

$$\|X_{T_{i+1}}\| \leq C(p, j, K, \tau, m, \alpha, i + 1) < \infty.$$

Since this is true for $1 \leq i \leq k$ and $T_k = T$, we have the result. \square

(3.25). LEMMA. Let $M, (M^n)_{n \geq 1}$ be semimartingales; let $J, (J^n)_{n \geq 1}$ be in \mathcal{C} ; let $F, (F^n)_{n \geq 1}$ be in $\text{Lip}(K)$. Let $X, (X^n)_{n \geq 1}$ be solutions respectively of:

$$X_t = J_t + \int_0^t (FX)_s - dM_s$$

$$X_t^n = J_t^n + \int_0^t (F^n X^n)_s - dM_s^n.$$

Assume that (1) $\|M\|_{\mathcal{H}^\omega} < \infty$ and $\lim_{n \rightarrow \infty} \|M^n - M\|_{\mathcal{H}^\omega} = 0$; (2) $\|J\|_{\mathcal{S}^p} < \infty$ and $\lim_{n \rightarrow \infty} \|J^n - J\|_{\mathcal{S}^p} = 0$; and (3) $\lim_{n \rightarrow \infty} \|F^n X - FX\|_{\mathcal{S}^p} = 0$. Let T be a stopping time such that $\|FO_-\|_{\mathcal{S}^p(T)} < \infty$ and $M^{T-} \in D^\omega(\alpha/2, k)$ for some $\alpha, 0 < \alpha < 1/s_p K$, where s_p is given in (2.10). Then $\inf_N \sup_{n > N} \|X^n\|_{\mathcal{S}^p(T)} < \infty$.

PROOF. By Proposition (2.16) we know that there exists an N such that for $n > N$ we have $M^n \in D^\omega(\alpha, k)$ with k not depending on n . We can assume without loss of generality that $\|M^n\|_{\mathcal{H}^\omega} \leq 2m$ for $n > N$. Note that

$$\|F^n O_-\|_{\mathcal{S}^p(T)} \leq \|F^n O - F^n X\|_{\mathcal{S}^p(T-)} + \|F^n X - FX\|_{\mathcal{S}^p(T-)} + \|FX - FO\|_{\mathcal{S}^p(T-)}$$

$$+ \|FO\|_{\mathcal{S}^p(T-)},$$

$$\leq 2K \|X\|_{\mathcal{S}^p(T-)} + \|F^n X - FX\|_{\mathcal{S}^p(T-)} + \|FO\|_{\mathcal{S}^p(T-)}$$

hence $\sup_{n > N} \|F^n O_-\|_{\mathcal{S}^p(T)} = \tau < \infty$ provided that $\|X\|_{\mathcal{S}^p(T-)} < \infty$. But $\|X\|_{\mathcal{S}^p(T)} < \infty$ by Lemma (3.14). Applying Lemma (3.14) to X^n as well, we see that for $n > N$:

$$\|X^n\|_{\mathcal{S}^p(T)} \leq C(p, j_n, K, \tau, 2m, \alpha, k)$$

where $j_n = \|J^n\|_{\mathcal{S}^p(T)}$, which is bounded as well by hypothesis (2). This completes the proof. \square

4. An amusing application. Recently Garcia, Maillard, and Peltraut [4] have shown that, given a totally inaccessible stopping time T and a constant K , there exists a martingale L with precisely one jump (at T) such that $L_T/L_{T-} = K$. We extend this result in Lemma (4.2) to allow random jumps in L^1 and \mathcal{F}_{T-} measurable, and in Theorem (4.4) we establish a “continuity” result for local martingales with (random) “multiplicative jumps”.

Let T denote a finite totally inaccessible stopping time, let $A_t = 1_{[T, \infty)}$, and let (M_t) denote the BMO martingale $M_t = A_t - \tilde{A}_t$, where \tilde{A} is the dual predictable projection of A (also called the compensator of A). Let $\wedge \in L^1(\mathcal{F}_{T-})$, and $N_t = E\{\wedge - 1 \mid \mathcal{F}_t\}$. Define F mapping \mathcal{C} to \mathcal{C} by $(FC)_t = N_t C_t$. Suppose X is the unique solution of

$$(4.1) \quad X_t = 1 + \int_0^t FX_s - dM_s.$$

Then X is the stochastic exponential of the local martingale $Y_t = \int_0^t N_s - dM_s$, and so

$$\frac{X_T}{X_{T-}} = 1 + N_{T-}(M_T - M_{T-})$$

which implies that $X_T/X_{T-} = \wedge$.

(4.2). LEMMA. *The local martingale X of (4.1) is locally in \mathcal{H}^p for $1 < p < \infty$.*

PROOF. Let $S^l = \inf\{t: |N_t| \geq l\} 1_{\{|N_0| \leq l\}}$. Define F^l by $F^l C = (FC)^{S^l}$. Then $F^l \in \text{Lip}(l)$. Since M is in BMO, it is also in \mathcal{H}^∞ when considered as a semimartingale. By Lemma (2.15) we can find a stopping time T^l with $P(T^l < S^l) < 1/2^l$ and such that (1) $M^{T^l} \in D^\infty(\alpha, k)$ for some $k < \infty$ and $0 < \alpha < 1/s_p l$; and (2) $\|F^l O_{-}\|_{\mathcal{S}^p(T^l)} < \infty$. Thus by Lemma (3.14) we conclude

$$\|X\|_{\mathcal{S}^p(T^l)} < \infty.$$

Since $\lim_{l \rightarrow \infty} S^l = \infty$ a.s., also $\lim_{l \rightarrow \infty} T^l = \infty$, and the proof is complete. \square

Let \wedge^n be a sequence of random variables in $L^1(\mathcal{F}_{T-})$ and let $N_t^n = E\{\wedge^n - 1 \mid \mathcal{F}_t\}$. Define $(F^n C)_t = N_t^n C_t$ for $C \in \mathcal{C}$. Suppose X^n are solutions of

$$(4.3) \quad X_t^n = 1 + \int_0^t F^n X_s^n - dM_s \quad n \geq 1.$$

Then each X^n has multiplicative jump of size \wedge^n at time T .

(4.4). THEOREM. *If \wedge^n converges to \wedge in $L^q(1 < q < \infty)$ with $|\wedge^n| \leq Y \in L^1$ for $n \geq 1$, then X^n converges locally in $\mathcal{H}^p(1 \leq p < q)$ to X .*

PROOF. Let $Y_t = E(Y \mid \mathcal{F}_t)$ and let $S^k = \inf\{t > 0: Y_t \geq k\}$. Then by stopping M at S^k we can assume without loss of generality that F and $F^n \in \text{Lip}(k)$ for all n . By Hölder's inequality we have $\|F^n X - FX\|_{\mathcal{S}^p} = \|(N^n - N)X\|_{\mathcal{S}^p} \leq \|N^n - N\|_{\mathcal{S}^q} \|X\|_{\mathcal{S}^r}$ for $(1/q) + (1/r) = (1/p)$. By Lemma (3.14) we know there exists an arbitrarily large stopping time R such that $\|X\|_{\mathcal{S}^r(R)} < \infty$. By Doob's inequality $\|N^n - N\|_{\mathcal{S}^q} \leq q(q-1)^{-1} \|\wedge^n - \wedge\|_{\mathcal{S}^q}$ which tends to 0 as n tends to ∞ . An application of Theorem (3.4) gives the result. \square

Acknowledgment. I wish to thank C. Doléans-Dade for her careful reading of an earlier

version of this paper. Her comments have led (I hope) to a much more readable and correct version. She also pointed out a gap in the proof of Theorem (3.4) which is now filled.

REFERENCES

- [1] EMERY, M. (1978). Stabilité des solutions des Equations Différentielles Stochastiques: Applications aux Intégrales Multiplicatives Stochastiques. *Z. Wahrscheinlichkeitstheorie und verw. Gebiete* **41** 241–262.
- [2] EMERY, M. (1979). Une Topologie sur L'Espace des Semimartingales. *Séminaire de Probabilités XIII. Springer Lecture Notes in Math.* **721** 260–280.
- [3] EMERY, M. (1979). Equations Différentielles Lipschitziennes: La Stabilité. *Séminaire de Probabilités XIII. Springer Lecture Notes in Math.* **721** 281–293.
- [4] GARCIA, M., MAILLARD, P. AND PELTRAUT, Y. (1978). Une Martingale de Saut Multiplicative Donné. *Springer Lecture Notes in Math.* **649** 51–52.
- [5] KAZAMAKI, N. (1980). A property of BMO-martingales. *Toyama Math. Report*. To appear.
- [6] KAZAMAKI, N. AND SEKIGUCHI, T. (1979). On the transformation of some classes of martingales by a change of law. *Tôhoku Math. J.* **31** 261–279.
- [7] MEYER, P. A. (1976). Un Cours sur les Intégrales Stochastiques. *Springer Lecture Notes in Math.* **511** 245–400.
- [8] MEYER, P. A. (1977). Le Théorème Fondamental sur les Martingales Locales. *Springer Lecture Notes in Math.* **581** 463–464.
- [9] MEYER, P. A. (1978). Inégalités de Normes pour les Intégrales Stochastiques. *Springer Lecture Notes in Math.* **649**, 757–762.
- [10] PROTTER, P. (1977a). On the existence, uniqueness, convergence and explosions of solutions of systems of stochastic integral equations. *Ann. Probability.* **5** 243–261.
- [11] PROTTER, P. (1977b). Markov solutions of stochastic differential equations. *Z. Wahrscheinlichkeitstheorie und verw. Gebiete* **41** 39–58.
- [12] PROTTER, P. (1978). H^p stability of solutions of stochastic differential equations. *Z. Wahrscheinlichkeitstheorie und verw. Gebiete* **44** 337–352.

DEPARTMENTS OF STATISTICS AND MATHEMATICS
PURDUE UNIVERSITY
W. LAFAYETTE, INDIANA 47907