

Reasonable Ultrafilters, Again

Andrzej Rosłanowski and Saharon Shelah

Abstract We continue investigations of *reasonable ultrafilters* on uncountable cardinals defined in previous work by Shelah. We introduce stronger properties of ultrafilters and we show that those properties may be handled in λ -support iterations of reasonably bounding forcing notions. We use this to show that consistently there are reasonable ultrafilters on an inaccessible cardinal λ with generating systems of size less than 2^λ . We also show how ultrafilters generated by small systems can be killed by forcing notions which have enough reasonable completeness to be iterated with λ -supports.

1 Introduction

Reasonable ultrafilters were introduced in Shelah [12] in order to suggest a line of research that would repeat in some sense the beautiful theory created around the notion of *P-points on ω* . Most of the generalizations of P-points to uncountable cardinals in the literature go in the direction of normal ultrafilters and large cardinals (see, e.g., Gitik [3]), but one may be interested in the opposite direction. If one wants to keep away from *normal ultrafilters on λ* , one may declare interest in ultrafilters which do not include some clubs and even demand that quotients by a closed unbounded subset of λ do not extend the club filter of λ . Such ultrafilters are called *weakly reasonable ultrafilters* (see Definition 2.1, Observation 2.2). But if we are interested in generalizing P-points, we have to consider also properties that would correspond to *any countable family of members of the ultrafilter has a pseudo-intersection in the ultrafilter*. The choice of the right property in the declared context of *very nonnormal ultrafilters* is not clear, and one of the goals of the present paper is to show that the *very reasonable ultrafilters* suggested in Shelah [12] (see Definition 2.3 here) are very reasonable indeed; that is, we may prove interesting theorems on them.

In Section 2 we recall some of the concepts and results presented in Shelah [12] and we introduce strong properties of generating systems (super and strong reasonability, see Definitions 2.11, 2.12) and we show that there may exist super reasonable systems which generate ultrafilters (Propositions 2.15, 2.16).

In Section 3 we recall from [7] some properties of forcing notions relevant for λ -support iterations. We also improve in some sense a result of [7] and we show a preservation theorem for *the nice double \mathbf{a} -bounding property* (Theorem 3.13).

Then in Section 4 we show that super reasonable families generating ultrafilters will be still at least strongly reasonable and will continue to generate ultrafilters after forcing with λ -support iterations of \mathbf{A} -bounding forcing notions. Therefore, for an inaccessible cardinal λ , it is consistent that $2^\lambda = \lambda^{++}$ and there is a very reasonable ultrafilter generated by a system of size λ^+ (Corollary 4.5). It should be stressed that “generating an ultrafilter” has the specific meaning stated in Definition 2.3(3). In particular, “having a small generating system” *does not* imply “having small ultrafilter base.”

Section 5 shows that some technical inconveniences of the proofs from Section 4 reflect the delicate nature of our concepts, not necessarily our lack of knowledge. We give an example of a nicely double \mathbf{a} -bounding forcing notion which kills ultrafilters generated by systems from the ground model. Then we show that for an inaccessible cardinal λ , it is consistent that $2^\lambda = \lambda^{++}$ and there is no ultrafilter generated by a system of size λ^+ (see Corollary 4.5).

Studies of ultrafilters generated according to the schema introduced in [12] are also carried out in Rosłanowski and Shelah [9].

Notation 1.1 Our notation is rather standard and compatible with that of classical textbooks (such as Jech [5]). In forcing we keep the older convention that *a stronger condition is the larger one*.

1. Ordinal numbers will be denoted by the lowercase initial letters of the Greek alphabet ($\alpha, \beta, \gamma, \delta \dots$) and also by i, j (with possible sub- and superscripts). Cardinal numbers will be called κ, λ, μ (with possible sub- and superscripts). λ is always assumed to be regular, sometimes even strongly inaccessible. By χ we will denote a sufficiently large regular cardinal; $\mathcal{H}(\chi)$ is the family of all sets hereditarily of size less than χ . Moreover, we fix a well ordering $<_\chi^*$ of $\mathcal{H}(\chi)$.
2. A sequence is a function with the domain being a set of ordinals. For two sequences η, ν we write $\nu \triangleleft \eta$ whenever ν is a proper initial segment of η , and $\nu \trianglelefteq \eta$ when either $\nu \triangleleft \eta$ or $\nu = \eta$. The length of a sequence η is the order type of its domain and it is denoted by $\text{lh}(\eta)$.
3. We will consider several games of two players. One player will be called GENERIC or COMPLETE or just COM, and we will refer to this player as “she”. Her opponent will be called ANTIGENERIC or INCOMPLETE or just INC and will be referred to as “he”.
4. For a forcing notion \mathbb{P} , all \mathbb{P} -names for objects in the extension via \mathbb{P} will be denoted with a tilde below (e.g., $\tilde{\tau}, \tilde{X}$). The canonical \mathbb{P} -name for the generic filter in \mathbb{P} is called $\tilde{G}_{\mathbb{P}}$. The weakest element of \mathbb{P} will be denoted by $\emptyset_{\mathbb{P}}$ (and we will always assume that there is one, and that there is no other condition equivalent to it). We will also assume that all forcing notions under consideration are atomless.

By “ λ -support iterations” we mean iterations in which domains of conditions are of size $\leq \lambda$. However, we will pretend that conditions in a λ -support iteration $\bar{\mathbb{Q}} = \langle \mathbb{P}_\zeta, \mathbb{Q}_\zeta : \zeta < \zeta^* \rangle$ are total functions on ζ^* and for $p \in \text{lim}(\bar{\mathbb{Q}})$ and $\alpha \in \zeta^* \setminus \text{Dom}(p)$ we will let $p(\alpha) = \emptyset_{\mathbb{Q}_\alpha}$.

5. For a filter D on λ , the family of all D -positive subsets of λ is called D^+ . (So $A \in D^+$ if and only if $A \subseteq \lambda$ and $A \cap B \neq \emptyset$ for all $B \in D$.) The club filter of λ is denoted by \mathcal{D}_λ .

2 More Reasonable Ultrafilters on λ

Here we recall some basic definitions and results from [12], and then we introduce even stronger properties of ultrafilters and/or generating systems. We also show that assumptions like $\diamond_{S_\lambda^+}$ imply the existence of such objects.

As explained in the introduction, we are interested in ultrafilters (on an uncountable cardinal λ) which are far from being normal. Weakly reasonable ultrafilters defined below do not contain some clubs even if we look at their quotients by a club.

Definition 2.1 ([12], Definition 1.4) We say that a uniform ultrafilter D on λ is *weakly reasonable* if for every function $f \in {}^\lambda \lambda$ there is a club C of λ such that

$$\bigcup \{[\delta, \delta + f(\delta)) : \delta \in C\} \notin D.$$

Observation 2.2 ([12], Observation 1.5) Let D be a uniform ultrafilter on λ . Then the following conditions are equivalent:

- (A) D is weakly reasonable,
- (B) for every increasing continuous sequence $\langle \delta_\xi : \xi < \lambda \rangle \subseteq \lambda$ there is a club C^* of λ such that

$$\bigcup \{[\delta_\xi, \delta_{\xi+1}) : \xi \in C^*\} \notin D.$$

We want to investigate ultrafilters on λ which are generated by systems defining “largeness in λ ” by giving a condition based on “largeness in intervals below λ .” The family \mathbb{Q}_λ^0 introduced below is a natural generalization of the approach used in [6, Sections 5, 6]. The directness of G^* is an easy way to guarantee that $\text{fil}(G^*)$ is a filter, and $(<\lambda^+)$ -directness has the flavor of P -pointness.

Definition 2.3 ([12], Definition 2.5)

1. Let \mathbb{Q}_λ^0 consist of all tuples

$$p = (C^p, \langle Z_\delta^p : \delta \in C^p \rangle, \langle d_\delta^p : \delta \in C^p \rangle)$$

such that

- (i) C^p is a club of λ consisting of limit ordinals only, and for $\delta \in C^p$:
 - (ii) $Z_\delta^p = [\delta, \min(C^p \setminus (\delta + 1)))$ and
 - (iii) $d_\delta^p \subseteq \mathcal{P}(Z_\delta^p)$ is a proper nonprincipal ultrafilter on Z_δ^p .
2. For $q \in \mathbb{Q}_\lambda^0$ we let

$$\text{fil}(q) \stackrel{\text{def}}{=} \{A \subseteq \lambda : (\exists \varepsilon < \lambda)(\forall \delta \in C^q \setminus \varepsilon)(A \cap Z_\delta^q \in d_\delta^q)\},$$

and for a set $G^* \subseteq \mathbb{Q}_\lambda^0$ we let $\text{fil}(G^*) \stackrel{\text{def}}{=} \bigcup \{\text{fil}(p) : p \in G^*\}$. We also define a binary relation \leq^0 on \mathbb{Q}_λ^0 by

$$p \leq^0 q \quad \text{if and only if} \quad \text{fil}(p) \subseteq \text{fil}(q).$$

3. We say that an ultrafilter D on λ is *reasonable* if it is weakly reasonable (see Definition 2.1) and there is a directed (with respect to \leq^0) set $G^* \subseteq \mathbb{Q}_\lambda^0$ such that $D = \text{fil}(G^*)$. The family G^* may be called *the generating system for D* .
4. An ultrafilter D on λ is said to be *very reasonable* if it is weakly reasonable and there is a $(<\lambda^+)$ -directed (with respect to \leq^0) set $G^* \subseteq \mathbb{Q}_\lambda^0$ such that $D = \text{fil}(G^*)$.

Definition 2.4 Suppose that

- (a) X is a nonempty set and e is an ultrafilter on X ,
- (b) d_x is an ultrafilter on a set Z_x (for $x \in X$).

We let

$$\bigoplus_{x \in X}^e d_x = \{A \subseteq \bigcup_{x \in X} Z_x : \{x \in X : Z_x \cap A \in d_x\} \in e\}.$$

(Clearly, $\bigoplus_{x \in X}^e d_x$ is an ultrafilter on $\bigcup_{x \in X} Z_x$.)

Proposition 2.5 ([12], Proposition 2.9) *Let $p, q \in \mathbb{Q}_\lambda^0$. Then the following are equivalent:*

- (a) $p \leq^0 q$,
- (b) *there is $\varepsilon < \lambda$ such that*

$$(\forall \alpha \in C^q \setminus \varepsilon)(\forall A \in d_\alpha^q)(\exists \beta \in C^p)(A \cap Z_\beta^p \in d_\beta^p),$$

- (c) *there is $\varepsilon < \lambda$ such that if $\alpha \in C^q \setminus \varepsilon$, $\beta_0 = \sup(C^p \cap (\alpha + 1))$, $\beta_1 = \min(C^p \setminus \min(C^q \setminus (\alpha + 1)))$, then there is an ultrafilter e on $[\beta_0, \beta_1) \cap C^p$ such that*

$$d_\alpha^q = \{A \cap Z_\alpha^q : A \in \bigoplus_{\beta \in [\beta_0, \beta_1) \cap C^p}^e \{d_\beta^p\}\}.$$

Observation 2.6 (Compare [12], Proposition 2.3(4)) If $p \in \mathbb{Q}_\lambda^0$, $A \subseteq \lambda$, then there is $q \in \mathbb{Q}_\lambda^0$ such that $p \leq^0 q$ and either $A \in \text{fil}(q)$ or $\lambda \setminus A \in \text{fil}(q)$.

Definition 2.7 ([12], Definition 2.10) Let $p \in \mathbb{Q}_\lambda^0$. Suppose that $X \in [C^p]^\lambda$ and $C \subseteq C^p$ is a club of λ such that

$$\text{if } \alpha < \beta \text{ are successive elements of } C, \text{ then } |[\alpha, \beta) \cap X| = 1.$$

(In this situation we say that p is *restrictable to $\langle X, C \rangle$* .) We define *the restriction of p to $\langle X, C \rangle$* as an element $q = p \upharpoonright \langle X, C \rangle \in \mathbb{Q}_\lambda^0$ such that $C^q = C$, and if $\alpha < \beta$ are successive elements of C , $x \in [\alpha, \beta) \cap X$, then $Z_\alpha^q = [\alpha, \beta)$ and $d_\alpha^q = \{A \subseteq Z_\alpha^q : A \cap Z_x^p \in d_x^p\}$.

Proposition 2.8 ([12], Proposition 2.11)

1. *If $G^* \subseteq \mathbb{Q}_\lambda^0$ is \leq^0 -directed and $|G^*| \leq \lambda$, then G^* has a \leq^0 -upper bound. (Hence, in particular, $\text{fil}(G^*)$ is not an ultrafilter.)*
2. *Assume that $G^* \subseteq \mathbb{Q}_\lambda^0$ is \leq^0 -directed and \leq^0 -downward closed, $p \in G^*$, $X \in [C^p]^\lambda$, and $C \subseteq C^p$ is a club of λ such that p is restrictable to $\langle X, C \rangle$. If $\bigcup_{x \in X} Z_x^p \in \text{fil}(G^*)$, then $p \upharpoonright \langle X, C \rangle \in G^*$.*

The following definition is used here to simplify our notation in Definition 2.11 only. However, these concepts play a more central role in [9].

Definition 2.9

1. Let \mathbb{Q}_λ^* be the family of all sets r such that
 - (a) members of r are triples (α, Z, d) such that $\alpha < \lambda$, $Z \subseteq [\alpha, \lambda)$, $\aleph_0 \leq |Z| < \lambda$, and d is a nonprincipal ultrafilter on Z ,
 - (b) $(\forall \zeta < \lambda)(|\{(\alpha, Z, d) \in r : \alpha = \zeta\}| < \lambda)$, and $|r| = \lambda$.
 For $r \in \mathbb{Q}_\lambda^*$ we define

$$\text{fil}^*(r) = \{A \subseteq \lambda : (\exists \varepsilon < \lambda)(\forall (\alpha, Z, d) \in r)(\varepsilon \leq \alpha \Rightarrow A \cap Z \in d)\},$$
 and we define a binary relation \leq^* on \mathbb{Q}_λ^* by

$$r_1 \leq^* r_2 \text{ if and only if } (r_1, r_2 \in \mathbb{Q}_\lambda^* \text{ and } \text{fil}^*(r_1) \subseteq \text{fil}^*(r_2)).$$
2. For a set $G_* \subseteq \mathbb{Q}_\lambda^*$ we let $\text{fil}^*(G_*) = \bigcup \{\text{fil}^*(r) : r \in G_*\}$.
3. We say that an $r \in \mathbb{Q}_\lambda^*$ is *strongly disjoint* if and only if
 - (a) $(\forall \zeta < \lambda)(|\{(\alpha, Z, d) \in r : \alpha = \zeta\}| < 2)$, and
 - (b) $(\forall (\alpha_1, Z_1, d_1), (\alpha_2, Z_2, d_2) \in r)(\alpha_1 < \alpha_2 \Rightarrow Z_1 \subseteq \alpha_2)$.
4. For $p \in \mathbb{Q}_\lambda^0$ we let $\#(p) = \{(\alpha, Z_\alpha^p, d_\alpha^p) : \alpha \in C^p\}$.

Observation 2.10

1. If $p \in \mathbb{Q}_\lambda^0$ then $\#(p) \in \mathbb{Q}_\lambda^*$ is strongly disjoint and $\text{fil}(p) = \text{fil}^*(\#(p))$.
Also, if $r \in \mathbb{Q}_\lambda^*$ is strongly disjoint, then $\text{fil}^*(r) = \text{fil}(p)$ for some $p \in \mathbb{Q}_\lambda^0$.
2. Let $r, s \in \mathbb{Q}_\lambda^*$. Then $r \leq^* s$ if and only if there is $\varepsilon < \lambda$ such that

$$(\forall (\alpha, Z, d) \in s)(\forall A \in d)(\alpha > \varepsilon \Rightarrow (\exists (\alpha', Z', d') \in r)(A \cap Z' \in d')).$$

The various definitions of super reasonable ultrafilters introduced in Definition 2.11 below are motivated by the proof of “the Sacks forcing preserves P -points.” In that proof, a fusion sequence is constructed so that at a stage $n < \omega$ of the construction one deals with *finitely many* nodes in a condition (the nodes that are declared to be kept). We would like to carry out this kind of argument, for example, for forcing notions used in [8, B.8.3, B.8.5], but now we have to deal with $< \lambda$ nodes in a tree, and the ultrafilter we try to preserve is not that complete. So what do we do? We deal with *finitely many* nodes at a time eventually taking care of everybody. One can think that in the definition below the set I_α is the set of nodes we have to keep and the finite sets $u_{\alpha,i}$ are the nodes taken care of at a substage i .

The technical aspects of Definition 2.11 are motivated by the iteration theorems in [7] and [10]: our games here are tailored to fit the games played on trees of conditions in λ -support iterations (see Theorems 4.2, 4.4 later). As said earlier, the main goal is to have a property of G^* which implies the preservation of “ $\text{fil}(G^*)$ is an ultrafilter” by many forcing notions. We would also love to preserve that property itself, but we failed to achieve it. The “super reasonability” is what we need to preserve the ultrafilter (see Theorem 4.2); “strong reasonability” is what we can prove about G^* in the extension (see Theorem 4.4).

Definition 2.11 Let $G^* \subseteq \mathbb{Q}_\lambda^0$ and let $\bar{\mu} = \langle \mu_\alpha : \alpha < \lambda \rangle$ be a sequence of cardinals, $2 \leq \mu_\alpha \leq \lambda$ for $\alpha < \lambda$.

1. We define a game $\mathcal{D}_{\bar{\mu}}^{\boxplus}(G^*)$ between two players, COM and INC. A play of $\mathcal{D}_{\bar{\mu}}^{\boxplus}(G^*)$ lasts λ steps and at a stage $\alpha < \lambda$ of the play the players choose $I_\alpha, i_\alpha, \bar{u}_\alpha$ and $\langle r_{\alpha,i}, r'_{\alpha,i}, (\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : i < i_\alpha \rangle$ applying the following procedure.

- (i) First, INC chooses a nonempty set I_α of cardinality $< \mu_\alpha$ and an enumeration $\bar{u}_\alpha = \langle u_{\alpha,i} : i < i_\alpha \rangle$ of $[I_\alpha]^{<\omega}$ (so $i_\alpha < \mu_\alpha \cdot \aleph_0$).
 - (ii) Next the two players play a subgame of length i_α . In the i th move of the subgame,
 - (a) COM chooses $r_{\alpha,i} \in G^*$, and then
 - (b) INC chooses $r'_{\alpha,i} \in G^*$ such that $r_{\alpha,i} \leq^0 r'_{\alpha,i}$, and finally
 - (c) COM picks $(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) \in \#(r'_{\alpha,i})$ such that $\beta_{\alpha,i} > \alpha$.
- In the end of the play COM wins if and only if

(\boxplus) there is $r \in G^*$ such that for every $\bar{j} = \langle j_\alpha : \alpha < \lambda \rangle \in \prod_{\alpha < \lambda} I_\alpha$ we have

$$\{(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : \alpha < \lambda, j_\alpha \in u_{\alpha,i} \text{ and } i < i_\alpha\} \leq^* \#(r).$$

A game $\mathfrak{D}_{\bar{\mu}}^{\boxplus}(G^*)$ is defined similarly to $\mathfrak{D}_{\bar{\mu}}^{\boxplus}(G^*)$ except (\boxplus) is weakened to

(\boxminus) for every $\bar{j} \in \prod_{\alpha < \lambda} I_\alpha$ the set $\bigcup \{Z_{\alpha,i} : \alpha < \lambda, i < i_\alpha \text{ and } j_\alpha \in u_{\alpha,i}\}$ belongs to $\text{fil}(G^*)$.

2. We say that the family G^* is $\bar{\mu}$ -super reasonable ($\bar{\mu}$ -super⁻ reasonable, respectively) if
 - (i) G^* is $(<\lambda^+)$ -directed (with respect to \leq^0), and
 - (ii) if $s \in G^*$, $r \in \mathbb{Q}_\lambda^0$ and for some $\alpha < \lambda$ we have $C^r = C^s \setminus \alpha$ and $d_\beta^r = d_\beta^s$ for $\beta \in C^r$, then $r \in G^*$, and
 - (iii) INC has no winning strategy in the game $\mathfrak{D}_{\bar{\mu}}^{\boxplus}(G^*)$ ($\mathfrak{D}_{\bar{\mu}}^{\boxminus}(G^*)$, respectively).
3. We say that a uniform ultrafilter D on λ is $\bar{\mu}$ -super reasonable ($\bar{\mu}$ -super⁻ reasonable, respectively) if there is a $\bar{\mu}$ -super reasonable ($\bar{\mu}$ -super⁻ reasonable, respectively) set $G^* \subseteq \mathbb{Q}_\lambda^0$ such that $D = \text{fil}(G^*)$.
4. If $\mu_\alpha = \lambda$ for all $\alpha < \lambda$, then we omit $\bar{\mu}$ and say just *super reasonable* or *super⁻ reasonable* (in reference to both ultrafilters on λ and families $G^* \subseteq \mathbb{Q}_\lambda^0$). Also in this case we may write \mathfrak{D}^{\boxplus} instead of $\mathfrak{D}_{\bar{\mu}}^{\boxplus}$.

Definition 2.12 Let $G^* \subseteq \mathbb{Q}_\lambda^0$ be directed with respect to \leq^0 and let $\bar{\mu} = \langle \mu_\alpha : \alpha < \lambda \rangle$ be a sequence of cardinals, $2 \leq \mu_\alpha \leq \lambda$ for $\alpha < \lambda$.

1. A game $\mathfrak{D}_{\bar{\mu}}^{\oplus}(G^*)$ between two players, COM and INC is defined as follows. A play of $\mathfrak{D}_{\bar{\mu}}^{\oplus}(G^*)$ lasts λ steps and at a stage $\alpha < \lambda$ of the play the players choose $I_\alpha, i_\alpha, \bar{u}_\alpha$ and $\langle r_{\alpha,i}, \delta_{\alpha,i}, (\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : i < i_\alpha \rangle$ applying the following procedure.
 - (i) First, INC chooses a nonempty set I_α of cardinality $< \mu_\alpha$, and then COM chooses $i_\alpha < \lambda$ and a sequence $\bar{u}_\alpha = \langle u_{\alpha,i} : i < i_\alpha \rangle$ of nonempty finite subsets of I_α such that $I_\alpha = \bigcup_{i < i_\alpha} u_{\alpha,i}$.
 - (ii) Next the two players play a subgame of length i_α . In the i th move of the subgame,
 - (a) COM chooses $r_{\alpha,i} \in G^*$, and then
 - (b) INC chooses $\delta_{\alpha,i} < \lambda$, and finally
 - (c) COM picks $(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) \in \#(r_{\alpha,i})$ such that $\beta_{\alpha,i}$ is above $\delta_{\alpha,i}$ and α .

In the end of the play COM wins if and only if

(\oplus) there is $r \in G^*$ such that for every $\bar{j} = \langle j_\alpha : \alpha < \lambda \rangle \in \prod_{\alpha < \lambda} I_\alpha$ we have

$$\{(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : \alpha < \lambda, j_\alpha \in u_{\alpha,i} \text{ and } i < i_\alpha\} \leq^* \#(r).$$

A game $\mathfrak{D}_{\bar{\mu}}^{\ominus}(G^*)$ is defined similarly to $\mathfrak{D}_{\bar{\mu}}^{\oplus}(G^*)$ except (\oplus) is weakened to

(\ominus) for every $\bar{j} \in \prod_{\alpha < \lambda} I_\alpha$ the set $\bigcup\{Z_{\alpha,i} : \alpha < \lambda, i < i_\alpha \text{ and } j_\alpha \in u_{\alpha,i}\}$ belongs to $\text{fil}(G^*)$.

2. If $G^* \subseteq \mathbb{Q}_\lambda^0$ is ($<\lambda^+$)-directed (with respect to \leq^0) and INC has no winning strategy in the game $\mathfrak{D}_{\bar{\mu}}^{\oplus}(G^*)$, then we say that G^* is $\bar{\mu}$ -strongly reasonable. Also, G^* is said to be $\bar{\mu}$ -strongly⁻ reasonable if it is ($<\lambda^+$)-directed and INC has no winning strategy in the game $\mathfrak{D}_{\bar{\mu}}^{\ominus}(G^*)$.
3. We say that a uniform ultrafilter D on λ is $\bar{\mu}$ -strongly reasonable ($\bar{\mu}$ -strongly⁻ reasonable, respectively) if there is a $\bar{\mu}$ -strongly reasonable ($\bar{\mu}$ -strongly⁻ reasonable, respectively) set $G^* \subseteq \mathbb{Q}_\lambda^0$ such that $D = \text{fil}(G^*)$. If $\mu_\alpha = \lambda$ for all $\alpha < \lambda$, then we omit $\bar{\mu}$ and say just strongly reasonable or strongly⁻ reasonable.

Observation 2.13 Assume that $2 \leq \mu_\alpha \leq \kappa_\alpha \leq \lambda$ for $\alpha < \lambda$ and $\bar{\mu} = \langle \mu_\alpha : \alpha < \lambda \rangle$, $\bar{\kappa} = \langle \kappa_\alpha : \alpha < \lambda \rangle$. Then for a family $G^* \subseteq \mathbb{Q}_\lambda^0$ and/or a uniform ultrafilter D on λ the following implications hold.

$$\begin{array}{ccccc} \bar{\kappa}\text{-super reasonable} & \Rightarrow & \bar{\mu}\text{-super reasonable} & \Rightarrow & \bar{\mu}\text{-strongly reasonable} \\ \downarrow & & \downarrow & & \downarrow \\ \bar{\kappa}\text{-super}^-\text{ reasonable} & \Rightarrow & \bar{\mu}\text{-super}^-\text{ reasonable} & \Rightarrow & \bar{\mu}\text{-strongly}^-\text{ reasonable} \end{array}$$

Proposition 2.14 Assume that $2 \leq \mu_\alpha \leq \lambda$ for $\alpha < \lambda$ and $\bar{\mu} = \langle \mu_\alpha : \alpha < \lambda \rangle$. If a uniform ultrafilter D on λ is $\bar{\mu}$ -strongly⁻ reasonable, then it is very reasonable.

Proof Pick a $\bar{\mu}$ -strongly⁻ reasonable family $G^* \subseteq \mathbb{Q}_\lambda^0$ such that $D = \text{fil}(G^*)$. Then G^* is ($<\lambda^+$)-directed and the proof will be completed once we show that D is weakly reasonable.

Let $f \in {}^\lambda\lambda$. We will argue that for some club $C = \{\gamma_\alpha : \alpha < \gamma\} \subseteq \lambda$ we have $\bigcup\{[\delta, \delta + f(\delta)) : \delta \in C\} \notin D$, where γ_α are given by the arguments below.

We consider the following strategy $\text{st}(f)$ for INC in $\mathfrak{D}_{\bar{\mu}}^{\ominus}(G^*)$. The strategy $\text{st}(f)$ instructs INC to construct on the side an increasing continuous sequence $\langle \gamma_\alpha : \alpha < \lambda \rangle \subseteq \lambda$ so that at a stage $\alpha < \lambda$ of the play, when

$$\langle I_\xi, i_\xi, \bar{u}_\xi, \langle r_{\xi,i}, \delta_{\xi,i}, (\beta_{\xi,i}, Z_{\xi,i}, d_{\xi,i}) : i < i_\xi \rangle : \xi < \alpha \rangle$$

is the result of the play so far, then

1. if α is limit, then $\gamma_\alpha = \sup(\gamma_\xi : \xi < \alpha)$,
2. if α is not limit, then $\gamma_\alpha = \sup(\bigcup\{Z_{\xi,i} : i < i_\xi, \xi < \alpha\}) + 1$.

Now (at the stage α) $\text{st}(f)$ instructs INC to choose $I_\alpha = \{0\}$ and then (after COM picks i_α, \bar{u}_α) he is instructed to play in the subgame of this stage as follows. At stage $i < i_\alpha$, after COM has picked $r_{\alpha,i}$, INC lets

$$\delta_{\alpha,i} = \gamma_\alpha + f(\gamma_\alpha) + \sup(\bigcup\{Z_{\alpha,j} : j < i\}) + 890.$$

(After this COM chooses $(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) \in \#(r_{\alpha,i})$ with $\beta_{\alpha,i} > \delta_{\alpha,i}$.)

The strategy $\text{st}(f)$ cannot be the winning one for INC, so there is a play

$$\langle I_\alpha, i_\alpha, \bar{u}_\alpha, \langle r_{\alpha,i}, \delta_{\alpha,i}, (\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : i < i_\alpha \rangle : \alpha < \lambda \rangle$$

of $\mathfrak{D}_{\bar{\mu}}^{\ominus}(G^*)$ in which INC follows $\mathbf{st}(f)$ but

$$A^* \stackrel{\text{def}}{=} \bigcup \{Z_{\alpha,i} : \alpha < \lambda, i < i_\alpha\} \in \text{fil}(G^*) = D$$

(note that necessarily $u_{\alpha,i} = I_\alpha = \{0\}$). It follows from the choice of $\gamma_\alpha, \delta_{\alpha,i}$ that for each $\alpha < \lambda$

$$[\gamma_\alpha, \gamma_\alpha + f(\gamma_\alpha)] \cap \bigcup \{Z_{\zeta,i} : \zeta < \lambda, i < i_\zeta\} = \emptyset,$$

and hence also $\bigcup \{[\gamma_\alpha, \gamma_\alpha + f(\gamma_\alpha)] : \alpha < \lambda\} \cap A^* = \emptyset$. Consequently, $\bigcup \{[\gamma_\alpha, \gamma_\alpha + f(\gamma_\alpha)] : \alpha < \lambda\} \notin D$ and one can easily finish the proof. \square

Proposition 2.15 *Assume $\lambda = \lambda^{<\lambda}$ and $\diamond_{S_\lambda^+}$ holds. There exists a sequence $\langle r_\zeta : \zeta < \lambda^+ \rangle \subseteq \mathbb{Q}_\lambda^0$ such that*

- (i) $(\forall \zeta < \zeta < \lambda^+)(r_\zeta \leq^0 r_\zeta)$, and
- (ii) *the family*

$$G^* \stackrel{\text{def}}{=} \{r \in \mathbb{Q}_\lambda^0 : (\exists \zeta < \lambda^+)(r \leq^0 r_\zeta)\}$$

is super reasonable and $\text{fil}(G^)$ is an ultrafilter on λ .*

Proof The sequence $\langle r_\zeta : \zeta < \lambda^+ \rangle$ will be constructed inductively. At successor stages we will use Observation 2.6 to make sure that $\text{fil}(G^*)$ is an ultrafilter. At limit stages we will use Proposition 2.8(1) to find upper bounds to the sequence constructed so far. Moreover, at (some) stages ζ of cofinality λ the element r_ζ will be chosen so that “it kills” a strategy for INC in $\mathfrak{D}^{\boxplus}(G^*)$ predicted by the diamond sequence.

For $\alpha < \lambda$ let X_α^1 be the set of all legal plays of $\mathfrak{D}^{\boxplus}(\mathbb{Q}_\lambda^0)$ of the form

$$(\odot)_\alpha^1 \quad \langle I_\gamma, i_\gamma, \bar{u}_\gamma, \langle r_{\gamma,i}, r'_{\gamma,i}, (\beta_{\gamma,i}, Z_{\gamma,i}, d_{\gamma,i}) : i < i_\gamma \rangle : \gamma < \alpha \rangle$$

where each I_γ (for $\gamma < \alpha$) is an ordinal below λ . Also let $X^1 = \bigcup_{\alpha < \lambda} X_\alpha^1$. Next, for $\alpha < \lambda, 0 < I < \lambda$ and an enumeration $\bar{u} = \langle u_j : j < i \rangle$ of $[I]^{<\omega}$ let $X_{\alpha,I,\bar{u}}^2$ be the set of all legal plays of $\mathfrak{D}^{\boxplus}(\mathbb{Q}_\lambda^0)$ of the form

$$(\odot)_{\alpha,I,\bar{u}}^2 \quad \bar{\sigma} \frown \langle (I, i, \bar{u}) \rangle \frown \langle r_j, r'_j, (\beta_j, Z_j, d_j) : j < j^* \rangle \frown \langle r \rangle,$$

where $\bar{\sigma} \in X_\alpha^1, j^* < i$ (and $\langle r_j, r'_j, (\beta_j, Z_j, d_j) : j < j^* \rangle \frown \langle r \rangle$ is a legal partial play of the subgame of level α ; in particular, $r_j, r'_j, r \in \mathbb{Q}_\lambda^0$). Also let

$$X^2 = \bigcup \{X_{\alpha,I,\bar{u}}^2 : \alpha < \lambda \text{ and } 0 < I < \lambda \text{ and } \bar{u} = \langle u_j : j < i \rangle \text{ is an enumeration of } [I]^{<\omega}\}.$$

Any strategy for INC in $\mathfrak{D}^{\boxplus}(\mathbb{Q}_\lambda^0)$ can be interpreted as a function \mathbf{st} such that

- (\odot)³ the domain of \mathbf{st} is $X^1 \cup X^2$,
- (\odot)⁴ if $\bar{\sigma} \in X_\alpha^1, \alpha < \lambda$, then $\mathbf{st}(\bar{\sigma}) = (I, i, \bar{u})$ for some $I < \lambda$ and an enumeration $\bar{u} = \langle u_j : j < i \rangle$ of $[I]^{<\omega}$,
- (\odot)⁵ if $\bar{\sigma} \in X_{\alpha,I,\bar{u}}^2, \alpha < \lambda, 0 < I < \lambda, \bar{u} = \langle u_j : j < i \rangle = [I]^{<\omega}$, and $\bar{\sigma} = \bar{\sigma}_0 \frown \langle (I, i, \bar{u}) \rangle \frown \langle r_j, r'_j, (\beta_j, Z_j, d_j) : j < j^* \rangle \frown \langle r \rangle$, then $\mathbf{st}(\bar{\sigma}) \in \mathbb{Q}_\lambda^0$ is such that $r \leq^0 \mathbf{st}(\bar{\sigma})$.

Below, whenever we say *a strategy for INC* we mean a function **st** satisfying conditions $(\odot)^3$ – $(\odot)^5$.

Since $|\mathbb{Q}_\lambda^0| = 2^{2^{<\lambda}} = \lambda^+$, we may pick a bijection $\pi_0 : \mathbb{Q}_\lambda^0 \xrightarrow{1-1} \lambda^+$ and for $\zeta < \lambda^+$ let \mathcal{X}_ζ consist of all $\bar{\sigma} \in X^1 \cup X^2$ such that $\pi_0(r) < \zeta$ for all elements $r \in \mathbb{Q}_\lambda^0$ involved in the representation of $\bar{\sigma}$ as in $(\odot)^1$, $(\odot)^2$. We also let \mathcal{Y}_ζ consist of all pairs $(\bar{\sigma}, a)$ such that

1. $\bar{\sigma} \in \mathcal{X}_\zeta$ and $a = \mathbf{st}(\bar{\sigma})$ for some strategy **st** of INC, and
2. if $\bar{\sigma} \in X^2$ (and so $a \in \mathbb{Q}_\lambda^0$) then $\pi_0(a) < \zeta$.

Note that $|\mathcal{X}_\zeta| \leq \lambda$ and $|\mathcal{Y}_\zeta| \leq \lambda$ (for each $\zeta < \lambda^+$). Put $\mathcal{Y} = \bigcup_{\zeta < \lambda^+} \mathcal{Y}_\zeta$. Plainly,

$|\mathcal{Y}| = \lambda^+$ so we may fix a bijection $\pi_1 : \lambda^+ \xrightarrow{\text{onto}} \mathcal{Y}$. Let $C = \{\zeta < \lambda^+ : \pi_1[\zeta] = \mathcal{Y}_\zeta\}$; it is a club of λ^+ .

Let $\langle A_\zeta : \zeta < \lambda^+ \rangle$ list all subsets of λ and let $\langle B_\zeta : \zeta \in S_\lambda^{\lambda^+} \rangle$ be a diamond sequence on $S_\lambda^{\lambda^+} = \{\zeta < \lambda^+ : \text{cf}(\zeta) = \lambda\}$. By induction on $\zeta < \lambda^+$ we choose a \leq^0 -increasing sequence $\langle r_\zeta : \zeta < \lambda^+ \rangle \subseteq \mathbb{Q}_\lambda^0$ applying the following procedure. Assume $\zeta < \lambda^+$ and we have constructed $\langle r_\zeta : \zeta < \zeta \rangle$.

Case 0 $\zeta = 0$. We let r_0 be the $<_\lambda^*$ -first member of \mathbb{Q}_λ^0 .

Case 1 $\zeta = \zeta + 1$. Pick $r_\zeta \in \mathbb{Q}_\lambda^0$ such that $r_\zeta \leq^0 r_\zeta$ and either $A_\zeta \in \text{fil}(r_\zeta)$ or $\lambda \setminus A_\zeta \in \text{fil}(r_\zeta)$ (remember Observation 2.6).

Case 2 ζ is a limit ordinal, $\text{cf}(\zeta) < \lambda$. Pick $r_\zeta \in \mathbb{Q}_\lambda^0$ such that $(\forall \zeta' < \zeta)(r_\zeta \leq^0 r_{\zeta'})$ (exists by Proposition 2.8(1)).

Case 3 ζ is a limit ordinal, $\text{cf}(\zeta) = \lambda$. Now we ask if

$(\odot)_\zeta^6$ $\zeta \in C$ and $(\forall \zeta' < \zeta)(\pi_0(r_{\zeta'}) < \zeta)$ and there is a strategy **st** for INC in $\mathfrak{D}^{\boxplus}(\mathbb{Q}_\lambda^0)$ such that $\pi_1[B_\zeta] = \mathbf{st} \cap \mathcal{Y}_\zeta = \mathbf{st} \upharpoonright \mathcal{X}_\zeta$.

If the answer to $(\odot)_\zeta^6$ is negative, then we choose $r_\zeta \in \mathbb{Q}_\lambda^0$ as in Case 2.

Suppose now that the answer to $(\odot)_\zeta^6$ is positive (so, in particular, $\zeta \in C$) and **st** is a strategy for INC such that $\pi_1[B_\zeta] = \mathbf{st} \cap \mathcal{Y}_\zeta = \mathbf{st} \upharpoonright \mathcal{X}_\zeta$. Let $\bar{\zeta} = \langle \zeta_\alpha : \alpha < \lambda \rangle$ be an increasing continuous sequence cofinal in ζ . Consider a play

$$\bar{\sigma} = \langle I_\alpha, i_\alpha, \bar{u}_\alpha, \langle r_{\alpha,i}, r'_{\alpha,i}, (\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : i < i_\alpha \rangle : \alpha < \lambda \rangle$$

of $\mathfrak{D}^{\boxplus}(\mathbb{Q}_\lambda^0)$ in which INC follows the strategy **st** and COM proceeds as follows. When playing $\mathfrak{D}^{\boxplus}(\mathbb{Q}_\lambda^0)$, at step $i < i_\alpha$ of the subgame of level $\alpha < \lambda$ (of $\mathfrak{D}^{\boxplus}(\mathbb{Q}_\lambda^0)$) COM chooses $r_{\alpha,i} = r_{\zeta_\alpha}$ and then, after INC determines $r'_{\alpha,i}$ by **st**, she picks the $<_\lambda^*$ -first $(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) \in \#(r'_{\alpha,i})$ satisfying

$(\odot)_{\zeta_\alpha, \alpha, i}^7$ $(\forall \gamma \leq \alpha)(\forall A \in d_{\alpha,i})(\exists \delta \in C^{r_{\zeta_\alpha}})(A \cap Z_\delta^{r_{\zeta_\alpha}} \in d_\delta^{r_{\zeta_\alpha}})$ (remember Proposition 2.5) and

$(\odot)_{\zeta_\alpha, \alpha, i}^8$ $(\forall \gamma < \alpha)(\forall j < i_\gamma)(Z_{\gamma,j} \subseteq \beta_{\alpha,i})$ and $(\forall j < i)(Z_{\alpha,j} \subseteq \beta_{\alpha,i})$.

The above rules fully determine the play $\bar{\sigma}$ and it should be clear that $\bar{\sigma} \upharpoonright \alpha \in \mathcal{X}_\zeta$ for each $\alpha < \lambda$. Note that $\bar{\sigma}$ depends on B_ζ and $\bar{\zeta}$ only (and not on **st**, provided it is as required by $(\odot)_\zeta^6$).

By the demands $(\odot)_{\xi, \alpha}^8$, we may choose an increasing continuous sequence $\langle \gamma_\alpha : \alpha < \lambda \rangle \subseteq \lambda$ such that $\gamma_0 = 0$ and $(\forall \alpha < \lambda)(\forall i < i_\alpha)(Z_{\alpha, i} \subseteq [\gamma_\alpha, \gamma_{\alpha+1}))$. Now, for $\alpha < \lambda$ choose an ultrafilter e_α on i_α such that

$$(\odot)_{\xi, \alpha}^9 \quad (\forall j \in I_\alpha)(\{i < i_\alpha : j \in u_{\alpha, i}\} \in e_\alpha)$$

and let d_α be an ultrafilter on $[\gamma_\alpha, \gamma_{\alpha+1})$ such that

$$(\odot)_{\xi, \alpha}^{10} \quad \bigoplus^{e_\alpha} \{d_{\alpha, i} : i < i_\alpha\} \subseteq d_\alpha.$$

Now let $r_\zeta \in \mathbb{Q}_\lambda^0$ be such that

1. $C^{r_\zeta} = \{\gamma_\alpha : \alpha < \lambda\}$, and
2. if $\delta = \gamma_\alpha$, then $Z_\delta^{r_\zeta} = [\gamma_\alpha, \gamma_{\alpha+1})$ and $d_\delta^{r_\zeta} = d_\alpha$.

One easily verifies that $r_{\zeta_\alpha} \leq^0 r_\zeta$ for all $\alpha < \lambda$ (remember $(\odot)^7$ and the choice of d_α ; use Proposition 2.5) and so $r_\zeta \leq^0 r_\zeta$ for all $\zeta < \xi$. It follows from $(\odot)_{\xi, \alpha}^9$ and $(\odot)_{\xi, \alpha}^{10}$ that

$$(\odot)_\xi^{11} \quad \text{for every } \bar{j} = \langle j_\alpha : \alpha < \lambda \rangle \in \prod_{\alpha < \lambda} I_\alpha \text{ we have}$$

$$\{(\beta_{\alpha, i}, Z_{\alpha, i}, d_{\alpha, i}) : \alpha < \lambda \ \& \ j_\alpha \in u_{\alpha, i} \ \& \ i < i_{\alpha, i}\} \leq^* \#(r_\xi).$$

After the construction of $\langle r_\zeta : \zeta < \lambda^+ \rangle$ is carried out we let

$$G^* = \{r \in \mathbb{Q}_\lambda^0 : (\exists \zeta < \lambda^+)(r \leq^0 r_\zeta)\}.$$

Plainly, G^* satisfies demands (i) and (ii) of Definition 2.11(2) and $\text{fil}(G^*)$ is an ultrafilter on λ (remember Case 1 of the construction). We should argue that INC has no winning strategy in $\mathfrak{D}^{\boxplus}(G^*)$. To this end suppose that \mathbf{st}^{\boxplus} is a strategy of INC in $\mathfrak{D}^{\boxplus}(G^*)$. Pick $\zeta \in S_\lambda^{\lambda^+} \cap C$ such that $(\forall \zeta < \xi)(\pi_0(r_\zeta) < \zeta)$ and $\pi_1[B_\zeta] = \mathbf{st}^{\boxplus} \cap \mathcal{Y}_\zeta = \mathbf{st}^{\boxplus} \upharpoonright \mathcal{X}_\zeta$. Then when choosing r_ζ we gave a positive answer to $(\odot)_\xi^6$ and we constructed a play $\bar{\sigma}$ of $\mathfrak{D}^{\boxplus}(\mathbb{Q}_\lambda^0)$. In that play, INC follows \mathbf{st}^{\boxplus} and COM chooses members of G^* , so it is a play of $\mathfrak{D}^{\boxplus}(G^*)$. Now the condition $(\odot)_\xi^{11}$ means that r_ζ witnesses that COM wins the play $\bar{\sigma}$ and consequently \mathbf{st}^{\boxplus} is not a winning strategy for INC. \square

Proposition 2.16 *Let $\mathbb{Q}_\lambda^0 = (\mathbb{Q}_\lambda^0, \leq^0)$.*

1. \mathbb{Q}_λ^0 is a $(< \lambda^+)$ -complete forcing notion of size $2^{2^{< \lambda}}$.
2. $\Vdash_{\mathbb{Q}_\lambda^0}$ “ $\mathcal{G}_{\mathbb{Q}_\lambda^0}$ is a super reasonable family and $\text{fil}(\mathcal{G}_{\mathbb{Q}_\lambda^0})$ is an ultrafilter”.

Proof (1) Should be clear; see also Proposition 2.8(1).

(2) By the completeness of \mathbb{Q}_λ^0 , forcing with it does not add new subsets of λ , and by Proposition 2.5

$$\Vdash_{\mathbb{Q}_\lambda^0} \text{“fil}(\mathcal{G}_{\mathbb{Q}_\lambda^0}) \text{ is a uniform ultrafilter on } \lambda\text{”}.$$

It should also be clear that $\mathcal{G}_{\mathbb{Q}_\lambda^0}$ satisfies the demands of Definition 2.11(2)(i+ii) (in $\mathbf{V}^{\mathbb{Q}_\lambda^0}$). Let us argue that

$$\Vdash_{\mathbb{Q}_\lambda^0} \text{“INC has no winning strategy in } \mathfrak{D}^{\boxplus}(\mathcal{G}_{\mathbb{Q}_\lambda^0})\text{”}$$

and to this end suppose $p \in \mathbb{Q}_\lambda^0$ and \mathbf{st} is a \mathbb{Q}_λ^0 -name such that

$$p \Vdash_{\mathbb{Q}_\lambda^0} \text{“}\mathbf{st} \text{ is a strategy of INC in } \mathfrak{D}^{\boxplus}(\mathcal{G}_{\mathbb{Q}_\lambda^0})\text{”}.$$

We are going to construct a condition $q \in \mathbb{Q}_\lambda^0$ stronger than p and a play $\bar{\sigma}$ of $\mathfrak{D}^{\boxplus}(\mathbb{Q}_\lambda^0)$ such that

$p \Vdash_{\mathbb{Q}_\lambda^0}$ “ $\bar{\sigma}$ is a play of $\mathfrak{D}^{\boxplus}(G_{\mathbb{Q}_\lambda^0})$ in which INC follows \mathfrak{st} but COM wins”.

Let X^1, X^2 be defined as in the proof of Proposition 2.15 (see $(\odot)_a^1, (\odot)_{a,I,\bar{u}}^2$ there). We may assume that

$p \Vdash_{\mathbb{Q}_\lambda^0}$ “ \mathfrak{st} is a function satisfying $(\odot)^3 - (\odot)^5$ of the proof of Proposition 2.15”.

By induction on $\alpha < \lambda$ we choose conditions $p_\alpha \in \mathbb{Q}_\lambda^0$ and partial plays $\bar{\sigma}_\alpha \in X_\alpha^1$ so that

(\square)₁ $p \leq^0 p_\alpha \leq^0 p_\beta$ and $\bar{\sigma}_\alpha \triangleleft \bar{\sigma}_\beta$ for $\alpha < \beta < \lambda$,

(\square)₂ $p_\alpha \Vdash_{\mathbb{Q}_\lambda^0}$ “ $\bar{\sigma}_\alpha$ is a partial play of $\mathfrak{D}^{\boxplus}(G_{\mathbb{Q}_\lambda^0})$ in which INC uses \mathfrak{st} ”,

(\square)₃ if $\bar{\sigma}_\alpha = \langle I_\gamma, i_\gamma, \bar{u}_\gamma, \langle r_{\gamma,i}, r'_{\gamma,i}, (\beta_{\gamma,i}, Z_{\gamma,i}, d_{\gamma,i}) : i < i_\gamma \rangle : \gamma < \alpha \rangle$, then for every $\gamma < \delta < \alpha$ and $j < i < i_\gamma$ we have

$$r'_{\gamma,i} \leq^0 p_\alpha \quad \text{and} \quad Z_{\gamma,j} \subseteq \beta_{\gamma,i} \quad \text{and} \quad Z_{\gamma,j} \subseteq \beta_{\delta,0}.$$

Suppose that $\alpha = \alpha^* + 1$ and we have determined $p_{\alpha^*}, \bar{\sigma}_{\alpha^*}$. Pick $p'_\alpha \geq^0 p_{\alpha^*}$ and $I_\alpha, i_\alpha, \bar{u}_\alpha$ such that $p'_\alpha \Vdash \mathfrak{st}(\bar{\sigma}_{\alpha^*}) = (I_\alpha, i_\alpha, \bar{u}_\alpha)$. Now choose inductively $p_{\alpha,i}^i, r_{\alpha,i}, r'_{\alpha,i}$ and $(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i})$ for $i < i_\alpha$ so that for each $i < j < i_\alpha$ we have

(\square)₄ (i) $p_\alpha^0 = p'_\alpha, p_\alpha^i \leq^0 p_\alpha^j, p_\alpha^i = r_{\alpha,i} \leq^0 r'_{\alpha,i} \leq^0 p_\alpha^{i+1}$, and

(ii) $p_\alpha^{i+1} \Vdash$ “ $r'_{\alpha,i}$ is the answer by \mathfrak{st} at stage i of the subgame”,

(iii) $\beta_{\alpha,i}$ satisfies the demand in (\square)₃ and $(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) \in \#(r'_{\alpha,i})$,

(iv) $(\forall A \in d_{\alpha,i})(\forall \gamma \leq \alpha^*)(\exists \delta \in C^{p_\gamma})(A \cap Z_\delta^{p_\gamma} \in d_\delta^{p_\gamma})$.

Then $p_{\alpha+1}$ is any \leq^0 -upper bound to $\{p_\alpha^i : i < i_\alpha\}$. The limit stages of the construction should be clear.

After the construction is carried out and we have $\bar{\sigma}_\lambda = \bigcup \{\bar{\sigma}_\alpha : \alpha < \lambda\}$, we define $r \in \mathbb{Q}_\lambda^0$ like r_ξ in the proof of Proposition 2.15 (see $(\odot)_{\xi,\alpha}^9 + (\odot)_\xi^{10}$ there). Then r is \leq^0 -stronger than all p_α (for $\alpha < \lambda$) and

$r \Vdash_{\mathbb{Q}_\lambda^0}$ “ $\bar{\sigma}_\lambda$ is a play of $\mathfrak{D}^{\boxplus}(G_{\mathbb{Q}_\lambda^0})$ in which INC uses \mathfrak{st} but COM wins”.

(Note that the respective version of $(\odot)_\xi^{11}$ of the proof of Proposition 2.15 holds. By the completeness it continues to hold in $\mathbf{V}^{\mathbb{Q}_\lambda^0}$.) \square

3 More on Reasonably Complete Forcing

Definition 3.1 Let \mathbb{P} be a forcing notion.

1. For a condition $r \in \mathbb{P}$, let $\mathfrak{D}_0^\lambda(\mathbb{P}, r)$ be the following game of two players, COMPLETE and INCOMPLETE:

the game lasts at most λ moves and during a play the players attempt to construct a sequence $\langle (p_i, q_i) : i < \lambda \rangle$ of pairs of conditions from \mathbb{P} in such a way that $(\forall j < i < \lambda)(r \leq p_j \leq q_j \leq p_i)$ and at the stage $i < \lambda$ of the game, first INCOMPLETE chooses p_i and then COMPLETE chooses q_i .

COMPLETE wins if and only if for every $i < \lambda$ there are legal moves for both players.

2. We say that the forcing notion \mathbb{P} is *strategically* ($<\lambda$)-*complete* if COMPLETE has a winning strategy in the game $\mathfrak{D}_0^\lambda(\mathbb{P}, p)$ for each condition $p \in \mathbb{P}$.
3. Let $N \prec (\mathcal{H}(\chi), \in, <_\chi^*)$ be a model such that ${}^{<\lambda}N \subseteq N$, $|N| = \lambda$, and $\mathbb{P} \in N$. We say that a condition $p \in \mathbb{P}$ is (N, \mathbb{P}) -*generic in the standard sense* (or just (N, \mathbb{P}) -*generic*) if for every \mathbb{P} -name $\tau \in N$ for an ordinal we have $p \Vdash \tau \in N$.
4. \mathbb{P} is λ -*proper in the standard sense* (or just λ -*proper*) if there is $x \in \mathcal{H}(\chi)$ such that for every model $N \prec (\mathcal{H}(\chi), \in, <_\chi^*)$ satisfying

$${}^{<\lambda}N \subseteq N, \quad |N| = \lambda, \quad \text{and} \quad \mathbb{P}, x \in N,$$

and every condition $p \in N \cap \mathbb{P}$ there is an (N, \mathbb{P}) -generic condition $q \in \mathbb{P}$ stronger than p .

Theorem 3.2 (See Shelah [11, Ch. III, Thm. 4.1], Abraham [1, §2], Eisworth [2, §3]) *Assume $2^\lambda = \lambda^+$, $\lambda^{<\lambda} = \lambda$. Let $\bar{\mathbb{Q}} = \langle \mathbb{P}_i, \bar{\mathbb{Q}}_i : i < \lambda^{++} \rangle$ be λ -support iteration such that for all $i < \lambda^{++}$ we have*

1. \mathbb{P}_i is λ -*proper*,
2. $\Vdash_{\mathbb{P}_i} \text{“}|\bar{\mathbb{Q}}_i| \leq \lambda^+ \text{”}$.

Then

1. for every $\delta < \lambda^{++}$, $\Vdash_{\mathbb{P}_\delta} 2^\lambda = \lambda^+$, and
2. the limit $\mathbb{P}_{\lambda^{++}}$ satisfies the λ^{++} -cc.

Proposition 3.3 ([8], Proposition A.1.6) *Suppose $\bar{\mathbb{Q}} = \langle \mathbb{P}_i, \bar{\mathbb{Q}}_i : i < \gamma \rangle$ is a λ -support iteration and, for each $i < \gamma$,*

$$\Vdash_{\mathbb{P}_i} \text{“}\bar{\mathbb{Q}}_i \text{ is strategically } (<\lambda)\text{-complete”}.$$

Then, for each $\varepsilon \leq \gamma$ and $r \in \mathbb{P}_\varepsilon$, there is a winning strategy $\mathbf{st}(\varepsilon, r)$ of COMPLETE in the game $\mathfrak{D}_0^\lambda(\mathbb{P}_\varepsilon, r)$ such that whenever $\varepsilon_0 < \varepsilon_1 \leq \gamma$ and $r \in \mathbb{P}_{\varepsilon_1}$ we have

- (i) if $\langle (p_i, q_i) : i < \lambda \rangle$ is a play of $\mathfrak{D}_0^\lambda(\mathbb{P}_{\varepsilon_0}, r \upharpoonright \varepsilon_0)$ in which COMPLETE follows the strategy $\mathbf{st}(\varepsilon_0, r \upharpoonright \varepsilon_0)$, then $\langle (p_i \hat{\smallfrown} r \upharpoonright [\varepsilon_0, \varepsilon_1], q_i \hat{\smallfrown} r \upharpoonright [\varepsilon_0, \varepsilon_1]) : i < \lambda \rangle$ is a play of $\mathfrak{D}_0^\lambda(\mathbb{P}_{\varepsilon_1}, r)$ in which COMPLETE uses $\mathbf{st}(\varepsilon_1, r)$;
- (ii) if $\langle (p_i, q_i) : i < \lambda \rangle$ is a play of $\mathfrak{D}_0^\lambda(\mathbb{P}_{\varepsilon_1}, r)$ in which COMPLETE plays according to the strategy $\mathbf{st}(\varepsilon_1, r)$, then $\langle (p_i \upharpoonright \varepsilon_0, q_i \upharpoonright \varepsilon_0) : i < \lambda \rangle$ is a play of $\mathfrak{D}_0^\lambda(\mathbb{P}_{\varepsilon_0}, r \upharpoonright \varepsilon_0)$ in which COMPLETE uses $\mathbf{st}(\varepsilon_0, r \upharpoonright \varepsilon_0)$;
- (iii) if $\langle (p_i, q_i) : i < i^* \rangle$ is a partial play of $\mathfrak{D}_0^\lambda(\mathbb{P}_{\varepsilon_1}, r)$ in which COMPLETE uses $\mathbf{st}(\varepsilon_1, r)$ and $p' \in \mathbb{P}_{\varepsilon_0}$ is stronger than all $p_i \upharpoonright \varepsilon_0$ (for $i < i^*$), then there is $p^* \in \mathbb{P}_{\varepsilon_1}$ such that $p' = p^* \upharpoonright \varepsilon_0$ and $p^* \geq p_i$ for $i < i^*$.

Definition 3.4 (Compare [7], Definition 2.2)

1. Let γ be an ordinal, $w \subseteq \gamma$. A *standard* $(w, 1)^\gamma$ -*tree* is a pair $\mathcal{T} = (T, \text{rk})$ such that
 - (a) $\text{rk} : T \longrightarrow w \cup \{\gamma\}$,
 - (b) if $t \in T$ and $\text{rk}(t) = \varepsilon$, then t is a sequence $\langle (t)_\zeta : \zeta \in w \cap \varepsilon \rangle$,
 - (c) (T, \triangleleft) is a tree with root $\langle \rangle$ and such that every chain in T has a \triangleleft -upper bound in T ,
 - (d) if $t \in T$, then there is $t' \in T$ such that $t \trianglelefteq t'$ and $\text{rk}(t') = \gamma$.

We will keep the convention that \mathcal{T}_y^x is (T_y^x, rk_y^x) .

2. Let $\bar{\mathbb{Q}} = \langle \mathbb{P}_i, \bar{\mathbb{Q}}_i : i < \gamma \rangle$ be a λ -support iteration. A *standard tree of conditions in $\bar{\mathbb{Q}}$* is a system $\bar{p} = \langle p_t : t \in T \rangle$ such that

- (a) (T, rk) is a standard $(w, 1)^\gamma$ -tree for some $w \subseteq \gamma$ and
 - (b) $p_t \in \mathbb{P}_{\text{rk}(t)}$ for $t \in T$ and
 - (c) if $s, t \in T, s \triangleleft t$, then $p_s = p_t \upharpoonright \text{rk}(s)$.
3. Let \bar{p}^0, \bar{p}^1 be standard trees of conditions in $\bar{\mathbb{Q}}, \bar{p}^i = \langle p_t^i : t \in T \rangle$. We write $\bar{p}^0 \leq \bar{p}^1$ whenever for each $t \in T$ we have $p_t^0 \leq p_t^1$.

Note that our standard trees and trees of conditions are a special case of that introduced in [8, Definition A.1.7] when $\alpha = 1$. Also, the rank function rk is essentially the function giving the level of a node, adjusted to have values in $w \cup \{\gamma\}$ via the canonical increasing bijection.

Proposition 3.5 (See [8], Proposition A.1.9) *Assume that $\bar{\mathbb{Q}} = \langle \mathbb{P}_i, \mathbb{Q}_i : i < \gamma \rangle$ is a λ -support iteration such that for all $i < \gamma$ we have*

$$\Vdash_{\mathbb{P}_i} \text{“}\mathbb{Q}_i \text{ is strategically } (<\lambda)\text{-complete”}.$$

Suppose that $\bar{p} = \langle p_t : t \in T \rangle$ is a standard tree of conditions in $\bar{\mathbb{Q}}, |T| < \lambda$, and $\mathcal{I} \subseteq \mathbb{P}_\gamma$ is open dense. Then there is a standard tree of conditions $\bar{q} = \langle q_t : t \in T \rangle$ such that $\bar{p} \leq \bar{q}$ and $(\forall t \in T)(\text{rk}(t) = \gamma \Rightarrow q_t \in \mathcal{I})$, and such that conditions q_{t_0}, q_{t_1} are incompatible whenever $t_0, t_1 \in T, \text{rk}(t_0) = \text{rk}(t_1)$ but $t_0 \neq t_1$.

Definition 3.6 (See [7], Definition 3.1) Let \mathbb{Q} be a forcing notion and let $\bar{\mu} = \langle \mu_\alpha : \alpha < \lambda \rangle$ be a sequence of regular cardinals such that $\aleph_0 \leq \mu_\alpha \leq \lambda$ for all $\alpha < \lambda$.

1. For a condition $p \in \mathbb{Q}$ we define a **reasonable A-completeness game** $\mathcal{D}_{\bar{\mu}}^{\text{rcA}}(p, \mathbb{Q})$ between two players, GENERIC and ANTIGENERIC, as follows. A play of $\mathcal{D}_{\bar{\mu}}^{\text{rcA}}(p, \mathbb{Q})$ lasts λ steps and during a play a sequence

$$\left\langle I_\alpha, \langle p_t^\alpha, q_t^\alpha : t \in I_\alpha \rangle : \alpha < \lambda \right\rangle$$

is constructed. Suppose that the players have arrived to a stage $\alpha < \lambda$ of the game. Now,

- (\aleph) $_\alpha$ first GENERIC chooses a nonempty set I_α of cardinality $< \mu_\alpha$ and a system $\langle p_t^\alpha : t \in I_\alpha \rangle$ of conditions from \mathbb{Q} ,
- (\sqsupset) $_\alpha$ then ANTIGENERIC answers by picking a system $\langle q_t^\alpha : t \in I_\alpha \rangle$ of conditions from \mathbb{Q} such that $(\forall t \in I_\alpha)(p_t^\alpha \leq q_t^\alpha)$.

At the end, GENERIC wins the play

$$\left\langle I_\alpha, \langle p_t^\alpha, q_t^\alpha : t \in I_\alpha \rangle : \alpha < \lambda \right\rangle$$

of $\mathcal{D}_{\bar{\mu}}^{\text{rcA}}(p, \mathbb{Q})$ if and only if

- (\otimes) $_{\bar{\mu}}^{\text{rcA}}$ there is a condition $p^* \in \mathbb{Q}$ stronger than p and such that

$$p^* \Vdash_{\mathbb{Q}} \text{“}(\forall \alpha < \lambda)(\exists t \in I_\alpha)(q_t^\alpha \in \mathcal{G}_{\mathbb{Q}})\text{”}.$$

2. We say that a forcing notion \mathbb{Q} is **reasonably A-bounding over $\bar{\mu}$** if
 - (a) \mathbb{Q} is strategically $(<\lambda)$ -complete, and
 - (b) for any $p \in \mathbb{Q}$, GENERIC has a winning strategy in the game $\mathcal{D}_{\bar{\mu}}^{\text{rcA}}(p, \mathbb{Q})$.

Definition 3.7 (See [7], Definition 3.2) Let $\bar{\mathbb{Q}} = \langle \mathbb{P}_\zeta, \mathbb{Q}_\zeta : \zeta < \gamma \rangle$ be a λ -support iteration and let $\bar{\mu} = \langle \mu_\alpha : \alpha < \lambda \rangle$ be a sequence of regular cardinals such that $\aleph_0 \leq \mu_\alpha \leq \lambda$ for all $\alpha < \lambda$.

1. For a condition $p \in \mathbb{P}_\gamma = \lim(\bar{\mathbb{Q}})$ we define a *tree A-completeness game* $\mathfrak{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ between two players, GENERIC and ANTIGENERIC, as follows. A play of $\mathfrak{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ lasts λ steps and in the course of a play a sequence $\langle \mathcal{T}_\alpha, \bar{p}^\alpha, \bar{q}^\alpha : \alpha < \lambda \rangle$ is constructed. Suppose that the players have arrived to a stage $\alpha < \lambda$ of the game. Now,

(\aleph) $_\alpha$ first GENERIC picks a standard $(w, 1)^\gamma$ -tree \mathcal{T}_α such that $|T_\alpha| < \mu_\alpha$ and a tree of conditions $\bar{p}^\alpha = \langle p_t^\alpha : t \in T_\alpha \rangle \subseteq \mathbb{P}_\gamma$ (so GENERIC, as a part of choosing \mathcal{T}_α , picks also $w = w_\alpha$),

(\beth) $_\alpha$ then ANTIGENERIC answers by choosing a tree of conditions $\bar{q}^\alpha = \langle q_t^\alpha : t \in T_\alpha \rangle \subseteq \mathbb{P}_\gamma$ such that $\bar{p}^\alpha \leq \bar{q}^\alpha$.

At the end, GENERIC wins the play $\langle \mathcal{T}_\alpha, \bar{p}^\alpha, \bar{q}^\alpha : \alpha < \lambda \rangle$ of $\mathfrak{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ if and only if

(\otimes) $_{\bar{A}}^{\text{tree}}$ there is a condition $p^* \in \mathbb{P}_\gamma$ stronger than p and such that

$$p^* \Vdash_{\mathbb{P}_\gamma} \text{“}(\forall \alpha < \lambda)(\exists t \in T_\alpha)(\text{rk}_\alpha(t) = \gamma \ \& \ q_t^\alpha \in G_{\mathbb{P}_\gamma})\text{”}.$$

2. We say that $\mathbb{P}_\gamma = \lim(\bar{\mathbb{Q}})$ is *reasonably* A($\bar{\mathbb{Q}}$)-bounding over $\bar{\mu}$* if GENERIC has a winning strategy in the game $\mathfrak{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ for every $p \in \mathbb{P}_\gamma$.

Theorem 3.8 (See [7], Theorem 3.2) *Assume that*

- (a) λ is a strongly inaccessible cardinal,
 (b) $\bar{\mu} = \langle \mu_\alpha : \alpha < \lambda \rangle$, each μ_α is a regular cardinal satisfying (for $\alpha < \lambda$)

$$\aleph_0 \leq \mu_\alpha \leq \lambda \quad \text{and} \quad (\forall f \in {}^\alpha \mu_\alpha)(\left| \prod_{\xi < \alpha} f(\xi) \right| < \mu_\alpha),$$

- (c) $\bar{\mathbb{Q}} = \langle \mathbb{P}_\xi, \mathbb{Q}_\xi : \xi < \gamma \rangle$ is a λ -support iteration such that for every $\xi < \gamma$,

$$\Vdash_{\mathbb{P}_\xi} \text{“}\mathbb{Q}_\xi \text{ is reasonably A-bounding over } \bar{\mu}\text{”}.$$

Then $\mathbb{P}_\gamma = \lim(\bar{\mathbb{Q}})$ is *reasonably* A($\bar{\mathbb{Q}}$)-bounding over $\bar{\mu}$* (and so \mathbb{P}_γ is also λ -proper).

In [7, §3], in addition to A-reasonable completeness game we considered its variant called *a-reasonable completeness game*. In that variant, at stage $\alpha < \lambda$ of the game, the players played a subgame to construct a sequence $\langle p_\xi^\alpha, q_\xi^\alpha : \xi < i_\alpha \rangle$ (corresponding to $\langle p_t^\alpha, q_t^\alpha : t \in I_\alpha \rangle$). In the following definition we introduce a further modification of that game. In the new game, the players will again play subgames, in some sense repeating several times the subgames from the *a-reasonable completeness game*.

Definition 3.9 Let \mathbb{Q} be a forcing notion and let $\bar{\mu} = \langle \mu_\alpha : \alpha < \lambda \rangle$ be a sequence of cardinals such that $\aleph_0 \leq \mu_\alpha < \lambda$ for all $\alpha < \lambda$. Suppose also that \mathcal{U} is a normal filter on λ .

1. For a condition $p \in \mathbb{Q}$ we define a *reasonable double-a-completeness game* $\mathfrak{D}_{\bar{\mu}}^{\text{rc2a}}(p, \mathbb{Q})$ between GENERIC and ANTIGENERIC as follows. A play of $\mathfrak{D}_{\bar{\mu}}^{\text{rc2a}}(p, \mathbb{Q})$ lasts at most λ steps and in the course of the play the players try to construct a sequence

$$(\boxtimes) \quad \langle \xi_\alpha, \langle p_\gamma^\alpha, q_\gamma^\alpha : \gamma < \mu_\alpha \cdot \xi_\alpha \rangle : \alpha < \lambda \rangle.$$

(Here μ_α is treated as an ordinal and $\mu_\alpha \cdot \xi_\alpha$ is the ordinal product of μ_α and ξ_α .) Suppose that the players have arrived to a stage $\alpha < \lambda$ of the game. First, ANTIGENERIC picks a nonzero ordinal $\xi_\alpha < \lambda$. Then the two players start a subgame of length $\mu_\alpha \cdot \xi_\alpha$ alternately choosing the terms of the sequence $\langle p_\gamma^\alpha, q_\gamma^\alpha : \gamma < \mu_\alpha \cdot \xi_\alpha \rangle$. At a stage $\gamma = \mu_\alpha \cdot i + j$ (where $i < \xi_\alpha, j < \mu_\alpha$) of the subgame, first GENERIC picks a condition $p_\gamma^\alpha \in \mathbb{Q}$ stronger than all conditions q_δ^α for $\delta < \gamma$ of the form $\delta = \mu_\alpha \cdot i' + j$ (where $i' < i$), and then ANTIGENERIC answers with a condition q_γ^α stronger than p_γ^α .

At the end, GENERIC wins the play (\boxtimes) of $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{Q})$ if and only if both players had always legal moves and

$(\otimes)_{2\mathbf{a}}^{\text{rc}}$ there is a condition $p^* \in \mathbb{Q}$ stronger than p and such that

$$p^* \Vdash_{\mathbb{Q}} \text{“}(\forall \alpha < \lambda)(\exists j < \mu_\alpha)(\{q_{\mu_\alpha \cdot i + j}^\alpha : i < \xi_\alpha\} \subseteq \mathcal{G}_{\mathbb{Q}})\text{”}.$$

2. Games $\mathcal{D}_{\bar{\mu}, \mathcal{U}}^{\text{rc}2\mathbf{b}}(p, \mathbb{Q})$ (for $p \in \mathbb{Q}$) are defined similarly; we only replace condition $(\otimes)_{2\mathbf{a}}^{\text{rc}}$ by

$(\otimes)_{2\mathbf{b}}^{\text{rc}}$ there is a condition $p^* \in \mathbb{Q}$ stronger than p and such that

$$p^* \Vdash_{\mathbb{Q}} \text{“}\{\alpha < \lambda : (\exists j < \mu_\alpha)(\{q_{\mu_\alpha \cdot i + j}^\alpha : i < \xi_\alpha\} \subseteq \mathcal{G}_{\mathbb{Q}})\} \in \mathcal{U}^{\mathbb{Q}}\text{”},$$

where $\mathcal{U}^{\mathbb{Q}}$ is the (\mathbb{Q} -name for the) normal filter generated by \mathcal{U} in $\mathbf{V}^{\mathbb{Q}}$.

3. A strategy \mathbf{st} for GENERIC in $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{Q})$ (or $\mathcal{D}_{\bar{\mu}, \mathcal{U}}^{\text{rc}2\mathbf{b}}(p, \mathbb{Q})$) is said to be *nice* if for every play $\langle \xi_\alpha, \langle p_\gamma^\alpha, q_\gamma^\alpha : \gamma < \mu_\alpha \cdot \xi_\alpha \rangle : \alpha < \lambda \rangle$ in which she uses \mathbf{st} , for every $\alpha < \lambda$, the conditions in $\{p_\gamma^\alpha : \gamma < \mu_\alpha\}$ are pairwise incompatible. (These are conditions played in the first “run” of the subgame. Note that then $p_\gamma^\alpha, p_{\gamma'}^{\alpha'}$ are incompatible whenever $\gamma \neq \gamma' \pmod{\mu_\alpha}$.)

4. Let $\mathbf{x} \in \{\mathbf{a}, \mathbf{b}\}$. A forcing notion \mathbb{Q} is *nicely double \mathbf{x} -bounding over $\bar{\mu}$* (and \mathcal{U} if $\mathbf{x} = \mathbf{b}$) if

(a) \mathbb{Q} is strategically $(< \lambda)$ -complete, and

(b) GENERIC has a nice winning strategy in the game $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{Q})$ ($\mathcal{D}_{\bar{\mu}, \mathcal{U}}^{\text{rc}2\mathbf{b}}(p, \mathbb{Q})$) if $\mathbf{x} = \mathbf{b}$) for every $p \in \mathbb{Q}$.

Remark 3.10

1. Reasonable double \mathbf{x} -boundedness (for $\mathbf{x} \in \{\mathbf{a}, \mathbf{b}\}$) is an iterable relative of reasonable \mathbf{x} -boundedness introduced in [7, Definition 3.1, pp. 206-7]. Technical differences in the definitions of suitable games are to achieve the preservation of the corresponding property in λ -support iterations (see Theorems 3.13, 3.14 below).

2. The game $\mathcal{D}_{\bar{\mu}, \mathcal{U}}^{\text{rc}2\mathbf{b}}(p, \mathbb{Q})$ is easier to win for GENERIC than $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{Q})$ (because the winning criterion is weaker). Therefore, if we are interested in λ -properness for λ -support iterations only, then Theorem 3.14 will cover a larger class of forcing notions than Theorem 3.13.

Definition 3.11 (See [7], Definition 6.1) Suppose that λ is inaccessible and $\bar{\kappa} = \langle \kappa_\alpha : \alpha < \lambda \rangle$ is a sequence of cardinals, $1 < \kappa_\alpha < \lambda$ for $\alpha < \lambda$. We define a forcing notion $\mathbb{P}^{\bar{\kappa}}$ as follows.

A condition in $\mathbb{P}^{\bar{\kappa}}$ is a pair $p = (f^p, C^p)$ such that

$$C^p \subseteq \lambda \text{ is a club of } \lambda \text{ and } f^p \in \prod \{\kappa_\iota : \iota \in \lambda \setminus C^p\}.$$

The order $\leq_{\mathbb{P}^{\bar{\kappa}}} = \leq$ of $\mathbb{P}^{\bar{\kappa}}$ is given by

$$p \leq_{\mathbb{P}^{\bar{\kappa}}} q \quad \text{if and only if} \quad C^q \subseteq C^p \text{ and } f^p \subseteq f^q.$$

Proposition 3.12

1. Assume that $\bar{\kappa}, \lambda$ are as in Definition 3.11 above and let a sequence $\bar{\mu} = \langle \mu_\alpha : \alpha < \lambda \rangle$ be chosen so that $\prod_{\beta < \alpha} \kappa_\beta \leq \mu_\alpha < \lambda$ (for $\alpha < \lambda$).
Then the forcing notion $\mathbb{P}^{\bar{\kappa}}$ is nicely double **b**-bounding over $\bar{\mu}, \mathcal{D}_\lambda$.
2. If $\kappa_\alpha = \kappa$ for all $\alpha < \lambda$ and $\mu_\alpha \geq \kappa^\alpha$, then $\mathbb{P}^{\bar{\kappa}}$ is nicely double **a**-bounding over $\bar{\mu}$.

Proof (1) A natural modification of the proof of [7, Proposition 6.1] works here. Note that if $\bar{\delta} = \langle \delta_\alpha : \alpha < \lambda \rangle$ is an increasing continuous sequence constructed as there during a play of $\mathcal{D}_{\bar{\mu}, \mathcal{D}_\lambda}^{\text{rc}2\mathbf{b}}(p, \mathbb{P}^{\bar{\kappa}})$, then the set $B \stackrel{\text{def}}{=} \{\alpha < \lambda : \prod_{\beta < \alpha} \kappa_{\delta_\beta} \leq \mu_\alpha\}$ is in the filter \mathcal{D}_λ . In the game, the stages $\alpha \in \lambda \setminus B$ are ignored and only those for $\alpha \in B$ are “active.” Also, at each stage α we may create μ_α “not active” steps at each run of the subgame by picking an antichain of conditions incompatible with p .

(2) Similar; we get double **a**-bounding here as at each stage $\alpha < \lambda$ of the game we know that $\prod_{\beta < \alpha} \kappa_{\delta_\beta} = \kappa^\alpha \leq \mu_\alpha$ (so all steps are “active”). \square

Theorem 3.13 Assume that

- (a) λ is a strongly inaccessible cardinal,
- (b) $\bar{\mu} = \langle \mu_\alpha : \alpha < \lambda \rangle$ is a sequence of cardinals below λ such that $(\forall \alpha < \lambda)(\aleph_0 \leq \mu_\alpha = \mu_\alpha^{|\alpha+1|})$,
- (c) $\bar{\mathbb{Q}} = \langle \mathbb{P}_\zeta, \mathbb{Q}_\zeta : \zeta < \zeta^* \rangle$ is a λ -support iteration such that for every $\zeta < \gamma$,

$$\Vdash_{\mathbb{P}_\zeta} \text{“}\mathbb{Q}_\zeta \text{ is nicely double } \mathbf{a}\text{-bounding over } \bar{\mu}\text{”}.$$

Then $\mathbb{P}_{\zeta^*} = \lim(\bar{\mathbb{Q}})$ is nicely double **a**-bounding over $\bar{\mu}$ (and so \mathbb{P}_{ζ^*} is also λ -proper).

Proof Our arguments refine those presented in the proof of [7, Theorem 3.2, p. 217], but the differences in the games involved eliminate the use of trees of conditions. However, trees of conditions are implicitly present here too. The tree at level δ of the argument is indexed by

$$T_\delta = \bigcup \left\{ \prod_{\zeta \in w_\delta \cap \zeta} \mu_\delta : \zeta \in w_\delta \cup \{\zeta^*\} \right\}$$

and it is formed in part by conditions played in the game for various $t \in \prod_{\zeta \in w_\delta} \mu_\delta = \{t \in T_\delta : \text{rk}(t) = \zeta^*\}$; note the coherence demand in $(\boxtimes)_7$.

Let $p \in \mathbb{P}_{\zeta^*}$. We will describe a strategy **st** for GENERIC in the game $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{P}_{\zeta^*})$. The strategy **st** instructs GENERIC to play the game $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}$ on each relevant coordinate $\zeta < \zeta^*$ using her winning strategy **st** $_\zeta$. At stage $\delta < \lambda$ GENERIC will be concerned with coordinates $\zeta \in w_\delta$ for some set w_δ of size $< \lambda$. If $\zeta_\delta < \lambda$ is the ordinal put by ANTIGENERIC in the play of $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{P}_{\zeta^*})$, then in the simulated plays on coordinates $\zeta \in w_\delta$ GENERIC pretends that her opponent put $\zeta_\delta^* = \mu_\delta \cdot \zeta_\delta$.

The innings of the two players, GENERIC and ANTIGENERIC, in the subgame of level δ on a coordinate ζ are

$$\bar{p}_{\delta,\zeta} = \langle p_{\delta,\zeta}^\gamma : \gamma < \mu_\delta \cdot \zeta_\delta^* \rangle \text{ and } \bar{q}_{\delta,\zeta} = \langle q_{\delta,\zeta}^\gamma : \gamma < \mu_\delta \cdot \zeta_\delta^* \rangle,$$

respectively. GENERIC's innings in the subgame of $\mathcal{D}_\mu^{\text{rc}2\mathbf{a}}(p, \mathbb{P}_{\zeta^*})$ will be associated with sequences $t \in \prod_{\zeta \in w_\delta} \mu_\delta = \langle t_j^\delta : j < \mu_\delta \rangle = \bar{t}^\delta$. The innings of the two players will be $p_\varepsilon^\delta, q_\varepsilon^\delta$ (for $\varepsilon < \mu_\delta \cdot \zeta_\delta$) and they will be related to what happens at coordinates $\zeta \in w_\delta$ as follows. If $t = t_j^\delta, \zeta \in w_\delta$, and $\beta = (t)_\zeta < \mu_\delta$, then in the subgame of $\mathcal{D}_\mu^{\text{rc}2\mathbf{a}}(p, \mathbb{P}_{\zeta^*})$ of level δ at stages of the form $\varepsilon = \mu_\delta \cdot i + j$ we will have $p_\varepsilon^\delta(\zeta) = p_{\delta,\zeta}^\gamma$ and $q_\varepsilon^\delta(\zeta) = q_{\delta,\zeta}^\gamma$, where $\gamma = \mu_\delta \cdot \varepsilon + \beta$. To keep track of what happens at coordinates $\zeta \notin w_\delta$ GENERIC will use conditions r_δ .

Let us note that the construction of \mathbf{st} presented in detail below would be somewhat simpler if we knew that all forcings \mathbb{Q}_ζ are $(<\lambda)$ -complete (and not only strategically $(<\lambda)$ -complete). Then $\underline{\mathbf{st}}_\zeta^0, r_\delta^-$ and $p_\varepsilon^{\delta,*}$ could be eliminated as their role is to make sure that some sequences of conditions (related to r_δ and/or p_ε^δ) have upper bounds. However, many natural forcing notions tend to have strategic completeness only (see [8, Part B]).

Let us formalize the ideas presented above. For each $\zeta < \zeta^*$, pick a \mathbb{P}_ζ -name $\underline{\mathbf{st}}_\zeta^0$ such that

$$\Vdash_{\mathbb{P}_\zeta} \text{“}\underline{\mathbf{st}}_\zeta^0 \text{ is a winning strategy for COMPLETE in } \mathcal{D}_0^\lambda(\mathbb{Q}_\zeta, \mathcal{Q}_{\mathbb{Q}_\zeta}) \text{ such that if INCOMPLETE plays } \mathcal{Q}_{\mathbb{Q}_\zeta}, \text{ then COMPLETE answers with } \mathcal{Q}_{\mathbb{Q}_\zeta} \text{ as well”}.$$

In the course of a play of $\mathcal{D}_\mu^{\text{rc}2\mathbf{a}}(p, \mathbb{P}_{\zeta^*})$, at a stage $\delta < \lambda$, GENERIC will be instructed to construct on the side

$$(\otimes)_\delta \quad w_\delta, \bar{t}^\delta, \zeta_\delta^*, \underline{\mathbf{st}}_\zeta \text{ (for } \zeta \in w_{\delta+1} \setminus w_\delta), \bar{p}_{\delta,\zeta}, \bar{q}_{\delta,\zeta}, p_\varepsilon^{\delta,*} \text{ (for } \varepsilon < \mu_\delta \cdot \zeta_\delta), \text{ and } r_\delta^-, r_\delta.$$

These objects will be chosen so that if

$$\langle \zeta_\delta, \langle p_\gamma^\delta, q_\gamma^\delta : \gamma < \mu_\delta \cdot \zeta_\delta \rangle : \delta < \lambda \rangle$$

is a play of $\mathcal{D}_\mu^{\text{rc}2\mathbf{a}}(p, \mathbb{P}_{\zeta^*})$ in which GENERIC follows \mathbf{st} , and the additional objects constructed at stage $\delta < \lambda$ are listed in $(\otimes)_\delta$, then the following conditions are satisfied (for each $\delta < \lambda$).

- (\boxtimes)₁ $r_\delta^-, r_\delta \in \mathbb{P}_{\zeta^*}, r_0^-(0) = r_0(0) = p(0), w_\delta \subseteq \zeta^*, |w_\delta| = |\delta + 1|,$
 $\bigcup_{\alpha < \lambda} \text{Dom}(r_\alpha) = \bigcup_{\alpha < \lambda} w_\alpha, w_0 = \{0\}, w_\delta \subseteq w_{\delta+1}$ and if δ is limit then
 $w_\delta = \bigcup_{\alpha < \delta} w_\alpha.$
- (\boxtimes)₂ For each $\alpha < \delta < \lambda$ we have $(\forall \zeta \in w_{\alpha+1})(r_\alpha(\zeta) = r_\delta^-(\zeta) = r_\delta(\zeta))$ and
 $p \leq r_\alpha^- \leq r_\alpha \leq r_\delta^- \leq r_\delta$, and $p_\varepsilon^{\delta,*} \in \mathbb{P}_{\zeta^*}$ (for $\varepsilon < \mu_\delta \cdot \zeta_\delta$).
- (\boxtimes)₃ If $\zeta \in \zeta^* \setminus w_\delta$, then
 $r_\delta \upharpoonright \zeta \Vdash_{\mathbb{P}_\zeta}$ “the sequence $\langle r_\alpha^-(\zeta), r_\alpha(\zeta) : \alpha \leq \delta \rangle$ is a legal partial play of
 $\mathcal{D}_0^\lambda(\mathbb{Q}_\zeta, \mathcal{Q}_{\mathbb{Q}_\zeta})$ in which COMPLETE follows $\underline{\mathbf{st}}_\zeta^0$ ”,

and if $\zeta \in w_{\delta+1} \setminus w_\delta$, then \mathbf{st}_ζ is a \mathbb{P}_ζ -name for a nice winning strategy for **GENERIC** in $\mathcal{D}_{\bar{\mu}}^{\text{rc2a}}(r_\delta(\zeta), \mathbb{Q}_\zeta)$. (And \mathbf{st}_0 is a nice winning strategy of **GENERIC** in $\mathcal{D}_{\bar{\mu}}^{\text{rc2a}}(p(0), \mathbb{Q}_0)$.)

- (\boxtimes)₄ $\bar{t}^\delta = \langle t_j^\delta : j < \mu_\delta \rangle$ is an enumeration of $\prod_{\zeta \in w_\delta} \mu_\delta = {}^{w_\delta} \mu_\delta$.
- (\boxtimes)₅ $\zeta_\delta^* = \mu_\delta \cdot \zeta_\delta$ (the ordinal product) and $\bar{p}_{\delta, \zeta}^\gamma = \langle p_{\delta, \zeta}^\gamma : \gamma < \mu_\delta \cdot \zeta_\delta^* \rangle$ and $\bar{q}_{\delta, \zeta}^\gamma = \langle q_{\delta, \zeta}^\gamma : \gamma < \mu_\delta \cdot \zeta_\delta^* \rangle$ are \mathbb{P}_ζ -names for sequences of conditions in \mathbb{Q}_ζ of length $\mu_\delta \cdot \zeta_\delta^*$ (for $\zeta \in \bigcup_{\alpha < \lambda} w_\alpha$).
- (\boxtimes)₆ If $\zeta \in w_{\beta+1} \setminus w_\beta$, $\beta < \delta$ (or $\zeta = \beta = 0$), then $\Vdash_{\mathbb{P}_\zeta}$ “ $\langle \zeta_\alpha^*, \langle p_{\alpha, \zeta}^\gamma, q_{\alpha, \zeta}^\gamma : \gamma < \mu_\alpha \cdot \zeta_\alpha^* \rangle : \alpha \leq \delta \rangle$ is a partial play of $\mathcal{D}_{\bar{\mu}}^{\text{rc2a}}(r_\beta(\zeta), \mathbb{Q}_\zeta)$ in which **GENERIC** uses \mathbf{st}_ζ ”.
- (\boxtimes)₇ If $\varepsilon = \mu_\delta \cdot i + j$, $i < \zeta_\delta$, $j < \mu_\delta$, then $\text{Dom}(p_\varepsilon^{\delta, *}) = \text{Dom}(p_\varepsilon^\delta) = w_\delta \cup \text{Dom}(p) \cup \bigcup_{\alpha < \delta} \text{Dom}(r_\alpha) \cup \bigcup_{\varepsilon' < \varepsilon} \text{Dom}(q_\varepsilon^{\delta, *})$, and for each $\zeta \in w_\delta \cup \{\zeta^*\}$ the condition $p_\varepsilon^{\delta, *} \upharpoonright \zeta$ is an upper bound to $\{p \upharpoonright \zeta\} \cup \{r_\alpha \upharpoonright \zeta : \alpha < \delta\} \cup \{q_\varepsilon^{\delta, *} \upharpoonright \zeta : \varepsilon' = \mu_\delta \cdot i' + j' < \varepsilon \ \& \ i' < \zeta_\delta \ \& \ j' < \mu_\delta \ \& \ t_{j'}^\delta \upharpoonright \zeta = t_j^\delta \upharpoonright \zeta\}$.
- (\boxtimes)₈ If $j < \mu_\delta$, $i < \zeta_\delta$, $\zeta \in w_\delta$, $(t_j^\delta)_\zeta = \beta$, and $\varepsilon = \mu_\delta \cdot i + j$, $\gamma = \mu_\delta \cdot \varepsilon + \beta$, then $p_\varepsilon^{\delta, *}(\zeta) = p_\varepsilon^\delta(\zeta) = p_{\delta, \zeta}^\gamma$ and $q_\varepsilon^{\delta, *} \upharpoonright \zeta \Vdash_{\mathbb{P}_\zeta} q_\varepsilon^\delta(\zeta) = q_{\delta, \zeta}^\gamma$.
- (\boxtimes)₉ If $\varepsilon = \mu_\delta \cdot i + j$, $i < \zeta_\delta$, $j < \mu_\delta$, $\zeta \in \zeta^* \setminus w_\delta$, and $t \in \prod \{\mu_\delta : \zeta \in w_\delta \cap \zeta\}$, $t \leq t_j^\delta$, then $p_\varepsilon^{\delta, *} \upharpoonright \zeta \Vdash_{\mathbb{P}_\zeta}$ “the sequence $\langle p_\varepsilon^{\delta, *}(\zeta), p_\varepsilon^\delta(\zeta) : \varepsilon' = \mu_\delta \cdot i' + j' \leq \varepsilon \ \& \ i' < \zeta_\delta \ \& \ j' < \mu_\delta \ \& \ t \leq t_{j'}^\delta \rangle$ is a legal partial play of $\mathcal{D}_0^\lambda(\mathbb{Q}_\zeta, p(\zeta))$ in which **COMPLETE** follows \mathbf{st}_ζ^0 ”.
- (\boxtimes)₁₀ $\text{Dom}(r_\delta^-) = \text{Dom}(r_\delta) = \bigcup \{\text{Dom}(q_\varepsilon^\delta) : \varepsilon < \mu_\delta \cdot \zeta_\delta\}$ and if $\zeta \in \zeta^* \setminus w_\delta$, $t \in \prod \{\mu_\delta : \zeta \in w_\delta \cap \zeta\}$, and $q \in \mathbb{P}_\zeta$, $q \geq r_\delta^- \upharpoonright \zeta$ and $q \geq q_\varepsilon^\delta \upharpoonright \zeta$ whenever $\varepsilon = \mu_\delta \cdot i + j$, $i < \zeta_\delta$, $j < \mu_\delta$, and $t \leq t_j^\delta$, then $q \Vdash_{\mathbb{P}_\zeta}$ “if the set $\{p(\zeta)\} \cup \{r_\alpha(\zeta) : \alpha < \delta\} \cup \{q_\varepsilon^\delta(\zeta) : \varepsilon = \mu_\delta \cdot i + j \ \& \ i < \zeta_\delta \ \& \ j < \mu_\delta \ \& \ t \leq t_j^\delta\}$ has an upper bound in \mathbb{Q}_ζ , then $r_\delta^-(\zeta)$ is such an upper bound; otherwise, $r_\delta^-(\zeta)$ is just an upper bound to $\{p(\zeta)\} \cup \{r_\alpha(\zeta) : \alpha < \delta\}$ ”.

Assume that the two players arrived to stage δ of $\mathcal{D}_{\bar{\mu}}^{\text{rc2a}}(p, \mathbb{P}_{\zeta^*})$ and

$$\langle \zeta_\alpha, \langle p_\varepsilon^\alpha, q_\varepsilon^\alpha : \varepsilon < \mu_\alpha \cdot \zeta_\alpha \rangle : \alpha < \delta \rangle$$

is the play constructed so far, and that **GENERIC** followed \mathbf{st} and determined objects listed in (\boxtimes) _{α} (for $\alpha < \delta$) with properties (\boxtimes)₁–(\boxtimes)₁₀.

Below, whenever we say **GENERIC chooses x such that** we mean **GENERIC chooses the $<^*_\chi$ -first x such that**, and so on. First, **GENERIC** uses her favorite bookkeeping device to determine w_δ so that the demands of (\boxtimes)₁ are satisfied (and that at the end we will have $\bigcup_{\alpha < \lambda} \text{Dom}(r_\alpha) = \bigcup_{\alpha < \lambda} w_\alpha$). If $\beta < \delta$ and $\zeta \in w_\beta$, then we already have $\bar{p}_{\alpha, \zeta}, \bar{q}_{\alpha, \zeta}$ for $\alpha < \delta$ (see (\boxtimes)₆), but we have not yet defined those objects

when $\delta = \delta_0 + 1$ and $\zeta \in w_\delta \setminus w_{\delta_0}$. So if $\delta = \delta_0 + 1$ and $\zeta \in w_\delta \setminus w_{\delta_0}$ then let $\bar{p}_{\alpha,\zeta} = \langle p_{\alpha,\zeta}^\gamma : \gamma < \mu_\alpha \cdot \zeta_\alpha^* \rangle$ and $\bar{q}_{\alpha,\zeta} = \langle q_{\alpha,\zeta}^\gamma : \gamma < \mu_\alpha \cdot \zeta_\alpha^* \rangle$ (for $\alpha < \delta$) be such that

$\Vdash_{\mathbb{P}_\zeta}$ “ $\langle \zeta_\alpha^*, \langle p_{\alpha,\zeta}^\gamma, q_{\alpha,\zeta}^\gamma : \gamma < \mu_\alpha \cdot \zeta_\alpha^* \rangle : \alpha < \delta \rangle$ is a partial play of $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(r_{\delta_0}(\zeta), \mathbb{Q}_\zeta)$ in which GENERIC uses \mathbf{st}_ζ and $p_{\alpha,\zeta}^\gamma = q_{\alpha,\zeta}^\gamma$ for all $\alpha < \delta, \gamma < \mu_\alpha \cdot \zeta_\alpha^*$ ”.

Condition $(\boxtimes)_4$ and our rule of taking “the $<^*$ -first” determine the enumeration $\bar{t}^\delta = \langle t_j^\delta : j < \mu_\delta \rangle$ of $\prod_{\zeta \in w_\delta} \mu_\delta$. Now ANTIGENERIC picks ζ_δ and the two players start a subgame of length $\mu_\delta \cdot \zeta_\delta$. During the subgame GENERIC will simulate subgames of level δ at coordinates $\zeta \in w_\delta$ pretending that ANTIGENERIC played $\zeta_\delta^* = \mu_\delta \cdot \zeta_\delta$ there. Each step in the subgame of $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{P}_{\zeta^*})$ will correspond to μ_δ steps in the subgames of $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(r_\beta(\zeta), \mathbb{Q}_\zeta)$ (when $\zeta \in w_{\beta+1} \setminus w_\beta, \beta < \delta$). So suppose that the two opponents have arrived to a stage $\varepsilon = \mu_\delta \cdot i + j$ of the subgame, $i < \zeta_\delta, j < \mu_\delta$, and assume also that GENERIC (playing according to \mathbf{st}) has already defined $p_{\delta,\zeta}^\gamma, q_{\delta,\zeta}^\gamma$ for $\zeta \in w_\delta, \gamma < \mu_\delta \cdot \varepsilon$, and $p_{\varepsilon'}^{\delta,*}$ for $\varepsilon' < \varepsilon$, so that the requirements of $(\boxtimes)_6$ – $(\boxtimes)_9$ are satisfied. Note that (by $(\boxtimes)_7$ – $(\boxtimes)_9$)

(*) if $\varepsilon > \varepsilon' = \mu_\delta \cdot i' + j' > \varepsilon'' = \mu_\delta \cdot i'' + j'', \zeta \in w_\delta \cup \{\zeta^*\}$ and $t_{j'}^\delta \upharpoonright \zeta = t_{j''}^\delta \upharpoonright \zeta$, then $p_{\varepsilon''}^{\delta,*} \upharpoonright \zeta \leq q_{\varepsilon''}^\delta \upharpoonright \zeta \leq p_{\varepsilon'}^{\delta,*} \upharpoonright \zeta \leq p_\varepsilon^\delta \upharpoonright \zeta$.

For each $\zeta \in w_\delta$ and $\beta < (t_j^\delta)_\zeta$, let $p_{\delta,\zeta}^{\mu_\delta \cdot \varepsilon + \beta} = q_{\delta,\zeta}^{\mu_\delta \cdot \varepsilon + \beta}$ be \mathbb{P}_ζ -names for conditions in \mathbb{Q}_ζ such that (the relevant part of) $(\boxtimes)_6$ holds. The same clause determines also $p_{\delta,\zeta}^{\mu_\delta \cdot \varepsilon + \beta}$ for $\beta = (t_j^\delta)_\zeta, \zeta \in w_\delta$. Then the requirements in $(\boxtimes)_7 + (\boxtimes)_8$ essentially describe what $p_\varepsilon^{\delta,*}$ is. Note that the “upper bound demands” in $(\boxtimes)_7$ can be satisfied because of $(\boxtimes)_9 + (\boxtimes)_3$ and (*) above. Next, GENERIC’s inning p_ε^δ in $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{P}_{\zeta^*})$ is chosen so that $\text{Dom}(p_\varepsilon^\delta) = \text{Dom}(p_\varepsilon^{\delta,*})$ and clauses $(\boxtimes)_8 + (\boxtimes)_9$ hold. After this ANTIGENERIC answers with a condition $q_\varepsilon^\delta \geq p_\varepsilon^\delta$, and GENERIC picks for the construction on the side names $q_{\delta,\zeta}^{\mu_\delta \cdot \varepsilon + \beta}$ for $\zeta \in w_\delta$ and $\beta = (t_j^\delta)_\zeta$ by the demand in $(\boxtimes)_8$. She also picks $p_{\delta,\zeta}^{\mu_\delta \cdot \varepsilon + \beta} = q_{\delta,\zeta}^{\mu_\delta \cdot \varepsilon + \beta}$ for $\zeta \in w_\delta$ and $(t_j^\delta)_\zeta < \beta < \mu_\delta$ so that $(\boxtimes)_6$ holds.

This completes the description of what happens during the $\mu_\delta \cdot \zeta_\delta$ steps of the subgame. After the subgame is over and the sequence $\langle p_\gamma^\delta, q_\gamma^\delta : \gamma < \mu_\delta \cdot \zeta_\delta \rangle$ is constructed, GENERIC chooses conditions $r_\delta^-, r_\delta \in \mathbb{P}_{\zeta^*}$ by $(\boxtimes)_1$ – $(\boxtimes)_3$ and $(\boxtimes)_10$. (Note: since \mathbf{st}_ζ are names for nice strategies, if $\zeta \in \zeta^* \setminus w_\delta, i_0, i_1 < \zeta_\delta, j_0, j_1 < \mu_\delta, \varepsilon_0 = \mu_\delta \cdot i_0 + j_0, \varepsilon_1 = \mu_\delta \cdot i_1 + j_1, t_0, t_1 \in \prod \{\mu_\delta : \zeta \in w_\delta \cap \zeta\}, t_0 \leq t_{j_0}^\delta, t_1 \leq t_{j_1}^\delta$ and $t_0 \neq t_1$, then the conditions $q_{\varepsilon_0}^\delta \upharpoonright \zeta, q_{\varepsilon_1}^\delta \upharpoonright \zeta$ are incompatible.) This finishes the description of the strategy \mathbf{st} .

Let us argue that \mathbf{st} is a winning strategy for GENERIC. Suppose that

$$\langle \zeta_\delta, \langle p_\gamma^\delta, q_\gamma^\delta : \gamma < \mu_\delta \cdot \zeta_\delta \rangle : \delta < \lambda \rangle$$

is a play of $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{P}_{\zeta^*})$ in which GENERIC followed \mathbf{st} and she constructed the side objects listed in $(\boxtimes)_\delta$ (for $\delta < \lambda$) so that demands $(\boxtimes)_1$ – $(\boxtimes)_10$ are satisfied. We define a condition $r \in \mathbb{P}_{\zeta^*}$ as follows. Let $\text{Dom}(r) = \bigcup_{\delta < \lambda} \text{Dom}(r_\delta)$. For $\zeta \in \text{Dom}(r)$ let $r(\zeta)$ be a \mathbb{P}_ζ -name for a condition in \mathbb{Q}_ζ such that

$(\boxtimes)_11$ if $\zeta \in w_{\alpha+1} \setminus w_\alpha, \alpha < \lambda$ (or $\zeta = \alpha = 0$), then

$\Vdash_{\mathbb{P}_\zeta}$ “ $r(\zeta) \geq r_\alpha(\zeta)$ and $r(\zeta) \Vdash_{\mathbb{Q}_\zeta} (\forall \delta < \lambda) (\exists j < \mu_\delta) (\forall \varepsilon < \zeta_\delta^*) (q_{\delta, \zeta}^{\mu_\delta \cdot \varepsilon + j} \in \mathcal{G}_{\mathbb{Q}_\zeta})$ ”.

Clearly, r is well defined (remember $(\boxtimes)_6$) and $(\forall \delta < \lambda)(r_\delta \leq r)$ and $p \leq r$.

Suppose now that $\delta < \lambda$ and $r' \geq r$. We are going to find $j < \mu_\delta$ and a condition $r'' \geq r'$ such that $(\forall i < \zeta_\delta)(q_{\mu_\delta \cdot i + j}^\delta \leq r'')$. To this end let $\langle \zeta_\alpha : \alpha \leq \alpha^* \rangle$ be the increasing enumeration of $w_\delta \cup \{\zeta^*\}$. For $\zeta \leq \zeta^*$ and $q \in \mathbb{P}_\zeta$, let $\mathbf{st}(\zeta, q)$ be a winning strategy of COMPLETE in $\mathcal{D}_0^{\lambda}(\mathbb{P}_\zeta, q)$ with the coherence properties given in Proposition 3.3.

By induction on $\alpha \leq \alpha^*$ we will choose conditions $r_\alpha^*, r_\alpha^{**} \in \mathbb{P}_{\zeta_\alpha}$ and $(t)_{\zeta_\alpha} < \mu_\delta$ such that

$$(\boxtimes)_{12} \quad r' \upharpoonright \zeta_\alpha \leq r_\alpha^*,$$

$$(\boxtimes)_{13} \quad \text{if } i < \zeta_\delta, j < \mu_\delta \text{ and } (t_j^\delta)_{\zeta_\beta} = (t)_{\zeta_\beta} \text{ for } \beta < \alpha, \text{ then } q_{\mu_\delta \cdot i + j}^\delta \upharpoonright \zeta_\alpha \leq r_\alpha^*,$$

$$(\boxtimes)_{14} \quad \langle r_\beta^* \widehat{\smile} r' \upharpoonright [\zeta_\beta, \zeta^*), r_\beta^{**} \widehat{\smile} r' \upharpoonright [\zeta_\beta, \zeta^*) : \beta < \alpha \rangle \text{ is a partial legal play of } \mathcal{D}_0^{\lambda}(\mathbb{P}_{\zeta^*}, r') \text{ in which COMPLETE uses her winning strategy } \mathbf{st}(\zeta^*, r').$$

Suppose that $\alpha \leq \alpha^*$ is a limit ordinal and we have already defined $(t)_{\zeta_\beta} < \mu_\delta$ and $r_\beta^*, r_\beta^{**} \in \mathbb{P}_{\zeta_\beta}$ for $\beta < \alpha$. Let $\zeta = \sup(\zeta_\beta : \beta < \alpha)$. It follows from $(\boxtimes)_{14}$ that we may pick a condition $s \in \mathbb{P}_\zeta$ stronger than all r_β^{**} for $\beta < \alpha$. Put $r_\alpha^* = s \widehat{\smile} r' \upharpoonright [\zeta, \zeta_\alpha) \in \mathbb{P}_{\zeta_\alpha}$. Then plainly $r' \upharpoonright \zeta_\alpha \leq r_\alpha^*$ and $q_{\mu_\delta \cdot i + j}^\delta \upharpoonright \zeta \leq r_\alpha^* \upharpoonright \zeta$ whenever

$$(\boxtimes)_{15}^{i,j,\alpha} \quad i < \zeta_\delta, j < \mu_\delta, \text{ and } (t_j^\delta)_{\zeta_\beta} = (t)_{\zeta_\beta} \text{ for all } \beta < \alpha.$$

Now by induction on $\zeta \leq \zeta_\alpha$ we show that $q_{\mu_\delta \cdot i + j}^\delta \upharpoonright \zeta \leq r_\alpha^* \upharpoonright \zeta$ whenever $(\boxtimes)_{15}^{i,j,\alpha}$ holds. For $\zeta \leq \zeta$ we are already done, so assume $\zeta \in [\zeta, \zeta_\alpha)$ and we have shown that $q_{\mu_\delta \cdot i + j}^\delta \upharpoonright \zeta \leq r_\alpha^* \upharpoonright \zeta$ whenever $(\boxtimes)_{15}^{i,j,\alpha}$ holds. It follows from $(\boxtimes)_7$ + $(\boxtimes)_9$ that the condition $r_\alpha^* \upharpoonright \zeta$ forces in \mathbb{P}_ζ that

$$\text{“the set } \{p(\zeta)\} \cup \{r_\alpha(\zeta) : \alpha < \delta\} \cup \{q_\varepsilon^\delta(\zeta) : \varepsilon = \mu_\delta \cdot i + j \ \& \ i < \zeta_\delta \ \& \ j < \mu_\delta \ \& \ (\forall \beta < \alpha)((t_j^\delta)_{\zeta_\beta} = (t)_{\zeta_\beta})\} \text{ has an upper bound in } \mathbb{Q}_\zeta\text{”}.$$

and therefore we may use $(\boxtimes)_{10}$ to conclude that

$$r_\alpha^* \upharpoonright \zeta \Vdash \text{“if } (\boxtimes)_{15}^{i,j,\alpha} \text{ holds, then } q_{\mu_\delta \cdot i + j}^\delta(\zeta) \leq r_\delta(\zeta) \leq r'(\zeta) = r_\alpha^*(\zeta)\text{”}.$$

The limit stages are trivial and we may claim that $q_{\mu_\delta \cdot i + j}^\delta \upharpoonright \zeta_\alpha \leq r_\alpha^*$ whenever $(\boxtimes)_{15}^{i,j,\alpha}$ holds. Next, r_α^{**} is determined by $(\boxtimes)_{14}$.

Now suppose that $\alpha = \beta + 1 \leq \alpha^*$ and we have already defined $r_\beta^*, r_\beta^{**} \in \mathbb{P}_{\zeta_\beta}$ and $\langle (t)_{\zeta_\gamma} : \gamma < \beta \rangle$. It follows from $(\boxtimes)_{11}$ that

$$r_\beta^{**} \Vdash_{\mathbb{P}_{\zeta_\beta}} \text{“} r(\zeta_\beta) \Vdash_{\mathbb{Q}_{\zeta_\beta}} (\exists \rho < \mu_\delta) (\forall \varepsilon < \zeta_\delta^*) (q_{\delta, \zeta_\beta}^{\mu_\delta \cdot \varepsilon + \rho} \in \mathcal{G}_{\mathbb{Q}_{\zeta_\beta}})\text{”},$$

so we may pick $\rho = (t)_{\zeta_\beta}$ and a condition $s \in \mathbb{P}_{\zeta_{\beta+1}}$ such that $r_\beta^{**} \leq s \upharpoonright \zeta_\beta$ and

$$s \upharpoonright \zeta_\beta \Vdash_{\mathbb{P}_{\zeta_\beta}} (\forall \varepsilon < \zeta_\delta^*) (q_{\delta, \zeta_\beta}^{\mu_\delta \cdot \varepsilon + \rho} \leq s(\zeta_\beta)).$$

It follows from $(\boxtimes)_{13}$ + $(\boxtimes)_8$ that then also $q_{\mu_\delta \cdot i + j}^\delta \upharpoonright (\zeta_\beta + 1) \leq s$ whenever $i < \zeta_\delta$, $j < \mu_\delta$, and $(t_j^\delta)_{\zeta_\gamma} = (t)_{\zeta_\gamma}$ for $\gamma \leq \beta$. We let $r_\alpha^* = s \widehat{\smile} r' \upharpoonright (\zeta_\beta, \zeta_\alpha)$ and exactly like in the limit case we argue that $r' \upharpoonright \zeta_\alpha \leq r_\alpha^*$ and $q_{\mu_\delta \cdot i + j}^\delta \upharpoonright \zeta_\alpha \leq r_\alpha^*$ whenever $i < \zeta_\delta$, $j < \mu_\delta$, and $(t_j^\delta)_{\zeta_\gamma} = (t)_{\zeta_\gamma}$ for $\gamma \leq \beta$. Again, r_α^{**} is determined by $(\boxtimes)_{14}$.

After the induction is completed look at $r'' = r_\alpha^*$ and $j < \mu_\delta$ such that $t_j^\delta = \langle (t)_{\zeta_\alpha} : \alpha < \alpha^* \rangle$. \square

Theorem 3.14 Assume (a), (b) of Theorem 3.13. Suppose that \mathcal{U} is a normal filter on λ and

(c) $\bar{\mathbb{Q}} = \langle \mathbb{P}_\zeta, \mathbb{Q}_\zeta : \zeta < \zeta^* \rangle$ is a λ -support iteration such that for every $\zeta < \gamma$,

$$\Vdash_{\mathbb{P}_\zeta} \text{“}\mathbb{Q}_\zeta \text{ is nicely double } \mathbf{b}\text{-bounding over } \bar{\mu}, \mathcal{U}^{\mathbb{P}_\zeta}\text{”}.$$

Then $\mathbb{P}_{\zeta^*} = \lim(\bar{\mathbb{Q}})$ is nicely double \mathbf{b} -bounding over $\bar{\mu}, \mathcal{U}$.

Proof The proof essentially repeats that of Theorem 3.13 with the following modifications in the arguments that \mathbf{st} is a winning strategy for **GENERIC** in $\mathfrak{D}_{\bar{\mu}, \mathcal{U}}^{\text{rc}2\mathbf{b}}(p, \mathbb{P}_{\zeta^*})$.

We assume that $\langle \zeta_\delta, \langle p_\gamma^\delta, q_\gamma^\delta : \gamma < \mu_\delta \cdot \zeta_\delta \rangle : \delta < \lambda \rangle$ is a play in which **GENERIC** follows \mathbf{st} and the objects listed in $(\otimes)_\delta$ were constructed on a side. A condition $r \in \mathbb{P}_{\zeta^*}$ is chosen so that $\text{Dom}(r) = \bigcup_{\delta < \lambda} \text{Dom}(r_\delta) = \bigcup_{\delta < \lambda} w_\delta$ and for each $\zeta \in w_{\alpha+1} \setminus w_\alpha$, $\alpha < \lambda$, we have

$$\begin{aligned} &\Vdash_{\mathbb{P}_\zeta} \text{“}r(\zeta) \geq r_\alpha(\zeta) \text{ and} \\ &r(\zeta) \Vdash_{\mathbb{Q}_\zeta} \{ \delta < \lambda : (\exists j < \mu_\delta)(\forall \varepsilon < \zeta_\delta^*)(q_{\delta, \zeta}^{\mu_\delta \cdot \varepsilon + j} \in \underline{G}_{\mathbb{Q}_\zeta}) \} \in \mathcal{U}^{\mathbb{P}_{\zeta+1}} \text{”}. \end{aligned}$$

Then, for each $\zeta \in \text{Dom}(r)$, we choose $\mathbb{P}_{\zeta+1}$ -names A_i^ζ for elements of \mathcal{U} such that

$$\Vdash_{\mathbb{P}_\zeta} \text{“}r(\zeta) \Vdash_{\mathbb{Q}_\zeta} (\forall \delta \in \Delta_{\delta < \lambda} A_i^\zeta)(\exists j < \mu_\delta)(\forall \varepsilon < \zeta_\delta^*)(q_{\delta, \zeta}^{\mu_\delta \cdot \varepsilon + j} \in \underline{G}_{\mathbb{Q}_\zeta})\text{”}.$$

Finally, we show that for each limit ordinal $\delta < \lambda$,

$$r \Vdash_{\mathbb{P}_{\zeta^*}} \text{“}(\forall \zeta \in w_\delta)(\delta \in \Delta_{\delta < \lambda} A_i^\zeta) \Rightarrow (\exists j < \mu_\delta)(\forall i < \zeta_\delta)(q_{\mu_\delta \cdot i + j}^\delta \in \underline{G}_{\mathbb{P}_{\zeta^*}})\text{”}.$$

For this we start with arbitrary condition $r' \geq r$ such that

$$r \Vdash_{\mathbb{P}_{\zeta^*}} \text{“}(\forall \zeta \in w_\delta)(\delta \in \Delta_{\delta < \lambda} A_i^\zeta)\text{”}$$

and we repeat the arguments from the end of the proof of Theorem 3.13 to find $j < \mu_\delta$ and $r'' \geq r'$ such that $(\forall i < \zeta_\delta)(q_{\mu_\delta \cdot i + j}^\delta \in \underline{G}_{\mathbb{P}_{\zeta^*}})$. \square

4 Reasonable Ultrafilters with Small Generating Systems

Our aim here is to show that, consistently, there may exist a very reasonable ultrafilter on an inaccessible cardinal λ with generating system of size less than 2^λ .

Lemma 4.1 Assume that $G^* \subseteq \mathbb{Q}_\lambda^0$ is directed (with respect to \leq^0) and $\text{fil}(G^*)$ is an ultrafilter on λ , $r \in G^*$. Let \mathbb{P} be a forcing notion not adding bounded subsets of λ , $p \in \mathbb{P}$ and let \underline{A} be a \mathbb{P} -name for a subset of λ such that $p \Vdash_{\mathbb{P}} \underline{A} \in (\text{fil}(G^*))^+$. Then

$$Y \stackrel{\text{def}}{=} \bigcup \{ Z_\delta^r : \delta \in C^r \text{ and } p \Vdash_{\mathbb{P}} \text{“}\underline{A} \cap Z_\delta^r \notin d_\delta^r\text{”} \} \in \text{fil}(G^*).$$

Proof Assume toward contradiction that $Y \notin \text{fil}(G^*)$. Then we may find $s \in G^*$ such that $r \leq^0 s$ and $\lambda \setminus Y \in \text{fil}(s)$. Take $\varepsilon < \lambda$ such that

$$\text{if } \alpha \in C^s \setminus \varepsilon, \text{ then } Z_\alpha^s \setminus Y \in d_\alpha^s \text{ and } (\forall A \in d_\alpha^s)(\exists \beta \in C^r)(A \cap Z_\beta^r \in d_\beta^r).$$

(Remember Proposition 2.5.) Now take a generic filter $G \subseteq \mathbb{P}$ over \mathbf{V} such that $p \in G$ and work in $\mathbf{V}[G]$. Since $\underline{A}^G \in \text{fil}(s)^+$, we may pick $\alpha \in C^s$ such that $\varepsilon < \alpha$ and $\underline{A}^G \cap Z_\alpha^s \in d_\alpha^s$. Then also $Z_\alpha^s \cap \underline{A}^G \setminus Y \in d_\alpha^s$ and thus we may find $\beta \in C^r$ such that $Z_\alpha^s \cap \underline{A}^G \cap Z_\beta^r \setminus Y \in d_\beta^r$. In particular, $Z_\beta^r \setminus Y \neq \emptyset$, so $p \Vdash_{\mathbb{P}} \underline{A} \cap Z_\beta^r \notin d_\beta^r$, and thus $\underline{A}^G \cap Z_\beta^r \notin d_\beta^r$. Consequently, $Z_\alpha^s \cap \underline{A}^G \cap Z_\beta^r \setminus Y \notin d_\beta^r$ giving a contradiction. \square

Theorem 4.2 *Assume that*

- (i) λ is strongly inaccessible, $\bar{\mu} = \langle \mu_\alpha : \alpha < \lambda \rangle$, each μ_α is a regular cardinal, $\aleph_0 \leq \mu_\alpha \leq \lambda$, and $(\forall f \in {}^\alpha \mu_\alpha) (|\prod_{\xi < \alpha} f(\xi)| < \mu_\alpha)$ for $\alpha < \lambda$;
- (ii) $\bar{\mathbb{Q}} = \langle \mathbb{P}_\xi, \mathbb{Q}_\xi : \xi < \gamma \rangle$ is a λ -support iteration such that for every $\xi < \gamma$,
 $\Vdash_{\mathbb{P}_\xi} \text{“}\mathbb{Q}_\xi \text{ is reasonably } A\text{-bounding over } \bar{\mu}\text{”}$;
- (iii) $G^* \subseteq \mathbb{Q}_\lambda^0$ is a \leq^0 -downward closed $\bar{\mu}$ -super reasonable family such that $\text{fil}(G^*)$ is an ultrafilter on λ .

Then

$$\Vdash_{\mathbb{P}_\gamma} \text{“}\text{fil}(G^*) \text{ is an ultrafilter on } \lambda\text{”}.$$

Proof The proof is by induction on the length γ of the iteration $\bar{\mathbb{Q}}$. So we assume that (i)–(iii) hold and for each $\xi < \gamma$

$$(\odot)_\xi \Vdash_{\mathbb{P}_\xi} \text{“}\text{fil}(G^*) \text{ is an ultrafilter on } \lambda\text{”}.$$

Note that (by the strategic $(< \lambda)$ -completeness of \mathbb{P}_γ) forcing with \mathbb{P}_γ does not add bounded subsets of λ , and therefore $(\mathbb{Q}_\lambda^0)^{\mathbf{V}} \subseteq (\mathbb{Q}_\lambda^0)^{\mathbf{V}^{\mathbb{P}_\gamma}}$.

Claim 4.3 *Assume that*

- (a) \underline{A} is a \mathbb{P}_γ -name for a subset of λ such that $\Vdash_{\mathbb{P}_\gamma} \underline{A} \in (\text{fil}(G^*))^+$,
- (b) $w \in [\gamma]^{<\omega}$ and \mathcal{T} is a finite standard $(w, 1)^\gamma$ -tree, and
- (c) $\bar{p} = \langle p_t : t \in T \rangle$ is a (finite) tree of conditions in $\bar{\mathbb{Q}}$, and
- (d) $r \in G^*$ and X is the set of all $\alpha \in C^r$ for which there is a tree of conditions $\bar{q} = \langle q_t : t \in T \rangle$ such that $\bar{q} \geq \bar{p}$ and

$$(\forall t \in T)(\text{rk}(t) = \gamma \Rightarrow q_t \Vdash \underline{A} \cap Z_\alpha^r \in d_\alpha^r).$$

Then $\bigcup \{Z_\alpha^r : \alpha \in X\} \in \text{fil}(G^*)$.

Proof of the Claim Induction on $|w|$. If $w = \emptyset$ and so $T = \{\emptyset\}$, then the assertion follows directly from Lemma 4.1 (with p, \mathbb{P} there standing for $p_\emptyset, \mathbb{P}_\gamma$ here). Assume that $|w| = n + 1$, $\zeta^* = \max(w)$, $w' = w \setminus \{\zeta^*\}$ and the claim is true for w' (in place of w) and any \underline{A}, \bar{p} . Let $\mathbb{P}_{\zeta^* \gamma}$ be a \mathbb{P}_{ζ^*} -name for a forcing notion with the universe $P_{\zeta^* \gamma} = \{p \restriction [\zeta^*, \gamma) : p \in \mathbb{P}_\gamma\}$ and the order relation $\leq_{\mathbb{P}_{\zeta^* \gamma}}$ such that

if $G \subseteq \mathbb{P}_{\zeta^*}$ is generic over \mathbf{V} and $f, g \in P_{\zeta^* \gamma}$,

$$\text{then } \mathbf{V}[G] \models f \leq_{\mathbb{P}_{\zeta^* \gamma}[G]} g \text{ if and only if } (\exists p \in G)(p \cup f \leq_{\mathbb{P}_\gamma} p \cup g).$$

Note that $P_{\zeta^* \gamma}$ is from \mathbf{V} but the relation $\leq_{\mathbb{P}_{\zeta^* \gamma}[G]}$ is defined in $\mathbf{V}[G]$ only. Also \mathbb{P}_γ is isomorphic with a dense subset of the composition $\mathbb{P}_{\zeta^*} * \mathbb{P}_{\zeta^* \gamma}$.

We are going to define a \mathbb{P}_{ζ^*} -name \underline{Y} for a subset of λ . Suppose that $G \subseteq \mathbb{P}_{\zeta^*}$ is generic over \mathbf{V} and work in $\mathbf{V}[G]$. For $t \in T$ such that $\text{rk}(t) = \gamma$ let X_t consist of all $\alpha \in C^r$ for which there is $f \in P_{\zeta^* \gamma}$ such that

$$p_t \restriction [\zeta^*, \gamma) \leq_{\mathbb{P}_{\zeta^* \gamma}[G]} f \quad \text{and} \quad f \Vdash_{\mathbb{P}_{\zeta^* \gamma}[G]} \underline{A} \cap Z_\alpha^r \in d_\alpha^r.$$

Let $Y_t = \bigcup \{Z_\alpha^r : \alpha \in X_t\}$ (for $t \in T$ such that $\text{rk}(t) = \gamma$). It follows from Lemma 4.1 that each Y_t belongs to $\text{fil}(G^*)$ (remember that $\Vdash_{\mathbb{P}_{\zeta^*}} \text{“}\text{fil}(G^*) \text{ is an ultrafilter”}$ by $(\odot)_{\zeta^*}$). Hence

$$Y^* \stackrel{\text{def}}{=} \bigcap \{Y_t : t \in T \ \& \ \text{rk}(t) = \gamma\} \in \text{fil}(G^*).$$

Note that for each $\alpha \in C^r$, either $Z'_\alpha \cap Y^* = \emptyset$ or $Z'_\alpha \subseteq Y^*$.

Going back to \mathbf{V} , let $\underline{Y}^*, \underline{Y}_t, \underline{X}_t$ be \mathbb{P}_{ζ^*} -names for the objects described as Y^*, Y_t, X_t above. Thus $\Vdash_{\mathbb{P}_{\zeta^*}} \underline{Y}^* \in \text{fil}(G^*)$ and we may apply the inductive hypothesis to w' , $T' = \{t \upharpoonright \zeta^* : t \in T\}$, and $\bar{p}' = \langle p_{t'} : t' \in T' \rangle \subseteq \mathbb{P}_{\zeta^*}$. Thus, if X^* is the set of all $\alpha \in C^r$ for which there is a tree of conditions $\bar{q}' = \langle q'_{t'} : t' \in T' \rangle \subseteq \mathbb{P}_{\zeta^*}$ such that $\bar{q}' \geq \bar{p}'$ and

$$(\forall t' \in T)(\text{rk}(t') = \zeta^* \Rightarrow q'_{t'} \Vdash_{\mathbb{P}_{\zeta^*}} \underline{Y}^* \cap Z'_\alpha \in d'_\alpha),$$

then $\bigcup \{Z'_\alpha : \alpha \in X^*\} \in \text{fil}(G^*)$.

Now suppose that $\alpha \in X^*$ is witnessed by \bar{q}' and let $t' \in T$ be such that $\text{rk}(t') = \zeta^*$. Then $q'_{t'} \Vdash_{\mathbb{P}_{\zeta^*}} Z'_\alpha \subseteq \underline{Y}^*$ and hence $q'_{t'} \Vdash_{\mathbb{P}_{\zeta^*}} \alpha \in \underline{X}_t$ for all $t \in T$ with $\text{rk}(t) = \gamma$, so we have \mathbb{P}_{ζ^*} -names \underline{f}'_t for elements of $\mathbb{P}_{\zeta^* \gamma}$ such that

$$q'_{t'} \Vdash_{\mathbb{P}_{\zeta^*}} \text{“} p_t \upharpoonright \zeta^*, \gamma \leq_{\mathbb{P}_{\zeta^* \gamma}} \underline{f}'_t \ \& \ \underline{f}'_t \Vdash_{\mathbb{P}_{\zeta^* \gamma}} \underline{A} \cap Z'_\alpha \in d'_\alpha \text{”}.$$

Now use Proposition 3.5 (or just finite induction) to get a tree of conditions

$$\bar{q}'' = \langle q''_{t'} : t' \in T' \rangle \subseteq \mathbb{P}_{\zeta^*}$$

and objects $g^t_{t'}$ (for $t' \in T', t \in T, \text{rk}(t') = \zeta^*, \text{rk}(t) = \gamma$) such that $\bar{q}' \leq \bar{q}''$ and $q''_{t'} \Vdash_{\mathbb{P}_{\zeta^*}} \underline{f}'_t = g^t_{t'}$. Now, for $t \in T$, put

1. $q_t = q''_{t \upharpoonright \zeta^*}$ if $\text{rk}(t) \leq \zeta^*$, and
2. $q_t = q''_{t \upharpoonright \zeta^*} \frown g^t_{t \upharpoonright \zeta^*}$ if $\text{rk}(t) = \gamma$.

It should be clear that $\bar{q} = \langle q_t : t \in T \rangle$ is a tree of conditions in $\bar{\mathbb{Q}}$, $\bar{p} \leq \bar{q}$ and for every $t \in T$ with $\text{rk}(t) = \gamma$ we have $q_t \Vdash_{\mathbb{P}_\gamma} \underline{A} \cap Z'_\alpha \in d'_\alpha$. This shows that X^* is included in the set X defined in the assumption (d), and hence $\bigcup \{Z'_\alpha : \alpha \in X\} \in \text{fil}(G^*)$. \square

Let \underline{A} be a \mathbb{P}_γ -name for a subset of λ such that $\Vdash_{\mathbb{P}_\gamma} \underline{A} \in (\text{fil}(G^*))^+$ and let $p \in \mathbb{P}_\gamma$. We will find a condition $p^* \geq p$ such that $p^* \Vdash_{\mathbb{P}_\gamma} \underline{A} \in \text{fil}(G^*)$. It will be provided by the winning criterion $(\otimes)_{\bar{\mu}}^{\text{treeA}}$ of the game $\mathfrak{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ (see Definition 3.7; remember \mathbb{P}_γ is reasonably* $A(\bar{\mathbb{Q}})$ -bounding over $\bar{\mu}$ by Theorem 3.8).

Let \mathbf{st} be a winning strategy of GENERIC in $\mathfrak{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$, and for $\varepsilon \leq \gamma$ and $q \in \mathbb{P}_\varepsilon$ let us fix a winning strategy $\mathbf{st}(\varepsilon, q)$ of COMPLETE in $\mathfrak{D}_0^\lambda(\mathbb{P}_\varepsilon, q)$ so that the coherence demands (i)–(iii) of Proposition 3.3 are satisfied.

We are going to describe a strategy \mathbf{st}^{\boxplus} of INC in the game $\mathfrak{D}_{\bar{\mu}}^{\boxplus}(G^*)$. In the course of a play of $\mathfrak{D}_{\bar{\mu}}^{\boxplus}(G^*)$, INC will construct on the side a play of $\mathfrak{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ in which GENERIC plays according to \mathbf{st} . So suppose that INC and COM arrived to a stage $\alpha < \lambda$ of a play of $\mathfrak{D}_{\bar{\mu}}^{\boxplus}(G^*)$, and they have constructed

$$(\otimes)_1^\alpha \ \langle I_\gamma, i_\gamma, \bar{u}_\gamma, \langle r_{\gamma,i}, r'_{\gamma,i}, \langle \beta_{\gamma,i}, Z_{\gamma,i}, d_{\gamma,i} \rangle : i < i_\gamma \rangle : \gamma < \alpha \rangle.$$

Also, let us assume that INC (playing according to \mathbf{st}^{\boxplus}) has written on the side a partial play

$$(\otimes)_2^\alpha \ \langle \mathcal{T}_\gamma, \bar{p}^\gamma, \bar{q}^\gamma : \gamma < \alpha \rangle$$

of $\mathfrak{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ (in which GENERIC plays according to \mathbf{st}). Let a standard tree \mathcal{T}_α and a tree of conditions $\bar{p}^\alpha = \langle p_t^\alpha : t \in T_\alpha \rangle$ be given to GENERIC by the strategy \mathbf{st} in answer to $(\otimes)_2^\alpha$ (so $|T_\alpha| < \mu_\alpha$).

On the board of $\mathfrak{D}_{\mu}^{\boxplus}(G^*)$, the strategy \mathbf{st}^{\boxplus} instructs INC to play the set

$$I_{\alpha} \stackrel{\text{def}}{=} \{t \in T_{\alpha} : \text{rk}_{\alpha}(t) = \gamma\}$$

and the \langle^*_{χ} -first enumeration $\bar{u}_{\alpha} = \langle u_{\alpha,i} : i < i_{\alpha} \rangle$ of $[I_{\alpha}]^{<\omega}$ (so $i_{\alpha} < \mu_{\alpha}$). Now the two players start playing a subgame of length i_{α} to determine a sequence $\langle r_{\alpha,i}, r'_{\alpha,i}, (\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : i < i_{\alpha} \rangle$. During the subgame INC will construct on the side a sequence $\langle \bar{q}_i^0, \bar{q}_i^1 : i < i_{\alpha} \rangle$ of trees of conditions in \mathbb{P}_{γ} so that

- (*)₃ $\bar{q}_i^{\ell} = \langle q_{t,i}^{\ell} : t \in T_{\alpha} \rangle$ (for $\ell < 2, i < i_{\alpha}$) and for each $t \in T_{\alpha}$, the sequence $\langle q_{t,i}^0, q_{t,i}^1 : i < i_{\alpha} \rangle$ is a legal play of $\mathfrak{D}_0^{\lambda}(\mathbb{P}_{\text{rk}_{\alpha}(t)}, p_t^{\alpha})$ in which COMPLETE uses her winning strategy $\mathbf{st}(\text{rk}_{\alpha}(t), p_t^{\alpha})$.

Suppose that COM and INC arrive at level $i < i_{\alpha}$ of the subgame (of $\mathfrak{D}_{\mu}^{\boxplus}(G^*)$) and

- (*)₄ $\langle r_{\alpha,j}, r'_{\alpha,j}, (\beta_{\alpha,j}, Z_{\alpha,j}, d_{\alpha,j}) : j < i \rangle$ and $\langle \bar{q}_j^0, \bar{q}_j^1 : j < i \rangle$

have been determined and COM has chosen $r_{\alpha,i} \in G^*$. INC's answer is given by \mathbf{st}^{\boxplus} as follows. First, INC takes the \langle^*_{χ} -first tree of conditions \bar{q}^{\diamond} in \mathbb{Q} such that

- (*)₅^a $\bar{q}^{\diamond} = \langle q_t^{\diamond} : t \in T_{\alpha} \rangle$ and $q_t^{\diamond} \in \mathbb{P}_{\text{rk}(t)}$ is an upper bound to the set $\{p_t^{\alpha}\} \cup \{q_{t,j}^1 : j < i\}$ (for each $t \in T_{\alpha}$)

(remember (*₃)). Then INC lets $X \subseteq C^{r_{\alpha,i}}$ be the set of all $\beta \in C^{r_{\alpha,i}}$ greater than $\sup \left(\bigcup_{\gamma < \alpha} \bigcup_{j < i_{\gamma}} Z_{\gamma,j} \cup \bigcup_{j < i} Z_{\alpha,j} \right) + 890$ and such that

- (*)₅^b there is a tree of conditions \bar{q}' in $\bar{\mathbb{Q}}$ such that $q^{\diamond} \leq \bar{q}'$ and

$$\text{if } t \in u_{\alpha,i}, \text{ then } q'_t \Vdash_{\mathbb{P}_{\gamma}} \bar{A} \cap Z_{\beta}^{r_{\alpha,i}} \in d_{\beta}^{r_{\alpha,i}}.$$

Since $u_{\alpha,i}$ is finite, it follows from Claim 4.3 that $\bigcup \{Z_{\beta}^{r_{\alpha,i}} : \beta \in X\} \in \text{fil}(G^*)$. Then INC picks also the club C of λ such that $C \subseteq C^{r_{\alpha,i}}$ and $r_{\alpha,i}$ is restrictable to $\langle X, C \rangle$ (see Definition 2.7) and $\min(C) = \min(X)$, and his inning at the stage i of the subgame of $\mathfrak{D}_{\mu}^{\boxplus}(G^*)$ is $r'_{\alpha,i} = r_{\alpha,i} \upharpoonright \langle X, C \rangle$ (again, see Definition 2.7; note that $r'_{\alpha,i} \in G^*$ by Proposition 2.8).

After this COM answers with $(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) \in \#(r'_{\alpha,i})$, and then INC chooses (for the construction on the side) the \langle^*_{χ} -first tree of conditions \bar{q}_i^0 in $\bar{\mathbb{Q}}$ such that $\bar{q}^{\diamond} \leq \bar{q}_i^0$ and

- (*)₆ if $t \in u_{\alpha,i}$, then $q_{t,i}^0 \Vdash_{\mathbb{P}_{\gamma}} \bar{A} \cap Z_{\alpha,i} \in d_{\alpha,i}$.

Then $\bar{q}_i^1 = \langle q_{t,i}^1 : t \in T_{\alpha} \rangle$ is a tree of conditions determined by the demand in (*₃) and the strategies $\mathbf{st}(\text{rk}_{\alpha}(t), p_t^{\alpha})$ (for $t \in T_{\alpha}$); remember the coherence conditions of Proposition 3.3.

This completes the description of how INC plays in the subgame of stage α . After the subgame is finished, INC determines the move \bar{q}^{α} of ANTIGENERIC in the play of $\mathfrak{D}_{\mu}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ which he is constructing on the side:

- (*)₇ \bar{q}^{α} is the \langle^*_{χ} -first tree of conditions $\langle q_t^{\alpha} : t \in T_{\alpha} \rangle$ such that $\bar{q}_i^0 \leq \bar{q}_i^1 \leq \bar{q}^{\alpha}$ for all $i < i_{\alpha}$.

(There is such a tree of conditions by $(*)_3$; remember $i_\alpha < \mu_\alpha \leq \lambda$.)

This completes the description of the strategy \mathbf{st}^\boxplus . Since G^* is $\bar{\mu}$ -super reasonable, \mathbf{st}^\boxplus cannot be a winning strategy, so there is a play

$$(*)_8 \quad \langle I_\alpha, i_\alpha, \bar{u}_\alpha, \langle r_{\alpha,i}, r'_{\alpha,i}, (\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : i < i_\alpha \rangle : \alpha < \lambda \rangle$$

of $\mathcal{D}_{\bar{\mu}}^\boxplus(G^*)$ in which INC follows \mathbf{st}^\boxplus , but

$$(*)_9 \quad \text{for some } r \in G^*, \text{ for every } \langle j_\alpha : \alpha < \lambda \rangle \in \prod_{\alpha < \lambda} I_\alpha \text{ we have}$$

$$\{(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : \alpha < \lambda \ \& \ i < i_\alpha \ \& \ j_\alpha \in u_{\alpha,i}\} \leq^* \#(r).$$

Let $\langle \mathcal{T}_\alpha, \bar{p}^\alpha, \bar{q}^\alpha : \alpha < \lambda \rangle$ be the play of $\mathcal{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ constructed on the side by INC (so this is a play in which GENERIC uses her winning strategy \mathbf{st}). Since GENERIC won that play, there is a condition $p^* \in \mathbb{P}_\gamma$ stronger than p and such that for each $\alpha < \lambda$ the set $\{q_t^\alpha : t \in T_\alpha \ \& \ \text{rk}_\alpha(t) = \gamma\}$ is pre-dense above p^* . Note that if we show that

$$(*)_{10} \quad \text{it is forced in } \mathbb{P}_\gamma \text{ that for every } \bar{j} = \langle j_\alpha : \alpha < \lambda \rangle \in \prod_{\alpha < \lambda} I_\alpha \text{ we have}$$

$$\{(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : \alpha < \lambda \ \& \ i < i_\alpha \ \& \ j_\alpha \in u_{\alpha,i}\} \leq^* \#(r),$$

then we will be able to conclude that $p^* \Vdash \dot{A} \in \text{fil}(r)$ (remember $(*)_6$ + $(*)_7$ and Observation 2.10), finishing the proof of the theorem. So let us argue that $(*)_{10}$ holds true.

It follows from the description of \mathbf{st}^\boxplus (see the description of X after $(*)_5^a$) that we may choose a continuous increasing sequence $\langle \delta_\alpha : \alpha < \lambda \rangle \subseteq \lambda$ such that

$$(\forall \alpha < \lambda) (\delta_\alpha \leq \beta_{\alpha,0} \leq \sup \{ \bigcup_{i < i_\alpha} Z_{\alpha,i} \} < \delta_{\alpha+1}).$$

Now, we will say that $\beta \in C^r$ is a sick case whenever there are $\alpha_0 < \alpha_1 < \lambda$ and $B \in d_\beta^r$ such that $Z_\beta^r \subseteq [\delta_{\alpha_0}, \delta_{\alpha_1})$ and

$$(\forall \alpha \in [\alpha_0, \alpha_1)) (\exists t \in I_\alpha) (\forall i < i_\alpha) (t \notin u_{\alpha,i} \text{ or } B \cap Z_{\alpha,i} \notin d_{\alpha,i}).$$

Using Observation 2.10(2) one can easily verify that the following two conditions are equivalent:

$$(*)_{11}^{\text{one}} \quad \text{there is } \langle j_\alpha : \alpha < \lambda \rangle \in \prod_{\alpha < \lambda} I_\alpha \text{ such that}$$

$$\{(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : \alpha < \lambda \ \& \ i < i_\alpha \ \& \ j_\alpha \in u_{\alpha,i}\} \not\leq^* \#(r);$$

$$(*)_{11}^{\text{two}} \quad \text{there are } \lambda \text{ many sick cases of } \beta \in C^r.$$

Since the forcing with \mathbb{P}_γ does not add bounded subsets of λ , being a sick case is absolute between \mathbf{V} and $\mathbf{V}^{\mathbb{P}_\gamma}$. So we may conclude (from $(*)_9$) that $(*)_{10}$ is true and thus the proof of Theorem 4.2 is complete. \square

Theorem 4.4 Assume (i) and (ii) of Theorem 4.2 and

(α) $\bar{\kappa} = \langle \kappa_\alpha : \alpha < \lambda \rangle$ is a sequence of regular cardinals such that for each $\alpha < \lambda$,

$$\mu_\alpha \leq \kappa_\alpha \leq \lambda \quad \text{and} \quad (\forall \mu < \mu_\alpha) (2^\mu < \kappa_\alpha),$$

(β) $G^* \subseteq \mathbb{Q}_\lambda^0$ is $\bar{\kappa}$ -super reasonable.

Then $\Vdash_{\mathbb{P}_\gamma}$ “ G^* is $\bar{\mu}$ -strongly reasonable”.

Proof First of all note that the forcing notion \mathbb{P}_γ is reasonably^{*} $A(\bar{\mathbb{Q}})$ -bounding over $\bar{\mu}$ and λ -proper (see Theorem 3.8). Therefore $\Vdash_{\mathbb{P}_\gamma} “([G^*]^{\leq \lambda})^\mathbf{V}$ is cofinal in $[G^*]^{\leq \lambda}$ ”, and consequently $\Vdash_{\mathbb{P}_\gamma} “G^*$ is $(< \lambda^+)$ -directed (with respect to \leq^0)”.

Suppose that \mathbf{st}^\oplus is a \mathbb{P}_γ -name, $p \in \mathbb{P}_\gamma$, and

$\Vdash_{\mathbb{P}_\gamma}$ “ \mathbf{st}^\oplus is a strategy of INC in $\mathfrak{D}_{\bar{\mu}}^\oplus(G^*)$ such that
all values given by it are from \mathbf{V} ”.

We are going to find a condition $p^* \geq p$ and a \mathbb{P}_γ -name g_λ such that

$p^* \Vdash_{\mathbb{P}_\gamma}$ “ g_λ is a play of $\mathfrak{D}_{\bar{\mu}}^\oplus(G^*)$ in which INC uses \mathbf{st}^\oplus but COM wins the play”.

The condition p^* will be provided by the winning criterion $(\otimes)_{\mathbf{A}}^{\text{tree}}$ of the game $\mathfrak{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ (see Definition 3.7).

In the rest of the proof whenever we say “INC chooses/picks x such that” we mean “INC chooses/picks the $<_\chi^*$ -first x such that”. Let us fix

- (i) a winning strategy \mathbf{st} of GENERIC in $\mathfrak{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$,
- (ii) winning strategies $\mathbf{st}(\varepsilon, q)$ of COMPLETE in $\mathfrak{D}_0^\lambda(\mathbb{P}_\varepsilon, q)$ (for $\varepsilon \leq \gamma$, $q \in \mathbb{P}_\varepsilon$) such that the coherence conditions of Proposition 3.3 are satisfied.

We are going to describe a strategy \mathbf{st}^\boxplus of INC in the game $\mathfrak{D}_{\bar{\kappa}}^\boxplus(G^*)$. In the course of a play of $\mathfrak{D}_{\bar{\kappa}}^\boxplus(G^*)$, INC will simulate a play of $\mathfrak{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ and he will consider names for partial plays of $\mathfrak{D}_{\bar{\mu}}^\oplus(G^*)$ in which INC uses \mathbf{st}^\oplus . Thus players INC/COM will appear in the play of $\mathfrak{D}_{\bar{\kappa}}^\boxplus(G^*)$ in \mathbf{V} and in the play of $\mathfrak{D}_{\bar{\mu}}^\oplus(G^*)$ in $\mathbf{V}^{\mathbb{P}_\gamma}$. To avoid confusion we will refer to them as $\text{COM}^\mathbf{V}$, $\text{INC}^\mathbf{V}$ for $\mathfrak{D}_{\bar{\kappa}}^\boxplus(G^*)$ (in \mathbf{V}) and $\text{COM}^{\mathbf{V}^{\mathbb{P}_\gamma}}$, $\text{INC}^{\mathbf{V}^{\mathbb{P}_\gamma}}$ for $\mathfrak{D}_{\bar{\mu}}^\oplus(G^*)$ (in $\mathbf{V}^{\mathbb{P}_\gamma}$).

So suppose that $\text{INC}^\mathbf{V}$ and $\text{COM}^\mathbf{V}$ arrived at a stage $\alpha < \lambda$ of the play of $\mathfrak{D}_{\bar{\kappa}}^\boxplus(G^*)$ (in \mathbf{V}), and $\text{INC}^\mathbf{V}$ (playing according to \mathbf{st}^\boxplus) has written on the side,

- $(\oplus)_1^\alpha$ a partial play $\langle \mathcal{T}_\beta, \bar{p}^\beta, \bar{q}^\beta : \beta < \alpha \rangle$ of $\mathfrak{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ in which GENERIC plays according to \mathbf{st} , and
- $(\oplus)_2^\alpha$ a \mathbb{P}_γ -name $\underline{g}_\alpha = \langle \underline{I}_\beta, \underline{i}_\beta, \underline{\bar{u}}_\beta, \underline{\bar{x}}_\beta : \beta < \alpha \rangle$ of a partial play of $\mathfrak{D}_{\bar{\mu}}^\oplus(G^*)$ (in $\mathbf{V}^{\mathbb{P}_\gamma}$) in which $\text{INC}^{\mathbf{V}^{\mathbb{P}_\gamma}}$ uses the strategy \mathbf{st}^\oplus ,
- $(\oplus)_3^\alpha$ ordinals $i_\beta < \mu_\beta$ such that $q_t^\beta \Vdash \underline{i}_\beta = i_\beta$ for every $t \in T_\beta$ with $\text{rk}_\beta(t) = \gamma$ (for $\beta < \alpha$).

Note that \underline{I}_β is a \mathbb{P}_γ -name for a set of size $< \mu_\beta$ from \mathbf{V} , $\underline{\bar{u}}_\beta$ is a \mathbb{P}_γ -name for an \underline{i}_β -sequence of finite subsets of \underline{I}_β and $\underline{\bar{x}}_\beta$ is a \mathbb{P}_γ -name for the result of the subgame of length \underline{i}_β of level β .

Let \underline{I}_α be a \mathbb{P}_γ -name for the answer by \mathbf{st}^\oplus to the play \underline{g}_α of $\mathfrak{D}_{\bar{\mu}}^\oplus(G^*)$ (in $\mathbf{V}^{\mathbb{P}_\gamma}$). Let \mathcal{T}_α and $\bar{p}^\alpha = \langle p_t^\alpha : t \in T_\alpha \rangle$ be given to GENERIC by the strategy \mathbf{st} as an answer to $(\oplus)_1^\alpha$. Let $\bar{q}^\alpha = \langle q_t^\alpha : t \in T_\alpha \rangle$ be a tree of conditions in $\bar{\mathbb{Q}}$ such that

- $(\oplus)_4^a$ $\bar{p}^\alpha \leq \bar{q}^\alpha$ and $q_{t_0}^\alpha, q_{t_1}^\alpha$ are incompatible for distinct $t_0, t_1 \in T_\alpha$ with $\text{rk}_\alpha(t_0) = \text{rk}_\alpha(t_1)$,
- $(\oplus)_4^b$ for every $t \in T_\alpha$ with $\text{rk}_\alpha(t) = \gamma$ the condition q_t^α decides the value of \underline{I}_α , say, $q_t^\alpha \Vdash_{\mathbb{P}_\gamma} “\underline{I}_\alpha = I_\alpha^t”$.

(Note that $\Vdash_{\mathbb{P}_\gamma} \underline{I}_\alpha \in \mathbf{V}$ by the choice of \mathbf{st}^\oplus ; remember Proposition 3.5.)

In the play of $\mathcal{D}_{\bar{\kappa}}^{\boxplus}(G^*)$, the strategy \mathbf{st}^{\boxplus} instructs $\text{INC}^{\mathbf{V}}$ to choose the set

$$I_\alpha = \prod \{I_\alpha^t : t \in T_\alpha \ \& \ \text{rk}_\alpha(t) = \gamma\}$$

and an enumeration $\bar{u}_\alpha = \langle u_{\alpha,i} : i < i_\alpha \rangle$ of $[I_\alpha]^{<\omega}$. Note that $|I_\alpha^t| < \mu_\alpha$ for all relevant $t \in T_\alpha$ and $|T_\alpha| < \mu_\alpha$, so by our assumptions on μ_α and κ_α we know that $|I_\alpha| < \kappa_\alpha$ (so also $i_\alpha < \kappa_\alpha$).

Then, in the play of $\mathcal{D}_{\bar{\mu}}^\oplus(G^*)$, $\text{INC}^{\mathbf{V}^{\mathbb{P}_\gamma}}$ pretends that $\text{COM}^{\mathbf{V}^{\mathbb{P}_\gamma}}$ played an ordinal $\underline{i}_\alpha \in [i_\alpha, \lambda)$ and $\bar{u}_\alpha = \langle u_{\alpha,i} : i < \underline{i}_\alpha \rangle$ such that

$$\Vdash_{\mathbb{P}_\gamma} \quad \text{“}\bar{u}_\alpha \subseteq [I_\alpha]^{<\omega} \text{ and } \bigcup \{u_{\alpha,i} : i < \underline{i}_\alpha\} = \underline{I}_\alpha\text{”}$$

and for each $t \in T_\alpha$ with $\text{rk}_\alpha(t) = \gamma$ we have

$$q_t^\diamond \Vdash_{\mathbb{P}_\gamma} \quad \text{“}\underline{i}_\alpha = i_\alpha \text{ and } u_{\alpha,i} = \{c(t) : c \in u_{\alpha,i}\} \text{ for } i < i_\alpha\text{”}.$$

Now, both in $\mathcal{D}_{\bar{\mu}}^\oplus(G^*)$ of $\mathbf{V}^{\mathbb{P}_\gamma}$ and in $\mathcal{D}_{\bar{\kappa}}^{\boxplus}(G^*)$ of \mathbf{V} the two players start a subgame. The length of the subgame in $\mathbf{V}^{\mathbb{P}_\gamma}$ may be longer than i_α , but we will restrict our attention to the first i_α steps of that subgame. In our *active case* we will have $\underline{i}_\alpha = i_\alpha$; see the choice of \underline{i}_α above. When playing the subgame, $\text{INC}^{\mathbf{V}}$ will build a sequence $\langle \bar{q}_i^0, \bar{q}_i^1 : i < i_\alpha \rangle$ of trees of conditions in $\bar{\mathbb{Q}}$ such that (in addition to demands stated later),

- (\oplus)₅^a $\bar{q}_j^\ell = \langle q_{t,j}^\ell : t \in T_\alpha \rangle$, $\bar{q}^\diamond \leq \bar{q}_j^1 \leq \bar{q}_j^0 \leq \bar{q}_i^0$ for $\ell < 2$, $j < i < i_\alpha$, and
- (\oplus)₅^b for each $t \in T_\alpha$, the sequence $\langle q_{t,i}^0, q_{t,i}^1 : i < i_\alpha \rangle$ is a legal play of the game $\mathcal{D}_0^\lambda(\mathbb{P}_{\text{rk}_\alpha(t)}, q_t^\diamond)$ in which COMPLETE uses her winning strategy $\mathbf{st}(\text{rk}_\alpha(t), q_t^\diamond)$.

He (as $\text{INC}^{\mathbf{V}^{\mathbb{P}_\gamma}}$) will also construct a name for a play of a subgame of $\mathcal{D}_{\bar{\mu}}^\oplus(G^*)$ of $\mathbf{V}^{\mathbb{P}_\gamma}$ for this stage.

Suppose that $\text{INC}^{\mathbf{V}}$ and $\text{COM}^{\mathbf{V}}$ have arrived to a stage $i < i_\alpha$ of the subgame and $\text{INC}^{\mathbf{V}}$ has determined on the side \bar{q}_j^ℓ for $j < i$, $\ell < 2$, and a \mathbb{P}_γ -name $\langle z_j^\alpha : j < i \rangle$ for a partial play of the subgame of $\mathcal{D}_{\bar{\mu}}^\oplus(G^*)$ of $\mathbf{V}^{\mathbb{P}_\gamma}$. Now $\text{COM}^{\mathbf{V}}$ chooses $r_{\alpha,i} \in G^*$ which $\text{INC}^{\mathbf{V}}$ passes to $\text{INC}^{\mathbf{V}^{\mathbb{P}_\gamma}}$ as an inning of $\text{COM}^{\mathbf{V}^{\mathbb{P}_\gamma}}$ at the i th step of the subgame of level α of $\mathcal{D}_{\bar{\mu}}^\oplus(G^*)$ in $\mathbf{V}^{\mathbb{P}_\gamma}$. There the strategy \mathbf{st}^\oplus gives $\text{INC}^{\mathbf{V}^{\mathbb{P}_\gamma}}$ an answer $\underline{\delta}_{\alpha,i} < \lambda$.

Next, $\text{INC}^{\mathbf{V}}$ picks a tree of conditions $\bar{q}_i^0 = \langle q_{t,i}^0 : t \in T_\alpha \rangle$ in $\bar{\mathbb{Q}}$ such that

- (\oplus)₆^a $(\forall j < i)(\bar{q}_j^1 \leq \bar{q}_i^0)$ and $\bar{q}^\diamond \leq \bar{q}_i^0$, and
- (\oplus)₆^b for every $t \in T_\alpha$ with $\text{rk}_\alpha(t) = \gamma$, the condition $q_{t,i}^0$ decides the value of $\underline{\delta}_{\alpha,i}$, say, $q_{t,i}^0 \Vdash_{\mathbb{P}_\gamma} \underline{\delta}_{\alpha,i} = \delta_{\alpha,i}^t$.

Then $\text{INC}^{\mathbf{V}}$ lets

$$\delta_{\alpha,i}^* = \sup \left(\{ \delta_{\alpha,i}^t : t \in T_\alpha \ \& \ \text{rk}_\alpha(t) = \gamma \} \cup \bigcup_{\beta < \alpha} \bigcup_{j < i_\beta} Z_{\beta,j} \cup \bigcup_{j < i} Z_{\alpha,j} \right) + 890$$

and in the subgame of $\mathcal{D}_{\bar{\kappa}}^{\boxplus}(G^*)$ (in \mathbf{V}) he is instructed to put $r'_{\alpha,i}$ such that

$$C^{r'_{\alpha,i}} = C^{r_{\alpha,i}} \setminus \delta_{\alpha,i}^* \quad \text{and} \quad d_\beta^{r'_{\alpha,i}} = d_\beta^{r_{\alpha,i}} \text{ for } \beta \in C^{r'_{\alpha,i}}.$$

(Note that $r'_{\alpha,i} \in G^*$ by Definition 2.11(2)(ii).)

After this $\text{COM}^{\mathbf{V}}$ chooses $(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) \in \#(r'_{\alpha,i})$, so $\beta_{\alpha,i} \in C^{r_{\alpha,i}}$, $\beta_{\alpha,i} \geq \delta_{\alpha,i}^*$ and $d_{\alpha,i} = d_{\beta_{\alpha,i}}^{r_{\alpha,i}}$. Next $\text{INC}^{\mathbf{V}}$ lets

1. \bar{q}_i^1 be the tree of conditions in $\bar{\mathbb{Q}}$ fully determined by demand $(\oplus)_5^b$ and
2. \bar{z}_i^α be a \mathbb{P}_γ -name for a legal result of stage i of the subgame of level α of $\mathcal{D}_{\bar{\mu}}^\oplus(G^*)$ in $\mathbf{V}^{\mathbb{P}_\gamma}$ such that for each $t \in T_\alpha$ with $\text{rk}_\alpha(t) = \gamma$ we have

$$q_{t,i}^0 \Vdash_{\mathbb{P}_\gamma} \bar{z}_i^\alpha = (r_{\alpha,i}, \bar{d}_{\alpha,i}, (\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i})).$$

Then the subgame continues.

After all i_α steps of the subgame are completed, $\text{INC}^{\mathbf{V}}$ chooses a tree of conditions $\bar{q}^\alpha = \langle q_t^\alpha : t \in T_\alpha \rangle$ in $\bar{\mathbb{Q}}$ such that $(\forall i < i_\alpha)(\bar{q}_i^1 \leq \bar{q}^\alpha)$ and he also lets \bar{x}_α be a \mathbb{P}_γ -name for the result of the subgame of level α of $\mathcal{D}_{\bar{\mu}}^\oplus(G^*)$ in $\mathbf{V}^{\mathbb{P}_\gamma}$ such that $\bar{x}_\alpha \upharpoonright i_\alpha = \langle \bar{z}_i^\alpha : i < i_\alpha \rangle$. Note that all the objects described by $(\oplus)_1^{\alpha+1} - (\oplus)_3^{\alpha+1}$ are determined now.

This completes the description of the strategy st^\boxplus of INC (i.e., $\text{INC}^{\mathbf{V}}$) in $\mathcal{D}_{\bar{\kappa}}^\boxplus(G^*)$. Since G^* is $\bar{\kappa}$ -super reasonable, this strategy cannot be a winning one, so there is a play

$$(\oplus)_7 \quad \langle I_\alpha, i_\alpha, \bar{u}_\alpha, \langle r_{\alpha,i}, r'_{\alpha,i}, (\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : i < i_\alpha \rangle : \alpha < \lambda \rangle$$

of $\mathcal{D}_{\bar{\kappa}}^\boxplus(G^*)$ in which INC follows st^\boxplus , but

$$(\oplus)_8 \quad \text{for some } r \in G^*, \text{ for every } \langle j_\alpha : \alpha < \lambda \rangle \in \prod_{\alpha < \lambda} I_\alpha \text{ we have}$$

$$\{(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : \alpha < \lambda \ \& \ i < i_\alpha \ \& \ j_\alpha \in u_{\alpha,i}\} \leq^* \#(r).$$

Exactly as in the proof of Theorem 4.2 we may argue that then also

$$(\oplus)_9 \quad \text{it is forced in } \mathbb{P}_\gamma \text{ that}$$

$$(\forall \bar{j} \in \prod_{\alpha < \lambda} I_\alpha) (\{(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : \alpha < \lambda \ \& \ i < i_\alpha \ \& \ j_\alpha \in u_{\alpha,i}\} \leq^* \#(r)).$$

(See $(\otimes)_{10}$ in the proof of Theorem 4.2.)

Let $(\mathcal{T}_\alpha, \bar{p}^\alpha, \bar{q}^\alpha : \alpha < \lambda)$ be the play of $\mathcal{D}_{\bar{\mu}}^{\text{treeA}}(p, \bar{\mathbb{Q}})$ constructed on the side by INC . GENERIC won that play, so there is a condition $p^* \in \mathbb{P}_\gamma$ stronger than p and such that for each $\alpha < \lambda$ the set $\{q_t^\alpha : t \in T_\alpha \ \& \ \text{rk}_\alpha(t) = \gamma\}$ is pre-dense above p^* . Also, let g_λ be the \mathbb{P}_γ -name of a play of $\mathcal{D}_{\bar{\mu}}^\oplus(G^*)$ (in $\mathbf{V}^{\mathbb{P}_\gamma}$) constructed on the side in the same run of $\mathcal{D}_{\bar{\kappa}}^\boxplus(G^*)$ (see $(\oplus)_2$). We are going to argue that

$$(\oplus)_{10} \quad \text{the condition } p^* \text{ forces (in } \mathbb{P}_\gamma) \text{ that}$$

$$(\forall \bar{j} \in \prod_{\alpha < \lambda} I_\alpha) (\{(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : \alpha < \lambda \ \& \ i < i_\alpha \ \& \ j_\alpha \in \underline{u}_{\alpha,i}\} \leq^* \#(r)),$$

that is,

$$p^* \Vdash_{\mathbb{P}_\gamma} \text{“COM}^{\mathbf{V}^{\mathbb{P}_\gamma}} \text{ wins the play } g_\lambda \text{ as witnessed by } r\text{”}.$$

Suppose that $G \subseteq \mathbb{P}_\gamma$ is generic over \mathbf{V} , $p^* \in G$ and let us work in $\mathbf{V}[G]$. For every $\alpha < \lambda$ there is a unique $t = t(\alpha) \in T_\alpha$ such that $\text{rk}_\alpha(t) = \gamma$ and $q_t^\alpha \in G$, and thus $(I_\alpha)^G = I_\alpha^t$, $(i_\alpha)^G = i_\alpha$ and $(\bar{u}_\alpha)^G = \langle (u_{\alpha,i})^G : i < i_\alpha \rangle$, where $(u_{\alpha,i})^G = \{c(t) : c \in u_{\alpha,i}\} \subseteq I_\alpha^t$. Suppose that $\bar{j} = \langle j_\alpha : \alpha < \lambda \rangle \in \prod_{\alpha < \lambda} I_\alpha^{t(\alpha)}$. For

each $\alpha < \lambda$, fix $j_\alpha^* \in I_\alpha = \prod\{I_\alpha^t : t \in T_\alpha \text{ \& } \text{rk}_\alpha(t) = \gamma\}$ such that $j_\alpha^*(t(\alpha)) = j_\alpha$. Note that if $j_\alpha^* \in u_{\alpha,i}$, $i < i_\alpha$, then $j_\alpha \in (u_{\alpha,i})^G$ and, therefore,

$$\begin{aligned} \{(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : \alpha < \lambda \text{ \& } i < i_\alpha \text{ \& } j_\alpha \in (u_{\alpha,i})^G\} &\leq^* \\ \{(\beta_{\alpha,i}, Z_{\alpha,i}, d_{\alpha,i}) : \alpha < \lambda \text{ \& } i < i_\alpha \text{ \& } j_\alpha^* \in u_{\alpha,i}\} &\leq^* \#(r) \end{aligned}$$

(remember $(\oplus)_9$). Now $(\oplus)_{10}$ follows and the proof of the theorem is complete. \square

Corollary 4.5 *Assume that λ is a strongly inaccessible cardinal. Then there is a forcing notion \mathbb{P} such that*

$\Vdash_{\mathbb{P}}$ “ λ is strongly inaccessible and $2^\lambda = \lambda^{++}$ and there is a strongly reasonable family $G^* \subseteq \mathbb{Q}_\lambda^0$ such that $\text{fil}(G^*)$ is an ultrafilter on λ and $|G^*| = \lambda^+$; in particular, there is a very reasonable ultrafilter on λ with a generating system of size $< 2^\lambda$ ”.

Proof We may start with a universe \mathbf{V} in which $\diamond_{S_\lambda^{i+}}$ holds (and λ is strongly inaccessible). It follows from Proposition 2.15 that (in \mathbf{V}) there is a \leq^0 -increasing sequence $\langle r_\alpha : \alpha < \lambda^+ \rangle \subseteq \mathbb{Q}_\lambda^0$ such that $G^* \stackrel{\text{def}}{=} \{r \in \mathbb{Q}_\lambda^0 : (\exists \alpha < \lambda^+)(r \leq^0 r_\alpha)\}$ is super reasonable and $\text{fil}(G^*)$ is an ultrafilter on λ .

Let $\bar{\mathbb{Q}} = \langle \mathbb{P}_\alpha, \bar{\mathbb{Q}}_\alpha : \alpha < \lambda^{++} \rangle$ be a λ -support iteration of the forcing notion $\mathbb{Q}_{\mathcal{D}_\lambda}^{\text{tree}}(K_1, \Sigma_1)$ defined in the proof of [8, Proposition B.8.5]. This forcing is reasonably \mathbf{A} -bounding (by [7, Proposition 4.1, p. 221] and [8, Theorem B.6.5]), so we may use Theorems 4.2 and 4.4 to conclude that

$\Vdash_{\mathbb{P}_{\lambda^{++}}}$ “ G^* is strongly reasonable, $|G^*| = \lambda^+ < 2^\lambda$, and $\text{fil}(G^*)$ is ultrafilter on λ ”.

If one analyzes the proof of Theorem 4.4, one may notice that even

$\Vdash_{\mathbb{P}_{\lambda^{++}}}$ “ $\{r_\alpha : \alpha < \lambda^+\}$ is strongly reasonable”.

5 A Feature, Not a Bug

One may wonder if Theorems 4.2 and 4.4 could be improved by replacing the assumption that we are working with the iteration of reasonably \mathbf{A} -bounding forcings by, say, just dealing with a nicely double \mathbf{a} -bounding forcing. A result of that sort would be more natural and the fact that we had to refer to an iteration-specific property could be seen as some lack of knowledge. However, this is a *feature, not a bug* as nicely double \mathbf{a} -bounding forcing notions may cause that $\text{fil}(G^*)$ is not an ultrafilter anymore.

In this section we assume that λ is a strongly inaccessible cardinal.

Definition 5.1

- Let \mathbb{P}^* consist of all pairs $p = (\eta^p, C^p)$ such that $\eta^p : \lambda \longrightarrow \{-1, 1\}$ and C^p is a club of λ . A binary relation $\leq_{\mathbb{P}^*}$ on \mathbb{P}^* is defined by letting $p \leq q$ if and only if
 - $C^q \subseteq C^p$, $\eta^q \upharpoonright \min(C^p) = \eta^p \upharpoonright \min(C^p)$, and
 - for all successive members $\alpha < \beta$ of C^p we have

$$(\forall \gamma \in [\alpha, \beta])(\eta^q(\gamma) = \frac{\eta^p(\alpha)}{\eta^q(\alpha)} \cdot \eta^p(\gamma)).$$

2. For $p \in \mathbb{P}^*$ and $\alpha \in C^p$ let

$$\text{pos}(p, \alpha) \stackrel{\text{def}}{=} \{\eta^q \upharpoonright \alpha : q \in \mathbb{P}^* \text{ \& } p \leq q\}.$$

3. For $p \in \mathbb{P}^*$, $\alpha < \lambda$, and $\nu : \alpha \rightarrow \{-1, 1\}$ we define

$$\nu *_{\alpha} p = (\nu \frown \eta^p \upharpoonright [\alpha, \lambda), C^p \setminus \alpha).$$

(Plainly, $\nu *_{\alpha} p \in \mathbb{P}^*$.)

Remark 5.2 \mathbb{P}^* is a natural generalization of the forcing notion used by Goldstern and Shelah [4] to the context of uncountable cardinals.

Proposition 5.3 Let $\bar{\mu} = \langle \mu_{\alpha} : \alpha < \lambda \rangle$, $\mu_{\alpha} = 2^{|\alpha| + \aleph_0}$ (for $\alpha < \lambda$). Then \mathbb{P}^* is a nicely double \mathbf{a} -bounding over $\bar{\mu}$ forcing notion. Also $|\mathbb{P}^*| = 2^{\lambda}$.

Proof One easily verifies that the relation $\leq_{\mathbb{P}^*}$ is transitive and reflexive; also plainly $|\mathbb{P}^*| = 2^{\lambda}$.

Claim 5.4 \mathbb{P}^* is $(< \lambda)$ -complete.

Proof of the Claim Suppose that $\delta < \lambda$ and $\langle p_{\xi} : \xi < \delta \rangle$ is a $\leq_{\mathbb{P}^*}$ -increasing sequence of conditions in \mathbb{P}^* . Let $C = \bigcap_{\xi < \delta} C^{p_{\xi}}$ (it is a club of λ) and let

$\eta : \lambda \rightarrow \{-1, 1\}$ be defined by

(i) if $\gamma < \min(C)$ and $\zeta = \min(\varepsilon < \delta : \gamma < \min(C^{p_{\varepsilon}}))$,
then $\eta(\gamma) = \eta^{p_{\zeta}}(\gamma)$;

(ii) if $\alpha < \beta$ are successive members of the club C , $\alpha \leq \gamma < \beta$ and $\zeta = \min(\varepsilon < \delta : \gamma < \min(C^{p_{\varepsilon}} \setminus (\alpha + 1)))$, then $\eta(\gamma) = \eta^{p_{\zeta}}(\alpha) \cdot \eta^{p_{\zeta}}(\gamma)$.

Plainly, η is well defined and $q \stackrel{\text{def}}{=} (\eta, C) \in \mathbb{P}^*$. We claim that $(\forall \xi < \delta)(p_{\xi} \leq q)$. To this end, suppose $\zeta < \delta$. Clearly, $C \subseteq C^{p_{\zeta}}$. Now, if $\gamma < \min(C^{p_{\zeta}})$, then $\eta(\gamma) = \eta^{p_{\zeta}}(\gamma)$ for some $\zeta \leq \xi$ such that $\gamma < \min(C^{p_{\zeta}})$. Since $p_{\zeta} \leq p_{\xi}$, we have $\eta^{p_{\zeta}}(\gamma) = \eta^{p_{\xi}}(\gamma)$ and thus $\eta^{p_{\xi}}(\gamma) = \eta(\gamma)$.

Next, suppose that $\alpha < \beta$ are successive members of $C^{p_{\xi}}$ and $\alpha \leq \gamma < \beta$. If $\gamma < \min(C)$ and $\zeta = \min(\varepsilon < \delta : \gamma < \min(C^{p_{\varepsilon}}))$, then $\zeta > \xi$, $\eta(\alpha) = \eta^{p_{\zeta}}(\alpha)$ and

$$(*)^1 \quad \eta(\gamma) = \eta^{p_{\zeta}}(\gamma) = \frac{\eta^{p_{\xi}}(\alpha)}{\eta^{p_{\zeta}}(\alpha)} \cdot \eta^{p_{\xi}}(\gamma) = \frac{\eta^{p_{\xi}}(\alpha)}{\eta(\alpha)} \cdot \eta^{p_{\xi}}(\gamma).$$

So assume $C \cap \beta \neq \emptyset$ and let $\alpha' < \beta'$ be successive members of C such that $\alpha' \leq \alpha \leq \gamma < \beta \leq \beta'$. Let $\zeta = \min(\varepsilon < \delta : \gamma < \min(C^{p_{\varepsilon}} \setminus (\alpha' + 1)))$. If $\alpha = \alpha'$, then $\zeta \leq \xi$ and

$$(*)^2 \quad \eta(\gamma) = \eta^{p_{\zeta}}(\alpha) \cdot \eta^{p_{\zeta}}(\gamma) = \eta^{p_{\zeta}}(\alpha) \cdot \frac{\eta^{p_{\xi}}(\alpha)}{\eta^{p_{\zeta}}(\alpha)} \cdot \eta^{p_{\xi}}(\gamma) = \eta^{p_{\xi}}(\alpha) \cdot \eta^{p_{\xi}}(\gamma) = \frac{\eta^{p_{\xi}}(\alpha)}{\eta(\alpha)} \cdot \eta^{p_{\xi}}(\gamma)$$

(as $\eta(\alpha) = \eta(\alpha') = 1$). If $\alpha' < \alpha$, then $\zeta < \xi$ and $\eta(\alpha) = \eta^{p_{\zeta}}(\alpha') \cdot \eta^{p_{\zeta}}(\alpha)$, and hence,

$$(*)^3 \quad \eta(\gamma) = \eta^{p_{\zeta}}(\alpha') \cdot \eta^{p_{\zeta}}(\gamma) = \frac{\eta(\alpha)}{\eta^{p_{\zeta}}(\alpha)} \cdot \eta^{p_{\zeta}}(\gamma) = \frac{\eta(\alpha)}{\eta^{p_{\zeta}}(\alpha)} \cdot \frac{\eta^{p_{\xi}}(\alpha)}{\eta^{p_{\zeta}}(\alpha)} \cdot \eta^{p_{\xi}}(\gamma) = \frac{\eta^{p_{\xi}}(\alpha)}{\eta(\alpha)} \cdot \eta^{p_{\xi}}(\gamma).$$

Clearly, $(*)^1$ – $(*)^3$ are what we need to justify Definition 5.1(1 β) and conclude $p_{\xi} \leq q$. \square

Claim 5.5 *Let $p \in \mathbb{P}^*$. Then GENERIC has a nice winning strategy in the game $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{P}^*)$ (see Definition 3.9).*

Proof of the Claim We will describe a strategy **st** for GENERIC in $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{P}^*)$. Whenever we say GENERIC chooses x such that we mean GENERIC chooses the $<_{\chi}^*$ -first x such that (and likewise for other variants).

During a play of $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{P}^*)$ GENERIC constructs on the side sequences $\langle p_\alpha : \alpha < \lambda \rangle$ and $\bar{\delta} = \langle \delta_\alpha : \alpha < \lambda \rangle$ so that for each $\alpha < \lambda$:

- (a) $\bar{\delta}$ is a strictly increasing continuous sequence of ordinals below λ , $p_\alpha \in \mathbb{P}^*$ and $\{\delta_\xi : \xi \leq \omega + \alpha\} = C^{p_\alpha} \cap (\delta_{\omega+\alpha} + 1)$,
- (b) if $\beta < \alpha$, then $p_\beta \leq p_\alpha$ and $\eta^{p_\alpha} \upharpoonright \delta_{\omega+\beta} = \eta^{p_\beta} \upharpoonright \delta_{\omega+\beta}$,
- (c) $\{\delta_\xi : \xi \leq \omega\} = \{\delta \in C^p : \text{otp}(\delta \cap C^p) \leq \omega\}$ and $p_0 = p$,
- (d) $\delta_{\omega+\alpha+1}$ and $p_{\alpha+1}$ are determined right after stage α of $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{P}^*)$.

So suppose that the two players have arrived to a stage $\alpha < \lambda$ of a play of $\mathcal{D}_{\bar{\mu}}^{\text{rc}2\mathbf{a}}(p, \mathbb{P}^*)$, and GENERIC has constructed on the side $\delta_{\omega+\beta+1}$ and $p_{\beta+1}$ for $\beta < \alpha$. If $\alpha = 0$ or α is a limit ordinal, then conditions (a)–(c) and our rule of taking “the $<_{\chi}^*$ -first” fully determine $\{\delta_\xi : \xi \leq \omega + \alpha\}$ and p_α (the suitable bounds exists essentially by Claim 5.4).

Now GENERIC chooses an enumeration (without repetition) $\bar{\rho} = \langle \rho_j^\alpha : j < \mu_\alpha \rangle$ of $\text{pos}(p_\alpha, \delta_{\omega+\alpha})$ such that $\rho_0^\alpha = \eta^{p_\alpha} \upharpoonright \delta_{\omega+\alpha}$. ANTIGENERIC picks a nonzero ordinal $\zeta_\alpha < \lambda$ and the two players start a subgame of length $\mu_\alpha \cdot \zeta_\alpha$. In the course of the subgame, in addition to her innings p_γ^α , GENERIC will also choose ordinals $\varepsilon_\gamma^\alpha = \varepsilon_\gamma < \lambda$ and sequences $\varphi_\gamma^\alpha = \varphi_\gamma : \varepsilon_\gamma \rightarrow \{-1, 1\}$. These objects will satisfy the following demands (letting q_γ^α be the innings of ANTIGENERIC):

- (e) $\delta_{\omega+\alpha} < \varepsilon_{\gamma'} < \varepsilon_\gamma \in C^{q_\gamma^\alpha}$ and $\varphi_{\gamma'} \upharpoonright [\delta_{\omega+\alpha}, \varepsilon_{\gamma'}) = \varphi_\gamma \upharpoonright [\delta_{\omega+\alpha}, \varepsilon_{\gamma'})$ for $\gamma' < \gamma < \mu_\alpha \cdot \zeta_\alpha$,
- (f) if $\gamma = \mu_\alpha \cdot i + 2j$, $i < \zeta_\alpha$ and $j < \mu_\alpha$, then
 - (i) $\rho_j^\alpha \triangleleft \varphi_\gamma \triangleleft \varphi_{\gamma+1}$, $\varphi_\gamma = \eta^{q_\gamma^\alpha} \upharpoonright \varepsilon_\gamma$, and $\varphi_{\gamma+1}(\delta) = -\eta^{q_{\gamma+1}^\alpha}(\delta)$ for $\delta \in [\delta_{\omega+\alpha}, \varepsilon_{\gamma+1})$,
 - (ii) $p_0^\alpha \geq \rho_0^\alpha *_{\delta_{\omega+\alpha}} p_\alpha$, $\min(C^{p_0^\alpha}) > \delta_{\omega+\alpha}$, and $(\varphi_\gamma \upharpoonright \varepsilon_{\gamma'}) *_{\varepsilon_{\gamma'}} q_{\gamma'}^\alpha \leq p_\gamma^\alpha$ for $\gamma' < \gamma$, and
 - (iii) $q_\gamma^\alpha \leq \varphi_\gamma *_{\varepsilon_\gamma} p_{\gamma+1}^\alpha \leq \varphi_{\gamma+1} *_{\varepsilon_{\gamma+1}} q_{\gamma+1}^\alpha$.

So suppose that the two players have arrived to a stage $\gamma = \mu_\alpha \cdot i + 2j$ ($i < \zeta_\alpha$, $j < \mu_\alpha$) of the subgame and $p_{\gamma'}^\alpha, q_{\gamma'}^\alpha, \varphi_{\gamma'}^\alpha, \varepsilon_{\gamma'}$ have been determined for $\gamma' < \gamma$. Let $\varphi = \rho_j^\alpha \frown \bigcup_{\gamma' < \gamma} \varphi_{\gamma'} \upharpoonright [\delta_{\omega+\alpha}, \varepsilon_{\gamma'})$. It follows from (f) that the sequence

$\langle \langle \varphi \upharpoonright \varepsilon_{\gamma'} *_{\varepsilon_{\gamma'}} q_{\gamma'}^\alpha : \gamma' < \gamma \rangle \rangle$ is $\leq_{\mathbb{P}^*}$ -increasing, so GENERIC may choose an upper bound $p_\gamma^\alpha \in \mathbb{P}^*$ to it. (Note that necessarily $\varphi \triangleleft \eta^{p_\gamma^\alpha}$, $\text{sup}(\varepsilon_{\gamma'} : \gamma' < \gamma) \leq \min(C^{p_\gamma^\alpha})$.) She plays p_γ^α in the subgame and ANTIGENERIC answers with $q_\gamma^\alpha \geq p_\gamma^\alpha$. Now GENERIC lets $\varepsilon_\gamma \in C^{q_\gamma^\alpha}$ be such that $|C^{q_\gamma^\alpha} \cap \varepsilon_\gamma| = 1$ and she puts $\varphi_\gamma = \eta^{q_\gamma^\alpha} \upharpoonright \varepsilon_\gamma$ and she lets $\psi : \varepsilon_\gamma \rightarrow \{-1, 1\}$ be defined by $\psi \upharpoonright \delta_{\omega+\alpha} = \rho_j^\alpha$ and $\psi(\delta) = -\varphi_\gamma(\delta)$ for $\delta \in [\delta_{\omega+\alpha}, \varepsilon_\gamma)$. Then GENERIC plays $p_{\gamma+1}^\alpha = \psi *_{\varepsilon_\gamma} q_\gamma^\alpha$ as her inning at stage $\gamma + 1$ of the subgame and ANTIGENERIC answers with $q_{\gamma+1}^\alpha \geq p_{\gamma+1}^\alpha$. Finally, GENERIC picks $\varepsilon_{\gamma+1} \in C^{q_{\gamma+1}^\alpha}$ such that $|C^{q_{\gamma+1}^\alpha} \cap \varepsilon_{\gamma+1}| = 1$ and she takes $\varphi_{\gamma+1} : \varepsilon_{\gamma+1} \rightarrow \{-1, 1\}$

such that $\varphi_\gamma \triangleleft \varphi_{\gamma+1}$ and $\varphi_{\gamma+1}(\delta) = -\eta^{q_{\gamma+1}^\alpha}(\delta)$ for $\delta \in [\varepsilon_\gamma, \varepsilon_{\gamma+1})$. Plainly, if $i' < i$, $\gamma' = \mu_\alpha \cdot i' + 2j$ then $q_{\gamma'}^\alpha \leq p_\gamma^\alpha$ and $q_{\gamma'+1}^\alpha \leq p_{\gamma+1}^\alpha$ so both p_γ^α and $p_{\gamma+1}^\alpha$ are legal innings in $\mathcal{D}_{\bar{\mu}}^{\text{rc}2a}(p, \mathbb{P}^*)$. Also easily the demands in (e)+(f) are satisfied. Moreover, if $j' < j < \mu_\alpha$ then the conditions $p_{\mu_\alpha+j'}^\alpha$ and $p_{\mu_\alpha+j}^\alpha$ are incompatible.

After the subgame is over, GENERIC lets

$$\varphi = \rho_0^\alpha \frown \bigcup \{\varphi_\gamma \upharpoonright [\delta_{\omega+\alpha}, \varepsilon_\gamma) : \gamma < \mu_\alpha \cdot \xi_\alpha\},$$

and she picks a $\leq_{\mathbb{P}^*}$ -upper bound $p'_{\alpha+1}$ to the increasing sequence

$$\langle (\varphi \upharpoonright \varepsilon_\gamma) *_{\varepsilon_\gamma} q_\gamma^\alpha : \gamma < \mu_\alpha \cdot \xi_\alpha \rangle.$$

Note that $\varepsilon_\gamma \leq \min(C^{p'_{\alpha+1}})$ and $\varphi \upharpoonright \varepsilon_\gamma \triangleleft \eta^{p'_{\alpha+1}}$ for all $\gamma < \mu_\alpha \cdot \xi_\alpha$, so also $\rho_0^\alpha = \eta^{p_\alpha} \upharpoonright \delta_{\omega+\alpha} \triangleleft \eta^{p'_{\alpha+1}}$. Also

$$(g) \quad (\eta^{p'_{\alpha+1}} \upharpoonright \varepsilon_\gamma) *_{\varepsilon_\gamma} q_\gamma^\alpha \leq p'_{\alpha+1} \text{ for all } \gamma < \mu_\alpha \cdot \xi_\alpha.$$

Let $p_{\alpha+1} \in \mathbb{P}^*$ be such that $C^{p_{\alpha+1}} = \{\delta_\xi : \xi \leq \omega + \alpha\} \cup C^{p'_{\alpha+1}}$ and $\eta^{p_{\alpha+1}} = \eta^{p'_{\alpha+1}}$ (plainly $p_\alpha \leq p_{\alpha+1}$) and let $\delta_{\omega+\alpha+1} = \min(C^{p'_{\alpha+1}})$.

This finishes the description of the strategy **st**. Let us argue that **st** is a winning strategy for GENERIC. To this end suppose that

$$(\boxplus) \quad \langle \xi_\alpha, \langle p_\gamma^\alpha, q_\gamma^\alpha : \gamma < \mu_\alpha \cdot \xi_\alpha \rangle : \alpha < \lambda \rangle$$

is a result of a play of $\mathcal{D}_{\bar{\mu}}^{\text{rc}2a}(p, \mathbb{P}^*)$ in which GENERIC follows **st** and the objects constructed on the side are

$$(\boxplus)_\alpha^* \quad \langle p'_\alpha, p_\alpha, \delta_\xi, \langle \varepsilon_\gamma^\alpha, \varphi_\gamma^\alpha : \gamma < \mu_\alpha \cdot \xi_\alpha \rangle, \langle \rho_j^\alpha : j < \mu_\alpha \rangle \rangle$$

(and the demands in (a)–(g) are satisfied). Let $C = \{\delta_\xi : \xi < \lambda\}$ (so it is a club of λ) and $\eta = \bigcup_{\alpha < \lambda} \eta^{p_\alpha} \upharpoonright \delta_{\omega+\alpha}$ (clearly, $\eta : \lambda \longrightarrow \{-1, 1\}$; remember (b)), and let $p^* = (\eta, C)$. It is a condition in \mathbb{P}^* and it is stronger than all p_α (for $\alpha < \lambda$) so also $p^* \geq p$. Suppose that $\alpha < \lambda$ and $p' \geq p^*$. We will show that there is $p'' \geq p'$ such that for some $j < \mu_\alpha$, the condition p'' is stronger than all $q_{\mu_\alpha \cdot i + j}^\alpha$ for all $i < \xi_\alpha$. Without loss of generality, $\min(C^{p'}) \geq \delta_{\omega+\alpha+1}$. Let $j' < \mu_\alpha$ be such that $\eta^{p'} \upharpoonright \delta_{\omega+\alpha} = \rho_{j'}^\alpha$. We consider two cases now.

Case 1 $\eta^{p'}(\delta_{\omega+\alpha}) = \eta^{p^*}(\delta_{\omega+\alpha}) = \eta^{p_{\alpha+1}}(\delta_{\omega+\alpha})$. Then $\eta^{p'} \upharpoonright [\delta_{\omega+\alpha}, \delta_{\omega+\alpha+1}) = \eta^{p_{\alpha+1}} \upharpoonright [\delta_{\omega+\alpha}, \delta_{\omega+\alpha+1})$. Let $j = 2 \cdot j' < \mu_\alpha$, and we will argue that $q_{\mu_\alpha \cdot i + j}^\alpha \leq p'$ for all $i < \xi_\alpha$. So let $i < \xi_\alpha$, $\gamma = \mu_\alpha \cdot i + j$. By the choice of j' we know that $\eta^{p'} \upharpoonright \delta_{\omega+\alpha} = \rho_{j'}^\alpha = \eta^{q_{\gamma'}^\alpha} \upharpoonright \delta_{\omega+\alpha}$ and also

$$\varphi_\gamma^\alpha \upharpoonright [\delta_{\omega+\alpha}, \varepsilon_\gamma^\alpha) = \eta^{p_{\alpha+1}} \upharpoonright [\delta_{\omega+\alpha}, \varepsilon_\gamma^\alpha) = \eta^{p'} \upharpoonright [\delta_{\omega+\alpha}, \varepsilon_\gamma^\alpha).$$

Hence (by (f)(i)) $\eta^{p'} \upharpoonright \varepsilon_\gamma^\alpha = \eta^{q_\gamma^\alpha} \upharpoonright \varepsilon_\gamma^\alpha$ and now

$$\begin{aligned} q_\gamma^\alpha &\leq (\eta^{p'} \upharpoonright \varepsilon_\gamma^\alpha) *_{\varepsilon_\gamma^\alpha} q_\gamma^\alpha \leq (\eta^{p'} \upharpoonright \delta_{\omega+\alpha}) *_{\delta_{\omega+\alpha}} p'_{\alpha+1} \\ &= (\eta^{p'} \upharpoonright \delta_{\omega+\alpha+1}) *_{\delta_{\omega+\alpha+1}} p'_{\alpha+1} = (\eta^{p'} \upharpoonright \delta_{\omega+\alpha+1}) *_{\delta_{\omega+\alpha+1}} p_{\alpha+1} \\ &\leq (\eta^{p'} \upharpoonright \delta_{\omega+\alpha+1}) *_{\delta_{\omega+\alpha+1}} p^* \leq (\eta^{p'} \upharpoonright \delta_{\omega+\alpha+1}) *_{\delta_{\omega+\alpha+1}} p' = p' \end{aligned}$$

(for the second inequality remember (g)).

Case 2 $\eta^{p'}(\delta_{\omega+\alpha}) = -\eta^{p^*}(\delta_{\omega+\alpha}) = -\eta^{p_{\alpha+1}}(\delta_{\omega+\alpha})$. Then $\eta^{p'}(\delta) = -\eta^{p^*}(\delta) = -\eta^{p_{\alpha+1}}(\delta)$ for all $\delta \in [\delta_{\omega+\alpha}, \delta_{\omega+\alpha+1})$. Let $j = 2 \cdot j' + 1$ and let us argue that $q_{\mu_\alpha \cdot i + j}^\alpha \leq p'$ for all $i < \xi_\alpha$. So let $i < \xi_\alpha$, $\gamma = \mu_\alpha \cdot i + j$. As in the previous case we show that $\eta^{p'} \upharpoonright \varepsilon_\gamma^\alpha = \eta^{q_\gamma^\alpha} \upharpoonright \varepsilon_\gamma^\alpha$ and then easily

$$q_\gamma^\alpha \leq (\eta^{p'} \upharpoonright \varepsilon_\gamma) *_{\varepsilon_\gamma^\alpha} q_\gamma^\alpha \leq (\eta^{p'} \upharpoonright \delta_{\omega+\alpha+1}) *_{\delta_{\omega+\alpha+1}} p'_{\alpha+1} \leq (\eta^{p'} \upharpoonright \delta_{\omega+\alpha+1}) *_{\delta_{\omega+\alpha+1}} p' = p'. \quad \square$$

Proposition 5.6 *Let η be a \mathbb{P}^* -name such that*

$$\Vdash_{\mathbb{P}^*} \eta = \bigcup \{ \eta^p \upharpoonright \min(C^p) : p \in \mathcal{G}_{\mathbb{P}^*} \}.$$

Then $\Vdash_{\mathbb{P}^*} \eta : \lambda \longrightarrow \{-1, 1\}$ and for every $s \in \mathbb{Q}_\lambda^0 \cap \mathbf{V}$,

$$\Vdash_{\mathbb{P}^*} \{ \alpha < \lambda : \eta(\alpha) = -1 \} \in \text{fil}(s)^+ \text{ and } \{ \alpha < \lambda : \eta(\alpha) = 1 \} \in \text{fil}(s)^+.$$

Proof It should be clear that $\Vdash_{\mathbb{P}^*} \eta : \lambda \longrightarrow \{-1, 1\}$, so let us show the second statement. Assume $p \in \mathbb{P}^*$, $s \in \mathbb{Q}_\lambda^0$. Choose a continuous increasing sequence $\langle \delta_\xi : \xi < \lambda \rangle \subseteq C^p$ such that for every $\xi < \lambda$ there is $\alpha = \alpha(\xi) \in C^s$ such that $Z_\alpha^s \subseteq [\delta_\xi, \delta_{\xi+1})$. Then let $C = \{ \delta_\xi : \xi < \lambda \text{ is even} \}$ (it is a club of λ) and let $\eta : \lambda \longrightarrow \{-1, 1\}$ be such that

1. $\eta \upharpoonright [\delta_\xi, \delta_{\xi+1}) \in \{ \eta^p \upharpoonright [\delta_\xi, \delta_{\xi+1}), -\eta^p \upharpoonright [\delta_\xi, \delta_{\xi+1}) \}$;
2. if $\xi < \lambda$ is even, then $\{ \delta \in Z_{\alpha(\xi)}^s : \eta(\delta) = 1 \} \in d_{\alpha(\xi)}^s$;
3. if $\xi < \lambda$ is odd, then $\{ \delta \in Z_{\alpha(\xi)}^s : \eta(\delta) = -1 \} \in d_{\alpha(\xi)}^s$.

Now note that $(\eta, C) \in \mathbb{P}^*$ is a condition stronger than p and it forces in \mathbb{P}^* that

$$\{ \alpha < \lambda : \eta(\alpha) = 1 \} \in \text{fil}(s)^+ \text{ and } \{ \alpha < \lambda : \eta(\alpha) = -1 \} \in \text{fil}(s)^+. \quad \square$$

Corollary 5.7 *Assume λ is a strongly inaccessible cardinal. Then there is a forcing notion \mathbb{P} such that*

$$\Vdash_{\mathbb{P}} \text{“} \lambda \text{ is strongly inaccessible and } 2^\lambda = \lambda^{++} \text{ and there is no very reasonable ultrafilter on } \lambda \text{ with a generating system of size } < 2^\lambda \text{”}.$$

Proof We may start with the universe \mathbf{V} in which $2^\lambda = \lambda^+$. Let $\bar{\mathbb{Q}} = \langle \mathbb{P}_\alpha, \mathbb{Q}_\alpha : \alpha < \lambda^{++} \rangle$ be a λ -support iteration of the forcing notion \mathbb{P}^* (see Definition 5.1). This forcing is nicely double **a**-bounding over $\bar{\mu}$ (where $\mu_\alpha = 2^{|\alpha| + \aleph_0}$; remember Proposition 5.3) and hence $\mathbb{P}_{\lambda^{++}}$ is nicely double **a**-bounding over $\bar{\mu}$ (by Theorem 3.13). Using Theorem 3.2 we conclude that $\mathbb{P}_{\lambda^{++}}$ does not collapse any cardinals and forces that $2^\lambda = \lambda^{++}$. Proposition 5.6 implies that

$$\Vdash_{\mathbb{P}_{\lambda^{++}}} \text{“for no family } G^* \subseteq \mathbb{Q}_\lambda^0 \text{ of size } < 2^\lambda, \text{fil}(G^*) \text{ is an ultrafilter on } \lambda \text{”}.$$

□

Problem 5.8

1. Is it consistent that for some uncountable regular cardinal λ we have that there is no super reasonable ultrafilter on λ ? Or even no very reasonable one?

2. In particular, are there super reasonable ultrafilters on λ in the model constructed for Corollary 5.7?
3. Do we need the inaccessibility of λ for the assertions of Corollaries 4.5 and 5.7 (concerning ultrafilters on λ)?

References

- [1] Abraham, U., “Lectures on proper forcing,” pp. 333–94 in *Handbook of Set Theory*, edited by A. Kanamori and M. Foreman, Springer, Dordrecht, 2010. [Zbl 1197.03001](#). [124](#)
- [2] Eisworth, T., “On iterated forcing for successors of regular cardinals,” *Fundamenta Mathematicae*, vol. 179 (2003), pp. 249–66. [Zbl 1066.03051](#). [MR 2029324](#). [124](#)
- [3] Gitik, M., “On nonminimal p -points over a measurable cardinal,” *Annals of Mathematical Logic*, vol. 20 (1981), pp. 269–88. [Zbl 0481.03032](#). [MR 631564](#). [113](#)
- [4] Goldstern, M., and S. Shelah, “Ramsey ultrafilters and the reaping number-Con($\mathfrak{r} < \mathfrak{u}$),” *Annals of Pure and Applied Logic*, vol. 49 (1990), pp. 121–42. [Zbl 0721.03032](#). [MR 1077075](#). [142](#)
- [5] Jech, T., *Set Theory*, 3d millennium edition, Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. [Zbl 1007.03002](#). [MR 1940513](#). [114](#)
- [6] Rosłanowski, A., and S. Shelah, “Norms on possibilities. I. Forcing with trees and creatures,” *Memoirs of the American Mathematical Society*, vol. 141 (1999). [Zbl 0940.03059](#). [MR 1613600](#). [115](#)
- [7] Rosłanowski, A., and S. Shelah, “Reasonably complete forcing notions,” pp. 195–239 in *Set Theory: Recent Trends and Applications*, edited by A. Andretta, vol. 17 of *Quaderni di Matematica*, Dipartimento di Matematica, Seconda Università degli Studi di Napoli, Napoli, 2006. [Zbl 1133.03024](#). [MR 2374767](#). [114](#), [117](#), [124](#), [125](#), [126](#), [127](#), [128](#), [141](#)
- [8] Rosłanowski, A., and S. Shelah, “Sheva-Sheva-Sheva: Large creatures,” *Israel Journal of Mathematics*, vol. 159 (2007), pp. 109–74. [Zbl 1130.03035](#). [MR 2342475](#). [117](#), [124](#), [125](#), [129](#), [141](#)
- [9] Rosłanowski, A., and S. Shelah, “Generating ultrafilters in a reasonable way,” *Mathematical Logic Quarterly*, vol. 54 (2008), pp. 202–20. [Zbl 1145.03025](#). [MR 2402629](#). [114](#), [116](#)
- [10] Rosłanowski, A., and S. Shelah, “Lords of the iteration,” pp. 287–330 in *Set Theory and Its Applications*, vol. 533 of *Contemporary Mathematics (CONM)*, American Mathematical Society, 2011. [117](#)
- [11] Shelah, S., *Proper and Improper Forcing*, 2d edition, Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 1998. [Zbl 0889.03041](#). [MR 1623206](#). [124](#)
- [12] Shelah, S., “The combinatorics of reasonable ultrafilters,” *Fundamenta Mathematicae*, vol. 192 (2006), pp. 1–23. [Zbl 1115.03058](#). [MR 2283626](#). [113](#), [114](#), [115](#), [116](#)

Acknowledgments

The first author would like to thank the Hebrew University of Jerusalem and the Lady Davis Fellowship Trust for awarding him a *Schönbrunn Visiting Professorship* under which this research was carried out. Both authors acknowledge support from the United States-Israel Binational Science Foundation (Grant no. 2006108). This is publication 890 of the second author.

Department of Mathematics
University of Nebraska at Omaha
Omaha NE 68182-0243
USA
roslanow@member.ams.org
<http://www.unomaha.edu/logic>

Einstein Institute of Mathematics
Edmond J. Safra Campus, Givat Ram
The Hebrew University of Jerusalem
Jerusalem 91904
ISRAEL

and
Department of Mathematics
Rutgers University
New Brunswick NJ 08854
USA
shelah@math.huji.ac.il
<http://shelah.logic.at>