

OBSERVATION OF VOLTERRA SYSTEMS WITH SCALAR KERNELS

BERNHARD H. HAAK AND BIRGIT JACOB

Communicated by Stig-Olof Londen

ABSTRACT. Volterra observation systems with scalar kernels are studied. New sufficient conditions for admissibility of observation operators are developed. The results obtained are applied to time-fractional diffusion equations of distributed order.

1. Introduction. Consider the following scalar abstract Volterra system

$$(1) \quad x(t) = x_0 + \int_0^t a(t-s) Ax(s) ds, \quad t \geq 0.$$

Here, the operator A is supposed to be a closed operator with dense domain on a Banach space X , $x_0 \in X$, the kernel function $a \in L^1_{\text{loc}}$ is supposed to be of sub-exponential growth so that its Laplace transform $\hat{a}(\lambda)$ exists for all λ with positive real part, and it is assumed that (1) is *parabolic* in the sense of Prüss [26], that is,

(P1) $\hat{a}(\lambda) \neq 0$ and $\frac{1}{\hat{a}(\lambda)} \in \varrho(A)$ for all λ with positive real part,

(P2) there exists a constant $M \geq 0$ such that $\|(1 - \hat{a}(\lambda)A)^{-1}\| \leq M$ for all λ with positive real part.

In addition, we always assume that the kernel function a is *1-regular*, that is, there is a constant $K > 0$ such that

$$(2) \quad |\lambda \hat{a}'(\lambda)| \leq K |\hat{a}(\lambda)|$$

for all λ with positive real part.

In Prüss [26, Theorem I.3.1] it is shown that under these assumptions, equation (1) admits a unique *solution family*, i.e., a family of bounded linear operators $(S(t))_{t \geq 0}$ on X , such that

Received by the editors on July 8, 2009.

DOI:10.1216/JIE-2011-23-3-421 Copyright ©2011 Rocky Mountain Mathematics Consortium

- (a) $S(0) = I$ and $S(\cdot)$ is strongly continuous on \mathbf{R}_+ .
- (b) $S(t)$ commutes with A , which means $S(t)(D(A)) \subset D(A)$ for all $t \geq 0$, and $AS(t)x = S(t)Ax$ for all $x \in D(A)$ and $t \geq 0$.
- (c) For all $x \in D(A)$ and all $t \geq 0$ the resolvent equations hold:

$$(3) \quad S(t)x = x + \int_0^t a(t-s)AS(s)x ds.$$

Moreover, $S \in C((0, \infty), \mathcal{B}(X))$ and $\|S(t)\| \leq K$.

For some results we need in addition that $-A$ be a sectorial operator of type $\omega \in (0, \pi)$ or that the kernel a is *sectorial of angle* $\theta \in (0, \pi)$. Recall that $-A$ is called a *sectorial operator of type* $\omega \in (0, \pi)$, if the operator A is a closed operator with dense domain on X having its spectrum contained in some open sectorial region of the complex plane, symmetric to the real axis and open to the left:

$$\sigma(A) \subseteq -\Sigma_\omega \quad \text{where} \quad \Sigma_\omega = \{z \in \mathbf{C} : |\arg(z)| < \omega\}$$

for some $\omega \in (0, \pi)$. Moreover, the resolvent of A is supposed to satisfy a growth condition of the type $\|\lambda R(\lambda, A)\| \leq M$ uniformly on each sector $\Sigma_{\pi-\omega-\varepsilon}$. Typical examples of such operators are generators of bounded strongly continuous semigroups, where $\omega \leq \pi/2$. We mention that ‘sectoriality’ may have different meanings for different authors in the literature.

The kernel a is called *sectorial of angle* $\theta \in (0, \pi)$ if

$$\hat{a}(\lambda) \in \Sigma_\theta \quad \text{for all } \lambda \text{ with positive real part.}$$

In particular, when $-A$ and a are both sectorial in the respective sense with angles that sum up to a constant strictly inferior to π , the Volterra equation is parabolic.

The purpose of this article is to present conditions for the admissibility of observation operators to parabolic Volterra equations, that is, we consider the ‘observed’ system

$$(V) \quad \begin{cases} x(t) = x_0 + \int_0^t a(t-s) Ax(s) ds & t \geq 0, \\ y(t) = Cx(t). \end{cases}$$

The operator C in the second line is supposed to be an operator from X into another Banach space Y that acts as a bounded operator from $X_1 \rightarrow Y$ where $X_1 = \mathcal{D}(A)$ is endowed by the graph norm of A . In order to guarantee that the output function lies locally in L_2 , we are interested in the following property.

Definition 1.1. A bounded linear operator $C : X_1 \rightarrow Y$ is called *finite-time admissible* for the Volterra equation (1) if there are constants $\eta, K > 0$ such that

$$\left(\int_0^t \|CS(r)x\|^2 dr \right)^{1/2} \leq Ke^{\eta t} \|x\|$$

for all $t \geq 0$ and all $x \in \mathcal{D}(A)$.

The notion of admissible observation operators is well studied in the literature for Cauchy systems, that is, $a \equiv 1$, see for example [17, 27, 28]. Admissible observation operators for Volterra systems are studied in [12, 18, 19, 22].

The Laplace transform of S , denoted by H , is given by

$$H(\lambda)x = \frac{1}{\lambda}(I - \hat{a}(\lambda)A)^{-1}x, \quad \operatorname{Re} \lambda > 0.$$

The following necessary condition for admissibility was shown in [19].

Proposition 1.2. *If C is a finite-time admissible observation operator for the Volterra equation (1), then there is a constant $M > 0$ such that*

$$(4) \quad \|\sqrt{\operatorname{Re} \lambda}CH(\lambda)\| \leq M, \quad \operatorname{Re} \lambda > 0.$$

In [19] it is shown that (4) is also sufficient for admissibility if X is a Hilbert space, Y is finite-dimensional and A generates a contraction semigroup. However, in general this condition is not sufficient (see, e.g., [17]).

We show that the slightly stronger growth condition on the resolvent

$$\sup_{r>0} \left\| (1 + \log^+ r)^\alpha r^{1/2} CH(r) \right\| < \infty,$$

is sufficient for admissibility if $\alpha > 1/2$ (see Theorem 3.6). This result generalizes the sufficient condition of [29] for Cauchy systems to general Volterra systems (1).

Our second main result, Theorem 3.1 provides a subordination argument to obtain admissibility for the observed Volterra equation from the admissibility of the observation operator for the underlying Cauchy problem. In the particular case of diagonal semigroups and one-dimensional output spaces Y , this improves a direct Carleson measure criterion from [12].

We proceed as follows. In Section 2 we obtain an integral representation for the solution family $(S(t))_{t \geq 0}$ and several regularity results of the corresponding kernel. Section 3 is devoted to sufficient conditions for admissibility of observation operators. A subordination result as well as a general sufficient condition are obtained. Several examples are included as well.

To enhance readability of the calculations, for the rest of this article, K denotes some positive constant that may change from one line to the other unless explicitly quantified.

2. Regularity transfer. The main result of this section is formulated in the following proposition. Let $s(t, \mu)$ denote the solution of the scalar equation

$$s(t, \mu) + \mu \int_0^t a(t-r)s(r, \mu) dr = 1, \quad t > 0, \mu \in \mathbf{C}.$$

We denote by $\mathcal{L}u$ or \widehat{u} the Laplace transform of the function u .

Proposition 2.1. *In addition to the general assumptions, we suppose that A generates a bounded C_0 -semigroup $(T(t))_{t \geq 0}$ and that the kernel a is sectorial of angle $\theta < \pi/2$. Then there exists a family of functions $(v_t)_{t > 0}$ such that*

$$\mathcal{L}(v_t)(\mu) = s(t, \mu) \quad \text{and} \quad S(t) = \int_0^\infty v_t(s)T(s) ds$$

satisfying

$$(a) \sup_{t>0} \|v_t\|_{L^1(\mathbf{R}_+)} < \infty.$$

(b) $\|v_t\|_{L^2(\mathbf{R}_+)} \leq K(t^{-\theta/\pi} + t^{+\theta/\pi})$ where K depends only on θ and the constant in (2).

$$(c) \|v_t\|_{W^{1,1}(\mathbf{R}_+)} \leq K(1 + t^{-2\theta/\pi} + t^{+2\theta/\pi}).$$

Moreover, for $\gamma \in [0, 1]$, $|\mu^\gamma s(t, \mu)| \leq Kt^{-2\gamma\theta/\pi}$.

For the proof of this proposition the following two lemmas are needed.

Lemma 2.2. *In addition to the general assumption on the kernel a , we suppose that a is sectorial of angle $\theta \leq \pi$. Let $\rho_0 := 2\theta/\pi$. Then there exists a constant $c > 0$ such that*

$$|\widehat{a}(\lambda)| \geq \begin{cases} c|\lambda|^{-\rho_0} & |\lambda| \geq 1 \\ c|\lambda|^{\rho_0} & |\lambda| \leq 1 \end{cases}$$

for all $\lambda \in \mathbf{C}$ with $\operatorname{Re} \lambda > 0$.

Proof. We borrow the argument from the proof of [25, Proposition 1]: we start with the analytic completion of the Poisson formula for the harmonic function $H(\lambda) = \arg \widehat{a}(\lambda)$, that is,

$$\log \widehat{a}(\lambda) = \kappa_0 + \frac{i}{\pi} \int_{-\infty}^{\infty} \left[\frac{1 - i\rho\lambda}{\lambda - i\rho} \right] h(i\rho) \frac{d\rho}{1 + \rho^2},$$

where $\kappa_0 \in \mathbf{R}$ is a constant. An easy calculation shows

$$|\operatorname{Re} \log \widehat{a}(\lambda)| \leq \kappa_0 + \rho_0 |\log \lambda|$$

for real $\lambda > 0$, and thus

$$|\widehat{a}(\lambda)| = e^{\log(|\widehat{a}(\lambda)|)} = e^{\Re \log \widehat{a}(\lambda)} \geq \begin{cases} c\lambda^{-\rho_0} & \lambda \geq 1 \\ c\lambda^{\rho_0} & 0 \leq \lambda \leq 1, \end{cases}$$

where $c := e^{-\kappa_0} > 0$. This estimate, together with [26, Lemma 8.1] stating the existence of a constant $c > 0$ such that $c^{-1} \leq |\widehat{a}(|\lambda|)/\widehat{a}(\lambda)| \leq c$ for all $\lambda \in \mathbf{C}$ with $\operatorname{Re} \lambda > 0$ completes the proof. \square




FIGURE 1. Illustration of (5).

Lemma 2.3. *Let $\theta \in (0, \pi)$. Then there exists a $c_\theta > 0$ such that*

$$(5) \quad 1 + |\lambda| \leq C_\theta |1 + \lambda|$$

for all $\lambda \in \Sigma_{\pi-\theta}$.

Proof. Clearly, $\alpha > \tilde{\alpha}$, see Figure 1. Since $\tilde{\alpha} = \frac{\theta}{2}$, the assertion then follows from the fact that $\frac{|1+\lambda|}{1+|\lambda|} = \frac{\sin(\alpha)}{\sin(\theta-\alpha)} \geq \sin(\alpha) \geq \sin(\theta/2)$. \square

Proof of Proposition 2.1. (a) is [26, Proposition I.3.5]. This latter result is also the principal inspiration of the next part:

(b) Let $\sigma(\lambda, \mu) = (\mathcal{L}s(\cdot, \mu))(\lambda)$, i.e., $\sigma(\lambda, \mu) = \frac{1}{\lambda(1+\mu\hat{a}(\lambda))}$. Fix $t > 0$ and $\varepsilon > 0$. Then

$$s(t, \mu) = \frac{1}{2\pi i} \int_{\varepsilon-i\infty}^{\varepsilon+i\infty} e^{\lambda t} \sigma(\lambda, \mu) d\lambda.$$

Then, by partial integration

$$\begin{aligned} s(t, \mu) &= \lim_{R \rightarrow \infty} \frac{1}{2\pi i} \left[\frac{1}{t} e^{\lambda t} \sigma(\lambda, \mu) \right]_{\lambda=\varepsilon-iR}^{\lambda=\varepsilon+iR} \\ &\quad - \frac{1}{2\pi i} \int_{\varepsilon-iR}^{\varepsilon+iR} \frac{1}{t} e^{\lambda t} \frac{d}{d\lambda} \sigma(\lambda, \mu) d\lambda \\ &= -\frac{1}{2\pi i} \int_{\varepsilon-i\infty}^{\varepsilon+i\infty} \frac{1}{t} e^{\lambda t} \frac{d}{d\lambda} \sigma(\lambda, \mu) d\lambda. \end{aligned}$$

An elementary calculation gives

$$\frac{d}{d\lambda} \frac{1}{\lambda(1 + \mu\hat{a}(\lambda))} = -\frac{1 + \mu\hat{a}(\lambda) \left(1 + \left(\frac{\lambda\hat{a}'(\lambda)}{\hat{a}(\lambda)} \right) \right)}{\lambda^2 (1 + \mu\hat{a}(\lambda))^2}.$$

By 1-regularity of the kernel, $\left| \frac{\lambda\hat{a}'(\lambda)}{\hat{a}(\lambda)} \right| \leq C$, and so Lemma 2.3 yields, for any $\delta > 0$,

$$\begin{aligned} &\left(\int_{-\infty}^{\infty} |s(t, \delta + iy)|^2 dy \right)^{1/2} \\ &\leq C_\theta(1 + C) \frac{e^{\varepsilon t}}{2\pi t} \left(\int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} \frac{1}{(\varepsilon^2 + x^2)(1 + |\delta + iy|\hat{a}(\varepsilon + ix))} dx \right)^2 dy \right)^{1/2} \\ &\leq \sqrt{2}C_\theta(1 + C) \frac{e^{\varepsilon t}}{2\pi t} \left(\int_0^{\infty} \left(\int_{-\infty}^{\infty} \frac{1}{(\varepsilon^2 + x^2)(1 + |y|\hat{a}(\varepsilon + ix))} dx \right)^2 dy \right)^{1/2} \\ &\leq \sqrt{2}C_\theta(1 + C) \frac{e^{\varepsilon t}}{2\pi t} \int_{-\infty}^{\infty} \left(\int_0^{\infty} \frac{1}{(\varepsilon^2 + x^2)^2 (1 + |y|\hat{a}(\varepsilon + ix))^2} dy \right)^{1/2} dx \\ &= \sqrt{2}C_\theta(1 + C) \frac{e^{\varepsilon t}}{2\pi t} \int_{-\infty}^{\infty} \frac{1}{(\varepsilon^2 + x^2)|\hat{a}(\varepsilon + ix)|^{1/2}} \left(\int_0^{\infty} \frac{1}{(1 + u)^2} du \right)^{1/2} dx \\ &= \sqrt{2}C_\theta(1 + C) \frac{e^{\varepsilon t}}{2\pi t} \int_{-\infty}^{\infty} \frac{1}{(\varepsilon^2 + x^2)|\hat{a}(\varepsilon + ix)|^{1/2}} dx \end{aligned}$$

Now we split the integral into two parts, by considering the cases $\varepsilon^2 + x^2 \geq 1$ and $\varepsilon^2 + x^2 < 1$ to apply Lemma 2.2 which is controlling $|1/\hat{a}|$. Substituting $x = \varepsilon t$ in both parts easily gives

$$\|s(t, \cdot)\|_{H^2} \leq \tilde{C}_\theta \frac{e^{\varepsilon t}}{t} (\varepsilon^{-1-(\theta/\pi)} + \varepsilon^{-1+(\theta/\pi)}),$$

which yields the assertion by letting $\varepsilon = 1/t$.

(c) We argue in the same spirit as above: by partial integration

$$\frac{d}{d\mu} (\mu s(t, \mu)) = \frac{1}{2\pi i} \int_{\varepsilon-i\infty}^{\varepsilon+i\infty} \frac{1}{t} e^{\lambda t} \frac{d^2}{d\mu d\lambda} (\mu \sigma(\lambda, \mu)) d\lambda.$$

An elementary calculation gives

$$\frac{d^2}{d\lambda d\mu} \frac{\mu \hat{a}(\lambda)}{(\lambda(1 + \mu \hat{a}(\lambda)))^2} = \frac{1 + \mu \hat{a}(\lambda) \left(1 + 2 \left(\frac{\lambda \hat{a}'(\lambda)}{\hat{a}(\lambda)}\right)\right)}{\lambda^2 (1 + \mu \hat{a}(\lambda))^3}.$$

By 1-regularity of the kernel, $\left|\frac{\lambda \hat{a}'(\lambda)}{\hat{a}(\lambda)}\right| \leq C$ and so Lemma 2.3 yields for any $\delta > 0$,

$$\begin{aligned} & \int_{-\infty}^{\infty} \left| \frac{d}{d\mu} (\mu s(t, \delta + iy)) \right| dy \\ & \leq C_\theta (1 + 2C) \frac{e^{\varepsilon t}}{2\pi t} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{1}{(\varepsilon^2 + x^2) (1 + |\delta + iy| |\hat{a}(\varepsilon + ix)|)^2} dx dy \\ & \leq C_\theta (1 + 2C) \frac{e^{\varepsilon t}}{2\pi t} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{1}{(\varepsilon^2 + x^2) (1 + |y| |\hat{a}(\varepsilon + ix)|)^2} dx dy \\ & = 2C_\theta (1 + 2C) \frac{e^{\varepsilon t}}{2\pi t} \int_{-\infty}^{\infty} \frac{1}{(\varepsilon^2 + x^2)} \frac{1}{|\hat{a}(\varepsilon + ix)|} \int_0^\infty \frac{1}{(1+u)^2} du dx \\ & = C_\theta (1 + 2C) \frac{e^{\varepsilon t}}{2\pi t} \int_{-\infty}^{\infty} \frac{1}{(\varepsilon^2 + x^2)} \frac{1}{|\hat{a}(\varepsilon + ix)|} dx \\ & \leq K(t^{-\frac{2\theta}{\pi}} + t^{+\frac{2\theta}{\pi}}) \end{aligned}$$

by choosing $\varepsilon = 1/t$. This shows that $f_t(\mu) = \frac{d}{d\mu} (\mu s(t, \mu)) \in H^1(\mathbf{C}_+)$. Note that $\mathbf{C}_+ := \Sigma_{\pi/2}$ is the open right half plane. We may apply Hardy's inequality (see, e.g., [8, page 198], [14, Theorem 4.2]),

$$\int_0^\infty \frac{|\check{f}_t(r)|}{r} dr \leq \frac{1}{2} \int_{-\infty}^\infty |f_t(i\omega)| d\omega,$$

so that $\frac{\check{f}_t(r)}{r} \in L^1(\mathbf{R}_+)$ is Laplace transformable for every $t > 0$. Since

$$\mathcal{L}\left(\frac{\check{f}_t(r)}{r}\right)(\sigma) = \int_{\sigma}^{\infty} f_t(\mu) d\mu = \sigma s(t, \sigma),$$

we find that $\mu \mapsto \mu s(t, \mu) \in H^\infty(\mathbf{C}_+)$ with a norm controlled by a multiple of $(t^{+(2\theta)/\pi} + t^{-(2\theta)/\pi})$. This implies that $v'_t \in L^1(\mathbf{R}_+)$. Together with (a) the claim follows.

Finally, the same technique gives an estimate for the growth of $s(t, \mu)$:

$$\begin{aligned} \mu^\gamma s(t, \mu) &\leq K \frac{|\mu|^\gamma e^{\varepsilon t}}{t} \int_{-\infty}^{\infty} \frac{1}{(\varepsilon^2 + r^2)(1 + |\mu| |\widehat{a}(\varepsilon + ir)|)} dr \\ &\leq K \frac{e^{\varepsilon t}}{t} \int_{-\infty}^{\infty} \frac{1}{(\varepsilon^2 + r^2) |\widehat{a}(\varepsilon + ir)|^\gamma} \frac{|\mu|^\gamma |\widehat{a}(\varepsilon + ir)|^\gamma}{(1 + |\mu| |\widehat{a}(\varepsilon + ir)|)} dr \\ &\leq K \frac{e^{\varepsilon t}}{t} \int_{-\infty}^{\infty} \frac{1}{(\varepsilon^2 + r^2) |\widehat{a}(\varepsilon + ir)|^\gamma} dr \\ &\stackrel{\varepsilon=1/t}{\leq} K (t^{-(2\gamma\theta)/\pi} + t^{+(2\gamma\theta)/\pi}). \quad \square \end{aligned}$$

3. Sufficient conditions for finite-time admissibility. In this section we present the two main results of this paper.

Theorem 3.1. *Let A generate an exponentially stable strongly continuous semigroup $(T(t))_{t \geq 0}$, and let $C : X_1 \rightarrow Y$ be bounded. Further, we assume that the kernel $a \in L^1_{\text{loc}}(\mathbf{R}_+)$ is of sub-exponential growth, 1-regular and sectorial of angle $\theta < \pi/2$. Then finite-time admissibility of C for the semigroup $(T(t))_{t \geq 0}$ implies that of C for the solution family $(S(t))_{t \geq 0}$.*

Proof. We first note that the assumptions of the theorem apply that equation (1) is parabolic. By Proposition 2.1 there exists a family of functions v_t such that $\|v_t\|_{L^2(\mathbf{R}_+)} \leq K(t^{-\theta/\pi} + t^{+\theta/\pi})$ for some constant $K > 0$ independent of $t > 0$ and

$$S(t) = \int_0^\infty v_t(r) T(r) dr, \quad t > 0.$$

For $x \in \mathcal{D}(A)$ we thus have

$$CS(t)x = \int_0^\infty v_t(r) CT(r)x dr.$$

Note that finite-time admissibility of C for $(T(t))_{t \geq 0}$ implies the existence of a constant $M > 0$ such that

$$\|CT(\cdot)x\|_{L^2(\mathbf{R}_+)} \leq M\|x\|, \quad x \in \mathcal{D}(A),$$

thanks to the exponential stability of $(T(t))_{t \geq 0}$. Thus the result follows from Cauchy-Schwarz inequality. \square

By replacing the Cauchy-Schwarz inequality by Hölder's inequality, similar arguments can be used to obtain sufficient conditions for L^p -admissibility.

Corollary 3.2. *Assume in addition to the hypotheses of the theorem that one of the following conditions is satisfied:*

- (a) *Y is finite-dimensional, X is a Hilbert space and A generates a contraction semigroup;*
- (b) *X is a Hilbert space and A generates a normal, analytic semigroup;*
- (c) *A generates an analytic semigroup and $(-A)^{1/2}$ is an finite-time admissible observation operator for $(T(t))_{t \geq 0}$.*

If there exists a constant $M > 0$ such that

$$(6) \quad \|C(\lambda - A)^{-1}\| \leq \frac{M}{\sqrt{\operatorname{Re} \lambda}}, \quad \operatorname{Re} \lambda > 0,$$

then C is a finite-time admissible observation operator for $(S(t))_{t \geq 0}$.

Proof. Under the assumption of the corollary, the inequality (6) implies that C is a finite-time admissible observation operator for $(S(t))_{t \geq 0}$, see [13, 16, 23]. Thus the result follows from Theorem 3.1. \square

The following corollary is an immediate consequence of the Carleson measure criterion of [15].

Corollary 3.3. *Assume in addition to the hypotheses of the theorem that A admits a Riesz basis of eigenfunctions (e_n) on a Hilbert space X with corresponding eigenvalues λ_n . If $Y = \mathbf{C}$ and if*

$$\mu = \sum_n |Ce_n|^2 \delta_{-\lambda_n}$$

is a Carleson measure on \mathbf{C}_+ , then C is finite-time admissible for the solution family $(S(t))_{t \geq 0}$.

A nice sufficient condition for admissibility for Cauchy problems is given by [29]. For the convenience of the reader we reproduce it here:

Theorem 3.4 [29]. *Let A be the infinitesimal generator of an exponentially stable C_0 -semigroup $(T(t))_{t \geq 0}$ on the Hilbert space H , and let $C : X_1 \rightarrow Y$ be bounded, where Y is another Hilbert space. If for some $\alpha > 1/2$,*

$$(7) \quad \sup_{\Re \lambda > 0} \left\| (1 + \log^+ \Re \lambda)^\alpha (\Re(\lambda))^{1/2} CR(\lambda, A) \right\| < \infty,$$

then C is a finite-time admissible observation operator.

Notice that the condition (7) can be reformulated by saying that in the sense of Evans, Opic and Pick (see [9, 10, 11])

$$\text{for all } x \in X : \|CR(\cdot, A)x\|_{(1/2), \infty, \mathbf{A}} < \infty$$

where $\mathbf{A} = (0, \alpha)$, see also [7] for logarithmic type interpolation functors.

Combining Theorem 3.1 and Theorem 3.4 we receive the following corollary.

Corollary 3.5. *Let, in addition to the assumptions of Theorem 3.4, the kernel a be of sub-exponential growth, 1-regular and sectorial of type $< \pi/2$. Then C is finite-time admissible for the solution family $(S(t))_{t \geq 0}$.*

In some situations, the condition of sectoriality of angle $< \pi/2$ in the above corollary may be inconvenient. Under much weaker assumptions one can also obtain admissibility by the following direct argument.

Theorem 3.6. *Assume that A is a closed operator with dense domain on X , the kernel function $a \in L^1_{\text{loc}}$ is of sub-exponential growth, 1-regular, and (1) is parabolic. Let $C : X_1 \rightarrow Y$ be bounded and assume that for some $\alpha > 1/2$,*

$$(8) \quad \sup_{r>0} \left\| (1 + \log^+ r)^\alpha r^{1/2} CH(r) \right\| < \infty.$$

Then C is finite-time admissible for $(S(t))_{t \geq 0}$.

Note that the exponent $\alpha > 1/2$ is optimal in the sense that for $\alpha < 1/2$ it is even wrong in the case $a \equiv 1$, see [20]. About the case $\alpha = 1/2$ nothing is known at the moment.

Proof. Let $\lambda \in \mathbf{C}_+$, and let φ be such that $\lambda = |\lambda|e^{2i\varphi}$. Then, by resolvent identity,

$$\begin{aligned} & 1 + (\log^+ (\operatorname{Re} \lambda))^\alpha |\lambda|^{1/2} CH(\lambda) \\ &= 1 + (\log^+ (\operatorname{Re} \lambda))^\alpha |\lambda|^{-1/2} C \frac{1}{\widehat{a}(\lambda)} R\left(\frac{1}{\widehat{a}(\lambda)}, A\right) \\ &= (1 + \log^+ |\lambda|)^\alpha |\lambda|^{1/2} CH(|\lambda|) \\ &\quad \times e^{-i\varphi} \frac{\widehat{a}(|\lambda|)}{\widehat{a}(\lambda)} \left[I + \left(\frac{1}{\widehat{a}(|\lambda|)} - \frac{1}{\widehat{a}(\lambda)} \right) R\left(\frac{1}{\widehat{a}(\lambda)}, A\right) \right] \\ &= (1 + \log^+ |\lambda|)^\alpha |\lambda|^{1/2} CH(|\lambda|) \\ &\quad \times e^{-i\varphi} \left[I + \left(\frac{\widehat{a}(\lambda)}{\widehat{a}(|\lambda|)} - 1 \right) (I - \widehat{a}(\lambda)A)^{-1} \right]. \end{aligned}$$

By [26, Lemma 8.1], $c^{-1} \leq |\widehat{a}(|\lambda|)/\widehat{a}(\lambda)| \leq c$ for some $c > 0$. This, together with the parabolicity of (1) yields uniform boundedness of expression in brackets and so the assumed estimate (8) gives

$$(9) \quad \|\lambda \mapsto CH(r+\lambda)\|_{H^\infty(\mathbf{C}_+)} \leq K(1 + \log^+ r)^{-\alpha} r^{-1/2}.$$

Since $(S(t))_{t \geq 0}$ is bounded,

$$\begin{aligned} & \|\lambda \mapsto H(r+\lambda)x\|_{H^2(\mathbf{C}_+)} \\ &= \|e^{-rt}S(t)x\|_{H^2(\mathbf{C}_+)} \leq Kr^{-1/2} \|x\| \quad \text{for all } r > 0 \end{aligned}$$

and together with (9), we infer

$$(10) \quad \|\lambda \mapsto CH(r+\lambda)^2 x\|_{H^2(\mathbf{C}_+)} \leq \frac{K}{(1 + \log^+ r)^\alpha r} \|x\| \quad \text{for all } r > 0.$$

Moreover, the estimate

$$\begin{aligned} \left\| \lambda \mapsto \frac{1}{r+\lambda} CH(r+\lambda)x \right\|_{H^2(\mathbf{C}_+)} \\ \leq \left\| \lambda \mapsto CH(r+\lambda)x \right\|_{H^\infty(\mathbf{C}_+)} \left\| \lambda \mapsto \frac{1}{r+\lambda} \right\|_{H^2(\mathbf{C}_+)} \end{aligned}$$

implies

$$(11) \quad \left\| \lambda \mapsto \frac{1}{r+\lambda} CH(r+\lambda)x \right\|_{H^2(\mathbf{C}_+)} \leq \frac{K}{(1 + \log^+ r)^\alpha r} \|x\| \quad \text{for all } r > 0.$$

Since $\frac{d}{d\lambda} H(\lambda) = \left(\frac{\lambda \widehat{a}'(\lambda)}{\widehat{a}(\lambda)} \right) H(\lambda)^2 - \frac{1}{\lambda} \left(1 + \frac{\lambda \widehat{a}'(\lambda)}{\widehat{a}(\lambda)} \right) H(\lambda)$, we infer from (10) and (11) that

$$\left\| \mu \mapsto \frac{d}{d\mu} CH(r+\mu)x \right\|_{H^2(\mathbf{C}_+)} \leq \frac{K}{(1 + \log^+ r)^\alpha r} \|x\| \quad \text{for all } r > 0.$$

Finally, (inverse) Laplace transform yields

$$\|t \mapsto r t e^{-rt} CS(t)x\|_{L^2(\mathbf{R}_+)} \leq \frac{K}{(1 + \log^+ r)^\alpha} \|x\| \quad \text{for all } r > 0$$

and that is the estimate we need in the following dyadic decomposition argument: notice that $x e^{-x} \geq 2e^{-2}$ for $x \in [1, 2]$. Fix some $t_0 > 0$. Then,

$$\begin{aligned} \int_0^{t_0} \|CS(t)x\|^2 dt &= \sum_{n=1}^{\infty} \int_{t_0 2^{-n}}^{t_0 2^{-n+1}} \|CS(t)x\|^2 dt \\ &\leq \frac{e^4}{4} \sum_{n=1}^{\infty} \int_{t_0 2^{-n}}^{t_0 2^{-n+1}} \|t 2^n t_0^{-1} e^{t 2^n t_0^{-1}} CS(t)x\|^2 dt \\ &\leq K \sum_{n=1}^{\infty} \frac{1}{(1 + \log^+(2^n t_0^{-1}))^{2\alpha}} \|x\|^2 \leq K \|x\|^2. \quad \square \end{aligned}$$

4. Example. In this section we apply the results obtained to time-fractional diffusion equations of distributed order.

Let A generate an exponentially stable strongly continuous semigroup $(T(t))_{t \geq 0}$. For $\omega > 0$ and $0 < \alpha < \beta \leq 1$, we study a time-fractional diffusion equation of distributed order of the form

$$(12) \quad \begin{aligned} \omega D_t^\alpha x(t) + D_t^\beta x(t) &= Ax(t), \quad t \geq 0, \\ x(0) &= x_0, \end{aligned}$$

where $D_t^\alpha x = (-\frac{\partial}{\partial t})^\alpha x$ denotes the Caputo derivative of x , given by the Phillips functional calculus of the right shift semigroup, that is,

$$D_t^\gamma x(t) = \frac{1}{\Gamma(1-\gamma)} \int_0^t (t-s)^{-\gamma} x'(s) ds.$$

for $\gamma \in (0, 1)$. Time-fractional diffusion equations of distributed order have attracted attention as a possible tool for the description of anomalous diffusion and relaxation phenomena in many areas such as turbulence, disordered medium, intermittent chaotic systems, mathematical finance and stochastic mechanics. For further information on time-fractional diffusion equations of distributed order, we refer the reader to [1–6, 21, 24].

Using the Laplace transform equation (12) is equivalent to

$$x(t) = x_0 + \int_0^t a(t-s) Ax(s) ds,$$

where

$$a(t) = a(t) = t^{\beta-1} E_{\beta-\alpha, \beta}(-\omega t^{\beta-\alpha})$$

Here $E_{\gamma, \delta}$, where $\gamma, \delta > 0$, denotes the Mittag-Leffler function

$$E_{\gamma, \delta}(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\gamma k + \delta)}.$$

The Laplace transformation of the kernel a is given by

$$\hat{a}(\lambda) = \frac{\lambda^{-\alpha}}{\omega + \lambda^{\beta-\alpha}}.$$

Thus the kernel a satisfies the assumption of Theorem 3.1.

We note that this example does, e.g., not satisfy the assumption of [12, Theorem 3.10] due to the ‘mixed’ growth conditions near infinity and the origin, such that, even when A is the Dirichlet Laplacian on a bounded domain, the latter result cannot be used to guarantee admissibility whereas a ‘standard’ Carleson measure criterion and the subordination result of Corollary 3.3 still applies.

Acknowledgments. This research was done at the Mathematisches Forschungsinstitut Oberwolfach during a stay within the Research in Pairs Programme from June 15 to June 28, 2008. We would like to thank the MFO for excellent working conditions.

REFERENCES

1. T.M. Atanackovic, *A generalized model for the uniaxial isothermal deformation of a viscoelastic body*, Acta Mech. **159** (2002), 77–86.
2. T.M. Atanackovic, M. Budincevic, and S. Pilipovic, *On a fractional distributed-order oscillator*, J. Phys. A, **38** (2005), 6703–6713.
3. R.L. Bagley and P.J. Torvik, *On the existence of the order domain and the solution of distributed order equations. I*, Inter. J. Appl. Math. **2** (2000), 865–882.
4. ———, *On the existence of the order domain and the solution of distributed order equations. II*, Inter. J. Appl. Math. **2** (2000), 965–987.
5. A. Carpinteri and F. Mainardi, *Fractals and fractional calculus in continuum mechanics*, CISM Courses and Lectures **378**, Springer, Wien, 1997.
6. A. Chechkin, R. Gorenflo and I. Sokolov, *Retarding sub- and accelerating superdiffusion governed by distributed order fractional diffusion equations*, arXiv:cond-mat/0202213v1, 2002.
7. F. Cobos, L.M. Fernández-Cabrera and H. Triebel, *Abstract and concrete logarithmic interpolation spaces*, J. London Math. Soc. **70** (2004), 231–243.
8. P.L. Duren, *Theory of H^p spaces*, Pure Appl. Math. **38**, Academic Press, New York, 1970.
9. W.D. Evans and B. Opic, *Real interpolation with logarithmic functors and reiteration*, Canad. J. Math. **52** (2000), 920–960.
10. W.D. Evans, B. Opic and L. Pick, *Interpolation of operators on scales of generalized Lorentz-Zygmund spaces*, Math. Nachr. **182** (1996), 127–181.
11. ———, *Real interpolation with logarithmic functors*, J. Inequal. Appl. **7** (2002), 187–269.
12. B. Haak, B. Jacob, J. Partington and S. Pott, *Admissibility and controllability of diagonal volterra equations with scalar inputs*, J. Differential Equations **246** (2009), 4423–4440.
13. S. Hansen and G. Weiss, *The operator Carleson measure criterion for admissibility of control operators for diagonal semigroups on l^2* , Syst. Contr. Lett. **16** (1991), 219–227.

14. E. Hille and J. Tamarkin, *On the absolute integrability of Fourier transforms*, Fund. Math. **25** (1935), 329–352.
15. L.F. Ho and D.L. Russell, *Admissible input elements for systems in Hilbert space and a Carleson measure criterion*, SIAM J. Control Optim. **21** (1983), 614–640; erratum **21**, 985–986.
16. B. Jacob and J.R. Partington, *The Weiss conjecture on admissibility of observation operators for contraction semigroups*, Integral Equations Operator Theory **40** (2001), 231–243.
17. ———, *Admissibility of control and observation operators for semigroups: A survey*, in *Current trends in operator theory and its applications*, Oper. Theory Adv. Appl. **149**, Birkhäuser, Basel, 2004.
18. ———, *Admissible control and observation operators for Volterra integral equations*, J. Evol. Equ. **4** (2004), 333–343.
19. ———, *A resolvent test for admissibility of Volterra observation operators*, J. Math. Anal. Appl. **332** (2007), 346–355.
20. B. Jacob, J.R. Partington and S. Pott, *Zero-class admissibility of observation operators*, Systems Control Lett. **58** (2009), 406–412.
21. G. Jumarie, *Fractionalization of the complex-valued Brownian motion of order n using Riemann-Liouville derivative. Applications to mathematical finance and stochastic mechanics*, Chaos Solitons Fractals **28** (2006), 1285–1305.
22. M. Jung, *Admissibility of control operators for solution families to Volterra integral equations*, SIAM J. Control Optim. **38** (2000), 1323–1333 (electronic).
23. C. Le Merdy, *The Weiss conjecture for bounded analytic semigroups*, J. London Math. Soc. **67** (2003), 715–738.
24. F. Mainardi and G. Pagnini, *The role of the Fox-Wright functions in fractional sub-diffusion of distributed order*, J. Comput. Appl. Math. **207** (2007), 245–257.
25. S. Monniaux and J. Prüss, *A theorem of the Dore-Venni type for noncommuting operators*, Trans. Amer. Math. Soc. **349** (1997), 4787–4814.
26. J. Prüss, *Evolutionary integral equations and applications*, Mono. Math. **87**, Birkhäuser Verlag, Basel, 1993.
27. O. Staffans, *Well-posed linear systems*, Encyc. Math. Appl. **103**, Cambridge University Press, Cambridge, 2005.
28. G. Weiss, *Admissible observation operators for linear semigroups*, Israel J. Math. **65** (1989), 17–43.
29. H. Zwart, *Sufficient conditions for admissibility*, Syst. Contr. Lett. **54** (2005), 973–979.

INSTITUT DE MATHÉMATIQUES DE BORDEAUX, 351 COURS DE LA LIBÉRATION,
33405 TALENCE CEDEX, FRANCE
Email address: Bernhard.Haak@math.u-bordeaux1.fr

DEPARTMENT OF MATHEMATICS AND INFORMATICS, GAUSS STRASSE 20, D-42119
WUPPERTAL, GERMANY
Email address: jacob@math.uni-wuppertal.de