

On Partial and Paraconsistent Logics

REINHARD MUSKENS

Abstract In this paper we consider the theory of predicate logics in which the principle of bivalence or the principle of noncontradiction or both fail. Such logics are partial or paraconsistent or both. We consider sequent calculi for these logics and prove model existence. For L_4 , the most general logic under consideration, we also prove a version of the Craig-Lyndon Interpolation Theorem. The paper shows that many techniques used for classical predicate logic generalize to partial and paraconsistent logics once the right setup is chosen. Our logic L_4 has a semantics that also underlies Belnap's logic and is related to the logic of bilattices. L_4 is in focus most of the time, but it is also shown how results obtained for L_4 can be transferred to several variants.

1 Introduction The principle of bivalence states that a sentence is either true or false; the principle of noncontradiction says that no sentence is both true and false. These two principles have been part and parcel of all standard formulations of logic since the subject began with Aristotle. But they need not be accepted and in fact if one of them is rejected (or if both are) we get a straightforward generalization of classical logic. Allowing the possibility that a sentence is neither true nor false yields a partial logic (see, e.g., Cleave [9], Blamey [5], Langholm [21]) and allowing sentences to be both true and false leads to a paraconsistent logic. The possibility of having partiality and paraconsistency at the same time is exemplified in Belnap [4]. Partial and paraconsistent logics have applications in database theory (see the motivation given in [4]), in treatments of the Liar paradox (see, e.g., Woodruff [30], Visser [29]), in knowledge representation (Thijsse [28], Jaspars [19]), in logic programming (Fitting [13], Bochman [6]), and in natural language semantics (see Barwise and Perry [3], Muskens [23], [24]). Apart from such applications there is another motivation to study them which derives from an interest in classical logic: how much of the latter's metatheory rests on bivalence and noncontradiction and how much remains if these principles are removed?

In this paper we shall consider predicate logics in which bivalence or noncontradiction or both fail. Our prime example will be the logic L_4 , a partial and paraconsistent predicate logic based on [4]. For this logic we shall give a simple Gentzen

Received November 30, 1998; revised April 20, 2000

sequent calculus and prove model existence (with useful corollaries such as completeness, compactness, and the Löwenheim-Skolem theorem) and Craig interpolation.

There will be logics other than \mathbf{L}_4 which we shall also consider briefly. Most of our methods are in fact independent from the way in which certain basic choices for setting up partial or paraconsistent logics are resolved. One such basic choice concerns the notion of *consequence*. In classical logic a set of premises Γ entails a set of conclusions Δ if and only if in each model in which all $\gamma \in \Gamma$ are true some $\delta \in \Delta$ is true. The same definition can be used for a partial or paraconsistent logic and in this case one obtains a notion \models^{tr} , *transmission of truth*. In classical systems this is indistinguishable from *transmission of nonfalsity*: We can define $\Gamma \models^{nf} \Delta$ to hold if and only if some $\delta \in \Delta$ is not false in each model in which no $\gamma \in \Gamma$ is false or, conversely, if and only if some $\gamma \in \Gamma$ is false whenever all $\delta \in \Delta$ are. A basic fact about the logics under consideration is that transmission of truth and transmission of nonfalsity in general are not equivalent. For example, if we have logical constants (zero-place connectives) \mathbf{t} and \mathbf{b} , with \mathbf{t} denoting the proposition that is always true and never false and \mathbf{b} the proposition that is always both true and false, we have $\mathbf{t} \models^{tr} \mathbf{b}$ but not $\mathbf{t} \models^{nf} \mathbf{b}$. Note that we needed the nonclassical connective \mathbf{b} for this example; it is well known that \models^{tr} and \models^{nf} are identical on formulas with only the classical \mathbf{t} , \neg , \wedge , \vee , \forall , and \exists (given the interpretations for these connectives considered below).

Which notion is the “right” notion of validity, \models^{tr} or \models^{nf} ? In \mathbf{L}_4 neither of these notions is taken, but the choice is resolved by requiring transmission of truth as well as nonfalsity. The relation of entailment is defined by letting $\Gamma \models \Delta$ if and only if $\Gamma \models^{tr} \Delta$ and $\Gamma \models^{nf} \Delta$. Thus, while \models^{tr} and \models^{nf} are duals in an obvious sense, the notion \models will be its own dual. This, we feel, is a strong argument in its favor. The choice for a “double-barreled” notion of consequence was also taken in [4], [5], and [24], but in the literature we find instantiations of the other possibilities as well. In [9], [21], [28], [19], and [6], for example, \models^{tr} is taken to be the basic notion of consequence, as it is in Hähnle [17] and Baaz, Fermüller, and Zach [2]. In Holden [18], on the other hand, we find its dual \models^{nf} . The results in this paper generalize over such variations in a simple way. As will be explained in some detail below, our basic notion will be \models ; but extra *structural elements* $\not\models$ and $\not\models^*$ can be present in our Gentzen sequents. Addition of $\not\models$ (but not $\not\models^*$) leads to the notion \models^{tr} , while adding $\not\models^*$ (but not $\not\models$) leads to \models^{nf} . The presence of $\not\models$ means that transmission of falsity from conclusions to premises need not obtain while the presence of $\not\models^*$ signals that there need be no transmission of truth.

The double-barreled notion of consequence distinguishes \mathbf{L}_4 from the approach taken in the tradition of *many-valued* logics (see, e.g., Schröter [26], Rousseau [25], Carnielli [7], [8], [2], and [17], and Zach [31]). Such logics are standardly associated with a set N of truth values and a set $D \subseteq N$ of *designated* truth values. A sentence φ follows from a set of sentences Γ in this approach if and only if φ evaluates to an element of D in every model in which each $\gamma \in \Gamma$ evaluates to an element of D . \models^{tr} and \models^{nf} easily fit within this scheme, as will be seen below, but \models is an animal of a different kind. Properties of \models cannot always be reduced to properties of \models^{tr} and \models^{nf} .

It is well known that partial and paraconsistent logics can usually be embedded into classical logic. Such embeddings (see Feferman [12], Gilmore [15], [21], and

[24]) give useful abstract information about the embedded logic, but for more concrete information direct methods are necessary. For instance, [24] observes that for \mathbf{L}_4 , the compactness theorem, the Löwenheim-Skolem theorem, and the recursive axiomatizability of \models all follow from a simple embedding into predicate logic and the corresponding theorems there. But this method of translation does not give a concrete axiomatization and cannot be used to obtain Interpolation.

Apart from their technical use, embeddings of partial and paraconsistent logics into the classical system give some intuitive guidance. The existence of such embeddings strongly suggests that many proofs for the classical theory will generalize to cases where bivalence or noncontradiction are not assumed to hold. One purpose of this paper, next to simply providing concrete syntactic characterizations of the consequence relation for various useful logics and studying properties of this consequence relation, is to show that this is indeed the case. The reader, therefore, should not be disappointed if our proofs turn out to be generalizations of similar proofs for the classical theory. Fascinating as partial and paraconsistent logics are, many of their properties can be studied with the same arsenal of methods that is used for the classical case.

The proof system in this paper will stay close to the sequent format introduced in Langholm [22]. For reasons that will be discussed below, Langholm's sequents are set up as 'quadrants', with *four* structural positions instead of the usual two (left and right). We found that this format helped to formulate Gentzen rules in a very concise way.

Our axiomatization of the \mathbf{L}_4 consequence relation with the help of Langholm's quadrants and the two structural elements $\not\vdash$ and $\not\vdash$ may seem strange at a first encounter. Is not a Gentzen calculus which depends on such unusual devices simply a "hack"? One way to test the quality of a calculus is to see whether it admits of Interpolation and indeed we shall find that a version of the Craig-Lyndon theorem can be proved in a very straightforward way. The result here should well be distinguished from the result in Langholm [21], where Interpolation is proved for a partial, but not paraconsistent, logic based on \models^{tr} . On the one hand, we have not been able to extend our interpolation result for \models to three-valued logics. On the other, [21] remarks that although \models "is perhaps a more worthy counterpart to the classical consequence relation," an interpolation theorem for this notion "does not seem to be as easily obtained as the interpolation theorem for \models_3 " (the partial but not paraconsistent version of our \models^{tr}). Langholm tends to emphasize the difference between partial logic and classical logic, arguing that the resemblances that people have noted between the two extend only to concepts (such as \models^{tr} and \models^{nf}) that concern only the truth or only the falsity behavior of sentences, while "the picture becomes considerably more complex when questions concerning the interaction between the two are brought into focus." If the results in this paper are right, such conclusions are at least not warranted for four-valued logics.

The setup of the rest of the paper will be as follows. In the next section we recall what happens when the classical connection between truth and falsity is given up: under reasonable assumptions we then arrive at the bilattice *FOUR* which was introduced in Belnap [4]. Section 3 describes the truth definition for \mathbf{L}_4 and shows the functional completeness of its basic set of connectives. Section 4 discusses semantic




Figure 1: The bilattice *FOUR*

consequence and introduces sequents, and Section 5 gives a sequent calculus. The completeness proof for this calculus is given in Section 6 via model existence. Section 7 discusses ways to base our logic on three instead of four values, and the logic’s interpolation theorem is proved in Section 8. A last section gives conclusions.

2 The elements of *FOUR* Let us introduce the basic notions that lead to the logic L_4 . If we give up both bivalence and noncontradiction, that is, if we sever the classical relation between truth and falsity completely, we arrive at Belnap’s four values *true and not false*, *false and not true*, *both true and false*, and *neither true nor false* (see [4]). These we shall abbreviate as **t**, **f**, **b**, and **n**, respectively. The first two of these values correspond to the classical possibilities; the third represents contradicting information; and the fourth no information at all. If we order $\mathbf{4} = \{\mathbf{t}, \mathbf{f}, \mathbf{b}, \mathbf{n}\}$ according to the information content of its elements, we arrive at the lattice ordering \leq_k depicted in Figure 1. If we order the same elements with respect to their degrees of truth and nonfalsity, we get the lattice ordering \leq_t . The structure $\langle \mathbf{4}, \leq_k \rangle$ was called an *approximation lattice* in [4], whereas $\langle \mathbf{4}, \leq_t \rangle$ was called a *logical lattice*.

Given that formulas φ and ψ take their values in $\mathbf{4}$, how can we compute values for $\neg\varphi$, $\varphi \wedge \psi$, $\varphi \vee \psi$? This can be answered in a very simple way by separating conditions for truth and conditions for falsity (see, e.g., Dunn [11]):

1. $\neg\varphi$ is true if and only if φ is false,
 $\neg\varphi$ is false if and only if φ is true;
2. $\varphi \wedge \psi$ is true if and only if φ is true and ψ is true,
 $\varphi \wedge \psi$ is false if and only if φ is false or ψ is false;
3. $\varphi \vee \psi$ is true if and only if φ is true or ψ is true,
 $\varphi \vee \psi$ is false if and only if φ is false and ψ is false.

So, for example, if φ receives the value **n** (neither true nor false) and ψ gets the value **t** (true and not false), then $\varphi \wedge \psi$ is evaluated as **n**: $\varphi \wedge \psi$ is not true since φ is not true and it is not false since neither φ nor ψ is false. Reasoning similarly in all other cases we arrive at the following tables.

\wedge	t	f	n	b	\vee	t	f	n	b	\neg	
t	t	f	n	b	t	t	t	t	t	t	f
f	f	f	f	f	f	t	f	n	b	f	t
n	n	f	n	f	n	t	n	n	t	n	n
b	b	f	f	b	b	t	b	t	b	b	b

It is easily seen that \wedge is meet and \vee is join in the lattice $\langle \mathbf{4}, \leq_t \rangle$. Note that the Strong Kleene truth tables are obtained if we restrict \wedge , \vee , and \neg to $\{\mathbf{t}, \mathbf{f}, \mathbf{n}\}$ or to $\{\mathbf{t}, \mathbf{f}, \mathbf{b}\}$. In fact, restricting values to $\{\mathbf{t}, \mathbf{f}, \mathbf{n}\}$ corresponds to accepting noncontradiction but not bivalence, while restricting to $\{\mathbf{t}, \mathbf{f}, \mathbf{b}\}$ corresponds to accepting bivalence but leaving open the possibility of paraconsistency. In Section 7 we shall show how the results from this paper can easily be adapted to logics that are either partial or paraconsistent, but not both.

The structure $FOUR = \langle \mathbf{4}, \leq_t, \leq_k, \neg \rangle$ is a prime example of what Ginzburg [16] has called a *bilattice*. For the general notion of a bilattice see [16] or one of Fitting’s papers on the subject (e.g., Fitting [13]). Here we shall content ourselves with considering predicate logics in which formulas can have their values only in $\mathbf{4}$.

\otimes	t	f	n	b	\oplus	t	f	n	b	\neg	
t	t	n	n	t	t	t	b	t	b	t	t
f	n	f	n	f	f	b	f	f	b	f	f
n	n	n	n	n	n	t	f	n	b	n	b
b	t	f	n	b	b	b	b	b	b	b	n

The truth functions \wedge , \vee , and \neg are *classical* in the sense that they always yield a value in $\{\mathbf{t}, \mathbf{f}\}$ when given arguments from $\{\mathbf{t}, \mathbf{f}\}$. The zero-place functions **t** and **f** are also classical in this sense, of course. Composition of classical functions can only give new classical functions and so it is clear that $\{\wedge, \vee, \neg, \mathbf{t}, \mathbf{f}\}$ cannot be functionally complete on $\mathbf{4}$. In fact many new interesting operators can be investigated once the classical connections between truth and falsity have been severed. For example, one can introduce connectives \otimes and \oplus , with truth tables as above, into the logical language. These connectives (called “consensus” and “gullibility” in the literature on bilattices) correspond to meet and join in the approximation lattice $\langle \mathbf{4}, \leq_k \rangle$, while \neg , *conflation*, does on $\langle \mathbf{4}, \leq_k \rangle$ what negation does on $\langle \mathbf{4}, \leq_t \rangle$. As was shown in [24] the set $\{\otimes, \wedge, \neg, \neg\}$ is in fact functionally complete.

Since \mathbf{L}_4 is based on the bilattice $FOUR$ there is a clear relation between this logic and what Arieli and Avron [1] have called *bilattice logics*. But there is also an important difference between \mathbf{L}_4 and the system presented in [1], as the latter’s consequence relation is a certain generalization of \models^{tr} adequate for arbitrary bilattices. Our preferred notion of consequence is \models , which directly reflects \leq_t . (Another difference is our restriction to $FOUR$, of course.) In Section 5 below we shall give more information about the difference between our setup and Arieli and Avron’s.

3 Satisfaction and functional completeness Having described the basic domain of truth values for \mathbf{L}_4 we may proceed with defining the syntax and semantics of the logic. The syntax is defined in the usual way with the help of function and relation symbols in some countable language \mathcal{L} , a countable set of variables, and the logical operators $\{\mathbf{n}, \approx, \neg, \neg, \wedge, \forall\}$. Of the latter, we have met **n**, \neg , \neg , and \wedge already; \approx is identity and \forall is universal quantification. The usual definitions of *free* and *bound* variables, *sentences*, and so on, obtain. *Constants* are zero-place function symbols. We write $[t_1/x_1, \dots, t_n/x_n]\varphi$ for the simultaneous substitution of t_1 for x_1 and \dots and t_n for x_n in φ . The function $[t_1/x_1, \dots, t_n/x_n]$ is called a *substitution*. φ is a *substitution instance* of ψ if $\varphi = \sigma\psi$ for some substitution σ . A *model* is a pair $\langle \mathcal{D}, I \rangle$ where

$\mathcal{D} \neq \emptyset$ and I is a function with domain \mathcal{L} such that $I(f)$ is an n -ary function on \mathcal{D} if $f \in \mathcal{L}$ is an n -ary function symbol and $I(R)$ is a pair of n -ary relations on \mathcal{D} if $R \in \mathcal{L}$ is an n -ary relation symbol. We denote the first element of this pair as $I^+(R)$, the second element as $I^-(R)$. Assignments and the notation $a_{d_1}^{x_1} \cdots a_{d_n}^{x_n}$ for assignments are defined as usual. The value of a term t in a model \mathcal{M} under an assignment a is written as $\llbracket t \rrbracket^{\mathcal{M},a}$, or $\llbracket t \rrbracket^{\mathcal{M}}$ if t is closed.

Definition 3.1 We define the three-place relations $\mathcal{M} \models \varphi[a]$ (formula φ is true in model \mathcal{M} under assignment a) and $\mathcal{M} \models \neg \varphi[a]$ (φ is false in \mathcal{M} under a) as follows.

1. $\mathcal{M} \models \mathbf{n}[a]$,
 $\mathcal{M} \models \mathbf{f}[a]$;
2. $\mathcal{M} \models Rt_1 \dots t_n[a] \iff \langle \llbracket t_1 \rrbracket^{\mathcal{M},a}, \dots, \llbracket t_n \rrbracket^{\mathcal{M},a} \rangle \in I^+(R)$,
 $\mathcal{M} \models \neg Rt_1 \dots t_n[a] \iff \langle \llbracket t_1 \rrbracket^{\mathcal{M},a}, \dots, \llbracket t_n \rrbracket^{\mathcal{M},a} \rangle \in I^-(R)$;
3. $\mathcal{M} \models t_1 \approx t_2[a] \iff \llbracket t_1 \rrbracket^{\mathcal{M},a} = \llbracket t_2 \rrbracket^{\mathcal{M},a}$,
 $\mathcal{M} \models \neg t_1 \approx t_2[a] \iff \llbracket t_1 \rrbracket^{\mathcal{M},a} \neq \llbracket t_2 \rrbracket^{\mathcal{M},a}$;
4. $\mathcal{M} \models \neg \varphi[a] \iff \mathcal{M} \models \varphi[a]$,
 $\mathcal{M} \models \varphi[a] \iff \mathcal{M} \models \neg \varphi[a]$;
5. $\mathcal{M} \models \neg \varphi[a] \iff \mathcal{M} \not\models \varphi[a]$,
 $\mathcal{M} \models \varphi[a] \iff \mathcal{M} \not\models \neg \varphi[a]$;
6. $\mathcal{M} \models \varphi \wedge \psi[a] \iff \mathcal{M} \models \varphi[a] \ \& \ \mathcal{M} \models \psi[a]$,
 $\mathcal{M} \models \neg \varphi \wedge \psi[a] \iff \mathcal{M} \models \varphi[a] \ \text{or} \ \mathcal{M} \models \psi[a]$;
7. $\mathcal{M} \models \forall x \varphi[a] \iff \mathcal{M} \models \varphi[a_d^x]$ for all $d \in \mathcal{D}$,
 $\mathcal{M} \models \neg \forall x \varphi[a] \iff \mathcal{M} \models \varphi[a_d^x]$ for some $d \in \mathcal{D}$.

We write $\mathcal{M} \models \varphi$ ($\mathcal{M} \models \neg \varphi$) if φ is a sentence and $\mathcal{M} \models \varphi[a]$ ($\mathcal{M} \models \neg \varphi[a]$) for some a .

Definition 3.1 uses the format of assigning truth conditions and falsity conditions separately, as discussed in the previous section. Alternatively, we can let formulas take their values directly in **4** by letting

$$\begin{aligned} \llbracket \varphi \rrbracket^{\mathcal{M},a} = \mathbf{t} & \quad \text{iff} \quad \mathcal{M} \models \varphi[a] \ \text{and} \ \mathcal{M} \not\models \neg \varphi[a], \\ \llbracket \varphi \rrbracket^{\mathcal{M},a} = \mathbf{f} & \quad \text{iff} \quad \mathcal{M} \not\models \varphi[a] \ \text{and} \ \mathcal{M} \models \neg \varphi[a], \\ \llbracket \varphi \rrbracket^{\mathcal{M},a} = \mathbf{n} & \quad \text{iff} \quad \mathcal{M} \not\models \varphi[a] \ \text{and} \ \mathcal{M} \not\models \neg \varphi[a], \\ \llbracket \varphi \rrbracket^{\mathcal{M},a} = \mathbf{b} & \quad \text{iff} \quad \mathcal{M} \models \varphi[a] \ \text{and} \ \mathcal{M} \models \neg \varphi[a]. \end{aligned}$$

Again we suppress superscripts where this may be done. It is easily verified that the connectives \mathbf{n} , \neg , $-$, and \wedge have a semantics as discussed in the previous section under this interpretation. The semantics of \forall is just what one would expect and bears the usual relation to that of \wedge . Note that

$$\llbracket \forall x \varphi \rrbracket^{\mathcal{M},a} = \bigwedge_{d \in \mathcal{D}} \llbracket \varphi \rrbracket^{\mathcal{M},a_d^x},$$

where \bigwedge denotes arbitrary meet in $(\mathbf{4}, \leq_t)$. This leaves it for us to motivate the semantics of \approx for which we need a short digression. One common way (see [4]) to motivate logics in which truth and nonfalsity are not the same concept is to point out the

existence of situations in which there is some form of distributed but fallible knowledge. Suppose we have a database which can be updated by more than one employee. Then it may occur that Tim enters that p , while Tom enters $\neg p$. If the reasoning system that comes with the database is based on classical logic this means that all future questions posed to the system will be answered with ‘yes’. A partial or paraconsistent logic can avoid this, for, as we shall see shortly, $p, \neg p \models q$ does not hold in such systems.

But this motivation does not preclude the possibility that the reasoning system decides for *some* sentences that they must take their values in $\{\mathbf{t}, \mathbf{f}\}$. For example, whatever information there is in the database system, it makes little sense for the computer to have doubts about statements it can decide itself, such as, say, $27 + 45 \approx 73$. The fact that *some* knowledge is distributed does not mean that *all* knowledge must be treated as such by the reasoning system.

For arbitrary formulas φ it is possible to state that the formula is true and not false by stating $\mathbf{t} \rightarrow \varphi$ (where \mathbf{t} abbreviates $\neg(\neg\mathbf{n} \wedge \mathbf{n})$ and \rightarrow is as below). We may imagine that an automated system which has expertise in a certain field simply asserts $\mathbf{t} \rightarrow \varphi$ for certain φ and overrules all employee attempts of entering potentially conflicting information. For identity statements it seems that bivalence is even the only possibility, provided that we wish to preserve two properties: (a) self-identity and (b) replacement of equals by equals. No respectable notion of identity can do without these. Suppose that some statement $t_1 \approx t_2$ could be both true and false. Then $\neg t_1 \approx t_2$ would also be both true and false. Given the definition of \models , self-identity, the property that $\models t_1 \approx t_1$, requires that $t_1 \approx t_1$ is true and not false and hence that $\neg t_1 \approx t_1$ is false and not true. Note that $t_1 \approx t_2, \neg t_1 \approx t_2 \models \neg t_1 \approx t_1$ is an instance of replacing equals by equals. But now we have a valid sequent with two premises which are both true (and false) but a conclusion which is not true. Contradiction. The assumption that $t_1 \approx t_2$ could be neither true nor false is dealt with in a similar way. In that case $t_1 \approx t_2, \neg t_1 \approx t_2 \models \neg t_1 \approx t_1$ has a false conclusion but no false premises. Again this is a contradiction. It follows that $t_1 \approx t_2$ must be bivalent and hence that the semantics as it is given is the only reasonable one.

We can introduce more connectives by means of abbreviation.

Definition 3.2 (Abbreviations) Write

$$\begin{aligned} \varphi \otimes \psi & \text{ for } (\varphi \wedge \psi) \vee ((\varphi \vee \psi) \wedge \mathbf{n}) \\ \varphi \oplus \psi & \text{ for } (\varphi \wedge \psi) \vee ((\varphi \vee \psi) \wedge \neg\mathbf{n}) \\ \varphi \rightarrow \psi & \text{ for } (\neg\varphi \vee \neg\psi) \wedge (\neg\neg\varphi \vee \psi). \end{aligned}$$

It is not difficult to check that \otimes and \oplus denote meet and join in the approximation lattice. The connective \rightarrow is related to \leq_t , for we have that

$$\begin{aligned} \llbracket \varphi \rightarrow \psi \rrbracket^{\mathcal{M},a} = \mathbf{t} & \text{ iff } \llbracket \varphi \rrbracket^{\mathcal{M},a} \leq_t \llbracket \psi \rrbracket^{\mathcal{M},a}, \\ \llbracket \varphi \rightarrow \psi \rrbracket^{\mathcal{M},a} = \mathbf{f} & \text{ iff } \llbracket \varphi \rrbracket^{\mathcal{M},a} \not\leq_t \llbracket \psi \rrbracket^{\mathcal{M},a}. \end{aligned}$$

Suitable definitions of \mathbf{f} , \mathbf{b} , \vee , and \exists are left to the reader.

Theorem 3.3 (Functional Completeness) *Every truth function is expressed by a formula.*

Proof: Directly from the functional completeness of $\{\otimes, \wedge, \neg, -\}$, shown in [24], and the definability of \otimes . \square

4 Consequence When we study the consequence relation \models it immediately becomes apparent that the usual rules for negation are no longer valid: $\Gamma, \neg\varphi \models \Delta$ does not follow from $\Gamma \models \varphi, \Delta$ (for example, we have $p \models p$, but not $p, \neg p \models$) and $\Gamma, \varphi \models \Delta$ does not entail $\Gamma \models \neg\varphi, \Delta$ (since $\not\models p, \neg p$). This means that such rules can no longer appear in a syntactic characterization of the consequence relation and that we must find something weaker. One solution is to give mixed rules for negation and other connectives as it is done, for example, in [9]. We can split the left rule for \wedge in two as follows.

$$\frac{\Gamma, \neg\varphi \vdash \Delta \quad \Gamma, \neg\psi \vdash \Delta}{\Gamma, \neg(\varphi \wedge \psi) \vdash \Delta} \qquad \frac{\Gamma, \varphi, \psi \vdash \Delta}{\Gamma, \varphi \wedge \psi \vdash \Delta}$$

We may also split the right rule:

$$\frac{\Gamma \vdash \varphi, \Delta \quad \Gamma \vdash \psi, \Delta}{\Gamma \vdash \varphi \wedge \psi, \Delta} \qquad \frac{\Gamma \vdash \neg\varphi, \neg\psi, \Delta}{\Gamma \vdash \neg(\varphi \wedge \psi), \Delta}$$

And other rules may be split in a similar way.

While it is possible to arrive at a sound and complete characterization of consequence in \mathbf{L}_4 in this manner, it may be thought less than nice that some of these rules are of a mixed character and combine two connectives. These rules do not conform to the so-called subformula property, as neither $\neg\varphi$ nor $\neg\psi$ is a subformula of $\neg(\varphi \wedge \psi)$. Notice, moreover, the similarity between the combined left rule for negation and conjunction and the right rule for conjunction without negation. A further similarity obtains between the left rule for unnegated conjunction and the combined right rule. We would do better if we could let such similar rules be instantiations of a single one.

In order to obtain such a more compact characterization we follow [22] in taking sequents to have *four* structural positions instead of the usual two and in letting these positions be arranged in a so-called quadrant. [22] also considers the various directions in which transmissions of truth and falsity may go and obtains sequents such as the following.

$$\frac{\Gamma \mid \Delta}{\Pi \mid \Sigma} \qquad \frac{\Gamma \mid \Delta}{\Pi \mid \Sigma} \qquad \frac{\Gamma \mid \Delta}{\Pi \mid \Sigma}$$

Here the two *north* positions correspond to the two positions in a normal Gentzen sequent $\Gamma \vdash \Delta$ and the two other positions are added for a convenient treatment of negation: having φ in a *southern* position will be equivalent to having $\neg\varphi$ in the corresponding *northern* position and vice versa. The idea of using sequents with multiple components dates back to [26] and [25], but Langholm's setup is different from these approaches, as will become apparent below. The 'biconsequence relations' of [6], on the other hand, are very similar to Langholm's quadrants.

We linearize notation by attaching two *signs* i and j to formulas. i can be n (*north*) or s (*south*), j can be e (*east*) or w (*west*). Instead of the rightmost sequent

displayed above we write

$$\{\varphi^{n,w} \mid \varphi \in \Gamma\} \cup \{\varphi^{n,e} \mid \varphi \in \Delta\} \cup \{\varphi^{s,w} \mid \varphi \in \Pi\} \cup \{\varphi^{s,e} \mid \varphi \in \Sigma\}.$$

While [22] considers the graphical representations shown above as different species of sequents, we let them be manifestations of a single variety. In order to distinguish the different kinds, we introduce the two *structural elements* \dashv and \dashv mentioned in the introduction. With the help of these we can define our basic data structure.

Definition 4.1 A *sequent* is a set of signed sentences and structural elements.

The usual notation for sequents will be employed. In particular, we write (as we did before) Γ, ϑ instead of $\Gamma \cup \{\vartheta\}$ whenever ϑ is a signed sentence or a structural element. We do not require sequents to be finite. The leftmost representation above corresponds to

$$\{\varphi^{n,w} \mid \varphi \in \Gamma\} \cup \{\varphi^{n,e} \mid \varphi \in \Delta\} \cup \{\varphi^{s,w} \mid \varphi \in \Pi\} \cup \{\varphi^{s,e} \mid \varphi \in \Sigma\} \cup \{\dashv\}.$$

The idea is that the direction from right to left in a sequent Γ is not considered if $\dashv \in \Gamma$ and that the direction from left to right is not considered if $\dashv \in \Gamma$. The situation that $\{\dashv, \dashv\} \subseteq \Gamma$ is a limiting case; Γ will then be an axiom.

We say that a signed sentence $\varphi^{i,j}$ is a *north* sentence if $i = n$, otherwise it is a *south* sentence. Similarly, $\varphi^{i,j}$ is a *west* sentence if $j = w$ and an *east* sentence if $j = e$. A model \mathcal{M} *accepts* a north sentence φ if $\mathcal{M} \models \varphi$; it accepts a south sentence φ if $\mathcal{M} \models \varphi$. \mathcal{M} *rejects* a north sentence φ if $\mathcal{M} \models \varphi$; it rejects a south sentence φ if $\mathcal{M} \models \varphi$.

Definition 4.2 \mathcal{M} *refutes* Γ if $\dashv \notin \Gamma$ and \mathcal{M} accepts all west sentences but no east sentence in Γ ; \mathcal{M} *refutes* Γ if $\dashv \notin \Gamma$ and \mathcal{M} rejects all east sentences but no west sentence in Γ ; and \mathcal{M} *refutes* Γ if it refutes Γ or refutes Γ . A sequent Γ is *valid* if no \mathcal{M} refutes Γ .

The notions of consequence considered in the introduction clearly are specializations of the notion of a valid sequent, as we have that

$$\begin{aligned} \Pi \models \Sigma & \text{ iff } \{\varphi^{n,w} \mid \varphi \in \Pi\} \cup \{\varphi^{n,e} \mid \varphi \in \Sigma\} \text{ is valid,} \\ \Pi \models^{tr} \Sigma & \text{ iff } \{\varphi^{n,w} \mid \varphi \in \Pi\} \cup \{\varphi^{n,e} \mid \varphi \in \Sigma\} \cup \{\dashv\} \text{ is valid,} \\ \Pi \models^{nf} \Sigma & \text{ iff } \{\varphi^{n,w} \mid \varphi \in \Pi\} \cup \{\varphi^{n,e} \mid \varphi \in \Sigma\} \cup \{\dashv\} \text{ is valid.} \end{aligned}$$

Remark 4.3 This place is as good as any to emphasize the fact that in general there is no unique way to associate quadrant positions with truth values in our system. It is true that for the notions *refute* Γ and *refute* Γ the following pictures emerge.

$$\begin{array}{c} \text{true} \mid \text{not true} \\ \hline \text{false} \mid \text{not false} \end{array} \quad \begin{array}{c} \text{not false} \mid \text{false} \\ \hline \text{not true} \mid \text{true} \end{array}$$

(See also the tableau system of D'Agostino [10] which is based on the values *true*, *false*, *nontrue*, and *nonfalse*.) But the general notion of refutation is a combination of *refute* Γ and *refute* Γ and there is no similar picture corresponding to it. For this

reason our main notion of logical consequence \models is not the kind of consequence relation that is studied in the tradition of many-valued logics, where an argument is valid whenever the conclusion gets a designated truth value if all premises get a designated truth value. The relations \models^{tr} and \models^{nf} , on the other hand, do fall within this realm. \models^{tr} is the relation we get when \mathbf{t} and \mathbf{b} are designated, while \models^{nf} is the relation we obtain when \mathbf{t} and \mathbf{n} are.

Definition 4.4 Let Γ be a sequent. We define the *dual* of Γ , $dual(\Gamma)$, to be the sequent which results from Γ by simultaneously replacing every superscript n in Γ by s , every s by n , every w by e , every e by w , \wedge by ∇ , and ∇ by \wedge .

Lemma 4.5 \mathcal{M} refutes $^{\rightarrow}$ Γ if and only if \mathcal{M} refutes $^{\leftarrow}$ $dual(\Gamma)$.

Proof: Immediate from the definitions. □

5 A sequent calculus We turn to the proof theory of our system and provide the notion of validity defined in the previous section with a corresponding notion of provability.

Definition 5.1 A sequent is *provable* if it follows in the usual way from the following sequent rules. (Here and elsewhere we shall let $-n = s$, $-s = n$, $-e = w$, $-w = e$.)

- | | | |
|-------------------------|------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|
| (R) | $\frac{}{\Gamma, \varphi^{i,w}, \varphi^{i,e}}$ | if φ is atomic |
| (∇ , \wedge) | $\frac{}{\Gamma, \nabla, \wedge}$ | |
| (\mathbf{n}^w) | $\frac{\Gamma, \nabla}{\Gamma, \mathbf{n}^{i,w}}$ | (\mathbf{n}^e) $\frac{\Gamma, \nabla}{\Gamma, \mathbf{n}^{i,e}}$ |
| (\neg) | $\frac{\Gamma, \varphi^{i,j}}{\Gamma, \neg \varphi^{-i,j}}$ | |
| ($-$) | $\frac{\Gamma, \varphi^{i,j}}{\Gamma, -\varphi^{-i,-j}}$ | |
| (\wedge_{sw}^{ne}) | $\frac{\Gamma, \varphi^{i,j} \quad \Gamma, \psi^{i,j}}{\Gamma, (\varphi \wedge \psi)^{i,j}}$ | where $\langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\}$ |
| (\wedge_{se}^{nw}) | $\frac{\Gamma, \varphi^{i,j}, \psi^{i,j}}{\Gamma, (\varphi \wedge \psi)^{i,j}}$ | where $\langle i, j \rangle \in \{\langle n, w \rangle, \langle s, e \rangle\}$ |
| (\forall_{sw}^{ne}) | $\frac{\Gamma, [c/x]\varphi^{i,j}}{\Gamma, \forall x \varphi^{i,j}}$ | where c is not in Γ or φ and $\langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\}$ |
| (\forall_{se}^{nw}) | $\frac{\Gamma, [t/x]\varphi^{i,j}}{\Gamma, \forall x \varphi^{i,j}}$ | where $\langle i, j \rangle \in \{\langle n, w \rangle, \langle s, e \rangle\}$ |
| (id) | $\frac{}{\Gamma, t \approx t^{i,j}}$ | where $\langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\}$ |
| (L) | $\frac{\Gamma, [t_2/x]\varphi^{i',j'}}{\Gamma, t_1 \approx t_2^{i,j}, [t_1/x]\varphi^{i',j'}}$ | where $\langle i, j \rangle \in \{\langle n, w \rangle, \langle s, e \rangle\}$, $i' \in \{n, s\}$, $j' \in \{e, w\}$ and φ is atomic. |

It is clear that this cut-free calculus obeys the modularity constraint that only one logical operator is dealt with in each rule. We also have a version of the *subformula property* as it is not difficult to show that, for each sentence φ occurring somewhere in a proof Π , φ is a substitution instance of some ψ occurring as a subformula of a sentence in the last sequent of Π .

Just as in the case of validity we can specialize our notion of provability and, letting Π and Σ vary over sets of sentences, write

$$\begin{aligned} \Pi \vdash \Sigma & \text{ iff } \{ \varphi^{n,w} \mid \varphi \in \Pi \} \cup \{ \varphi^{n,e} \mid \varphi \in \Sigma \} \text{ is provable,} \\ \Pi \vdash^{tr} \Sigma & \text{ iff } \{ \varphi^{n,w} \mid \varphi \in \Pi \} \cup \{ \varphi^{n,e} \mid \varphi \in \Sigma \} \cup \{ \not\vdash \} \text{ is provable,} \\ \Pi \vdash^{nf} \Sigma & \text{ iff } \{ \varphi^{n,w} \mid \varphi \in \Pi \} \cup \{ \varphi^{n,e} \mid \varphi \in \Sigma \} \cup \{ \not\vdash \} \text{ is provable.} \end{aligned}$$

Example 5.2 The following proof shows that $\neg(\varphi \wedge \psi) \vdash \neg\varphi \wedge \neg\psi$.

$$\frac{\frac{\frac{\varphi^{s,e}, \psi^{s,e}, \varphi^{s,w}}{\varphi^{s,e}, \psi^{s,e}, -\varphi^{n,e}} (R) \quad \frac{\varphi^{s,e}, \psi^{s,e}, \psi^{s,w}}{\varphi^{s,e}, \psi^{s,e}, -\psi^{n,e}} (R)}{\varphi^{s,e}, \psi^{s,e}, -\varphi \wedge -\psi^{n,e}} (-) \quad \frac{\varphi^{s,e}, \psi^{s,e}, -\psi^{n,e}}{\varphi^{s,e}, \psi^{s,e}, -\psi^{n,e}} (\wedge_{sw}^{ne})}{\varphi^{s,e}, \psi^{s,e}, -\varphi \wedge -\psi^{n,e}} (\wedge_{se}^{nw})}{\varphi \wedge \psi^{s,e}, -\varphi \wedge -\psi^{n,e}} (\wedge_{se}^{nw})}{\neg(\varphi \wedge \psi)^{n,w}, -\varphi \wedge -\psi^{n,e}} (-)$$

Remark 5.3 Consider a calculus with rules as above except that $(\not\vdash, \not\vdash)$, (\mathbf{n}^w) , and (\mathbf{n}^e) are replaced by the single

$$\overline{\Gamma, \mathbf{n}^{i,w}}.$$

Call a sequent *tr-provable* if it follows from this calculus. It is not difficult to show that, if Γ does not contain $\not\vdash$, Γ is tr-provable if and only if $\Gamma, \not\vdash$ is provable. This gives an alternative characterization of \vdash^{tr} . An alternative characterization of \vdash^{nf} is obtained by proceeding dually. For Γ not containing structural elements, we have that Γ is provable if and only if Γ is provable in the system resulting from the present one with $(\not\vdash, \not\vdash)$, (\mathbf{n}^w) , and (\mathbf{n}^e) replaced by

$$\overline{\Gamma, \mathbf{n}^{i,w}, \mathbf{n}^{i',e}}.$$

We conclude that the structural elements are not strictly necessary for the setup. But see Remark 5.4 below.

It is easy to check that the following are derived rules of our calculus.

$$\begin{aligned} (\mathbf{f}_{se}^{nw}) & \quad \overline{\Gamma, \mathbf{f}^{i,j}}, & \text{ where } \langle i, j \rangle \in \{ \langle n, w \rangle, \langle s, e \rangle \} \\ (\vee_{se}^{nw}) & \quad \frac{\Gamma, \varphi^{i,j} \quad \Gamma, \psi^{i,j}}{\Gamma, (\varphi \vee \psi)^{i,j}}, & \text{ where } \langle i, j \rangle \in \{ \langle n, w \rangle, \langle s, e \rangle \} \\ (\vee_{sw}^{ne}) & \quad \frac{\Gamma, \varphi^{i,j}, \psi^{i,j}}{\Gamma, (\varphi \vee \psi)^{i,j}}, & \text{ where } \langle i, j \rangle \in \{ \langle n, e \rangle, \langle s, w \rangle \} \\ (\rightarrow_{sw}^{ne}) & \quad \frac{\Gamma, \varphi^{n,w}, \psi^{n,e} \quad \Gamma, \varphi^{s,e}, \psi^{s,w}}{\Gamma, (\varphi \rightarrow \psi)^{i,j}}, & \text{ where } \langle i, j \rangle \in \{ \langle n, e \rangle, \langle s, w \rangle \} \end{aligned}$$

$$\begin{array}{l}
(\rightarrow_{se}^{nw}) \quad \frac{\Gamma, \varphi^{n,e}, \psi^{s,e} \quad \Gamma, \varphi^{n,e}, \varphi^{s,w} \quad \Gamma, \psi^{n,w}, \psi^{s,e} \quad \Gamma, \psi^{n,w}, \varphi^{s,w}}{\Gamma, (\varphi \rightarrow \psi)^{i,j}}, \\
\text{where } \langle i, j \rangle \in \{\langle n, w \rangle, \langle s, e \rangle\} \\
(\exists_{se}^{nw}) \quad \frac{\Gamma, [c/x]\varphi^{i,j}}{\Gamma, \exists x\varphi^{i,j}}, \quad \text{where } c \text{ is not in } \Gamma \text{ or } \varphi \text{ and } \langle i, j \rangle \in \\
\{\langle n, w \rangle, \langle s, e \rangle\} \\
(\exists_{sw}^{ne}) \quad \frac{\Gamma, [t/x]\varphi^{i,j}}{\Gamma, \exists x\varphi^{i,j}}, \quad \text{where } \langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\}
\end{array}$$

An inspection of the rules shows that if we restrict ourselves to sentences in which **n** does not occur our various notions of provability collapse, that is, for Π and Σ in which no signed sentence contains **n** we have that $\Pi \vdash \Sigma$ if and only if $\Pi \vdash^{tr} \Sigma$ if and only if $\Pi \vdash^{nf} \Sigma$, since no application of a rule can create $\not\vdash$ or $\not\vdash$. But as soon as **n** enters the picture it is important to keep track of the extra structural elements. Here are derived rules for **b**, \otimes , and \oplus in which $\not\vdash$ and $\not\vdash$ play an important role.

$$\begin{array}{l}
(\mathbf{b}^e) \quad \frac{\Gamma, \not\vdash}{\Gamma, \mathbf{b}^{i,e}} \quad (\mathbf{b}^w) \quad \frac{\Gamma, \not\vdash}{\Gamma, \mathbf{b}^{i,w}} \\
(\otimes^e) \quad \frac{\Gamma, \varphi^{i,e}, \psi^{i,e} \quad \Gamma, \varphi^{i,e}, \not\vdash \quad \Gamma, \psi^{i,e}, \not\vdash}{\Gamma, (\varphi \otimes \psi)^{i,e}}, \quad \text{where } i \in \{n, s\} \\
(\otimes^w) \quad \frac{\Gamma, \varphi^{i,w}, \psi^{i,w} \quad \Gamma, \varphi^{i,w}, \not\vdash \quad \Gamma, \psi^{i,w}, \not\vdash}{\Gamma, (\varphi \otimes \psi)^{i,w}}, \quad \text{where } i \in \{n, s\} \\
(\oplus^e) \quad \frac{\Gamma, \varphi^{i,e}, \psi^{i,e} \quad \Gamma, \varphi^{i,e}, \not\vdash \quad \Gamma, \psi^{i,e}, \not\vdash}{\Gamma, (\varphi \oplus \psi)^{i,e}}, \quad \text{where } i \in \{n, s\} \\
(\oplus^w) \quad \frac{\Gamma, \varphi^{i,w}, \psi^{i,w} \quad \Gamma, \varphi^{i,w}, \not\vdash \quad \Gamma, \psi^{i,w}, \not\vdash}{\Gamma, (\varphi \oplus \psi)^{i,e}}, \quad \text{where } i \in \{n, s\}
\end{array}$$

Remark 5.4 Note that the use of our structural elements here makes it possible to formulate these rules without any violation of the subformula property.

Remark 5.5 Arieli and Avron [1] offer the following sequent rules for \otimes and \oplus .

$$\begin{array}{ll}
[\otimes \Rightarrow] \quad \frac{\Gamma, \varphi, \psi \Rightarrow \Delta}{\Gamma, \varphi \otimes \psi \Rightarrow \Delta} & [\Rightarrow \otimes] \quad \frac{\Gamma \Rightarrow \Delta, \varphi \quad \Gamma \Rightarrow \Delta, \psi}{\Gamma \Rightarrow \Delta, \varphi \otimes \psi} \\
[\neg \otimes \Rightarrow] \quad \frac{\Gamma, \neg \varphi, \neg \psi \Rightarrow \Delta}{\Gamma, \neg(\varphi \otimes \psi) \Rightarrow \Delta} & [\Rightarrow \neg \otimes] \quad \frac{\Gamma \Rightarrow \Delta, \neg \varphi \quad \Gamma \Rightarrow \Delta, \neg \psi}{\Gamma \Rightarrow \Delta, \neg(\varphi \otimes \psi)} \\
[\oplus \Rightarrow] \quad \frac{\Gamma, \varphi \Rightarrow \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \varphi \oplus \psi \Rightarrow \Delta} & [\Rightarrow \oplus] \quad \frac{\Gamma \Rightarrow \Delta, \varphi, \psi}{\Gamma \Rightarrow \Delta, \varphi \oplus \psi} \\
[\neg \oplus \Rightarrow] \quad \frac{\Gamma, \neg \varphi \Rightarrow \Delta \quad \Gamma, \neg \psi \Rightarrow \Delta}{\Gamma, \neg(\varphi \oplus \psi) \Rightarrow \Delta} & [\Rightarrow \neg \oplus] \quad \frac{\Gamma \Rightarrow \Delta, \neg \varphi, \neg \psi}{\Gamma \Rightarrow \Delta, \neg(\varphi \oplus \psi)}
\end{array}$$

It is not difficult to show that these rules are derivable in our calculus plus an additional thinning rule, provided we interpret \Rightarrow as \vdash^{tr} . In view of the soundness result below, this also means that they are sound with respect to \models^{tr} . However, soundness with respect to \models does not obtain. Consider $[\otimes \Rightarrow]$, for example. Since $\mathbf{f}, \mathbf{t} \Rightarrow \mathbf{f}$, it follows from this rule that $\mathbf{f} \otimes \mathbf{t} \Rightarrow \mathbf{f}$ is derivable. But clearly, $\mathbf{f} \otimes \mathbf{t} \not\models \mathbf{f}$, as $\mathbf{f} \otimes \mathbf{t}$

evaluates as **n**. We now prove the soundness of our calculus after stating a useful lemma.

Lemma 5.6 $\Gamma_1, \dots, \Gamma_n / \Gamma$ is a sequent rule if and only if $dual(\Gamma_1), \dots, dual(\Gamma_n) / dual(\Gamma)$ is a sequent rule.

Proof: By a simple inspection of the sequent rules. \square

Theorem 5.7 (Soundness) *If a sequent is provable then it is valid.*

Proof: It can be shown for any sequent rule that some condition of the rule is refutable $^{\rightarrow}$ if the conclusion of the rule is refutable $^{\rightarrow}$. We prove this statement for (\forall_{sw}^{ne}) , leaving the other cases to the reader. Suppose that $\mathcal{M} = \langle \mathcal{D}, I \rangle$ and that \mathcal{M} refutes $^{\rightarrow} \Gamma, \forall x \varphi^{n,e}$. Then $\mathcal{M} \not\models \forall x \varphi[a]$ for some a , so that there is some $d \in \mathcal{D}$ such that $\mathcal{M} \not\models \varphi[a_d^x]$. Let I' be the interpretation function which is just like I with the possible exception that $I'(c) = d$ and let $\mathcal{M}' = \langle \mathcal{D}, I' \rangle$. Then $\mathcal{M}' \not\models [c/x]\varphi$ and, since c does not occur in $\Gamma \cup \{\varphi\}$, \mathcal{M}' refutes $^{\rightarrow} \Gamma, [c/x]\varphi^{n,e}$. For the case that $\langle i, j \rangle = \langle s, w \rangle$, repeat this argument, but uniformly replace $\not\models$ with \models .

Other cases are proved in a similar vein and this settles that, for any sequent rule, some condition of the rule is refutable $^{\rightarrow}$ if the conclusion is refutable $^{\rightarrow}$. For the dual case, suppose that the conclusion Γ of a rule $\Gamma_1, \dots, \Gamma_n / \Gamma$ is refutable $^{\leftarrow}$. Then, by Lemma 4.5, $dual(\Gamma)$ is refutable $^{\rightarrow}$. By Lemma 5.6, $dual(\Gamma_1), \dots, dual(\Gamma_n) / dual(\Gamma)$ is a sequent rule, whence, by the previous reasoning, $dual(\Gamma_k)$ is refutable $^{\rightarrow}$ for some k . A second application of Lemma 4.5 gives that Γ_k is refutable $^{\leftarrow}$.

We conclude that the conclusion of any rule is valid if all its conditions are valid and the theorem follows by an induction on the complexity of proofs. \square

6 Elementary model theory The purpose of this section is to prove a model existence theorem for our logic. From this some useful corollaries in the form of a compactness theorem, a Löwenheim-Skolem theorem and a Completeness theorem will follow.

Definition 6.1 A sequent Γ is called a *Hintikka sequent* if and only if

1. $\{\varphi^{i,w}, \varphi^{i,e}\} \not\subseteq \Gamma$, if φ is atomic;
2. $\{\not\wedge, \not\vee\} \not\subseteq \Gamma$;
3. $\mathbf{n}^{i,w} \in \Gamma \implies \not\wedge \in \Gamma$,
 $\mathbf{n}^{i,e} \in \Gamma \implies \not\vee \in \Gamma$;
4. $\neg\varphi^{i,j} \in \Gamma \implies \varphi^{-i,j} \in \Gamma$;
5. $-\varphi^{i,j} \in \Gamma \implies \varphi^{-i,-j} \in \Gamma$;
6. $\varphi \wedge \psi^{i,j} \in \Gamma \implies \varphi^{i,j} \in \Gamma$ or $\psi^{i,j} \in \Gamma$, if $\langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\}$,
 $\varphi \wedge \psi^{i,j} \in \Gamma \implies \{\varphi^{i,j}, \psi^{i,j}\} \subseteq \Gamma$, if $\langle i, j \rangle \in \{\langle n, w \rangle, \langle s, e \rangle\}$;
7. $\forall x \varphi^{i,j} \in \Gamma \implies [t/x]\varphi^{i,j} \in \Gamma$, for all closed terms t , if $\langle i, j \rangle \in \{\langle n, w \rangle, \langle s, e \rangle\}$,

- $\forall x\varphi^{i,j} \in \Gamma \implies [c/x]\varphi^{i,j} \in \Gamma$, for some constant c , if $\langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\}$;
8. $t \approx t^{i,j} \notin \Gamma$, if $\langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\}$,
 $\{t_1 \approx t_2^{i,j}, [t_1/x]\varphi^{i',j'}\} \subseteq \Gamma \implies [t_2/x]\varphi^{i',j'} \in \Gamma$, if $\langle i, j \rangle \in \{\langle n, w \rangle, \langle s, e \rangle\}$, $i' \in \{n, s\}$, $j' \in \{e, w\}$ and φ is atomic.

Lemma 6.2 *If Γ is a Hintikka sequent then $\text{dual}(\Gamma)$ is a Hintikka sequent.*

Proof: By inspection. □

Lemma 6.3 (Hintikka Lemma) *Each Hintikka sequent is refutable by a countable model.*

Proof: Let Γ be a Hintikka sequent. We first consider the case that $\not\vdash \notin \Gamma$ and construct a model \mathcal{M} which accepts all west sentences but no east sentence in Γ . Define the relation \sim between closed terms by setting

$$t_1 \sim t_2 \iff (t_1 \approx t_2^{n,w} \in \Gamma \text{ or } t_1 \approx t_2^{s,e} \in \Gamma).$$

It is easily verified that \sim is an equivalence relation. For each term t , let \tilde{t} be the equivalence class $\{t' \mid t' \sim t\}$ and let \mathcal{D} be the set $\{\tilde{t} \mid t \text{ is a closed term}\}$. Define, for each n -ary function symbol $f \in \mathcal{L}$ and each n -ary relation symbol $R \in \mathcal{L}$:

$$I(f)(\tilde{t}_1, \dots, \tilde{t}_n) = \widetilde{ft_1 \dots t_n},$$

$$I(R) = \{\langle \tilde{t}_1, \dots, \tilde{t}_n \rangle \mid Rt_1, \dots, t_n^{n,w} \in \Gamma\}, \{\langle \tilde{t}_1, \dots, \tilde{t}_n \rangle \mid Rt_1, \dots, t_n^{s,w} \in \Gamma\}.$$

The last clause of Definition 6.1 ensures that this definition does not depend on the choice of t_1, \dots, t_n . Now let $\mathcal{M} = \langle \mathcal{D}, I \rangle$. Clearly, \mathcal{M} is a countable model. An induction on term complexity shows that, for each t , $\llbracket t \rrbracket^{\mathcal{M}} = \tilde{t}$. Another induction on the number of connectives occurring in a sentence establishes that, for each φ ,

- A. $\varphi^{n,e} \in \Gamma \implies \mathcal{M} \not\models \varphi$
- B. $\varphi^{s,w} \in \Gamma \implies \mathcal{M} \models \varphi$
- C. $\varphi^{n,w} \in \Gamma \implies \mathcal{M} \models \varphi$
- D. $\varphi^{s,e} \in \Gamma \implies \mathcal{M} \not\models \varphi$

We work out the \forall case of the induction.

- A. Assume that $\forall x\varphi^{n,e} \in \Gamma$. Then, by the definition of a Hintikka sequent, $[c/x]\varphi^{n,e} \in \Gamma$ for some constant c . By the induction hypothesis, $\mathcal{M} \not\models [c/x]\varphi$, so that $\mathcal{M} \not\models \varphi[a_c^x]$, where a is arbitrary, follows by the usual substitution lemma. From this we have that $\mathcal{M} \not\models \forall x\varphi$.
- B. $\forall x\varphi^{s,w} \in \Gamma \implies [c/x]\varphi^{s,w} \in \Gamma$ for some $c \implies \mathcal{M} \models [c/x]\varphi \implies \mathcal{M} \models \varphi[a_c^x] \implies \mathcal{M} \models \forall x\varphi$.
- C. $\forall x\varphi^{n,w} \in \Gamma \implies [t/x]\varphi^{n,w} \in \Gamma$ for all closed terms $t \implies \mathcal{M} \models [t/x]\varphi$ for all $t \implies \mathcal{M} \models \varphi[a_t^x]$ for all $t \implies \mathcal{M} \models \varphi[a_d^x]$ for all $d \in \mathcal{D} \implies \mathcal{M} \models \forall x\varphi$.
- D. $\forall x\varphi^{s,e} \in \Gamma \implies [t/x]\varphi^{s,e} \in \Gamma$ for all closed $t \implies \mathcal{M} \not\models [t/x]\varphi$ for all $t \implies \mathcal{M} \not\models \varphi[a_t^x]$ for all $t \implies \mathcal{M} \not\models \varphi[a_d^x]$ for all $d \in \mathcal{D} \implies \mathcal{M} \not\models \forall x\varphi$.

The other cases of this induction are similar and are left to the reader. It follows that \mathcal{M} refutes $\neg \Gamma$.

Now consider the case that $\not\vdash \notin \Gamma$. Since $dual(\Gamma)$ is a Hintikka sequent by Lemma 6.2 and since $\not\vdash \notin dual(\Gamma)$ we have that there is a countable \mathcal{M} which refutes $\neg dual(\Gamma)$ and hence, by Lemma 4.5, refutes $\neg \Gamma$. Since $\not\vdash$ and $\not\vdash$ cannot both be elements of Γ , we have established the theorem. \square

Definition 6.4 Let \mathcal{P} be a set of sequents in the language \mathcal{L} . \mathcal{P} is a *provability property* with respect to \mathcal{L} if and only if

1. If $\{\Gamma_1, \dots, \Gamma_n\} \subseteq \mathcal{P}$ and $\Gamma_1, \dots, \Gamma_n / \Gamma$ is a sequent rule, then $\Gamma \in \mathcal{P}$;
2. If $\Gamma \in \mathcal{P}$ and $\Gamma \subseteq \Gamma'$, then $\Gamma' \in \mathcal{P}$, for each $\Gamma' \in \mathcal{L}$.

Theorem 6.5 (Model Existence) *Let \mathcal{L} be a language and let C be a countably infinite set of constants such that $\mathcal{L} \cap C = \emptyset$. Assume that \mathcal{P} is a provability property with respect to $\mathcal{L} \cup C$ and that Γ is a sequent in the language \mathcal{L} . If $\Gamma \notin \mathcal{P}$ then Γ is refutable by a countable model.*

Proof: Let \mathcal{P} and Γ be as described. We construct a Hintikka sequent Γ^* such that $\Gamma \subseteq \Gamma^*$. Let $\vartheta_1, \dots, \vartheta_n, \dots$ be an enumeration of all signed sentences in $\mathcal{L} \cup C$ plus the structural elements. Write $\iota(\vartheta)$ for the index that the signed sentence or structural element ϑ obtains in this enumeration. Define

$$\Gamma_0 = \Gamma$$

$$\Gamma_{n+1} = \begin{cases} \Gamma_n, & \text{if } \Gamma_n \cup \{\vartheta_n\} \in \mathcal{P} \\ \Gamma_n \cup \{\vartheta_n\}, & \text{if } \Gamma_n \cup \{\vartheta_n\} \notin \mathcal{P} \text{ and } \vartheta_n \text{ is not of the} \\ & \text{form } \forall x \varphi^{n,e} \text{ or } \forall x \varphi^{s,w} \\ \Gamma_n \cup \{\vartheta_n, [c/x]\varphi^{i,j}\}, & \text{if } \Gamma_n \cup \{\vartheta_n\} \notin \mathcal{P}, \langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\} \\ & \text{and } \vartheta_n \text{ is of the form } \forall x \varphi^{i,j}, \text{ where } c \text{ is} \\ & \text{the first constant in } C \text{ which does not} \\ & \text{occur in } \Gamma_n \cup \{\vartheta_n\} \end{cases}$$

This is well defined since each Γ_n contains only a finite number of constants from C . That $\Gamma_n \notin \mathcal{P}$ for each n follows by a simple induction which uses the definition of a provability property and the fact that (\forall_{sw}^{ne}) is a sequent rule. Define $\Gamma^* = \bigcup_n \Gamma_n$. We prove that, for all finite sets $\{\vartheta_{k_1}, \dots, \vartheta_{k_n}\}$ and for all $k \geq \max\{k_1, \dots, k_n\}$,

$$\{\vartheta_{k_1}, \dots, \vartheta_{k_n}\} \subseteq \Gamma^* \iff \Gamma_k \cup \{\vartheta_{k_1}, \dots, \vartheta_{k_n}\} \notin \mathcal{P}. \quad (1)$$

In order to show that this holds, let $k \geq \max\{k_1, \dots, k_n\}$ and suppose that $\{\vartheta_{k_1}, \dots, \vartheta_{k_n}\} \subseteq \Gamma^*$. Then there is some ℓ such that $\{\vartheta_{k_1}, \dots, \vartheta_{k_n}\} \subseteq \Gamma_\ell$. Let $m = \max\{k, \ell\}$. We have that $\Gamma_k \cup \{\vartheta_{k_1}, \dots, \vartheta_{k_n}\} \subseteq \Gamma_m$. Since $\Gamma_m \notin \mathcal{P}$ and \mathcal{P} is closed under supersets it follows that $\Gamma_k \cup \{\vartheta_{k_1}, \dots, \vartheta_{k_n}\} \notin \mathcal{P}$. For the reverse direction, suppose that $\Gamma_k \cup \{\vartheta_{k_1}, \dots, \vartheta_{k_n}\} \notin \mathcal{P}$. Then, since \mathcal{P} is closed under supersets, $\Gamma_{k_i} \cup \{\vartheta_{k_i}\} \notin \mathcal{P}$, for each of the k_i . By the construction of Γ^* each $\vartheta_{k_i} \in \Gamma^*$ and $\{\vartheta_{k_1}, \dots, \vartheta_{k_n}\} \subseteq \Gamma^*$.

With the help of (1) we verify that Γ^* is a Hintikka sequent. Here we check only a few conditions of Definition 6.1.

1. In order to check the first part of condition 6 of Definition 6.1, let $\langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\}$ and suppose $\varphi \wedge \psi^{i,j} \in \Gamma^*$. Let k be the maximum of $\iota(\varphi \wedge \psi^{i,j})$, $\iota(\varphi^{i,j})$, and $\iota(\psi^{i,j})$. (1) entails that $\Gamma_k \cup \{\varphi \wedge \psi^{i,j}\} \notin \mathcal{P}$. Since \mathcal{P} is closed under sequent rules, it follows with (\wedge_{sw}^{ne}) that $\Gamma_k \cup \{\varphi^{i,j}\} \notin \mathcal{P}$ or $\Gamma_k \cup \{\psi^{i,j}\} \notin \mathcal{P}$. By (1) this implies that $\varphi^{i,j} \in \Gamma^*$ or $\psi^{i,j} \in \Gamma^*$.
2. We verify that the seventh condition of Definition 6.1 holds for Γ^* . First suppose that $\forall x \varphi^{i,j} \in \Gamma^*$, that $\langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\}$, and that t is an arbitrary closed term. Let $k = \max\{\iota(\forall x \varphi^{i,j}), \iota([t/x]\varphi^{i,j})\}$. (1) gives that $\Gamma_k \cup \{\forall x \varphi^{i,j}\} \notin \mathcal{P}$ and by the closure of \mathcal{P} under sequent rules we find that $\Gamma_k \cup \{[t/x]\varphi^{i,j}\} \notin \mathcal{P}$. This in its turn, by (1), has as a consequence that $[t/x]\varphi^{i,j} \in \Gamma^*$. The construction of Γ^* ensures that $[c/x]\varphi^{i,j} \in \Gamma^*$ for some c if $\forall x \varphi^{n,e} \in \Gamma^*$ if $\langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\}$, so that the second part of clause 7 of Definition 6.1 is satisfied.

Checking the other conditions in the definition of a Hintikka sequent gives rise to considerations that are very similar to those already encountered and is left to the reader. Since $\Gamma \subseteq \Gamma^*$ and Γ^* is refutable by a countable model, Γ is refutable by that model. \square

In the following corollaries Γ will always be a sequent in some language \mathcal{L} while Δ ranges over sequents in $\mathcal{L} \cup C$ where \mathcal{L} and C are as in the formulation of Theorem 6.5.

Corollary 6.6 (Compactness) *If Γ is valid then there is some finite $\Gamma_0 \subseteq \Gamma$ which is valid.*

Proof: The set $\{\Delta \mid \text{some finite } \Delta_0 \subseteq \Delta \text{ is valid}\}$ is easily seen to be a provability property. It follows by Theorem 6.5 that Γ is refutable if no finite $\Gamma_0 \subseteq \Gamma$ is valid. By contraposition we find that some finite $\Gamma_0 \subseteq \Gamma$ must be valid if Γ is valid. \square

Corollary 6.7 (Löwenheim–Skolem) *If Γ is not valid then Γ is refutable by a countable model.*

Proof: $\{\Delta \mid \Delta \text{ is valid}\}$ is a provability property. \square

Corollary 6.8 (Completeness) *If Γ is valid then Γ is provable.*

Proof: The set $\{\Delta \mid \Delta \text{ is provable}\}$ is a provability property. It follows that Γ is refutable if Γ is not provable. \square

7 Three values We have obtained our results for a logic that was both partial and paraconsistent. What if we do not want to allow paraconsistency or do not want partiality?

The solution is simple. For a logic without paraconsistency we must remove— from our syntax and in order to make up for this loss we must add **f** and \rightarrow as primitive connectives. Next we additionally require in the definition of a *model* that, for any relation symbol R , $I^+(R) \cap I^-(R) = \emptyset$ (*no gluts*). This removes the possibility of paraconsistency. The truth definition should give the new primitives the semantics they previously obtained by expansion of their definitions:

1. $\mathcal{M} \not\models \mathbf{f}$
 $\mathcal{M} \models \mathbf{f}$
2. $\mathcal{M} \models \varphi \rightarrow \psi[a] \iff (\mathcal{M} \models \varphi[a] \implies \mathcal{M} \models \psi[a]) \ \& \ (\mathcal{M} \models \psi[a] \implies \mathcal{M} \models \varphi[a])$
 $\mathcal{M} \models \varphi \rightarrow \psi[a] \iff (\mathcal{M} \models \varphi[a] \ \& \ \mathcal{M} \not\models \psi[a]) \ \text{or} \ (\mathcal{M} \models \psi[a] \ \& \ \mathcal{M} \not\models \varphi[a])$

That the set $\{\mathbf{f}, \mathbf{n}, \neg, \wedge, \rightarrow\}$ is functionally complete for the new setup is easily seen to hold on the basis of a minor variant of the proof of Theorem 3.3. Apart from (\mathbf{f}_{sw}^{ne}) , (\rightarrow_{sw}^{ne}) and (\rightarrow_{se}^{nw}) two extra rules are added to the sequent calculus to counterbalance our restriction of the class of models (the rule for $-$ will be omitted, of course).

$$(3^w) \quad \frac{\Gamma, \not\wedge}{\Gamma, \varphi^{n,w}, \varphi^{s,w}} \quad (3^e) \quad \frac{\Gamma, \not\wedge}{\Gamma, \varphi^{n,e}, \varphi^{s,e}}$$

The five new rules bring five new conditions in Definition 6.1 with them.

1. $\mathbf{f}^{i,j} \notin \Gamma$, if $\langle i, j \rangle \in \{\langle n, w \rangle, \langle s, e \rangle\}$
2. $\{\varphi^{n,w}, \varphi^{s,w}\} \subseteq \Gamma \implies \not\wedge \in \Gamma$
3. $\{\varphi^{n,e}, \varphi^{s,e}\} \subseteq \Gamma \implies \not\wedge \in \Gamma$
4. $\varphi \rightarrow \psi^{i,j} \in \Gamma \implies \{\varphi^{n,w}, \psi^{n,e}\} \subseteq \Gamma$ or $\{\varphi^{s,e}, \psi^{s,w}\} \subseteq \Gamma$, if $\langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\}$
5. $\varphi \rightarrow \psi^{i,j} \in \Gamma \implies \{\varphi^{n,e}, \psi^{s,e}\} \subseteq \Gamma$ or $\{\varphi^{n,e}, \varphi^{s,w}\} \subseteq \Gamma$ or $\{\psi^{n,w}, \psi^{s,e}\} \subseteq \Gamma$ or $\{\psi^{n,w}, \varphi^{s,w}\} \subseteq \Gamma$, if $\langle i, j \rangle \in \{\langle n, w \rangle, \langle s, e \rangle\}$

It must then be checked in the proof of the Hintikka Lemma that the model which is constructed satisfies our new requirement that $I^+(R) \cap I^-(R) = \emptyset$ for each R . But, in fact, this is trivial from the construction. The reader will have no difficulty in seeing that, under these new conditions, all our previous proofs will go through.

In order to get logics that are not partial (but may be paraconsistent), we proceed dually. First, we require \mathbf{n} and $-$ not to be in the language but introduce \mathbf{b}, \mathbf{f} , and \rightarrow . We put the additional constraint on models that $\mathcal{D}^n \subseteq I^+(R) \cup I^-(R)$ for all n -ary relation symbols R (*no gaps*). The extra rules which need to be added are (\mathbf{b}^e) , (\mathbf{b}^w) , (\mathbf{f}_{sw}^{ne}) , (\rightarrow_{sw}^{ne}) , and (\rightarrow_{se}^{nw}) plus the following:

$$(3'^w) \quad \frac{\Gamma, \not\wedge}{\Gamma, \varphi^{n,w}, \varphi^{s,w}} \quad (3'^e) \quad \frac{\Gamma, \not\wedge}{\Gamma, \varphi^{n,e}, \varphi^{s,e}} .$$

Again we must add conditions corresponding to these rules to Definition 6.1 and we must check that this causes the new requirement on models to be satisfied by the model constructed in the Hintikka Lemma.

This shows that we can easily trade our four-valued logics for three-valued ones. The choice of which notion of logical consequence should be used, \models , \models^{tr} , or \models^{nf} , is independent from the choice which truth-values should be accepted.

To obtain classical logic add the *no gluts* and the *no gaps* requirements on models, remove \mathbf{n} , and $-$, but add \mathbf{f} and \rightarrow . A sequent calculus is obtained by adopting extra rules (\mathbf{f}_{sw}^{ne}) , (\rightarrow_{sw}^{ne}) , (\rightarrow_{se}^{nw}) , (3^w) , (3^e) , $(3'^w)$, and $(3'^e)$.

8 Interpolation in L_4 The purpose of this section is to prove a Craig-Lyndon interpolation theorem for L_4 . The theorem will be restricted to languages without function symbols other than individual constants, as a consideration of complex terms leads to certain complications orthogonal to the main concerns of this paper. The Lyndon part of the theorem can be formulated in a way that is slightly subtler than is possible in classical logic. Due to the fact that we have *two* negations (\neg and $-$) in our setup, there will also be two notions of positive and negative occurrences of formulas. It will turn out that the theorem holds for both these notions. The proof of the theorem was inspired by the proof in Kleene [20] for the classical case, but also bears resemblance to the Maehara method discussed in Takeuti [27]. Proofs of interpolation for \vdash^r and \vdash^f have been known before (see [21]), but since it is unclear how interpolation for \vdash could be obtained from these, we give a direct proof here.

We *color* signed sentences in proofs in order to be able to keep track of them. A colored signed sentence is a signed sentence which carries an additional subscript r (red) or b (blue) and a coloring of a sequent proof Π is obtained by coloring the signed sentences in the sequents of Π in such a way that a signed sentence shares its color with its ancestors in the proof (because of our representation of sequents as sets this may require that some signed sentences $\varphi^{i,j}$ now get two representations, $\varphi_r^{i,j}$ and $\varphi_b^{i,j}$). It is clear that, given any initial coloring of the end sequent of any proof Π , Π itself can be colored.

If Γ is a sequent in which the signed sentences are colored, we write Γ^b (Γ^r) for the sequent which results from Γ by removing all signed sentences colored red (blue) in Γ (note that this leaves the structural elements in place) and removing all subscripts from the remaining signed sentences.

A relation symbol R *occurs \neg -positively* (*\neg -negatively*) in a sentence φ if it occurs within the scope of an even (odd) number of negation symbols \neg . Similarly R *occurs $-$ -positively* (*$-$ -negatively*) in φ if it occurs within the scope of an even (odd) number of $-$ s. These notions of positive and negative occurrence within an unsigned sentence can be extended to signed colored sentences by counting certain combinations of signs and colors as extra negation symbols in the following way.

1. R occurs \neg -positively (\neg -negatively) in $\varphi_k^{i,j}$ iff R occurs \neg -positively (\neg -negatively) in φ and $\langle i, j, k \rangle \in \{\langle n, w, r \rangle, \langle n, e, b \rangle, \langle s, w, b \rangle, \langle s, e, r \rangle\}$.
2. R occurs \neg -negatively (\neg -positively) in $\varphi_k^{i,j}$ iff R occurs \neg -positively (\neg -negatively) in φ and $\langle i, j, k \rangle \in \{\langle s, w, r \rangle, \langle s, e, b \rangle, \langle n, w, b \rangle, \langle n, e, r \rangle\}$.
3. R occurs $-$ -positively ($-$ -negatively) in $\varphi_k^{i,j}$ iff R occurs $-$ -positively ($-$ -negatively) in φ and $\langle i, j, k \rangle \in \{\langle n, w, r \rangle, \langle n, e, b \rangle, \langle s, w, r \rangle, \langle s, e, b \rangle\}$.
4. R occurs $-$ -negatively ($-$ -positively) in $\varphi_k^{i,j}$ iff R occurs $-$ -positively ($-$ -negatively) in φ and $\langle i, j, k \rangle \in \{\langle n, w, b \rangle, \langle n, e, r \rangle, \langle s, w, b \rangle, \langle s, e, r \rangle\}$.

The main interpolation lemma can now be formulated and proved as follows.

Lemma 8.1 *Let Δ be a colored provable sequent in a language without function symbols other than individual constants. Then there is a sentence χ such that*

1. Δ^r , $\chi^{n,e}$ and Δ^b , $\chi^{n,w}$ are provable, and
2. each individual constant occurring in χ also occurs in Δ^r and in Δ^b ;

3. each relation symbol which occurs \neg -positively (\neg -negatively, $-$ -positively, $-$ -negatively) in χ also occurs \neg -positively (\neg -negatively, $-$ -positively, $-$ -negatively) in both Δ^r and Δ^b .

χ is called an interpolant of Δ and we write $\Delta : \chi$.

Proof: Let Π be a colored proof for Δ . The argument will proceed by induction on the complexity of Π . Consider the last rule ρ used in Π and suppose that ρ has premises $\Delta_1, \dots, \Delta_n$. Induction gives us χ_1, \dots, χ_n such that $\Delta_1 : \chi_1, \dots, \Delta_n : \chi_n$. In the following statements, which exhaust all possibilities for ρ , the abbreviation

$$\frac{\Delta_1 : \chi_1, \dots, \Delta_n : \chi_n}{\Delta : \chi}$$

means ‘ χ is an interpolant for Δ if each χ_i is an interpolant for Δ_i ($1 \leq i \leq n$).’

$$\begin{array}{l}
(\mathbf{R})^{int} \quad (a) \frac{}{\Gamma, \varphi_r^{i,w}, \varphi_r^{i,e} : \mathbf{f}}, \quad (b) \frac{}{\Gamma, \varphi_b^{i,w}, \varphi_b^{i,e} : \mathbf{t}}, \\
\quad (c) \frac{}{\Gamma, \varphi_r^{n,w}, \varphi_b^{n,e} : \varphi}, \quad (d) \frac{}{\Gamma, \varphi_b^{n,w}, \varphi_r^{n,e} : \neg - \varphi}, \\
\quad (e) \frac{}{\Gamma, \varphi_r^{s,w}, \varphi_b^{s,e} : \neg \varphi}, \quad (f) \frac{}{\Gamma, \varphi_b^{s,w}, \varphi_r^{s,e} : -\varphi}, \\
(\varphi \text{ atomic}) \\
(\neq, \neq)^{int} \quad \frac{}{\Gamma, \neq, \neq : \mathbf{t}} \\
(\mathbf{n}^w)^{int} \quad (a) \frac{\Gamma, \neq : \chi}{\Gamma, \mathbf{n}_r^{i,w} : \chi \wedge \mathbf{n}} \quad (b) \frac{\Gamma, \neq : \chi}{\Gamma, \mathbf{n}_b^{i,w} : \chi \vee \mathbf{b}} \\
(\mathbf{n}^e)^{int} \quad (a) \frac{\Gamma, \neq : \chi}{\Gamma, \mathbf{n}_r^{i,e} : \chi \wedge \mathbf{b}} \quad (b) \frac{\Gamma, \neq : \chi}{\Gamma, \mathbf{n}_b^{i,e} : \chi \vee \mathbf{n}} \\
(\neg)^{int} \quad \frac{\Gamma, \varphi_k^{i,j} : \chi}{\Gamma, \neg \varphi_k^{-i,j} : \chi} \\
(-)^{int} \quad \frac{\Gamma, \varphi_k^{i,j} : \chi}{\Gamma, -\varphi_k^{-i,-j} : \chi} \\
(\wedge_{sw}^{ne})^{int} \quad (a) \frac{\Gamma, \varphi_r^{i,j} : \chi_1 \quad \Gamma, \psi_r^{i,j} : \chi_2}{\Gamma, (\varphi \wedge \psi)_r^{i,j} : \chi_1 \vee \chi_2} \quad (b) \frac{\Gamma, \varphi_b^{i,j} : \chi_1 \quad \Gamma, \psi_b^{i,j} : \chi_2}{\Gamma, (\varphi \wedge \psi)_b^{i,j} : \chi_1 \wedge \chi_2} \\
((i, j) \in \{\langle n, e \rangle, \langle s, w \rangle\}) \\
(\wedge_{se}^{nw})^{int} \quad \frac{\Gamma, \varphi_k^{i,j}, \psi_k^{i,j} : \chi}{\Gamma, (\varphi \wedge \psi)_k^{i,j} : \chi}, \quad ((i, j) \in \{\langle n, w \rangle, \langle s, e \rangle\}) \\
(\forall_{sw}^{ne})^{int} \quad \frac{\Gamma, [c/x]\varphi_k^{i,j} : \chi}{\Gamma, \forall x \varphi_k^{i,j} : \chi}, \quad (k \in \{r, b\}, c \text{ not in } \Gamma, \varphi \text{ and } \langle i, j \rangle \in \{\langle n, e \rangle, \langle s, w \rangle\})
\end{array}$$

- (\forall_{se}^{nw})^{int}
- (a) $\frac{\Gamma, [c/x]\varphi_r^{i,j} : \chi}{\Gamma, \forall x\varphi_r^{i,j} : \chi}$, if c occurs in Γ^r or φ
- (b) $\frac{\Gamma, [c/x]\varphi_r^{i,j} : \chi}{\Gamma, \forall x\varphi_r^{i,j} : \forall x[x/c]\chi}$, if c does not occur in Γ^r or φ
- (c) $\frac{\Gamma, [c/x]\varphi_b^{i,j} : \chi}{\Gamma, \forall x\varphi_b^{i,j} : \chi}$, if c occurs in Γ^b or φ
- (d) $\frac{\Gamma, [c/x]\varphi_b^{i,j} : \chi}{\Gamma, \forall x\varphi_b^{i,j} : \exists x[x/c]\chi}$, if c does not occur in Γ^b or φ
- $((i, j) \in \{\langle n, w \rangle, \langle s, e \rangle\})$
- (id)^{int}
- (a) $\frac{}{\Gamma, d \approx d_r^{i,j} : \mathbf{f}}$, (b) $\frac{}{\Gamma, d \approx d_b^{i,j} : \mathbf{t}}$, $((i, j) \in \{\langle n, e \rangle, \langle s, w \rangle\})$
- (L)^{int}
- (a) $\frac{\Gamma, [c/x]\varphi_k^{i',j'} : \chi}{\Gamma, d \approx c_{k'}^{i,j}, [d/x]\varphi_k^{i',j'} : \chi}$,
if $k = k'$ ($k, k' \in \{r, b\}$) or x is not free in φ ;
- (b) $\frac{\Gamma, [c/x]\varphi_r^{i',j'} : \chi}{\Gamma, d \approx c_b^{i,j}, [d/x]\varphi_r^{i',j'} : \neg d \approx c \vee \chi}$,
if x is free in φ and c occurs in d, φ , or Γ^r ;
- (c) $\frac{\Gamma, [c/x]\varphi_r^{i',j'} : \chi}{\Gamma, d \approx c_b^{i,j}, [d/x]\varphi_r^{i',j'} : \forall y(\neg d \approx y \vee [y/c]\chi)}$,
if x is free in φ and c does not occur in d, φ , or Γ^r ;
- (d) $\frac{\Gamma, [c/x]\varphi_b^{i',j'} : \chi}{\Gamma, d \approx c_r^{i,j}, [d/x]\varphi_b^{i',j'} : d \approx c \wedge \chi}$,
if x is free in φ and c occurs in d, φ , or Γ^b ;
- (e) $\frac{\Gamma, [c/x]\varphi_b^{i',j'} : \chi}{\Gamma, d \approx c_r^{i,j}, [d/x]\varphi_b^{i',j'} : \exists y(d \approx y \wedge [y/c]\chi)}$,
if x is free in φ and c does not occur in d, φ , or Γ^b ;
- $((i, j) \in \{\langle n, w \rangle, \langle s, e \rangle\}, i' \in \{n, s\}, j' \in \{e, w\}$ and φ is atomic)

This ends the long list of possible cases for ρ . We shall prove one characteristic case, the (c) case of (L)^{int}, leaving the others to an interested reader. Suppose that $\Gamma, [c/x]\varphi_r^{i',j'} : \chi$, where x is free in φ and c does not occur in d, φ , or Γ^r . Then by definition $\Gamma^r, [c/x]\varphi^{i',j'}, \chi^{n,e}$ and $\Gamma^b, \chi^{n,w}$ are provable. But a proof of

$\Gamma^r, [c/x]\varphi^{i,j}, \chi^{n,e}$ can be extended as follows.

$$\frac{\frac{\frac{\Gamma^r, [c/x]\varphi^{i,j}, \chi^{n,e}}{\Gamma^r, [d/x]\varphi^{i,j}, d \approx c^{s,e}, \chi^{n,e}} \text{ (L)}}{\Gamma^r, [d/x]\varphi^{i,j}, \neg d \approx c^{n,e}, \chi^{n,e}} \text{ (\neg)}}{\Gamma^r, [d/x]\varphi^{i,j}, \neg d \approx c \vee \chi^{n,e}} \text{ (\vee}_{sw}^{ne})}$$

$$\frac{\Gamma^r, [d/x]\varphi^{i,j}, \neg d \approx c \vee \chi^{n,e}}{\Gamma^r, [d/x]\varphi^{i,j}, \forall y(\neg d \approx y \vee [y/c]\chi)^{n,e}} \text{ (\forall}_{sw}^{ne})}$$

The last step was possible because c does not occur in d , φ , or Γ^r . The proof of $\Gamma^b, \chi^{n,w}$ can be extended as follows.

$$\frac{\frac{\frac{\Gamma^b, c \approx c^{s,w}}{\Gamma^b, \neg c \approx c^{n,w}} \text{ (\neg)}}{\Gamma^b, \neg c \approx c \vee \chi^{n,w}} \text{ (\vee}_{se}^{nw})}}{\Gamma^b, d \approx c^{i,j}, \neg d \approx c \vee \chi^{n,w}} \text{ (L)}$$

$$\frac{\Gamma^b, d \approx c^{i,j}, \neg d \approx c \vee \chi^{n,w}}{\Gamma^b, d \approx c^{i,j}, \forall y(\neg d \approx y \vee [y/c]\chi)^{n,w}} \text{ (\forall}_{se}^{nw})}$$

Since x occurs free in φ , d will occur in $[d/x]\varphi$ as well as in $d \approx c$. Moreover, since $\Gamma, [c/x]\varphi_r^{i,j} : \chi$, each individual constant in $[y/c]\chi$ occurs in Γ^b and also either in Γ^r or in φ . This means that each individual constant in $\forall y(\neg d \approx y \vee [y/c]\chi)$ occurs both in $\Gamma^b, d \approx c^{i,j}$ and in $\Gamma^r, [d/x]\varphi^{i,j}$. If R is a nonlogical relation constant occurring \neg -positively (\neg -negatively, etc.) in $\forall y(\neg d \approx y \vee [y/c]\chi)$, then R occurs \neg -positively (\neg -negatively, etc.) in χ . Using the fact that $\Gamma, [c/x]\varphi_r^{i,j} : \chi$ we easily find that R occurs \neg -positively (\neg -negatively, $--$ -positively, $--$ -negatively) both in $\Gamma^b, d \approx c^{i,j}$ and in $\Gamma^r, [d/x]\varphi^{i,j}$. We conclude that $\forall y(\neg d \approx y \vee [y/c]\chi)$ is the required interpolant.

This concludes the construction of an interpolant for Δ . \square

Theorem 8.2 (Craig-Lyndon Interpolation Theorem) *Let Γ and Δ be sets of sentences in a language without function symbols other than individual constants. If $\Gamma \vdash \Delta$ there is a sentence χ such that*

1. $\Gamma \vdash \chi$ and $\chi \vdash \Delta$,
2. each individual constant which occurs in χ also occurs in Γ and in Δ ;
3. each relation symbol which occurs \neg -positively (\neg -negatively, $--$ -positively, $--$ -negatively) in χ also occurs \neg -positively (\neg -negatively, $--$ -positively, $--$ -negatively) in both Γ and Δ .

Here \vdash can uniformly be replaced with \vdash^{tr} or with \vdash^{nf} .

Proof: The \vdash case follows by applying the previous lemma to the colored sequent $\Theta = \{\varphi_r^{n,w} \mid \varphi \in \Gamma\} \cup \{\varphi_b^{n,e} \mid \varphi \in \Delta\}$. For the \vdash^{tr} and \vdash^{nf} cases consider $\Theta \cup \{\neq\}$ and $\Theta \cup \{\neq\}$, respectively. \square

A set of sentences Π is *inconsistent* if $\Pi \vdash$, that is, if $\{\varphi^{n,w} \mid \varphi \in \Pi\}$ is provable.

Corollary 8.3 (Robinson Joint Consistency Theorem) *Suppose Π_1 and Π_2 are sets of sentences in a language without function symbols and that $\Pi_1 \cup \Pi_2$ is inconsistent. Then there is a sentence χ , all whose nonlogical symbols also occur in $\Pi_1 \cup \Pi_2$, such that $\Pi_1 \vdash \chi$ and $\Pi_2, \chi \vdash$.*

Proof: Directly from Lemma 8.1. □

9 Conclusion We have generalized the sequent calculus for predicate logic to systems for a range of partial and paraconsistent logics. Our methods work for systems based on one-directional notions of logical consequence (\models^r and \models^f), but also for the bidirectional notion \models based on \leq_t , which to us seems more attractive from an esthetic point of view. The bidirectional notion differs from the one-directional notions only if connectives are considered that can only be defined in terms of \mathbf{n} , but among these are the important \oplus and \otimes . Other interesting connectives which are definable with \mathbf{n} , but not without this connective, are discussed in [5]. That the techniques we have used stay close to the techniques usually employed for predicate logic (e.g., in Fitting [14]) comes as a surprise in view of the remarks in the otherwise excellent [21] and [22]. The fact that we have been able to prove an interpolation theorem for \mathbf{L}_4 gives some support to the idea that a reasonable sequent formalization for this logic was found.

Acknowledgments I wish to thank Manfred Pinkal for inviting me to the stimulating environment of Computational Linguistics group at the Universität des Saarlandes (SFB 378–Ressourcenadaptive Kognitive Prozesse) and Hans Kamp for inviting me to the no less inspiring environment of the Institut für Maschinelle Sprachverarbeitung in Stuttgart. Most research for this paper was done during these two stays. I also wish to thank the anonymous referee for a list of useful comments and suggestions.

REFERENCES

- [1] Arieli, O., and A. Avron, “Reasoning with logical lattices,” *The Journal of Logic, Language, and Information*, vol. 5 (1996), pp. 25–63.
Zbl 0851.03017 MR 97c:03094 2, 2, 5.5
- [2] Baaz, M., C. G. Fermüller, and R. Zach, “Elimination of cuts in first-order finite-valued logics,” *The Journal of Information Processing and Cybernetics*, vol. 29 (1994), pp. 333–55. Zbl 0821.03013 1, 1
- [3] Barwise, J., and J. Perry, *Situations and Attitudes*, The MIT Press, Cambridge, 1983.
Zbl 0946.03007 MR 2001h:03051 1
- [4] Belnap, N. D., Jr., “A useful four-valued logic,” pp. 8–37 in *Modern Uses of Multiple-Valued Logic*, edited by J. M. Dunn and G. Epstein, Reidel, Dordrecht, 1977.
Zbl 0424.03012 MR 58:5021 1, 1, 1, 1, 2, 2, 3
- [5] Blamey, S., “Partial logic,” pp. 1–70 in *Handbook of Philosophical Logic*, vol. 3, edited by D. M. Gabbay and F. Guenther, Reidel, Dordrecht, 1986. Zbl 0875.03023 1, 1, 9
- [6] Bochman, A., “A logical foundation for logic programming I: biconsequence relations and nonmonotonic completion,” *The Journal of Logic Programming*, vol. 35 (1998), pp. 151–70. Zbl 0905.68032 MR 2000e:68028 1, 1, 4
- [7] Carnielli, W. A., “Systematization of finite many-valued logics through the method of tableaux,” *The Journal of Symbolic Logic*, vol. 52 (1987), pp. 473–93.
Zbl 0633.03008 MR 88g:03034 1

- [8] Carnielli, W. A., “On sequents and tableaux for many-valued logics,” *The Journal of Non-Classical Logic*, vol. 8 (1991), pp. 59–78. [Zbl 0774.03006](#) [MR 94b:03046](#) 1
- [9] Cleave, J. P., “The notion of logical consequence in the logic of inexact predicates,” *Zeitschrift für mathematische Logik und Grundlagen der Mathematik*, vol. 20 (1974), pp. 307–24. [Zbl 0299.02015](#) [MR 51:10028](#) 1, 1, 4
- [10] D’Agostino, M., “Investigations into the complexity of some propositional calculi,” Technical Report, Oxford University Computing Laboratory, 1990. [4.3](#)
- [11] Dunn, J. M., “Intuitive semantics for first-degree entailments and ‘coupled trees’,” *Philosophical Studies*, vol. 29 (1976), pp. 149–68. [MR 58:10311](#) 2
- [12] Feferman, S., “Toward useful type free theories I,” *The Journal of Symbolic Logic*, vol. 49 (1984), pp. 75–111. [Zbl 0574.03043](#) [MR 85i:03068](#) 1
- [13] Fitting, M., “Bilattices and the semantics of logic programming,” *The Journal of Logic Programming*, vol. 11 (1991), pp. 91–116. [Zbl 0757.68028](#) [MR 92c:68119](#) 1, 2
- [14] Fitting, M., *First-Order Logic and Automated Theorem Proving*, Springer, New York, 1996. [Zbl 0848.68101](#) [MR 97b:68197](#) 9
- [15] Gilmore, P. C., “The consistency of partial set theory without extensionality,” pp. 147–53 in *Axiomatic Set Theory, Proceedings of Symposia in Pure Mathematics 13, Part II*, American Mathematical Society, Providence, 1974. [Zbl 0309.02065](#) [MR 50:12721](#) 1
- [16] Ginzburg, M. L., “Multivalued logics: a uniform approach to reasoning in A,” *Computer Intelligence*, vol. 4 (1988), pp. 256–316. [2, 2](#)
- [17] Hähnle, R., *Automated Deduction in Multiple-valued Logics*, Oxford University Press, Oxford, 1993. [Zbl 0798.03010](#) [MR 96a:68107](#) 1, 1
- [18] Holden, M., “Weak logic theory,” *Theoretical Computer Science*, vol. 79 (1991), pp. 295–321. [Zbl 0724.03020](#) [MR 92f:68111](#) 1
- [19] Jaspars, J., *Calculi for Constructive Communication*, Ph.D thesis, Tilburg University, Tilburg, 1994. [1, 1](#)
- [20] Kleene, S. C., *Introduction to Metamathematics*, Van Nostrand, Amsterdam, 1952. [Zbl 0047.00703](#) [MR 14:525m](#) 8
- [21] Langholm, T., *Partiality, Truth and Persistence*, CSLI Lecture Notes, Stanford, 1988. [Zbl 0665.03024](#) 1, 1, 1, 1, 1, 8, 9
- [22] Langholm, T., “How different is partial logic?,” pp. 3–43 in *Partiality, Modality, and Nonmonotonicity*, edited by P. Doherty, CSLI, Stanford, 1996. [Zbl 0904.03012](#) [MR 1427283](#) 1, 4, 4, 4, 9
- [23] Muskens, R. A., “Going partial in Montague grammar,” pp. 175–220 in *Semantics and Contextual Expression, Proceedings of the Sixth Amsterdam Colloquium*, edited by R. Bartsch, J. van Benthem, and P. van Emde Boas, Foris, Dordrecht, 1989. [Zbl 0773.03020](#) 1
- [24] Muskens, R. A., *Meaning and Partiality*, CSLI, Stanford, 1995. [1, 1, 1, 1, 2, 3](#)
- [25] Rousseau, G., “Sequents in many-valued logic I,” *Fundamenta Mathematicæ*, vol. 60 (1967), pp. 23–33. [Zbl 0154.25504](#) [MR 35:1451](#) 1, 4
- [26] Schröter, K., “Methoden zur Axiomatisierung beliebiger Aussagen- und Prädikatenkalküle,” *Zeitschrift für mathematische Logik und Grundlagen der Mathematik*, vol. 1 (1955), pp. 241–51. [Zbl 0066.25604](#) [MR 17:1038g](#) 1, 4

- [27] Takeuti, G., *Proof Theory*, North-Holland, Amsterdam, 1975.
[Zbl 0681.03039](#) [MR 58:27366b](#) 8
- [28] Thijsse, E., *Partial Logic and Knowledge Representation*, Ph.D thesis, Tilburg University, Tilburg, 1992. 1, 1
- [29] Visser, A., “Four-valued semantics and the Liar,” *The Journal of Philosophical Logic*, vol. 13 (1984), pp. 181–212. [Zbl 0546.03007](#) [MR 86f:03042](#) 1
- [30] Woodruff, P. W., “Truth and logic, part I: paradox and truth,” *The Journal of Philosophical Logic*, vol. 13 (1984), pp. 213–32. 1
- [31] Zach, R., “Proof theory of finite-valued logics.” Technical Report TUW-E185.2-Z.1-93, Institut für Computersprachen, Technische Universität Wien, Vienna, 1993. 1

Department of Linguistics
Tilburg University
PO Box 90153
NL-5000 LE Tilburg
THE NETHERLANDS
email: R.A.Muskens@kub.nl