

## CORRIGENDUM TO MY PAPER

"A PROPOSITIONAL CALCULUS INTERMEDIATE BETWEEN THE  
MINIMAL CALCULUS AND THE CLASSICAL"

CHARLES PARSONS

Dr. R. A. Bull has pointed out to me that a step in the proof of theorem 2 of my paper, 'A propositional calculus intermediate between the minimal calculus and the classical' (this *Journal*, vol. 7 (1966), pp. 353-358) is fallacious. What I was entitled to infer from [iv] and  $-A_1 \dots -A_n \vdash S$  is that  $S$  is derivable from all formulae

$$-C_i \vee. - -C_j \supset C_j \quad ,$$

where  $C_1 \dots C_{m+n}$  are  $A_1 \dots A_n, B_1 \dots B_m$ . Thus  $S$  is provable in **MCC** if  $\bar{q} \vee. \bar{p} \supset p$  is.

The text shows  $\vdash \bar{p} \vee. \bar{p} \supset p$ , and  $\vdash \bar{p} \vee. \bar{p} \supset p$  by ax. 10. Hence  $\vdash \bar{p} \wedge \bar{p} \vee. \bar{p} \supset p$ . But  $\vdash \bar{p} \supset. \bar{p} \supset \bar{q}$  in **MC**. Hence  $\vdash \bar{q} \vee. \bar{p} \supset p$ , q.e.d.

Similarly, to justify the claim on p. 358 that every intuitionistically valid and pseudo-valid formula  $S$  is provable in **MC** +  $\bar{p} \vee. \bar{p} \supset. p \supset q$ , it is necessary to show  $\bar{r} \vee. \bar{p} \supset. p \supset q$  provable in that system. We have  $\vdash -(s \supset s) \vee. -(s \supset s) \supset q$  and hence

$$\vdash r \supset - (s \supset s) \vee. p \supset - (s \supset s) \supset. p \supset q.$$

But  $\bar{r} \equiv. r \supset -(s \supset s)$  and  $\bar{p} \equiv. p \supset -(s \supset s)$  are provable in **MC**.

*Columbia University*  
*New York City, New York*