

# TRANSLATION IN MEASURE ALGEBRAS AND THE CORRESPONDENCE TO FOURIER TRANSFORMS VANISHING AT INFINITY

Charles F. Dunkl and Donald E. Ramirez

Let  $G$  denote a locally compact (not necessarily abelian) group and  $M(G)$  the collection of finite regular Borel measures on  $G$ . The set  $M(G)$  is a semisimple Banach algebra with identity under convolution  $*$ . It can be identified with the dual space of  $C_0(G)$ , the space of continuous complex-valued functions on  $G$  that vanish at infinity, with the sup-norm. The group  $G$  has a left-invariant regular Borel measure  $dm(x)$  that is unique up to a constant and is called the left Haar measure of  $G$ . Let  $C^B(G)$  denote the space of bounded continuous functions on  $G$ . For each  $x \in G$ , we define on  $C^B(G)$  the left-translation operator by the relation

$$L(x)f(y) = f(x^{-1}y) \quad (f \in C^B(G)).$$

We say that  $f \in C^B(G)$  is right uniformly continuous if  $L(x_\alpha)f \xrightarrow{\alpha} L(x)f$  uniformly, whenever  $x_\alpha \xrightarrow{\alpha} x$ . Let  $C_{ru}^B(G)$  denote the subspace of  $C^B(G)$  of right uniformly continuous functions. For  $\mu \in M(G)$ , define  $L(x)\mu \in M(G)$  by the condition

$$\int_G f(t) dL(x)\mu(t) = \int_G L(x^{-1})f(t) d\mu(t),$$

where  $f \in C_0(G)$ . We wish to study for which  $\mu \in M(G)$  the map  $x \mapsto L(x)\mu$  is continuous from  $G$  into  $M(G)$ , where  $M(G)$  will be equipped with an  $L(x)$ -invariant metric topology. In particular, we shall characterize  $M_0(G)$ , the algebra of measures whose Fourier transform vanishes at infinity.

Let  $A \subset C_{ru}^B(G)$  be a linear subspace with sufficiently many elements to separate the points of  $M(G)$ ; in other words, if  $\mu \in M(G)$  and if

$$\int_G f(t) d\mu(t) = 0$$

for all  $f \in A$ , then  $\mu = 0$ . We are then able to pair  $A$  and  $M(G)$  by the relation

$$\langle f, \mu \rangle = \int_G f(t) d\mu(t) \quad (f \in A; \mu \in M(G)).$$

Let  $\sigma(A, M(G))$  denote the weak topology on  $A$  induced by this pairing. Suppose  $A$  can be written as  $\bigcup_{k=1}^{\infty} A_k$ , where each  $A_k$  is a subset of  $A$  that is  $L(x)$ -invariant for all  $x \in G$  and where each  $A_k$  is  $\sigma(A, M(G))$ -bounded. Note that  $A_k$  is

---

Received December 24, 1969.

This research was supported in part by NSF contract number GP 8981.

Michigan Math. J. 17 (1970).

$\sigma(A, M(G))$ -bounded if and only if  $A_k$  is bounded in sup-norm. We let  $\mathcal{F}(A_k)$  denote the topology on  $M(G)$  of uniform convergence on the sets  $A_k$ . Note that  $\mathcal{F}(A_k)$  gives an  $L(x)$ -invariant metric topology on  $M(G)$ . For  $k \geq 1$ , let

$$\tau_k(\mu) = \sup \{ |\langle f, \mu \rangle| : f \in A_k \}.$$

Then  $\tau_k$  is an  $L(x)$ -invariant seminorm on  $M(G)$ .

*Definition.* For  $\mu \in M(G)$ , we say that  $\mu$  has *separable orbit* in  $(M(G), \mathcal{F}(A_k))$  if there exists a sequence  $\{x_n\}_{n=1}^\infty \subset G$  such that for each  $x \in G$ ,  $k \geq 1$ , and  $\varepsilon > 0$ , there exists an  $x_n$  such that  $\tau_k(L(x)\mu - L(x_n)\mu) < \varepsilon$ .

**PROPOSITION 1.** *Let  $\mu \in M(G)$  have separable orbit in  $(M(G), \mathcal{F}(A_k))$ . Then  $s \mapsto L(s)\mu$  is continuous from  $G$  to  $(M(G), \mathcal{F}(A_k))$ .*

*Proof.* Let  $s_\alpha \xrightarrow{\alpha} s$ . Choose  $k \geq 1$  and  $\varepsilon > 0$ . We need to show there exists an  $\alpha_0$  such that for  $\alpha \geq \alpha_0$ , we have the inequality  $\tau_k(L(s_\alpha)\mu - L(s)\mu) < \varepsilon$ . Note that for  $f \in C_{ru}^B(G)$ ,  $L(y_\beta^{-1})f \xrightarrow{\beta} L(y^{-1})f$  uniformly as  $y_\beta \xrightarrow{\beta} y$  (and hence as  $y_\beta^{-1} \xrightarrow{\beta} y^{-1}$ ). Thus

$$\langle f, L(y_\beta)\mu \rangle = \langle L(y_\beta^{-1})f, \mu \rangle \xrightarrow{\beta} \langle L(y^{-1})f, \mu \rangle = \langle f, L(y)\mu \rangle.$$

Let  $S(n) = \{y \in G : \tau_k(L(y)\mu - L(x_n)\mu) \leq \varepsilon/3\}$ . We wish to show that  $S(n)$  is closed. Let  $y_\beta \in S(n)$  be such that  $y_\beta \xrightarrow{\beta} y$ . Thus

$$\tau_k(L(y)\mu - L(x_n)\mu) = \sup \{ \lim_{\beta} |\langle f, L(y_\beta)\mu - L(x_n)\mu \rangle| : f \in A_k \} \leq \varepsilon/3.$$

Hence  $S(n)$  is closed.

By hypothesis,  $G = \bigcup_{n=1}^\infty S(n)$ . By the Baire category theorem for locally compact groups, there exists  $n_0$  such that  $S(n_0)$  has an interior. Thus there exists an open set  $U$  about  $s$  such that  $t_0 s^{-1}U \subset S(n_0)$  for some  $t_0 \in S(n_0)$ . Let  $\alpha_0$  be such that  $s_\alpha \in U$  for  $\alpha \geq \alpha_0$ . We now show that for  $\alpha \geq \alpha_0$ , the inequality

$$\tau_k(L(s_\alpha)\mu - L(s)\mu) < \varepsilon$$

holds. For  $\alpha \geq \alpha_0$ , we have that

$$\begin{aligned} \tau_k(L(s_\alpha)\mu - L(s)\mu) &= \tau_k(L(t_0 s^{-1})L(s_\alpha)\mu - L(t_0 s^{-1})L(s)\mu) \\ &\leq \tau_k(L(t_0 s^{-1}s_\alpha)\mu - L(x_{n_0})\mu) + \tau_k(L(x_{n_0})\mu - L(t_0)\mu) \\ &\leq \frac{\varepsilon}{3} + \frac{\varepsilon}{3} < \varepsilon, \end{aligned}$$

since  $t_0, t_0 s^{-1}s_\alpha \in t_0 s^{-1}U \subset S(n_0)$ . ■

**PROPOSITION 2.** *Let  $G$  be  $\sigma$ -compact. If  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $(M(G), \mathcal{F}(A_k))$ , then  $\mu$  has separable orbit in  $(M(G), \mathcal{F}(A_k))$ .*

*Proof.* Note that  $(M(G), \mathcal{F}(A_k))$  is a metric space. Let  $G = \bigcup_{n=1}^\infty K_n$ , where  $K_n$  is compact. The image of  $K_n$  under  $x \mapsto L(x)\mu$  is a compact metric space and hence is separable. Thus the image of  $G$  is separable. ■

If  $G$  is not  $\sigma$ -compact and  $M(G)$  has the measure norm topology, then no non-zero measure has a separable orbit.

We now show that  $\mu \in M(G)$  has the property that  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $(M(G), \mathcal{T}(A_k))$  if and only if  $\mu$  is in the  $\mathcal{T}(A_k)$ -closure of  $L^1(G)$ , denoted by  $L^1(\overline{G})^A$ .

**THEOREM 3.** *Let  $\mu \in M(G)$  be such that  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $(M(G), \mathcal{T}(A_k))$ . Then  $\mu \in L^1(\overline{G})^A$ .*

*Proof.* Let  $\{f_\alpha\}$  be an approximate identity in  $L^1(G)$ , indexed over a neighborhood base of  $e$ ; in other words,  $\text{support}(f_\alpha) \subset \alpha$ ,  $f_\alpha \geq 0$ , and  $\|f_\alpha\|_1 = 1$ . Choose  $k_0 \geq 1$  and  $\varepsilon > 0$ . It suffices to show that  $\tau_k(f_\alpha * \mu - \mu) \leq \varepsilon$  for  $\alpha \geq \alpha_0$ , for some  $\alpha_0$ . Pick  $U$  to be a symmetric neighborhood of  $e$  in  $G$  such that

$$\tau_k(L(x)\mu - \mu) < \varepsilon$$

for  $x \in U$ . Choose  $\alpha_0$  such that the inequality  $\alpha \geq \alpha_0$  implies that  $\text{support}(f_\alpha) \subset U$ . Now for  $\alpha \geq \alpha_0$ ,

$$\begin{aligned} \tau_k(f_\alpha * \mu - \mu) &= \sup \left\{ \left| \langle \phi, f * \mu - \mu \rangle \right| : \phi \in A_k \right\} \\ &= \sup \left\{ \left| \int_G \phi(x) df_\alpha * \mu(x) - \int_G \phi(y) d\mu(y) \right| : \phi \in A_k \right\} \\ &= \sup \left\{ \left| \int_G \int_G \phi(xy) d\mu(y) f_\alpha(x) dx - \int_G \int_G f_\alpha(x) dx \phi(y) d\mu(y) \right| : \phi \in A_k \right\} \\ &= \sup \left\{ \left| \int_G f_\alpha(x) dx \left[ \int_G \phi(y) dL(x)\mu(y) - \int_G \phi(y) d\mu(y) \right] \right| : \phi \in A_k \right\} \\ &\leq \sup_{x \in U} \tau_k(L(x)\mu - \mu) \leq \varepsilon. \blacksquare \end{aligned}$$

**THEOREM 4.** *Let  $\mu \in L^1(\overline{G})^A$ . Then  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $(M(G), \mathcal{T}(A_k))$ .*

*Proof.* We note first that since  $A_k$  is  $\sigma(A, M(G))$ -bounded and  $L(x)$ -invariant,  $A_k$  is a sup-norm bounded set in  $C^B(G)$ ; in fact, for all  $x \in G$ , we have that

$$\sup_{f \in A_k} |f(x)| = \sup_{f \in A_k} \left| \int_G L(x^{-1})fd\delta_e \right| = \sup_{f \in A_k} \left| \int_G fd\delta_e \right| = M < \infty,$$

where  $\delta_e$  is the unit mass at  $e$ . Now  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $M(G)$  in the measure norm, for  $\mu \in L^1(G)$ . Thus, since  $A_k$  is a sup-norm bounded set,  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $(M(G), \mathcal{T}(A_k))$  for  $\mu \in L^1(G)$ . Choose

$\mu \in L^1(\overline{G})^A$ , and let  $x_\alpha \xrightarrow{\alpha} x$ . Let  $k \geq 1$  and  $\varepsilon > 0$ . We need to find an  $\alpha_0$  such that if  $\alpha_0 \leq \alpha$ , then  $\tau_k(L(x_\alpha)\mu - L(x)\mu) < \varepsilon$ . First pick  $f \in L^1(G)$  such that  $\tau_k(f - \mu) < \varepsilon/3$ . Now choose  $\alpha_0$  such that for  $\alpha \geq \alpha_0$ ,  $\tau_k(L(x_\alpha)f - L(x)f) < \varepsilon/3$ . Thus for  $\alpha \geq \alpha_0$ ,

$$\begin{aligned} \tau_k(L(x_\alpha)\mu - L(x)\mu) &\leq \tau_k(L(x_\alpha)\mu - L(x_\alpha)f) + \tau_k(L(x_\alpha)f - L(x)f) + \tau_k(L(x)f - L(x)\mu) \\ &< \tau_k(\mu - f) + \frac{\varepsilon}{3} + \tau_k(f - \mu) < \varepsilon. \blacksquare \end{aligned}$$

*Remark.* The two theorems above also hold if  $A$  is a space of bounded Borel functions, rather than a subspace of  $C_{ru}^B(G)$ .

For  $\mu \in M(G)$ , let  $\|\mu\|$  denote the measure norm of  $\mu$ , that is, the norm of  $\mu$  as a linear functional on  $C_0(G)$  with sup-norm  $\|f\|_\infty = \sup\{|f(x)|: x \in G\}$ . If we let  $A_k = \{f \in C_0(G): \|f\|_\infty < k\}$ , then  $\mathcal{F}(A_k)$  is the measure norm topology. Thus we have the following corollaries.

**COROLLARY 5.** *Let  $\mu \in M(G)$ . If  $\mu$  has separable orbit in  $(M(G), \|\cdot\|)$ , then  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $(M(G), \|\cdot\|)$ .*

*Suppose  $G$  is  $\sigma$ -compact. If  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $(M(G), \|\cdot\|)$ , then  $\mu$  has separable orbit in  $(M(G), \|\cdot\|)$ .*

**COROLLARY 6.** *Let  $\mu \in M(G)$ . The measure  $\mu$  is absolutely continuous if and only if  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $(M(G), \|\cdot\|)$ .*

*Remarks.* Propositions 1 and 2 are similar in spirit to a theorem of K. Shiga [8] in the compact case. Corollary 5 was obtained by R. Larsen [5] for the case where  $G$  is second countable and by K. W. Tam [9] in the general case. Corollary 6 was obtained by W. Rudin [7].

We now study  $M(G)$  under its sup-norm  $\|\cdot\|_\infty$ . We shall give first the abelian case for motivation. We then treat the compact nonabelian case and finally the general case.

Let  $G$  be abelian, and let  $\hat{G}$  denote the character group of  $G$ . For  $\mu \in M(G)$ , define  $\hat{\mu}(\gamma) = \int_G \overline{\gamma(x)} d\mu(x)$ , for  $\gamma \in \hat{G}$ . Then  $\hat{\mu}$  is the Fourier transform of  $\mu$ . For  $\mu \in M(G)$ , let

$$\|\mu\|_\infty = \sup\{|\hat{\mu}(\gamma)|: \gamma \in \hat{G}\}.$$

Let  $M_0(G) = \{\mu \in M(G): \mu \in C_0(\hat{G})\}$ .

**COROLLARY 7.** *Let  $G$  be abelian. The map  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $(M(G), \|\cdot\|_\infty)$  if and only if  $\mu \in M_0(G)$ .*

*Proof.* Let  $A_k = \{\hat{f}: f \in L^1(\hat{G}) \text{ with } \|f\|_1 < k\}$ . Then  $\mathcal{F}(A_k)$  is the topology of  $(M(G), \|\cdot\|_\infty)$ . ■

*Remark.* Corollary 7 was obtained by R. Goldberg and A. Simon [3]. They used the following result: If  $U$  is a relatively compact neighborhood of  $0$  in  $G$  (where  $G$  is abelian), there exists a compact subset  $K$  of  $\hat{G}$  such that for  $\gamma \in \hat{G} \setminus K$ , there exists an  $x \in U$  with  $\Re \gamma(x) \leq 0$ . To see this, let  $\sqrt{2} \leq \delta < \sqrt{3}$ , and define  $U^0 = \{\gamma \in \hat{G}: |\gamma(x) - 1| < \delta \text{ for all } x \in U\}$ . Note that  $U^0$  is relatively compact in  $\hat{G}$  (K. H. Hofmann and P. S. Mostert [4, p. 284] or Pontryagin [6, p. 237]). Let  $K$  be the closure of  $U^0$  in  $\hat{G}$ . We now prove the analogous result for the case where  $G$  is compact and nonabelian. This result is independent of the rest of this paper. We use the notation of Dunkl and Ramirez [1, Chapters 7 and 8], where proofs of unproved statements below may be found.

Let  $G$  be a compact, nonabelian group. We let  $\hat{G}$  denote the set of equivalence classes of continuous, unitary irreducible representations of  $G$ . For  $\alpha \in \hat{G}$ , let  $T_\alpha$  be an element of  $\alpha$ . Then  $T_\alpha$  is a homomorphism of  $G$  into  $U(n_\alpha)$ , the group of unitary  $n_\alpha \times n_\alpha$  matrices, where  $n_\alpha$  is the dimension of  $\alpha$ . We use  $T_\alpha(x)_{ij}$  to denote the matrix entries of  $T_\alpha(x)$  ( $1 \leq i, j \leq n$ ) and  $T_{\alpha ij}$  to denote the function  $x \mapsto T_\alpha(x)_{ij}$ . Clearly

$$T_\alpha(xy)_{ij} = \sum_{k=1}^{n_\alpha} T_\alpha(x)_{ik} T_\alpha(y)_{kj} \quad \text{and} \quad T_\alpha(y^{-1})_{ij} = \overline{T_\alpha(y)_{ji}}.$$

Furthermore,  $T_{\alpha ij} \in C(G)$ , where  $C(G)$  denotes the set of continuous functions on  $G$ . For  $\alpha \in \hat{G}$ , let

$$\chi_\alpha(x) = \text{trace}(T_\alpha(x)) = \sum_{i=1}^{n_\alpha} T_\alpha(x)_{ii}.$$

This trace  $\chi_\alpha$  is called the character of  $\alpha$ , and it is independent of the choice of  $T_\alpha$  in  $\alpha$ . Let  $X$  be an  $n$ -dimensional, complex inner-product space. Let  $\mathcal{B}(X)$  denote the space of linear maps from  $X$  into  $X$ . We define the operator norm of  $A \in \mathcal{B}(X)$  by

$$\|A\|_\infty = \sup \{ |A\xi| : \xi \in X, |\xi| \leq 1 \}.$$

For the trace of  $A$ , we find that  $\text{Tr } A = \sum_{i=1}^n (A\xi_i, \xi_i)$ , where  $\{\xi_i\}_{i=1}^n$  is some orthonormal basis for  $X$  and  $(\cdot, \cdot)$  denotes the inner product in  $X$ . Let  $|A|$  denote  $(A^*A)^{1/2}$ . The operator norm of  $A$  is  $\|A\|_\infty$ , that is,  $\max \{\lambda_i : 1 \leq i \leq n\}$ , where the  $\lambda_i$  are the eigenvalues of  $|A|$ . For each  $A \in \mathcal{B}(X)$ , we have the inequality  $|\text{Tr } A| \leq n \|A\|_\infty$ .

**PROPOSITION 8.** *Let  $G$  be a compact group. Suppose  $0 < \delta < \sqrt{3}$ , and let  $U$  be a neighborhood of  $e$  in  $G$ . Let  $U^0 = \{\alpha \in \hat{G} : \|T_\alpha(x) - I\|_\infty < \delta \text{ for all } x \in U\}$ . Then  $U^0$  is finite.*

*Proof.* We show that  $U^0$  is an equicontinuous set of representations of  $G$ . Choose  $\varepsilon > 0$ . Let  $K$  be a positive constant such that for  $0 \leq \theta \leq 2\pi/3$ , we have the inequality  $|e^{i\theta} - 1| \leq K\theta$  (for example, let  $K = 3\pi\sqrt{3}/2$ ). Define

$$V_m = \{x \in G : x, x^2, \dots, x^m \in U\}.$$

Clearly,  $V_m$  is a neighborhood of  $e$  in  $G$ . Pick  $m$  such that  $K\delta/m < \varepsilon$ . Then for  $x_1, x_2 \in G$  with  $x = x_1^{-1}x_2 \in V_m$ , we have that

$$\begin{aligned} \|T_\alpha(x_1) - T_\alpha(x_2)\|_\infty &= \|I - T_\alpha(x_1^{-1}x_2)\|_\infty = \|I - T_\alpha(x)\|_\infty \\ &= \sup \{ |1 - e^{i\theta_j}| : 1 \leq j \leq n_\alpha \} \quad (\alpha \in U^0), \end{aligned}$$

by diagonalizing  $T_\alpha(x)$ . Thus

$$\|I - T_\alpha(x^r)\|_\infty = \sup \{ |1 - e^{ir\theta_j}| : 1 \leq j \leq n_\alpha \} < \delta$$

for  $1 \leq r \leq m$ . Therefore

$$\|I - T_\alpha(x)\|_\infty = \sup \{ |1 - e^{i\theta_j}| : 1 \leq j \leq n_\alpha \} < \frac{K\delta}{m} < \varepsilon.$$

Thus  $U^0$  is an equicontinuous set of representation of  $G$ .

Let  $\chi_\alpha = \text{Tr } T_\alpha$ . We claim that  $\{\chi_\alpha/n_\alpha : \alpha \in U^0\}$  is an equicontinuous, uniformly bounded set of functions. This is the case since

$$|\text{Tr}(I - T_\alpha)| \leq n_\alpha \|I - T_\alpha\|_\infty.$$

Further  $\|\chi_\alpha/n_\alpha\|_\infty \leq 1$ , and hence  $\{\chi_\alpha/n_\alpha : \alpha \in U^0\}$  is relatively compact, by the Arzelà-Ascoli theorem. Since the  $\{\chi_\alpha/n_\alpha\}$  are orthogonal in  $L^2(G)$ , either  $U^0$  is finite or  $\{\chi_\alpha/n_\alpha : \alpha \in U^0\}$  has 0 as a uniform cluster point. This latter condition cannot happen, since  $\chi_\alpha(e)/n_\alpha = 1$ . ■

Let  $G$  be as above (that is, compact and nonabelian). We shall give the analogue to Corollary 7. Let the set  $\phi = \{\phi_\alpha : \alpha \in \hat{G}, \text{ where } \phi_\alpha \in \mathcal{B}(C^{n_\alpha})\}$  be such that  $\sup \{\|\phi_\alpha\|_\infty : \alpha \in \hat{G}\} < \infty$ . The set of all such  $\phi$  is denoted by  $\mathcal{L}^\infty(\hat{G})$ . It is a Banach algebra under the norm  $\|\phi\|_\infty = \sup \{\|\phi_\alpha\|_\infty : \alpha \in \hat{G}\}$  and under co-ordinatewise operations. Let

$$\mathcal{E}_0(\hat{G}) = \{ \phi \in \mathcal{L}^\infty(\hat{G}) : \lim_{\alpha \rightarrow \infty} \|\phi_\alpha\|_\infty = 0 \}.$$

For  $\mu \in M(G)$ , the Fourier transform  $\hat{\mu}$  of  $\mu$  is a matrix-valued function, defined for  $\alpha \in \hat{G}$  by the relation

$$\alpha \mapsto \hat{\mu}_\alpha = \int_G T_\alpha(x^{-1}) d\mu(x).$$

Note that  $\hat{\mu} \in \mathcal{L}^\infty(\hat{G})$ . Thus for  $\mu \in M(G)$ , let  $\|\mu\|_\infty = \sup \{\|\hat{\mu}_\alpha\|_\infty : \alpha \in \hat{G}\}$ . We define  $M_0(G)$  to be the set  $\{\mu \in M(G) : \hat{\mu} \in \mathcal{E}_0(\hat{G})\}$ .

Let  $A \in \mathcal{B}(X)$ , where  $X$  is a finite-dimensional, complex inner-product space. We define the dual norm to  $\|\cdot\|_\infty$  by  $\|A\|_1 = \sup \{ |\text{Tr}(AB)| : \|B\|_\infty \leq 1 \}$ . This norm can also be characterized by the condition  $\|A\|_1 = \text{Tr}(|A|)$ . For  $\phi \in \mathcal{L}^\infty(G)$ , we put

$$\|\phi\|_1 = \sum_{\alpha \in \hat{G}} n_\alpha \|\phi_\alpha\|_1.$$

Let  $\mathcal{L}^1(\hat{G}) = \{ \phi \in \mathcal{L}^\infty(\hat{G}) : \|\phi\|_1 < \infty \}$ . Then  $\mathcal{L}^1(\hat{G})$  is a Banach space under  $\|\cdot\|_1$ . For  $\phi \in \mathcal{L}^1(\hat{G})$ , let  $\text{Tr}(\phi) = \sum_{\alpha \in \hat{G}} n_\alpha \text{Tr}(\phi_\alpha)$ . For  $\psi \in \mathcal{L}^1(G)$  and  $\phi \in \mathcal{L}^\infty(G)$ , we obtain the inequality  $|\text{Tr}(\phi\psi)| \leq \|\phi\|_\infty \|\psi\|_1$ .

We now define  $A(G)$ , the Fourier algebra of  $G$ , and we pair  $A(G)$  and  $M(G)$  to get the compact analogue of Corollary 7. Let  $A(G)$  be the set of  $f \in C(G)$  for which  $\hat{f} \in \mathcal{L}^1(\hat{G})$ . We define a norm on  $A(G)$  by

$$\|f\|_A = \|\hat{f}\|_1 = \sum_{\alpha \in \hat{G}} n_\alpha \|\hat{f}_\alpha\|_1 < \infty.$$

Note that  $A(G)$  is isomorphic to  $\mathcal{L}^1(\hat{G})$ , because for each  $\phi \in \mathcal{L}^1(\hat{G})$ , the function  $f(x) = \sum_{\alpha \in \hat{G}} n_\alpha \operatorname{Tr}(\phi_\alpha T_\alpha(x))$  is in  $A(G)$ ; further,

$$\|f\|_\infty = \sup_{x \in G} \left| \sum_{\alpha \in \hat{G}} n_\alpha \operatorname{Tr}(\phi_\alpha T_\alpha(x)) \right| \leq \sum_{\alpha \in \hat{G}} n_\alpha \|\phi_\alpha\|_1 = \|\phi\|_1.$$

We note that for  $f \in A(G)$ ,  $\|L(x)f\|_A = \|f\|_A$ .

**THEOREM 9.** *Let  $G$  be a compact (nonabelian) group, and let  $\mu \in M(G)$ . Then  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $(M(G), \|\cdot\|_\infty)$  if and only if  $\mu \in M_0(G)$ .*

*Proof.* For  $\mu \in M(G)$  and  $f \in A(G)$ , we define

$$\langle f, \mu \rangle = \int_G f(t) d\mu(t) = \operatorname{Tr}(\hat{\mu}\hat{h}),$$

where  $h(t) = f(t^{-1})$ . If  $\check{f}$  is defined by  $\check{f}(t) = f(t^{-1})$ , then  $\|\check{f}\|_A = \|f\|_A$ . Thus  $\langle \check{f}, \mu \rangle = \operatorname{Tr}(\hat{\mu}\hat{f})$ . Let  $A_k = \{f \in A(G) : \|f\|_A < k\}$ , and let  $\mathcal{T}(A_k)$  be the topology on  $M(G)$  of uniform convergence on the sets  $A_k$ . Since

$$|\operatorname{Tr}(\hat{\mu}\hat{f})| \leq \|\hat{\mu}\|_\infty \|\hat{f}\|_1 = \|\hat{\mu}\|_\infty \|f\|_A = \|\mu\|_\infty \|f\|_A,$$

the topology  $\mathcal{T}(A_k)$  is weaker than the  $\|\cdot\|_\infty$ -topology on  $M(G)$ . However, since  $\mathcal{L}^\infty(\hat{G})$  is identified with the dual space of  $\mathcal{L}^1(\hat{G})$  by  $\psi \rightarrow \operatorname{Tr}(\phi\psi)$  for  $\phi \in \mathcal{L}^\infty(\hat{G})$  and  $\psi \in \mathcal{L}^1(\hat{G})$ ,  $\mathcal{T}(A_k)$  is the same as the  $\|\cdot\|_\infty$ -topology on  $M(G)$ . Furthermore,  $A_k$  is  $L(x)$ -invariant, since  $\|L(x)f\|_A = \|f\|_A$  for  $f \in A(G)$ . We now apply Theorems 3 and 4. ■

We conclude now with the general case. We shall use the machinery developed by P. Eymard [2], and we shall follow his conventions in the use of  $x$  in various formulae, where we used  $x^{-1}$  in the compact and abelian cases discussed above.

Let  $G$  be a locally compact group. Let  $\Sigma$  denote the equivalence classes of the continuous unitary representations on  $G$ . For  $\pi \in \Sigma$ , let  $\mathcal{H}_\pi$  denote the representation space. We define  $\hat{\mu}$  to be a function on  $\Sigma$  by  $\pi \mapsto \hat{\mu}(\pi) = \int_G \pi(x) d\mu(x)$ . For  $\mathcal{S} \subset \Sigma$ , let

$$\|\mu\|_{\mathcal{S}} = \sup \{ \|\hat{\mu}(\pi)\|_\infty : \pi \in \mathcal{S} \},$$

where  $\|\hat{\mu}(\pi)\|_\infty$  denotes the operator norm on  $\mathcal{H}_\pi$ . We define  $C^*(G)$  to be the completion of  $L^1(G)$  in  $\|\cdot\|_\Sigma$  (see [2, Section 1.14]). Let  $\{\rho\}$  denote the subset of  $\Sigma$  containing just the left regular representation of  $G$  on  $L^2(G)$ . Let  $C_\rho^*(G)$  denote the completion of  $L^1(G)$  in  $\|\cdot\|_\rho$  (see [2, Section 1.16]).

For  $\mu \in M(G)$ , we let  $\rho(\mu)$  denote the bounded operator on  $L^2(G)$ , defined by  $h \mapsto \mu * h$  ( $h \in L^2(G)$ ), with operator norm  $\|\rho(\mu)\|_\rho$ . Let  $\mathcal{B}(L^2(G))$  denote the set of bounded operators on  $L^2(G)$ . Then  $C_\rho^*(G)$  can be identified with the closure of  $\rho(L^1(G)) = \{\rho(f) : f \in L^1(G)\}$  in  $\mathcal{B}(L^2(G))$ . If  $G$  is abelian, then  $C_\rho^*(G) = C_0(G)$ . If  $G$  is compact, then  $C^*(G) = \mathcal{C}_0(\hat{G})$ .

Let  $VN(G)$  denote the von Neumann subalgebra of  $\mathcal{B}(L^2(G))$  generated by the left translation operators (see [2, Section 3.9]). For  $\mu \in M(G)$ , we have that  $\rho(\mu) \in VN(G)$ . Further,  $C_\rho^*(G) \subset VN(G)$ . If  $G$  is abelian, then  $VN(G) = L^\infty(\hat{G})$ . If  $G$  is compact, then  $VN(G) = \mathcal{L}^\infty(\hat{G})$ .

*Definition.*  $M_0(G) = \{\mu \in M(G): \rho(\mu) \in C_\rho^*(G)\}$ .

Let  $B(G)$  denote the linear subspace of  $C^B(G)$  generated by the continuous positive-definite functions. Then  $B(G)$  can be identified with the dual space of  $C^*(G)$  (see [2, Section 2.2]). For  $f \in B(G)$ , let  $\|f\|_B$  denote the norm of  $f$  as a linear functional on  $C^*(G)$ . Finally, let  $A(G)$  be the closed subalgebra of  $B(G)$  generated by the continuous positive-definite functions with compact support (see [2, Section 3.4]). If  $G$  is abelian, then  $A(G) = L^1(\hat{G})^\wedge$ . If  $G$  is compact, then our previous definitions and those of Eymard are consistent. We have the inclusion  $A(G) \subset C_{ru}^B(G)$ , since  $A(G) \subset C_0(G)$ . We let  $A_k = \{f \in A(G): \|f\|_B < k\}$ . Now for  $f \in A(G)$ ,  $\|L(x)f\|_B = \|f\|_B$ ; hence each  $A_k$  is  $L(x)$ -invariant. We pair  $A(G)$  and  $M(G)$  by the relation

$$\langle f, \mu \rangle = \int_G f(t) d\mu(t) \quad (f \in A(G) \text{ and } \mu \in M(G)).$$

Let  $\mathcal{T}(A_k)$  be the topology on  $M(G)$  of uniform convergence on the sets  $A_k$ . We wish to apply Theorems 3 and 4 as we did in Theorem 9. To do this, it remains only to observe that  $VN(G)$  can be identified as the dual space of  $A(G)$  (see [2, Section 3.10]), and for  $\mu \in M(G)$ , the identification is given by the relation

$$f \mapsto \int_G f(x) d\mu(x) = \langle f, \mu \rangle,$$

where  $f \in A(G)$ . It follows now by Theorems 3 and 4 that  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $(M(G), \|\cdot\|_\rho)$  if and only if  $\rho(\mu) \in \rho(L^1(\overline{G}))$  (the closure in  $\mathcal{B}(L^2(G))$ ). Hence we have the following result.

**THEOREM 10.** *Let  $G$  be a locally compact group. Let  $\mu \in M(G)$ . Then  $x \mapsto L(x)\mu$  is continuous from  $G$  to  $(M(G), \|\cdot\|_\rho)$  if and only if  $\mu \in M_0(G)$ .*

**THEOREM 11.** *Suppose  $A \subset C_{ru}^B(G)$  has the further property that  $A$  is dense in  $L^1(|\mu|)$  for each  $\mu \in M(G)$ , and that for each  $f \in A$  we have inclusions  $fA_k \subset CA_k$ , ( $k = 1, 2, \dots$ ), where the constants  $C$  and  $k'$  depend on  $f$  and on  $k$ . Then  $L^1(\overline{G})^A$  is a band; in other words, if  $\mu \in L^1(\overline{G})^A$  and  $\nu \ll \mu$ , then  $\nu \in L^1(\overline{G})^A$ .*

*Proof.* Let  $\mu \in L^1(\overline{G})^A$  and  $\nu \ll \mu$ ; then  $d\nu = g d\mu$ , for some Borel function  $g \in L^1(|\mu|)$ . Now there exist functions  $f_m \in A$  ( $m = 1, 2, \dots$ ) such that

$$\int_G |f_m - g| d|\mu| < 1/m,$$

that is,  $\|f_m d\mu - d\nu\|_{M(G)} \rightarrow 0$  as  $m \rightarrow \infty$ . We claim that each  $f_m d\mu$  belongs to  $L^1(\overline{G})^A$ . For if  $\{g_n\} \subset L^1(G)$  and  $g_n \xrightarrow{n} \mu$  in  $\mathcal{T}(A_k)$ , then  $f_m g_n \xrightarrow{n} f_m d\mu$  (note that  $f_m g_n \in L^1(G)$ ). In fact, for each  $k$ , we have the relations



$$\begin{aligned} \tau_k(f_m g_n - f_m d\mu) &= \sup \left\{ \left| \int_G \phi(x) f_m(x) [g_n(x) dx - d\mu(x)] \right| : \phi \in A_k \right\} \\ &\leq C \sup \left\{ \left| \int_G \phi(x) [g_n(x) dx - d\mu(x)] \right| : \phi \in A_{k'} \right\} = C \tau_{k'}(g_n - d\mu), \end{aligned}$$

where  $C$  and  $k'$  depend on  $k$  and  $f_m$ . Thus  $\tau_k(f_m g_n - f_m d\mu) \xrightarrow{n} 0$ , and  $f_m d\mu \in L^1(\overline{G})^A$ .

Since  $\mathcal{T}(A_k)$ -closed sets are closed in the measure norm topology ( $\sup \{ \|\phi\|_\infty : \phi \in A_k \} < \infty$ ), we have that  $\nu \in L^1(\overline{G})^A$ . ■

**COROLLARY 12.** *For every locally compact group  $G$ ,  $M_0(G)$  is a band.*

*Proof.* Let  $A = A(G)$  as before, and recall that  $A(G)$  is a dense subalgebra of  $C_0(G)$  (for the locally compact case, see [2, Section 3.4]). ■

#### REFERENCES

1. C. Dunkl and D. Ramirez, *Topics in harmonic analysis* (to appear).
2. P. Eymard, *L'algèbre de Fourier d'un groupe localement compact*. Bull. Soc. Math. France 92 (1964), 181-236.
3. R. R. Goldberg and A. B. Simon, *Characterization of some classes of measures*. Acta Sci. Math. (Szeged) 27 (1966), 157-161.
4. K. H. Hofmann and P. S. Mostert, *Elements of compact semigroups*. Charles E. Merrill Books, Columbus, Ohio, 1966.
5. R. Larsen, *Measures with separable orbits*. Proc. Amer. Math. Soc. 19 (1968), 569-572.
6. L. S. Pontryagin, *Topological groups*. Gordon and Breach, New York, 1966.
7. W. Rudin, *Measure algebras on abelian groups*. Bull. Amer. Math. Soc. 65 (1959), 227-247.
8. K. Shiga, *Representations of a compact group on a Banach space*. J. Math. Soc. Japan 7 (1955), 224-248.
9. K. W. Tam, *On measures with separable orbit*. Proc. Amer. Math. Soc. 23 (1969), 409-411.

University of Virginia  
Charlottesville, Virginia 22901

