



Banach J. Math. Anal. 1 (2007), no. 1, 85–90

**BANACH JOURNAL OF MATHEMATICAL ANALYSIS**

ISSN: 1735-8787 (electronic)

<http://www.math-analysis.org>

## A SPECIAL GAUSSIAN RULE FOR TRIGONOMETRIC POLYNOMIALS

GRADIMIR V. MILOVANOVIĆ<sup>1\*</sup>, ALEKSANDAR S. CVETKOVIĆ<sup>2</sup> AND MARIJA P. STANIĆ<sup>3</sup>

*This paper is dedicated to Professor Themistocles M. Rassias.*

Submitted by S. S. Dragomir

ABSTRACT. Abram Haimovich Turetzkii [*Uchenye Zapiski*, **1** (149) (1959), 31–55 (translation in English in *East J. Approx.* **11** (2005), 337–359)] considered interpolatory quadrature rules which have the following form  $\int_0^{2\pi} f(x)w(x)dx \approx \sum_{\nu=0}^{2n} w_{\nu}f(x_{\nu})$ , and which are exact for all trigonometric polynomials of degree less than or equal to  $n$ . Maximal trigonometric degree of exactness of such quadratures is  $2n$ , and such kind of quadratures are known as *quadratures of Gaussian type* or *Gaussian quadratures for trigonometric polynomials*. In this paper we prove some interesting properties of a special Gaussian quadrature with respect to the weight function  $w_m(x) = 1 + \sin mx$ , where  $m$  is a positive integer.

### 1. INTRODUCTION

This paper is dedicated to Themistocles M. Rassias for our long and fruitful collaboration in mathematical research, including the writing of our book [4], published in 1994 by World Scientific Publishing Co., jointly with Professor Dragoslav S. Mitrinović.

Let the weight function  $w(x)$  be integrable and nonnegative on the interval  $[0, 2\pi)$ , vanishing there only on a set of a measure zero.

*Date:* Received: 24 March 2007; Accepted: 24 October 2007.

\* Corresponding author.

2000 *Mathematics Subject Classification.* Primary 65D30; Secondary 42A05.

*Key words and phrases.* Trigonometric polynomials of semi-integer degree, Orthogonality, Gaussian type quadratures.

By  $\mathcal{T}_n$  we denote the linear space of all trigonometric polynomials of degree less than or equal to  $n$ , and by  $\mathcal{T}_n^{1/2}$  we denote the linear span of the following set  $\{\cos(\nu + 1/2)x, \sin(\nu + 1/2)x, \nu = 0, 1, \dots, n\}$ . Trigonometric functions from  $\mathcal{T}_n^{1/2}$ , i.e., trigonometric functions of the following form

$$\sum_{\nu=0}^n \left[ c_\nu \cos\left(\nu + \frac{1}{2}\right)x + d_\nu \sin\left(\nu + \frac{1}{2}\right)x \right], \quad (1.1)$$

where  $c_\nu, d_\nu \in \mathbb{R}$ ,  $|c_n| + |d_n| \neq 0$ , are known as trigonometric polynomials of semi-integer degree  $n + 1/2$  (see [5], [2]). This kind of trigonometric functions are important for construction of the Gaussian quadrature rules for trigonometric polynomials.

It is obvious that

$$A_{n+1/2}(x) = A \prod_{k=0}^{2n} \sin \frac{x - x_k}{2} \quad (A \text{ is a non-zero constant}), \quad (1.2)$$

is trigonometric polynomial of semi-integer degree  $n + 1/2$ . Also, every trigonometric polynomial of semi-integer degree  $n + 1/2$  of the form (1.1) can be represented in the form (1.2) (see [5, Lemma 1]).

For any positive integer  $n$ , the quadrature rule of the Gaussian type is the following one

$$\int_0^{2\pi} t(x)w(x)dx = \sum_{\nu=0}^{2n} w_\nu t(x_\nu), \quad t \in \mathcal{T}_{2n}, \quad (1.3)$$

where weights  $w_\nu$  are given by

$$w_\nu = \int_0^{2\pi} \frac{A_{n+1/2}(x)}{2 \sin \frac{x - x_\nu}{2} A'_{n+1/2}(x_\nu)} w(x) dx, \quad \nu = 0, 1, \dots, 2n, \quad (1.4)$$

and nodes are zeros of  $A_{n+1/2}(x)$ , which is orthogonal on  $[0, 2\pi)$  with respect to the weight function  $w(x)$  to every trigonometric polynomial of the semi-integer degree from  $\mathcal{T}_{n-1}^{1/2}$  (see [5], [2]). It is known that such orthogonal trigonometric polynomial of semi-integer degree  $A_{n+1/2}(x)$  has in  $[0, 2\pi)$  exactly  $2n + 1$  distinct simple zeros (see [5, Theorem 3]) and  $A_{n+1/2}$  with given leading coefficients  $c_n$  and  $d_n$  is uniquely determined (see [5, §3.]). In [2] and [3] two choices of leading coefficients were considered. For the choice  $c_n = 1, d_n = 0$ , we denote an orthogonal trigonometric polynomial of semi-integer degree by  $A_{n+1/2}^C$ , and for the choice  $c_n = 0$  and  $d_n = 1$  by  $A_{n+1/2}^S$ .

The Gaussian quadrature (1.3) is not given uniquely since it is possible to use any orthogonal polynomial of semi-integer degree  $A_{n+1/2} \in \mathcal{T}_n^{1/2}$ . In [2] a numerical method for constructing the Gaussian quadratures using  $A_{n+1/2}^C$  is given. That method is based on five-term recurrence relations for  $A_{n+1/2}^C$  and  $A_{n+1/2}^S$ . The main problem in that method is the calculation of five-term recurrence coefficients. Explicit formulas for five-term recurrence coefficients for some weight functions are given in [3].

In this paper, in Section 2, we prove some interesting properties for Gaussian quadratures (1.3) with respect to the weight function  $w_m(x) = 1 + \sin mx$ ,  $m \in \mathbb{N}$ , and in Section 3 we give a numerical example.

## 2. MAIN RESULTS

As an example, the nodes  $x_\nu$  and weight  $w_\nu$ ,  $\nu = 0, 1, \dots, 2n$ , for the Gaussian quadrature for  $n = 25$  and  $w_{15}(x) = 1 + \sin 15x$  are given in [2]. In that case we have seen some kind of symmetry, since  $w_\nu = w_{\nu+17j}$  and  $x_{\nu+17j} = x_\nu + 2j\pi/3$ ,  $j = 1, 2$ ,  $\nu = 0, 1, \dots, 16$ . This symmetry is not an isolated case, namely Gaussian quadratures (1.3) with respect to the weight functions  $w_m(x) = 1 + \sin mx$ ,  $m \in \mathbb{N}$ , have interesting properties, which are presented in Theorem 2.1. If we want to construct a Gaussian quadrature rule with  $2n_1 + 1$  nodes for the weight function  $w_{m_1}(x) = 1 + \sin m_1x$  where  $\gcd(m_1, 2n_1 + 1) = d \neq 1$  we can obtain nodes and weights for such quadrature directly from nodes and weights of the Gaussian quadrature rule with  $(2n_1 + 1)/d$  nodes with respect to the weight  $w_{m_1/d}$ . In a given example in [2] we have  $n_1 = 25$ ,  $m_1 = 15$  and  $\gcd(15, 51) = 3$ .

**Theorem 2.1.** *Let denote by  $x_\nu, w_\nu$ ,  $\nu = 1, \dots, 2n + 1$ , the nodes and weights for the Gaussian quadrature rule with respect to the weight function  $w_m(x) = 1 + \sin mx$ . Then*

$$\widehat{x}_{j(2n+1)+\nu} = \frac{x_\nu}{q} + \frac{2j\pi}{q}, \quad \widehat{w}_{j(2n+1)+\nu} = \frac{w_\nu}{q},$$

for  $j = 0, 1, \dots, q - 1$ ,  $\nu = 1, \dots, 2n + 1$ , are nodes and weights for the Gaussian quadrature rule with  $(2n + 1)q$  nodes with respect to the weight function  $w_{mq}(x) = 1 + \sin mqx$ , where  $q$  is an odd positive integer.

*Proof.* According to orthogonality conditions for trigonometric polynomials of semi-integer degree with respect to the weight function  $w_m(x) = 1 + \sin mx$ , we obtain

$$\int_0^{2\pi} A_{n+1/2}^C(x) \cos(k + 1/2)x(1 + \sin mx)dx = 0, \quad 0 \leq k \leq n - 1,$$

$$\int_0^{2\pi} A_{n+1/2}^C(x) \sin(k + 1/2)x(1 + \sin mx)dx = 0, \quad 0 \leq k \leq n - 1.$$

If we introduce  $x := qx$ , we have

$$\int_0^{2\pi/q} A_{n+1/2}^C(qx) \cos(k + 1/2)qx(1 + \sin mqx)dx = 0, \quad 0 \leq k \leq n - 1,$$

$$\int_0^{2\pi/q} A_{n+1/2}^C(qx) \sin(k + 1/2)qx(1 + \sin mqx)dx = 0, \quad 0 \leq k \leq n - 1.$$

Substituting  $t = x + 2j\pi/q$ ,  $j = 1, \dots, q - 1$ , we get

$$\int_{\frac{2j\pi}{q}}^{\frac{2(j+1)\pi}{q}} A_{n+1/2}^C(qt - 2j\pi) \cos(k + 1/2)(qt - 2j\pi)(1 + \sin m(qt - 2j\pi))dt = 0,$$

$$\int_{\frac{2j\pi}{q}}^{\frac{2(j+1)\pi}{q}} A_{n+1/2}^C(qt - 2j\pi) \sin(k + 1/2)(qt - 2j\pi)(1 + \sin m(qt - 2j\pi)) dt = 0,$$

for  $0 \leq k \leq n - 1$ , i.e.,

$$\int_{2j\pi/q}^{2(j+1)\pi/q} A_{n+1/2}^C(qt) \cos(k + 1/2)qt(1 + \sin mqt) dt = 0,$$

$$\int_{2j\pi/q}^{2(j+1)\pi/q} A_{n+1/2}^C(qt) \sin(k + 1/2)qt(1 + \sin mqt) dt = 0,$$

because of

$$\cos(k + 1/2)(qt - 2j\pi) = (-1)^j \cos(k + 1/2)qt$$

and

$$\sin(k + 1/2)(qt - 2j\pi) = (-1)^j \sin(k + 1/2)qt,$$

for  $j = 1, \dots, q - 1$ .

Thus, we have

$$\int_0^{2\pi} A_{n+1/2}^C(qx) \cos(k + 1/2)qx(1 + \sin mqx) dx = 0, \quad 0 \leq k \leq n - 1,$$

$$\int_0^{2\pi} A_{n+1/2}^C(qx) \sin(k + 1/2)qx(1 + \sin mqx) dx = 0, \quad 0 \leq k \leq n - 1.$$

Let denote by  $\mathcal{J}^q$  the linear span of the following functions

$$\cos(k + 1/2)qx = \cos(kq + (q - 1)/2 + 1/2)x, \quad k = 0, 1, \dots, n,$$

$$\sin(k + 1/2)qx = \sin(kq + (q - 1)/2 + 1/2)x, \quad k = 0, 1, \dots, n.$$

Obviously,  $\mathcal{J}^q \subseteq \mathcal{J}_{nq+(q-1)/2}^{1/2}$ . Using integrals

$$\int_0^{2\pi} \cos(\nu + 1/2)x \cos(\ell + 1/2)x w_{qm}(x) dx = \pi \delta_{\nu, \ell},$$

$$\int_0^{2\pi} \sin(\nu + 1/2)x \sin(\ell + 1/2)x w_{qm}(x) dx = \pi \delta_{\nu, \ell},$$

$$\int_0^{2\pi} \cos(\nu + 1/2)x \sin(\ell + 1/2)x w_{qm}(x) dx = \frac{\pi}{2}(\delta_{\nu, \ell - qm} + \delta_{\nu, qm - \ell - 1} - \delta_{\nu, \ell + qm}),$$

for  $\nu = kq + (q - 1)/2$ , we see that integrals might not be equal zero only under condition  $\ell = iq + (q - 1)/2$ ,  $i \in \mathbb{Z}$ . With this argument we obtain the following orthogonality  $\mathcal{J}^q \perp (\mathcal{J}_{nq+(q-1)/2}^{1/2} \ominus \mathcal{J}^q)$  with respect to the inner product

$$(f, g) = \int_0^{2\pi} f(x)g(x)w_{mq}(x)dx.$$

Then,  $A_{n+1/2}^C(qx)$  is orthogonal on  $[0, 2\pi)$  to all trigonometric polynomials of semi-integer degree  $t \in \mathcal{J}_{nq+(q-1)/2-1}^{1/2}$  with respect to the weight function  $w_{mq}(x)$ .

Since  $A_{n+1/2}^C(x)$  can be represented as

$$A_{n+1/2}^C(x) = A \prod_{\nu=1}^{2n+1} \sin \frac{x - x_\nu}{2}, \quad A \neq 0,$$

we get

$$A_{n+1/2}^C(qx) = A \prod_{\nu=1}^{2n+1} \sin \frac{qx - x_\nu}{2} = A \prod_{\nu=1}^{2n+1} \sin \frac{q(x - x_\nu/q)}{2},$$

so it is obvious that  $\hat{x}_{j(2n+1)+\nu}$ ,  $j = 0, 1, \dots, q-1$ ,  $\nu = 1, \dots, 2n+1$ , are zeros of  $A_{n+1/2}^C(qx)$ .

Formulas for weight coefficients  $\hat{w}_{j(2n+1)+\nu}$ ,  $j = 0, 1, \dots, q-1$ ,  $\nu = 1, \dots, 2n+1$ , can be easily obtained using (1.4).  $\square$

### 3. NUMERICAL EXAMPLE

We determine the nodes and weights for a Gaussian quadrature with 75 nodes and the weight function  $w_{45}(x) = 1 + \sin 45x$ .

Since  $\gcd(45, 75) = 15$ , according to Theorem 2.1, we can obtain parameters for this Gaussian quadrature directly from the nodes and weights of a Gaussian quadrature with 5 nodes and the weight function  $w_3(x) = 1 + \sin 3x$ .

The parameters  $x_\nu$  and weight  $w_\nu$ ,  $\nu = 0, 1, \dots, 2n$ , for  $n = 2$  and  $w(x) = 1 + \sin 3x$  are given in Table 1. (Numbers in parentheses indicate decimal exponents.) We calculate these parameters using a numerical method described in [2] with explicit formulas for five-term recurrence coefficients given in [3]. All computations are performed in double precision arithmetic (16 decimal digits mantissa) in MATHEMATICA, using the corresponding software package described in [1].

Now it is easy to obtain the nodes  $\hat{x}_\nu$  and the weights  $\hat{w}_\nu$ ,  $\nu = 0, 1, \dots, 74$ , of the Gaussian quadrature for the  $w_{45}(x) = 1 + \sin 45x$ . The nodes  $\hat{x}_\nu$ ,  $\nu = 0, 1, \dots, 4$ , and the weights  $\hat{w}_k$ ,  $k = 0, 1, \dots, 74$ , are given in Table 2. The other nodes  $x_k$ ,  $k = 5, 6, \dots, 74$ , are given by

$$\hat{x}_{5j+\nu} = \hat{x}_\nu + \frac{2j\pi}{15}, \quad j = 1, \dots, 14, \nu = 0, 1, \dots, 4.$$

TABLE 1. Nodes  $x_\nu$  and weights  $w_\nu$ ,  $\nu = 0, 1, \dots, 4$ , for  $w_3(x) = 1 + \sin 3x$

$\nu$	$x_\nu$	$w_\nu$
0	5.717322718269771(-1)	1.700722884785417
1	2.062284642241416	9.534684415662144(-1)
2	2.933402103642034	1.316125839231413
3	4.554306420567181	1.586274703648100
4	5.586237829671358	7.265934379484426(-1)

TABLE 2. Nodes  $\hat{x}_\nu$  and weights  $\hat{w}_{5j+\nu}$ ,  $\nu = 0, 1, \dots, 4$ ,  $j = 0, 1, \dots, 14$ , for  $n = 37$  and  $w_{45}(x) = 1 + \sin 45x$

$\nu$	$\hat{x}_\nu$	$\hat{w}_{5j+\nu}, j = 0, 1, \dots, 14$
0	0.3811548478846514(-1)	0.1133815256523611
1	0.1374856428160944	0.6356456277108096(-1)
2	0.1955601402428023	0.8774172261542751(-1)
3	0.3036204280378120	0.1057516469098733
4	0.3724158553114239	0.4843956252989617(-1)

**Acknowledgements:** The authors were supported in part by the Serbian Ministry of Science (Project: *Orthogonal Systems and Applications*, grant number #144004G) and the Swiss National Science Foundation (SCOPEs Joint Research Project No. IB7320–111079 “*New Methods for Quadrature*”)

#### REFERENCES

1. A.S. Cvetković and G.V. Milovanović, *The Mathematica Package “OrthogonalPolynomials”*, Facta Univ. Ser. Math. Inform. **19** (2004), 17–36.
2. G.V. Milovanović, A.S. Cvetković and M.P. Stanić, *Trigonometric orthogonal systems and quadrature formulae*, preprint.
3. G.V. Milovanović, A.S. Cvetković and M.P. Stanić, *Explicit formulas for five-term recurrence coefficients of orthogonal trigonometric polynomials of semi-integer degree*, Appl. Math. Comput. (to appear).
4. G.V. Milovanović, D.S. Mitrinović and Th.M. Rassias, *Topics in Polynomials: Extremal Problems, Inequalities, Zeros*, World Scientific, Singapore–New Jersey–London–Hong Kong, 1994.
5. A.H. Turetzkii, *On quadrature formulae that are exact for trigonometric polynomials*, East J. Approx. **11** (2005), 337–359 (translation in English from Uchenye Zapiski, Vypusk 1 (149), Seria math. Theory of Functions, Collection of papers, Izdatel'stvo Belgosuniversiteta imeni V.I. Lenina, Minsk (1959), 31–54).

<sup>1</sup> DEPARTMENT OF MATHEMATICS, FACULTY OF ELECTRONIC ENGINEERING, UNIVERSITY OF NIŠ, P.O. BOX 73, 18000 NIŠ, SERBIA.  
E-mail address: [grade@junis.ni.ac.yu](mailto:grade@junis.ni.ac.yu)

<sup>2</sup> DEPARTMENT OF MATHEMATICS, FACULTY OF ELECTRONIC ENGINEERING, UNIVERSITY OF NIŠ, P.O. BOX 73, 18000 NIŠ, SERBIA.  
E-mail address: [aca@elfak.ni.ac.yu](mailto:aca@elfak.ni.ac.yu)

<sup>3</sup> DEPARTMENT OF MATHEMATICS AND INFORMATICS, FACULTY OF SCIENCE, UNIVERSITY OF KRAGUJEVAC, P.O. BOX 60, 34000 KRAGUJEVAC, SERBIA.  
E-mail address: [stanicm@kg.ac.yu](mailto:stanicm@kg.ac.yu)