

# *Algebra & Number Theory*

Volume 12

2018

No. 2

**Sparsity of  $p$ -divisible unramified liftings for subvarieties of abelian varieties with trivial stabilizer**

Danny Scarponi



# Sparsity of $p$ -divisible unramified liftings for subvarieties of abelian varieties with trivial stabilizer

Danny Scarponi

By means of the theory of strongly semistable sheaves and the theory of the Greenberg transform, we generalize to higher dimensions a result on the sparsity of  $p$ -divisible unramified liftings which played a crucial role in Raynaud's proof of the Manin–Mumford conjecture for curves. We also give a bound for the number of irreducible components of the first critical scheme of subvarieties of an abelian variety which are complete intersections.

## 1. Introduction

The Manin–Mumford conjecture is a significant question concerning the intersection of a subvariety  $X$  of an abelian variety  $A$  with the group of torsion points of  $A$ . Raised independently by Manin and Mumford, the conjecture was originally formulated in the case of curves. Suppose that  $A$  is an abelian variety over a number field  $K$  and that  $C$  is a smooth subcurve of  $A$  of genus at least two. Then only finitely many torsion points of  $A(\bar{K})$  lie in  $C$ . In 1983, Raynaud proved this conjecture and generalized it to higher dimensions: if  $A/K$  is as above and  $X/K$  is a smooth subvariety of  $A$  which does not contain any translate of a nontrivial abelian subvariety, then the set of torsion points of  $A(\bar{K})$  lying in  $X$  is finite [Raynaud 1983b; 1983c].

Let us fix  $K$ ,  $X$  and  $A$  as above. Let  $U$  be a nonempty open subscheme of  $\text{Spec } \mathcal{O}_K$  not containing any ramified primes and such that  $A/K$  extends to an abelian scheme  $\mathcal{A}/U$  and  $X$  extends to a smooth closed integral subscheme  $\mathcal{X}$  of  $\mathcal{A}$ . For any  $\mathfrak{p} \in U$ , let  $R$  and  $R_n$  be the ring of Witt vectors and Witt vectors of length  $n + 1$ , respectively, with coordinates in the algebraic closure  $\overline{k(\mathfrak{p})}$  of the residue field of  $\mathfrak{p}$ . Recall that  $R$  is a DVR with maximal ideal generated by  $p$  such that  $R_0 = R/p = \overline{k(\mathfrak{p})}$ . Denote by  $X_{\mathfrak{p}^n}$  and  $A_{\mathfrak{p}^n}$  the  $R_n$ -schemes  $\mathcal{X} \times_U \text{Spec } R_n$  and  $\mathcal{A} \times_U \text{Spec } R_n$ , respectively, and consider the reduction map

$$pA_{\mathfrak{p}^1}(R_1) \cap X_{\mathfrak{p}^1}(R_1) \rightarrow X_{\mathfrak{p}^0}(R_0). \quad (1)$$

In [Raynaud 1983b] it was shown that, if  $X$  is a curve, the image of (1) is not Zariski dense in  $X_{\mathfrak{p}^0}$ , i.e., it is a finite set. This local result is crucial in Raynaud's proof of the Manin–Mumford conjecture for curves, since it easily implies that only finitely many prime-to- $p$  torsion points of  $A(\bar{K})$  lie on  $X$  [Raynaud 1983b, Théorème II].

*MSC2010:* primary 14K12; secondary 14K15.

*Keywords:* Manin–Mumford conjecture, number fields,  $p$ -divisible unramified liftings, Greenberg transform, strongly semistable sheaves.

It is quite natural to expect that a similar result also holds in higher dimensions. More explicitly, one can ask: is it true that, if a smooth subvariety  $X$  of  $A$  does not contain any translate of a nontrivial abelian subvariety, the image of (1) is not Zariski dense? In this paper we give a positive answer to this question (see [Theorem 5.3](#)).

**Theorem 1.1** (sparsity of  $p$ -divisible unramified liftings). *Suppose that  $X$  has trivial stabilizer. For all  $\mathfrak{p} \in U$  above a prime  $p > (\dim X)^2 \deg(\Omega_X)$  such that  $X_{\mathfrak{p}^0}$  has trivial stabilizer, the image of*

$$pA_{\mathfrak{p}^1}(R_1) \cap X_{\mathfrak{p}^1}(R_1) \rightarrow X_{\mathfrak{p}^0}(R_0)$$

is not Zariski dense in  $X_{\mathfrak{p}^0}$ .

Here  $\deg(\Omega_X)$  refers to the degree of the cotangent bundle  $\Omega_X$  computed with respect to any fixed very ample line bundle on  $X$ .

Notice that if  $X$  does not contain any translate of a nontrivial abelian subvariety, then it has finite stabilizer. Therefore, replacing  $A$  and  $X$  with their quotients by the stabilizer of  $X$ , one can assume the stabilizer is trivial (see the beginning of the next section for the definition of stabilizer).

A different generalization of Raynaud’s local result was given by Rössler [\[2013\]](#) who proved that, if the torsion points of  $\mathcal{A}(\overline{\text{Frac}(R)})$  are not dense in  $\mathcal{X}(\overline{\text{Frac}(R)})$ , then for  $m$  big enough the image of

$$p^m A_{\mathfrak{p}^m}(R_m) \cap X_{\mathfrak{p}^m}(R_m) \rightarrow X_{\mathfrak{p}^0}(R_0) \tag{2}$$

is not Zariski dense in  $X_{\mathfrak{p}^0}$  [\[Rössler 2013, Theorem 4.1\]](#). [Theorem 1.1](#) makes Rössler’s result effective, showing that if the stabilizer of  $X$  is trivial, then it is sufficient to consider the map (2) for  $m = 1$ .

The proof of [Theorem 1.1](#) strongly relies on Rössler’s paper [\[2016\]](#) and is done by contradiction. First we use some basic properties of the Greenberg transform to show that, if the image of (1) is Zariski dense in  $X_{\mathfrak{p}^0}$ , the absolute Frobenius  $F_{X_{\mathfrak{p}^0}} : X_{\mathfrak{p}^0} \rightarrow X_{\mathfrak{p}^0}$  lifts to an endomorphism of  $X_{\mathfrak{p}^1}$ . A well-known consequence of this liftability is the existence of a map of sheaves of differentials  $F_{X_{\mathfrak{p}^0}}^* \Omega_{X_{\mathfrak{p}^0}} \rightarrow \Omega_{X_{\mathfrak{p}^0}}$  which is nonzero. If  $X$  is a curve, such a map cannot exist, since  $\deg(F_{X_{\mathfrak{p}^0}}^* \Omega_{X_{\mathfrak{p}^0}})$  is strictly bigger than  $\deg(\Omega_{X_{\mathfrak{p}^0}})$ . This simple observation was in fact used by Raynaud to prove Lemma I.5.4 in [\[Raynaud 1983a\]](#). By means of the theory of strongly semistable sheaves developed by Rössler [\[2016\]](#), we show that when  $X$  has dimension higher than one, there are no nontrivial maps from  $F_{X_{\mathfrak{p}^0}}^* \Omega_{X_{\mathfrak{p}^0}}$  to  $\Omega_{X_{\mathfrak{p}^0}}$ . This gives us the wanted contradiction.

In the last section of this paper, we consider subvarieties of abelian varieties which are complete intersections. If  $\text{Gr}_1$  denotes the Greenberg transform of level 1 (see [Section 3](#)), then we know that the first critical scheme

$$\text{Crit}^1(\mathcal{X}, \mathcal{A}) := [p]_* \text{Gr}_1(A_{\mathfrak{p}^1}) \cap \text{Gr}_1(X_{\mathfrak{p}^1})$$

is a scheme over  $R_0$  such that

$$\text{Crit}^1(\mathcal{X}, \mathcal{A})(R_0) = pA_{\mathfrak{p}^1}(R_1) \cap X_{\mathfrak{p}^1}(R_1).$$

Using exactly the same technique that allowed Buium [1996] to give an effective form of the Manin–Mumford conjecture in the case of curves, we get a bound for the number of irreducible components of  $\text{Crit}^1(\mathcal{X}, \mathcal{A})$  when  $X$  is a complete intersection (not necessarily with trivial stabilizer).

**Theorem 1.2.** *Let  $K$  be a number field,  $A/K$  be an abelian variety of dimension  $n$  and let  $L$  be a very ample line bundle on  $A$ . Let  $c \in \mathbb{N}$  be positive and let  $H_1, H_2, \dots, H_c \in |L|$  be general. Suppose that  $X := H_1 \cap H_2 \cap \dots \cap H_c$  is smooth. There exists a nonempty open subscheme  $V \subseteq \text{Spec } \mathcal{O}_K$  (see the beginning of Section 6 for its definition) such that if  $\mathfrak{p} \in V$ , the number of irreducible components of  $\text{Crit}^1(\mathcal{X}, \mathcal{A})$  is bounded by*

$$p^{2n} \left( \sum_{h=0}^{n-c} \binom{2n-2c}{h} \binom{c}{n-c-h} p^{n-c-h} \right) (L^n)^2.$$

Here  $(L^n)$  denotes the intersection number of  $L$ .

We conclude the introduction with the following remark. Since the field of definition of points in the prime-to- $p$  torsion  $\text{Tor}^p(A(\bar{K}))$  is unramified at  $\mathfrak{p}$  and the specialization map  $\mathcal{A}(R) \rightarrow A_{\mathfrak{p}^1}(R_1)$  is injective on the prime-to- $p$  torsion, we have an injection

$$\text{Tor}^p(A(\bar{K})) \cap X(\bar{K}) \subseteq pA_{\mathfrak{p}^1}(R_1) \cap X_{\mathfrak{p}^1}(R_1).$$

This implies that, if  $X$  is a complete intersection such that  $\text{Crit}^1(\mathcal{X}, \mathcal{A})(R_0)$  is finite, then the bound in Theorem 1.2 is a bound for the cardinality of  $\text{Tor}^p(A(\bar{K})) \cap X(\bar{K})$ , i.e., an effective form of the Manin–Mumford conjecture for the prime-to- $p$  torsion.

## 2. Notations

We fix the following notations

- $K$  a number field,
- $\bar{K}$  an algebraic closure of  $K$ ,
- $A/K$  an abelian variety,
- $X \subseteq A$  a closed integral subscheme, smooth over  $K$ ,
- $\text{Stab}_A(X)$  the translation stabilizer of  $X$  in  $A$ , i.e., the closed subgroup scheme of  $A$  characterized uniquely by the fact that for any  $K$ -scheme  $S$  and any morphism  $b : S \rightarrow A$ , translation by  $b$  on the product  $A \times_K S$  maps the subscheme  $X \times_K S$  to itself if and only if  $b$  factors through  $\text{Stab}_A(X)$  (for its existence we refer the reader to [SGA 3 II 1970, Exemple 6.5(e), Expose VIII]),
- $U$  an open subscheme of  $\text{Spec } \mathcal{O}_K$  not containing any ramified prime and such that  $A/K$  extends to an abelian scheme  $\mathcal{A}/U$  and  $X$  extends to a smooth closed integral subscheme  $\mathcal{X}$  of  $\mathcal{A}$ .

For any prime number  $p$ , any unramified prime  $\mathfrak{p}$  of  $K$  above  $p$  and any  $n \geq 0$ , we denote by

- $k(\mathfrak{p})$  the residue field  $\mathcal{O}_K/\mathfrak{p}$  for  $\mathfrak{p}$ ,

- $K_{\mathfrak{p}}$  the completion of  $K$  with respect to  $\mathfrak{p}$ ,
- $\widehat{K}_{\mathfrak{p}}^{\text{unr}}$  the completion of the maximal unramified extension of  $K_{\mathfrak{p}}$ ,
- $R := W(\overline{k(\mathfrak{p})})$  and  $R_n := W_n(\overline{k(\mathfrak{p})})$  the ring of Witt vectors and the ring of Witt vectors of length  $n + 1$ , respectively, with coordinates in  $\overline{k(\mathfrak{p})}$ . We recall that  $R$  can be identified with the ring of integers of  $\widehat{K}_{\mathfrak{p}}^{\text{unr}}$  and  $R_0$  with  $\overline{k(\mathfrak{p})}$ ,
- $X_{\mathfrak{p}^n}$  the  $R_n$ -scheme  $\mathcal{X} \times_U \text{Spec } R_n$   $A_{\mathfrak{p}^n}$  the  $R_n$ -scheme  $\mathcal{A} \times_U \text{Spec } R_n$ .

### 3. The Greenberg transform and the critical schemes

Now we recall some basic facts about the Greenberg transform (for more details, see [Greenberg 1961; 1963; Bosch et al. 1990, pp. 276–277]).

Fix a prime number  $p$  and an unramified prime  $\mathfrak{p}$  of  $K$  above  $p$ .

For any  $n \geq 0$ , the Greenberg transform of level  $n$  is a covariant functor  $\text{Gr}_n$  from the category of  $R_n$ -schemes locally of finite type, to the category of  $R_0$ -schemes locally of finite type. If  $Y_n$  is an  $R_n$ -scheme locally of finite type,  $\text{Gr}_n(Y_n)$  is a  $R_0$ -scheme with the property

$$Y_n(R_n) = \text{Gr}_n(Y_n)(R_0).$$

More precisely, we can interpret  $R_n$  as the set of  $\overline{k(\mathfrak{p})}$ -valued points of a ring scheme  $\mathfrak{R}_n$  over  $\overline{k(\mathfrak{p})}$ . For any  $R_0$ -scheme  $T$ , we define  $\mathbb{W}_n(T)$  as the ringed space over  $R_n$  consisting of  $T$  as a topological space and of  $\text{Hom}_{R_0}(T, \mathfrak{R}_n)$  as a structure sheaf. By definition  $\text{Gr}_n(Y_n)$  represents the functor from the category of schemes over  $R_0$  to the category of sets given by

$$T \mapsto \text{Hom}_{R_n}(\mathbb{W}_n(T), Y_n)$$

where  $\text{Hom}$  stands for homomorphisms of ringed spaces. In other words, the functor  $\text{Gr}_n$  is right adjoint to the functor  $\mathbb{W}_n$ .

The functor  $\text{Gr}_n$  respects closed immersions, open immersions, fiber products, smooth, étale morphisms and is the identity for  $n = 0$ . Furthermore it sends group schemes over  $R_n$  to group schemes over  $R_0$ . The canonical morphism  $R_{n+1} \rightarrow R_n$  gives rise to a functorial transition morphism  $\pi_{n+1} : \text{Gr}_{n+1} \rightarrow \text{Gr}_n$ .

Let  $Y_n$  be a scheme over  $R_n$  locally of finite type. Then for any  $m < n$  we define

$$Y_m := Y_n \times_{R_n} R_m.$$

Let us call  $F_{Y_0} : Y_0 \rightarrow Y_0$  the absolute Frobenius endomorphism of  $Y_0$  and  $\Omega_{Y_0/R_0}$  the sheaf of relative differentials.

For any finite rank locally free sheaf  $\mathcal{F}$  over  $Y_0$  we will write

$$V(\mathcal{F}) := \underline{\text{Spec}}(\text{Sym}(\mathcal{F}^\vee))$$

for the vector bundle over  $Y_0$  associated to  $\mathcal{F}$ .

Suppose now that  $Y_n$  is smooth over  $R_n$ , so that  $\Omega_{Y_0/R_0}$  is locally free. A key result about the Greenberg transform is the following fact [Greenberg 1963, Section 2]:

$$\pi_1 : \mathrm{Gr}_1(Y_1) \rightarrow \mathrm{Gr}_0(Y_0) = Y_0$$

is a torsor under the Frobenius tangent bundle

$$V(F_{Y_0}^* \Omega_{Y_0/R_0}^\vee).$$

Let  $X, A, \mathcal{X}, \mathcal{A}$  and  $U$  be as fixed in the previous section and suppose that  $\mathfrak{p} \in U$ . We refer the reader to Section II.1 in [Raynaud 1983a] for more details on what we will recall from now till the end of the section. For any  $n \geq 0$ , the kernel of

$$\mathrm{Gr}_n(A_{\mathfrak{p}^n}) \rightarrow \mathrm{Gr}_0(A_{\mathfrak{p}^0}) = A_{\mathfrak{p}^0}$$

is unipotent, killed by  $p^n$ . Thus, the scheme-theoretic image  $[p^n]_* \mathrm{Gr}_n(A_{\mathfrak{p}^n})$  of multiplication by  $p^n$  in  $\mathrm{Gr}_n(A_{\mathfrak{p}^n})$  is the greatest abelian subvariety of  $\mathrm{Gr}_n(A_{\mathfrak{p}^n})$  and, since  $R_0$  is algebraically closed,  $[p^n]_* \mathrm{Gr}_n(A_{\mathfrak{p}^n})(R_0) = p^n \mathrm{Gr}_n(A_{\mathfrak{p}^n})(R_0)$ .

We define the  $n$ -critical scheme as

$$\mathrm{Crit}^n(\mathcal{X}, \mathcal{A}) := [p^n]_* \mathrm{Gr}_n(A_{\mathfrak{p}^n}) \cap \mathrm{Gr}_n(X_{\mathfrak{p}^n}).$$

Notice that  $\mathrm{Crit}^n(\mathcal{X}, \mathcal{A})$  is a scheme over  $R_0$  and that  $\mathrm{Crit}^0(\mathcal{X}, \mathcal{A}) = X_{\mathfrak{p}^0}$ .

The transition morphisms  $\pi_{n+1} : \mathrm{Gr}_{n+1}(A_{\mathfrak{p}^{n+1}}) \rightarrow \mathrm{Gr}_n(A_{\mathfrak{p}^n})$  lead to a projective system of  $R_0$ -schemes

$$\dots \rightarrow \mathrm{Crit}^2(\mathcal{X}, \mathcal{A}) \rightarrow \mathrm{Crit}^1(\mathcal{X}, \mathcal{A}) \rightarrow \mathrm{Crit}^0(\mathcal{X}, \mathcal{A}) = X_{\mathfrak{p}^0},$$

whose connecting morphisms are both affine and proper, hence finite. In fact, transition morphisms are affine and the subscheme  $[p^n]_* \mathrm{Gr}_n(A_{\mathfrak{p}^n})$  is proper, being the greatest abelian subvariety of  $\mathrm{Gr}_n(A_{\mathfrak{p}^n})$ .

We shall write  $\mathrm{Exc}^n(\mathcal{X}, \mathcal{A})$  for the scheme theoretic image of the morphism  $\mathrm{Crit}^n(\mathcal{X}, \mathcal{A}) \rightarrow X_{\mathfrak{p}^0}$ .

#### 4. The geometry of vector bundles in positive characteristic

In this section we recall some results on the geometry of vector bundles in positive characteristic by Langer [2004] and Rössler [2016]. These results will play a crucial role in the proof of Lemma 5.1 and Theorem 5.3.

Let us start with some basic definitions and facts regarding semistable sheaves in positive characteristic.

Let  $Y$  be a smooth projective variety over an algebraically closed field  $l_0$  of positive characteristic. We write as before  $\Omega_{Y/l_0}$  for the sheaf of differentials of  $Y$  over  $l_0$  and  $F_Y : Y \rightarrow Y$  for the absolute Frobenius endomorphism of  $Y$ . Now let  $L$  be a very ample line bundle on  $Y$ . If  $V$  is a torsion free coherent sheaf on  $Y$ , we shall write

$$\mu(V) = \mu_L(V) = \deg_L(V) / \mathrm{rk}(V)$$

for the slope of  $V$  (with respect to  $L$ ). Here  $\text{rk}(V)$  is the rank of  $V$ , i.e., the dimension of the stalk of  $V$  at the generic point of  $Y$ . Furthermore,

$$\text{deg}_L(V) := \int_Y c_1(V) \cdot c_1(L)^{\dim(Y)-1}$$

where  $c_1(\cdot)$  refers to the first Chern class with values in an arbitrary Weil cohomology theory and the integral  $\int_Y$  stands for the push-forward morphism to  $\text{Spec } l_0$  in that theory. Recall that  $V$  is called semistable (with respect to  $L$ ) if for every coherent subsheaf  $W$  of  $V$ , we have  $\mu(W) \leq \mu(V)$  and it is called strongly semistable if  $F_Y^{n,*}V$  is semistable for all  $n \geq 0$ .

In general, there exists a filtration

$$0 = V_0 \subseteq V_1 \subseteq \dots \subseteq V_{r-1} \subseteq V_r = V$$

of  $V$  by subsheaves, such that the quotients  $V_i/V_{i-1}$  are all semistable and such that the slopes  $\mu(V_i/V_{i-1})$  are strictly decreasing for  $i \geq 1$ . This filtration is unique and is called the Harder–Narasimhan (HN) filtration of  $V$ . We will say that  $V$  has a strongly semistable HN filtration if all the quotients  $V_i/V_{i-1}$  are strongly semistable. We shall write

$$\mu_{\min}(V) := \mu(V_r/V_{r-1}) \quad \text{and} \quad \mu_{\max}(V) := \mu(V_1).$$

By the very definition of HN filtration, we have

$$V \text{ is semistable} \Leftrightarrow \mu_{\min}(V) = \mu_{\max}(V).$$

An important consequence of the definitions is the following fact; if  $V$  and  $W$  are two torsion free sheaves on  $Y$  and  $\mu_{\min}(V) > \mu_{\max}(W)$ , then  $\text{Hom}_Y(V, W) = 0$ .

For more on the theory of semistable sheaves, see the monograph [Huybrechts and Lehn 2010].

The following two theorems are key results from Langer.

**Theorem 4.1** [Langer 2004, Theorem 2.7]. *If  $V$  is a torsion free coherent sheaf on  $Y$ , then there exists  $n_0 \geq 0$  such that  $F_Y^{n,*}V$  has a strongly semistable HN filtration for all  $n \geq n_0$ .*

If  $V$  is a torsion free coherent sheaf on  $Y$ , we now define

$$\bar{\mu}_{\min}(V) := \lim_{r \rightarrow \infty} \mu_{\min}(F_Y^{r,*}V) \text{char}(l_0)^r \quad \text{and} \quad \bar{\mu}_{\max}(V) := \lim_{r \rightarrow \infty} \mu_{\max}(F_Y^{r,*}V) / \text{char}(l_0)^r.$$

Note that Theorem 4.1 implies that the two sequences  $\mu_{\min}(F_Y^{r,*}V) / \text{char}(l_0)^r$  and  $\mu_{\max}(F_Y^{r,*}V) / \text{char}(l_0)^r$  become constant when  $r$  is sufficiently large, so the above definitions of  $\bar{\mu}_{\min}$  and  $\bar{\mu}_{\max}$  make sense. Furthermore the sequences  $\mu_{\min}(F_Y^{r,*}V) \text{char}(l_0)^r$  and  $\mu_{\max}(F_Y^{r,*}V) \text{char}(l_0)^r$  are respectively weakly decreasing and weakly increasing, therefore we have

$$\mu_{\min}(V) \geq \bar{\mu}_{\min}(V) \quad \text{and} \quad \bar{\mu}_{\max}(V) \geq \mu_{\max}(V).$$

Let us define

$$\alpha(V) := \max\{\mu_{\min}(V) - \bar{\mu}_{\min}(V), \bar{\mu}_{\max}(V) - \mu_{\max}(V)\}.$$

**Theorem 4.2** [Langer 2004, Cororollary 6.2]. *If  $V$  is of rank  $r$ , then*

$$\alpha(V) \leq \frac{r-1}{\text{char}(l_0)} \max\{\bar{\mu}_{\max}(\Omega_{Y/l_0}), 0\}.$$

*In particular, if  $\bar{\mu}_{\max}(\Omega_{Y/l_0}) \geq 0$  and  $\text{char}(l_0) \geq d = \dim Y$ ,*

$$\bar{\mu}_{\max}(\Omega_{Y/l_0}) \leq \frac{\text{char}(l_0)}{\text{char}(l_0) + 1 - d} \mu_{\max}(\Omega_{Y/l_0}).$$

We conclude this section with the following two lemmas from Rössler.

**Lemma 4.3** [Rössler 2016, Lemma 3.8]. *Suppose that there is a closed  $l_0$ -immersion  $i : Y \hookrightarrow B$ , where  $B$  is an abelian variety over  $l_0$ . Suppose that  $\text{Stab}_B(Y) = 0$ . Then  $\Omega_Y^\vee$  is globally generated and for any dominant proper morphism  $\phi : Y_0 \rightarrow Y$ , where  $Y_0$  is integral, we have  $H^0(Y_0, \phi^* \Omega_Y^\vee) = 0$ . Furthermore, we have  $\bar{\mu}_{\min}(\Omega_Y) > 0$ .*

**Lemma 4.4** [Rössler 2016, Cororollary 3.11]. *Let  $V$  be a finite rank, locally free sheaf over  $Y$ . Suppose that*

- *for any surjective finite map  $\phi : Y' \rightarrow Y$  with  $Y'$  integral, we have  $H^0(Y', \phi^* V) = 0$ ,*
- *$V^\vee$  is globally generated.*

*Then  $H^0(Y, F_Y^{n,*} V \otimes \Omega_{Y/l_0}) = 0$  for  $n$  sufficiently big.*

*Furthermore, let  $T \rightarrow Y$  be a torsor under  $V(F_Y^{n_0,*} V)$ , where  $n_0$  satisfies  $H^0(Y, F_Y^{n,*} V \otimes \Omega_{Y/l_0}) = 0$  for all  $n > n_0$ . Let  $\phi : Y' \rightarrow Y$  be a finite surjective morphism and suppose that  $Y'$  is integral. Then we have the implication*

$$\phi^* T \text{ is a trivial } V(\phi^*(F_Y^{n_0,*} V))\text{-torsor} \implies T \text{ is a trivial } V(F_Y^{n_0,*} V)\text{-torsor}.$$

The main ingredient of the proof of [Lemma 4.4](#) is a result by Szpiro and Lewin-Ménégaux which we will need later.

**Proposition 4.5** [Szpiro 1981, Expose 2, Proposition 1]. *If  $V$  is a vector bundle over  $Y$  such that  $H^0(Y, F_Y^* V \otimes \Omega_{Y/l_0}) = 0$ , then the map*

$$H^1(Y, V) \rightarrow H^1(Y, F_Y^* V)$$

*is injective.*

## 5. Sparsity of $p$ -divisible unramified liftings

In this section we prove our result on the sparsity of  $p$ -divisible unramified liftings (see [Theorem 5.3](#)).

Let  $K, A, X$  and  $U$  be as fixed in [Section 2](#) and let  $\text{Stab}_A(X)$  be trivial. The construction of the stabilizer commutes with the base change, so we have

$$\text{Stab}_A(X) = \text{Stab}_A(\mathcal{X}) \times_U \text{Spec } K.$$

Since  $\text{Stab}_A(X)$  is trivial, by generic flatness and finiteness, we can restrict the map  $\pi : \text{Stab}_A(\mathcal{X}) \rightarrow U$  to the inverse image of a nonempty open subscheme  $U' \subset U$  to obtain a finite flat commutative group scheme of degree one

$$\pi|_{\pi^{-1}(U')} : \pi^{-1}(U') \rightarrow U'.$$

This implies that  $\pi|_{\pi^{-1}(U')}$  is an isomorphism and for any  $\mathfrak{q} \in U'$  we have that  $\text{Stab}_{A_{\mathfrak{q}^0}}(X_{\mathfrak{q}^0})$  is trivial. We will denote by  $\tilde{U} \subseteq U$  the nonempty open subscheme

$$\tilde{U} := \{\mathfrak{q} \in U \mid \text{Stab}_{A_{\mathfrak{q}^0}}(X_{\mathfrak{q}^0}) \text{ is trivial}\}.$$

For any  $\mathfrak{p} \in U$  we denote by  $F_{\overline{k(\mathfrak{p})}}$  the Frobenius endomorphism on  $\overline{k(\mathfrak{p})}$  and by  $F_{R_1}$  the endomorphism of  $R_1$  induced by  $F_{\overline{k(\mathfrak{p})}}$  by functoriality. We define

$$X'_{\mathfrak{p}^0} := X_{\mathfrak{p}^0} \times_{F_{\overline{k(\mathfrak{p})}}} \overline{k(\mathfrak{p})} \quad \text{and} \quad X'_{\mathfrak{p}^1} := X_{\mathfrak{p}^1} \times_{F_{R_1}} R_1$$

and we write

$$F_{X_{\mathfrak{p}^0}/\overline{k(\mathfrak{p})}} : X_{\mathfrak{p}^0} \rightarrow X'_{\mathfrak{p}^0}$$

for the relative Frobenius on  $X_{\mathfrak{p}^0}$ . For brevity's sake, from now on we will write

$$\Omega_{X_{\mathfrak{p}^0}}, \Omega_{X'_{\mathfrak{p}^0}}, \Omega_{X_{\mathfrak{p}^1}}, \Omega_{X'_{\mathfrak{p}^1}} \text{ and } \Omega_X$$

instead of

$$\Omega_{X_{\mathfrak{p}^0}/\overline{k(\mathfrak{p})}}, \Omega_{X'_{\mathfrak{p}^0}/\overline{k(\mathfrak{p})}}, \Omega_{X_{\mathfrak{p}^1}/R_1}, \Omega_{X'_{\mathfrak{p}^1}/R_1} \text{ and } \Omega_{X/K}.$$

Observe that since  $U$  is normal,  $\mathcal{A}$  is projective over  $U$  [Raynaud 1970, Theorem XI 1.4]. Therefore there exists a  $U$ -very ample line bundle  $L$  on  $\mathcal{X}$ . For any  $\mathfrak{p} \in U$  different from the generic point  $\xi$ , let us denote by  $L_{\mathfrak{p}}$  the inverse image of  $L$  on  $X_{\mathfrak{p}^0}$ . Similarly we denote by  $L_{\xi}$  the inverse image of  $L$  on  $X$ . From now on, for any vector bundle  $G_{\mathfrak{p}}$  over  $X_{\mathfrak{p}^0}$ , we will write  $\text{deg}(G_{\mathfrak{p}})$  for the degree of  $G_{\mathfrak{p}}$  with respect to  $L_{\mathfrak{p}}$ . Analogously, if  $G_{\xi}$  is a vector bundle over  $X$ , we will write  $\text{deg}(G_{\xi})$  for the degree of  $G_{\xi}$  with respect to  $L_{\xi}$ . Now consider the vector bundle  $\Omega_{\mathcal{X}/U}$  over  $\mathcal{X}$ . For any natural number  $m$ , the map from  $U$  to  $\mathbb{Z}$  defined by

$$\mathfrak{p} \mapsto \chi((\Omega_{\mathcal{X}/U} \otimes L^m)_{\mathfrak{p}}) = \chi(\Omega_{X_{\mathfrak{p}^0}} \otimes L_{\mathfrak{p}}^m) \quad \text{and} \quad \xi \mapsto \chi((\Omega_{\mathcal{X}/U} \otimes L^m)_{\xi}) = \chi(\Omega_X \otimes L_{\xi}^m)$$

(here  $\chi$  refers to the Euler characteristic) is constant on  $U$  [Mumford 1970, Chapter II, Section 5]. Therefore we have the equality

$$\chi(\Omega_{X_{\mathfrak{p}^0}} \otimes L_{\mathfrak{p}}^m) = \chi(\Omega_X \otimes L_{\xi}^m)$$

for all  $m \in \mathbb{N}$  and for all  $\mathfrak{p} \in U$ . In other words, the Hilbert polynomial of  $\Omega_{X_{\mathfrak{p}^0}}$  with respect to  $L_{\mathfrak{p}}$  coincides with the Hilbert polynomial of  $\Omega_X$  with respect to  $L_{\xi}$ . Since the degree of a vector bundle we defined at the beginning of this section can be described in terms of its Hilbert polynomial [Huybrechts and Lehn 2010, Definition 1.2.11], we obtain that for every  $\mathfrak{p} \in U$  we have  $\text{deg}(\Omega_{X_{\mathfrak{p}^0}}) = \text{deg}(\Omega_X)$ .

The following lemma is a fundamental step to prove our sparsity [Theorem 5.3](#).

**Lemma 5.1.** *Let  $K$ ,  $A$ ,  $X$  and  $U$  be as fixed in [Section 2](#), let  $\text{Stab}_A(X)$  be trivial and let  $n$  be the dimension of  $X$  over  $K$ . Then*

$$\text{Hom}_{X_{p^0}}(F_{X_{p^0}}^{k,*} \Omega_{X_{p^0}}, \Omega_{X_{p^0}}) = 0$$

for any  $k \geq 1$  and any  $\mathfrak{p} \in \tilde{U}$  above a prime  $p > n^2 \deg(\Omega_X)$ .

*Proof.* Let us notice first that, if  $n = 1$ , then  $X$  is a curve of genus  $g$  at least 2 and

$$\text{Hom}_{X_{p^0}}(F_{X_{p^0}}^{k,*} \Omega_{X_{p^0}}, \Omega_{X_{p^0}}) = 0$$

is a simple consequence of the fact

$$\deg(F_{X_{p^0}}^{k,*} \Omega_{X_{p^0}}) = p^k(2g - 2) > 2g - 2 = \deg \Omega_{X_{p^0}}.$$

To treat the general case, let us fix  $\mathfrak{p} \in \tilde{U}$  above a prime  $p > n^2 \deg(\Omega_X)$ . We know that if

$$\mu_{\min}(F_{X_{p^0}}^{k,*} \Omega_{X_{p^0}}) > \mu_{\max}(\Omega_{X_{p^0}})$$

then  $\text{Hom}_{X_{p^0}}(F_{X_{p^0}}^{k,*} \Omega_{X_{p^0}}, \Omega_{X_{p^0}}) = 0$ . Since  $\mu_{\min} \geq \bar{\mu}_{\min}$  and  $\bar{\mu}_{\max} \geq \mu_{\max}$ , it is sufficient to show that, for every  $k \geq 1$

$$\bar{\mu}_{\min}(F_{X_{p^0}}^{k,*} \Omega_{X_{p^0}}) > \bar{\mu}_{\max}(\Omega_{X_{p^0}}). \quad (3)$$

Since  $\text{Stab}_{A_{p^0}}(X_{p^0})$  is trivial, we can apply [Lemma 4.3](#) to obtain  $\bar{\mu}_{\min}(\Omega_{X_{p^0}}) > 0$ . In particular  $\mu_{\min}(\Omega_{X_{p^0}}) > 0$  and  $\deg(\Omega_{X_{p^0}}) > 0$ . Using this and the equality  $\bar{\mu}_{\min}(F_{X_{p^0}}^{k,*} \Omega_{X_{p^0}}) = p^k \bar{\mu}_{\min}(\Omega_{X_{p^0}})$ , we see that (3) is implied by

$$p \bar{\mu}_{\min}(\Omega_{X_{p^0}}) > \bar{\mu}_{\max}(\Omega_{X_{p^0}}). \quad (4)$$

[Theorem 4.2](#) gives us the following inequality

$$p \bar{\mu}_{\min}(\Omega_{X_{p^0}}) \geq p \mu_{\min}(\Omega_{X_{p^0}}) + (1 - n) \bar{\mu}_{\max}(\Omega_{X_{p^0}}),$$

so that (4) is satisfied if

$$p \mu_{\min}(\Omega_{X_{p^0}}) > n \bar{\mu}_{\max}(\Omega_{X_{p^0}}). \quad (5)$$

Since  $p > n^2 \deg(\Omega_X) \geq n$ , we can apply the second part of [Theorem 4.2](#)

$$\bar{\mu}_{\max}(\Omega_{X_{p^0}}) \leq \frac{p}{p + 1 - n} \mu_{\max}(\Omega_{X_{p^0}}),$$

so that inequality (5) is implied by

$$(p + 1 - n) \mu_{\min}(\Omega_{X_{p^0}}) > n \mu_{\max}(\Omega_{X_{p^0}}). \quad (6)$$

If  $\Omega_{X_{p^0}}$  is semistable, (6) gives  $p > 2n - 1$ . Otherwise, we can estimate  $\mu_{\max}(\Omega_{X_{p^0}})$  and  $\mu_{\min}(\Omega_{X_{p^0}})$  in the following way. We know that

$$\mu_{\max}(\Omega_{X_{p^0}}) = \frac{\deg(M)}{\text{rk}(M)}$$

for some subsheaf  $0 \neq M \subsetneq \Omega_{X_{p^0}}$ . Therefore we have  $\mu_{\max}(\Omega_{X_{p^0}}) \leq \deg(M)$ . Furthermore, since  $\bar{\mu}_{\min}(\Omega_{X_{p^0}}) > 0$ , we have that  $\deg(\Omega_{X_{p^0}}/M) > 0$ . This and the additivity of the degree on short exact sequences gives us

$$\mu_{\max}(\Omega_{X_{p^0}}) \leq \deg(M) \leq \deg(\Omega_{X_{p^0}}) - 1.$$

Similarly,

$$\mu_{\min}(\Omega_{X_{p^0}}) = \frac{\deg(Q)}{\text{rk}(Q)}$$

for some  $Q$  quotient of  $\Omega_{X_{p^0}}$ , so  $\mu_{\min}(\Omega_{X_{p^0}}) \geq 1/n$ . Inequality (6) is then implied by

$$p > n^2 \deg(\Omega_{X_{p^0}}) + (n - 1 - n^2).$$

Since  $n - 1 - n^2$  is always negative, we are reduced to  $p > n^2 \deg(\Omega_{X_{p^0}})$ . Now  $\deg(\Omega_{X_{p^0}})$  is greater or equal to one, so  $n^2 \deg(\Omega_{X_{p^0}}) \geq 2n - 1$  for any  $n$ . This ensures us that the condition

$$p > n^2 \deg(\Omega_{X_{p^0}})$$

is sufficient to have  $\mu_{\min}(F_{X_{p^0}}^{k,*} \Omega_{X_{p^0}}) > \mu_{\max}(\Omega_{X_{p^0}})$  for every  $k \geq 1$  whether  $\Omega_{X_{p^0}}$  is semistable or not. To conclude it is enough to remember that  $\deg(\Omega_{X_{p^0}})$  coincides with  $\deg(\Omega_X)$ .  $\square$

**Corollary 5.2.** *The map*

$$H^1(X_{p^0}, F_{X_{p^0}}^* \Omega_{X_{p^0}}^\vee) \rightarrow H^1(X_{p^0}, F_{X_{p^0}}^{k,*} \Omega_{X_{p^0}}^\vee)$$

is injective for every  $k \geq 1$  and every  $\mathfrak{p} \in \tilde{U}$  above a prime  $p > n^2 \deg(\Omega_X)$ .

*Proof.* Lemma 5.1 and Proposition 4.5 imply that

$$H^1(X_{p^0}, F_{X_{p^0}}^{h,*} \Omega_{X_{p^0}}^\vee) \rightarrow H^1(X_{p^0}, F_{X_{p^0}}^{h+1,*} \Omega_{X_{p^0}}^\vee)$$

is injective for every  $h \geq 0$ . Therefore the composition

$$H^1(X_{p^0}, F_{X_{p^0}}^* \Omega_{X_{p^0}}^\vee) \hookrightarrow H^1(X_{p^0}, F_{X_{p^0}}^{2,*} \Omega_{X_{p^0}}^\vee) \hookrightarrow \dots \hookrightarrow H^1(X_{p^0}, F_{X_{p^0}}^{k,*} \Omega_{X_{p^0}}^\vee)$$

is an injective map.  $\square$

We are now ready to prove our sparsity result.

**Theorem 5.3.** *With the same hypotheses as in Lemma 5.1, for any  $\mathfrak{p} \in \tilde{U}$  above a prime  $p > n^2 \deg(\Omega_X)$ , the set*

$$\{P \in X_{p^0}(R_0) \mid P \text{ lifts to an element of } pA_{p^1}(R_1) \cap X_{p^1}(R_1)\}$$

is not Zariski dense in  $X_{p^0}$ .

*Proof.* Let us fix  $\mathfrak{p}$  as in the hypotheses. Since

$$\text{Crit}^1(\mathcal{X}, \mathcal{A})(R_0) = pA_{p^1}(R_1) \cap X_{p^1}(R_1),$$

we have that

$$\{P \in X_{p^0}(R_0) \mid P \text{ lifts to an element of } pA_{p^1}(R_1) \cap X_{p^1}(R_1)\}$$

coincides with the image of  $\text{Crit}^1(\mathcal{X}, \mathcal{A})(R_0) \rightarrow X_{p^0}(R_0)$ .

Let us assume by contradiction that this image is dense in  $X_{p^0}(R_0)$ . This implies that  $\pi_1 : \text{Gr}_1(X_{p^1}) \rightarrow X_{p^0}$  is a trivial torsor; the argument we use to show this is taken from Rössler (see the beginning of the proof of Theorem 2.2 in [Rössler 2016]). First of all the closed map  $\text{Crit}^1(\mathcal{X}, \mathcal{A}) \rightarrow X_{p^0}$  is surjective and so we can choose an irreducible component

$$\text{Crit}^1(\mathcal{X}, \mathcal{A})_0 \hookrightarrow \text{Crit}^1(\mathcal{X}, \mathcal{A})$$

which dominates  $X_{p^0}$ . Lemmas 4.3 and 5.1 allow us to apply the second part of Lemma 4.4 with  $V = \Omega_{X_{p^0}}^\vee$ ,  $Y = X_{p^0}$ ,  $n_0 = 1$ ,  $T = \text{Gr}_1(X_{p^1})$  and  $\phi$  equal to  $\text{Crit}^1(\mathcal{X}, \mathcal{A})_0 \rightarrow X_{p^0}$ . We have that  $\phi^* \text{Gr}_1(X_{p^1})$  is trivial as a  $V(\phi^* F_{X_{p^0}}^* \Omega_{X_{p^0}}^\vee)$ -torsor, since  $\text{Crit}^1(\mathcal{X}, \mathcal{A})_0$  is contained in  $\text{Gr}_1(X_{p^1})$ . Hence  $\pi_1 : \text{Gr}_1(X_{p^1}) \rightarrow X_{p^0}$  is trivial as a  $V(F_{X_{p^0}}^* \Omega_{X_{p^0}}^\vee)$ -torsor. Let us take a section  $\sigma : X_{p^0} \rightarrow \text{Gr}_1(X_{p^1})$ . By the definition of the Greenberg transform, the map  $\sigma$  over  $R_0$  corresponds to a map  $\bar{\sigma} : \mathbb{W}_1(X_{p^0}) \rightarrow X_{p^1}$  over  $R_1$ . We can precompose  $\bar{\sigma}$  with the morphism  $t : X_{p^1} \rightarrow \mathbb{W}_1(X_{p^0})$  corresponding to

$$\begin{aligned} W_1(\mathcal{O}_{X_{p^0}}) &\rightarrow \mathcal{O}_{X_{p^1}} \\ (a_0, a_1) &\mapsto \tilde{a}_0^p + \tilde{a}_1 p, \end{aligned}$$

where  $\tilde{a}_i$  lifts  $a_i$ . Consider the following diagram

$$\begin{array}{ccccc} X_{p^1} & \xrightarrow{t} & \mathbb{W}_1(X_{p^0}) & \xrightarrow{\bar{\sigma}} & X_{p^1} \\ \uparrow & & \uparrow & & \uparrow \\ X_{p^0} & \xrightarrow{F_{X_{p^0}}} & X_{p^0} & \xrightarrow{\text{Id}} & X_{p^0} \end{array}$$

Its left square is commutative, since the composition

$$X_{p^0} \longrightarrow X_{p^1} \xrightarrow{t} \mathbb{W}_1(X_{p^0})$$

simply corresponds to the map

$$\begin{aligned} W_1(\mathcal{O}_{X_{p^0}}) &\rightarrow \mathcal{O}_{X_{p^0}} \\ (a_0, a_1) &\mapsto a_0^p. \end{aligned}$$

For the commutativity of the right square, notice that by the very definition of the transition morphism  $\pi_1 : \text{Gr}_1(X_{p^1}) \rightarrow X_{p^0}$  we have a commutative diagram

$$\begin{array}{ccc} \text{Hom}_{R_1}(\mathbb{W}_1(X_{p^0}), X_{p^1}) & \longrightarrow & \text{Hom}_{R_0}(X_{p^0}, \text{Gr}_1(X_{p^1})) \\ \downarrow \text{reduction mod } p & & \swarrow (\pi_1 \circ -) \\ \text{Hom}_{R_0}(X_{p^0}, X_{p^0}) & & \end{array}$$

In particular,  $\text{Id}_{X_{p^0}} = \pi_1 \circ \sigma = (\text{reduction mod } p)(\bar{\sigma})$ , which is exactly what we wanted to verify. We obtain therefore that  $\bar{\sigma} \circ t : X_{p^1} \rightarrow X_{p^1}$  is a lift of the Frobenius  $F_{X_{p^0}}$ .

The diagram below is also commutative

$$\begin{array}{ccccc}
 X_{p^1} & \xrightarrow{t} & \mathbb{W}_1(X_{p^0}) & \xrightarrow{\bar{\sigma}} & X_{p^1} \\
 \downarrow & & \downarrow & & \downarrow \\
 \text{Spec}(R_1) & \xrightarrow{F_{R_1}} & \text{Spec}(R_1) & \xrightarrow{\text{Id}} & \text{Spec}(R_1)
 \end{array}$$

In fact, by definition,  $\bar{\sigma}$  is a morphism over  $R_1$ , so the right square is commutative. The commutativity of the left square is easy to check, since we know explicitly  $t$  and  $F_{R_1}$ . Therefore  $\bar{\sigma} \circ t$  is a lift of the Frobenius  $F_{X_{p^0}}$  compatible with  $F_{R_1}$ ; this implies the existence of a morphism of  $R_1$ -schemes

$$\tilde{F} : X_{p^1} \rightarrow X'_{p^1}$$

lifting the relative Frobenius  $F_{X_{p^0}/R_0}$ .

As shown in part (b) of the proof of Théorème 2.1 in [Deligne and Illusie 1987], since the image of  $\tilde{F}^* : \Omega_{X'_{p^1}} \rightarrow \tilde{F}_* \Omega_{X_{p^1}}$  is contained in  $p\tilde{F}_* \Omega_{X_{p^1}}$  and the multiplication by  $p$  induces an isomorphism  $p : F_{X_{p^0}/R_0,*} \Omega_{X_{p^0}} \xrightarrow{\sim} p\tilde{F}_* \Omega_{X_{p^1}}$ , there exists a unique map

$$f := p^{-1} \tilde{F}^* : \Omega_{X'_{p^1}} \rightarrow F_{X_{p^0}/R_0,*} \Omega_{X_{p^0}},$$

making the diagram below commutative.

$$\begin{array}{ccc}
 \Omega_{X'_{p^1}} & \xrightarrow{\tilde{F}^*} & p\tilde{F}_* \Omega_{X_{p^1}} \\
 \downarrow & & \uparrow p \\
 \Omega_{X'_{p^0}} & \xrightarrow{f} & F_{X_{p^0}/R_0,*} \Omega_{X_{p^0}}
 \end{array}$$

Proposition 3 in [Xin 2016] states that the adjoint of  $f$ ,

$$\bar{f} : F_{X_{p^0}}^* \Omega_{X_{p^0}} = F_{X_{p^0}/R_0}^* \Omega_{X'_{p^0}} \rightarrow \Omega_{X_{p^0}},$$

is generically bijective. This clearly contradicts Lemma 5.1. □

### 6. The number of irreducible components of the critical scheme of complete intersections

In this last section we provide an upper bound for the number of irreducible components of the critical scheme  $\text{Crit}^1(\mathcal{X}, \mathcal{A})$  in the case in which  $X$  is a smooth complete intersection.

Let  $A/K$  be an abelian variety of dimension  $n$  and let  $L$  be a very ample line bundle on  $A$ . Let  $c \in \mathbb{N}$  be positive and let  $H_1, H_2, \dots, H_c \in |L|$  be general. We define  $X := H_1 \cap H_2 \cap \dots \cap H_c$ . Suppose that  $X$  is smooth.

Let us take a sufficiently small open  $V \subseteq \text{Spec}(\mathcal{O}_K)$  such that  $A$  extends over  $V$  to an abelian scheme  $\mathcal{A}$ ,  $L$  extends to a  $V$ -very ample line bundle  $\mathcal{L}$ ,  $H_i$  extends to  $\mathcal{H}_i$  for every  $i$  and  $\mathcal{X} := \mathcal{H}_1 \cap \mathcal{H}_2 \cap \cdots \cap \mathcal{H}_c$  is smooth. We can restrict  $V$  if necessary and suppose  $K/\mathbb{Q}$  is unramified at  $\mathfrak{p}$ .

**Theorem 6.1.** *Let  $K$  be a number field,  $A/K$  be an abelian variety of dimension  $n$  and let  $L$  be a very ample line bundle on  $A$ . Let  $c \in \mathbb{N}$  be positive and let  $H_1, H_2, \dots, H_c \in |L|$  be general. Suppose that  $X := H_1 \cap H_2 \cap \cdots \cap H_c$  is smooth. If  $\mathfrak{p}$  is in the open subscheme  $V$  defined above, then the number of irreducible components of  $\text{Crit}^1(\mathcal{X}, \mathcal{A})$  is bounded by*

$$p^{2n} \left( \sum_{h=0}^{n-c} \binom{2n-2c}{h} \binom{c}{n-c-h} p^{n-c-h} \right) (L^n)^2.$$

Here  $(L^n)$  denotes the intersection number of  $L$ .

*Proof.* To obtain [Theorem 6.1](#), we follow the approach of [[Buium 1996](#), Theorem 1.11], proving the Manin–Mumford conjecture for curves; we first show that  $\text{Crit}^1(\mathcal{X}, \mathcal{A})$  can be realized as the intersection of two projective varieties (see  $\mathbb{P}(E_X)$  and  $[p]_* \text{Gr}_1(A_{\mathfrak{p}^1})$  below) and then use the product of their degrees to bound the number of its irreducible components. Since  $X$  is not necessarily of dimension one, the computation of the degree of  $\mathbb{P}(E_X)$  is slightly more demanding here than the corresponding one in Buium’s work.

Let us fix  $\mathfrak{p} \in V$ . The torsors  $\text{Gr}_1(X_{\mathfrak{p}^1}) \rightarrow X_{\mathfrak{p}^0}$  and  $\text{Gr}_1(A_{\mathfrak{p}^1}) \rightarrow A_{\mathfrak{p}^0}$  correspond to elements  $\eta_X \in H^1(X_{\mathfrak{p}^0}, F_{X_{\mathfrak{p}^0}}^* \Omega_{X_{\mathfrak{p}^0}}^\vee)$  and  $\eta_A \in H^1(A_{\mathfrak{p}^0}, F_{A_{\mathfrak{p}^0}}^* \Omega_{A_{\mathfrak{p}^0}}^\vee)$ , respectively. Under the natural isomorphisms

$$H^1(X_{\mathfrak{p}^0}, F_{X_{\mathfrak{p}^0}}^* \Omega_{X_{\mathfrak{p}^0}}^\vee) \simeq \text{Ext}^1(F_{X_{\mathfrak{p}^0}}^* \Omega_{X_{\mathfrak{p}^0}}, \mathcal{O}_{X_{\mathfrak{p}^0}}) \quad \text{and} \quad H^1(A_{\mathfrak{p}^0}, F_{A_{\mathfrak{p}^0}}^* \Omega_{A_{\mathfrak{p}^0}}^\vee) \simeq \text{Ext}^1(F_{A_{\mathfrak{p}^0}}^* \Omega_{A_{\mathfrak{p}^0}}, \mathcal{O}_{A_{\mathfrak{p}^0}}),$$

$\eta_X$  and  $\eta_A$  correspond to extensions of vector bundles

$$0 \rightarrow \mathcal{O}_{X_{\mathfrak{p}^0}} \rightarrow E_X \rightarrow F_{X_{\mathfrak{p}^0}}^* \Omega_{X_{\mathfrak{p}^0}} \rightarrow 0 \quad \text{and} \quad 0 \rightarrow \mathcal{O}_{A_{\mathfrak{p}^0}} \rightarrow E_A \rightarrow F_{A_{\mathfrak{p}^0}}^* \Omega_{A_{\mathfrak{p}^0}} \rightarrow 0.$$

For any locally free sheaf  $W$  over a base  $S$  of finite type over a field, we shall write  $\mathbb{P}(W)$  for the projective bundle associated to  $W$ , i.e., the  $S$ -scheme representing the functor on  $S$ -schemes

$$T \mapsto \{\text{isomorphism classes of surjective morphisms of } \mathcal{O}_T\text{-modules } W_T \rightarrow Q, \\ \text{where } Q \text{ is locally free of rank } 1\}.$$


As shown in paragraph 1 of [[Martin-Deschamps 1984](#)], the two extensions above give us two divisors

$$D_X := \mathbb{P}(F_{X_{\mathfrak{p}^0}}^* \Omega_{X_{\mathfrak{p}^0}}) \subseteq \mathbb{P}(E_X) \quad \text{and} \quad D_A := \mathbb{P}(F_{A_{\mathfrak{p}^0}}^* \Omega_{A_{\mathfrak{p}^0}}) \subseteq \mathbb{P}(E_A),$$

belonging respectively to the linear systems  $|\mathcal{O}_{\mathbb{P}(E_X)}(1)|$  and  $|\mathcal{O}_{\mathbb{P}(E_A)}(1)|$ , and

$$\text{Gr}_1(X_{\mathfrak{p}^1}) \simeq \mathbb{P}(E_X) \setminus D_X \quad \text{and} \quad \text{Gr}_1(A_{\mathfrak{p}^1}) \simeq \mathbb{P}(E_A) \setminus D_A.$$

If  $i$  denotes the closed immersion  $i : X_{p^0} \rightarrow A_{p^0}$ , then it is not difficult to show that there is a natural restriction homomorphism  $i^* E_A \rightarrow E_X$  prolonging the homomorphism  $i^* \Omega_{A_{p^0}} \rightarrow \Omega_{X_{p^0}}$ . The homomorphism  $i^* E_A \rightarrow E_X$  is clearly surjective, so it induces a closed immersion  $j : \mathbb{P}(E_X) \hookrightarrow \mathbb{P}(E_A)$  prolonging  $\text{Gr}_1(X_{p^1}) \hookrightarrow \text{Gr}_1(A_{p^1})$ . Therefore we have a commutative diagram



Let us denote by  $\mathcal{L}_p$  the base change of  $\mathcal{L}$  to  $A_{p^0}$ . It is standard to prove that

$$\mathcal{H} := \pi_A^* \mathcal{L}_p \otimes \mathcal{O}_{\mathbb{P}(E_A)}(1)$$

is very ample on  $\mathbb{P}(E_A)$  [Buium and Voloch 1996, p. 4]. We have

$$\mathcal{H}|_{\mathbb{P}(E_X)} = \pi_X^* i^* \mathcal{L}_p \otimes \mathcal{O}_{\mathbb{P}(E_X)}(1) \quad \text{and} \quad \mathcal{H}|_{[p]_* \text{Gr}_1(A_{p^1})} = T^* \mathcal{L}_p,$$

since  $D_A \in |\mathcal{O}_{\mathbb{P}(E_A)}(1)|$  and  $[p]_* \text{Gr}_1(A_{p^1}) \subseteq \text{Gr}_1(A_{p^1}) \simeq \mathbb{P}(E_A) \setminus D_A$ . We know that  $[p]_* \text{Gr}_1(A_{p^1})$  is the maximal abelian subvariety of  $\text{Gr}_1(A_{p^1})$  and we know that the multiplication by  $p$  map on  $\text{Gr}_1(A_{p^1})$  factors through the isogeny  $T$ . This implies that  $T$  has degree at most  $p^{2n}$ , so we have the following estimate

$$\text{deg}_{\mathcal{H}}([p]_* \text{Gr}_1(A_{p^1})) \leq p^{2n} (\mathcal{L}_p^n).$$

Let us now consider  $\text{deg}_{\mathcal{H}}(\mathbb{P}(E_X))$ . It coincides with

$$\int_{\mathbb{P}(E_X)} c_1(\mathcal{H}|_{\mathbb{P}(E_X)})^{2n-2c} \tag{7}$$

where  $c_1$  stands for the first Chern class in the Chow ring and  $\int_{\mathbb{P}(E_X)}$  stands for the push-forward morphism to  $\text{Spec}(R_0)$  in the Chow theory. Since

$$c_1(\mathcal{H}|_{\mathbb{P}(E_X)}) = c_1(\pi_X^* i^* \mathcal{L}_p) + c_1(\mathcal{O}_{\mathbb{P}(E_X)}(1))$$

we can rewrite (7) as

$$\int_{\mathbb{P}(E_X)} \sum_{h=0}^{2n-2c} \binom{2n-2c}{h} c_1(\pi_X^* i^* \mathcal{L}_p)^h \cdot c_1(\mathcal{O}_{\mathbb{P}(E_X)}(1))^{2n-2c-h}.$$

Equivalently

$$\int_{X_{p^0}} \sum_{h=0}^{2n-2c} \binom{2n-2c}{h} c_1(i^* \mathcal{L}_p)^h \cdot \pi_{X,*}(c_1(\mathcal{O}_{\mathbb{P}(E_X)}(1))^{2n-2c-h})$$

and by definition of Segre class this is

$$\int_{X_{p^0}} \sum_{h=0}^{2n-2c} \binom{2n-2c}{h} c_1(i^* \mathcal{L}_p)^h \cdot s_{n-c-h}(E_X^\vee).$$

Notice that the Segre classes of the dual of  $E_X$  appear in our formula; this is due to the fact that we are not using Fulton's geometric notation for the projective bundle associated to a vector bundle (see the note at the end of B.5.5 in [Fulton 1998]). Since  $s_k = 0$  if  $k < 0$ , we end up with

$$\int_{X_{p^0}} \sum_{h=0}^{n-c} \binom{2n-2c}{h} c_1(i^* \mathcal{L}_p)^h \cdot s_{n-c-h}(E_X^\vee).$$

Now the exact sequence

$$0 \rightarrow \mathcal{O}_{X_{p^0}} \rightarrow E_X \rightarrow F_{X_{p^0}}^* \Omega_{X_{p^0}} \rightarrow 0$$

implies

$$s_{n-c-h}(E_X^\vee) = s_{n-c-h}(F_{X_{p^0}}^* \Omega_{X_{p^0}}^\vee)$$

and so

$$s_{n-c-h}(E_X^\vee) = p^{n-c-h} s_{n-c-h}(\Omega_{X_{p^0}}^\vee)$$

(here we have used the following fact: the pullback of a cycle  $\eta$  of codimension  $j$  through the Frobenius map coincides with  $p^j \eta$ ). Therefore we have to study the following sum

$$\sum_{h=0}^{n-c} \binom{2n-2c}{h} p^{n-c-h} c_1(i^* \mathcal{L}_p)^h \cdot s_{n-c-h}(\Omega_{X_{p^0}}^\vee). \quad (8)$$

The short exact sequence

$$0 \rightarrow \Omega_{X_{p^0}}^\vee \rightarrow i^* \Omega_{A_{p^0}}^\vee \rightarrow N \rightarrow 0$$

(where  $N$  is the normal bundle for  $i$ ) gives

$$c_t(\Omega_{X_{p^0}}^\vee) c_t(N) = c_t(i^* \Omega_{A_{p^0}}^\vee) = 1,$$

so that  $c_t(N) = s_t(\Omega_{X_{p^0}}^\vee)$ . Recalling that

$$c_t(N) = (1 + c_1(i^* \mathcal{L}_p)t)^c$$

we obtain

$$s_{n-c-h}(\Omega_{X_{p^0}}^\vee) = c_{n-c-h}(N) = \binom{c}{n-c-h} c_1(i^* \mathcal{L}_p)^{n-c-h}.$$

Substituting in (8), we obtain

$$\left( \sum_{h=0}^{n-c} \binom{2n-2c}{h} \binom{c}{n-c-h} p^{n-c-h} \right) c_1(i^* \mathcal{L}_p)^{n-c}.$$

Therefore  $\text{deg}_{\mathcal{H}}(\mathbb{P}(E_X))$  is

$$\left( \sum_{h=0}^{n-c} \binom{2n-2c}{h} \binom{c}{n-c-h} p^{n-c-h} \right) \int_{X_{p^0}} c_1(i^* \mathcal{L}_p)^{n-c}.$$

Since  $X_{p^0} = H_{1,p} \cap \dots \cap H_{c,p}$  where  $H_{1,p}, \dots, H_{c,p}$  belong to  $|\mathcal{L}_p|$ , we have

$$\int_{X_{p^0}} c_1(i^* \mathcal{L}_p)^{n-c} = \int_{A_{p^0}} c_1(\mathcal{L}_p)^n = (\mathcal{L}_p^n)$$

and

$$\text{deg}_{\mathcal{H}}(\mathbb{P}(E_X)) = \left( \sum_{h=0}^{n-c} \binom{2n-2c}{h} \binom{c}{n-c-h} p^{n-c-h} \right) (\mathcal{L}_p^n).$$

Now Bézout’s theorem in Fulton’s form [1998, p. 148] says that the number of irreducible components in the intersection of two projective varieties of degrees  $d_1$  and  $d_2$  cannot exceed  $d_1 d_2$ . In particular, the number of irreducible components of  $\text{Crit}^1(\mathcal{X}, \mathcal{A})$  is less than or equal to

$$p^{2n} \left( \sum_{h=0}^{n-c} \binom{2n-2c}{h} \binom{c}{n-c-h} p^{n-c-h} \right) (\mathcal{L}_p^n)^2.$$

Notice that  $(\mathcal{L}_p^n) = (L^n)$ , by the same reasoning as before Lemma 5.1. □

**Remark 6.2.** One can consider any intersection  $X := H_1 \cap H_2 \cap \dots \cap H_c$  where  $H_i \in |L_i|$  for some very ample line bundles  $L_i$ . In this more general case, the computations in our proof become a bit more complex, but it is still possible to give an explicit bound for the number of irreducible components of  $\text{Crit}^1(\mathcal{X}, \mathcal{A})$ . We have

$$c_j(N) = \sum_{1 \leq i_1 < \dots < i_j \leq c} \prod_{k=i_1}^{i_j} c_1(i^* \mathcal{L}_{k,p})$$

which implies

$$s_{n-c-h}(\Omega_{X_{p^0}}^\vee) = \sum_{1 \leq i_1 < \dots < i_{n-c-h} \leq c} \prod_{k=i_1}^{i_{n-c-h}} c_1(i^* \mathcal{L}_{k,p}).$$

Therefore, defining  $\mathcal{H} := \pi_A^* \mathcal{L}_{1,p} \otimes \mathcal{O}_{\mathbb{P}(E_A)}(1)$ , then  $\text{deg}_{\mathcal{H}}(\mathbb{P}(E_X))$  is

$$\sum_{h=0}^{n-c} \binom{2n-2c}{h} p^{n-c-h} \sum_{1 \leq i_1 < \dots < i_{n-c-h} \leq c} \left( \int_{X_{p^0}} c_1(i^* \mathcal{L}_{1,p})^h \prod_{k=i_1}^{i_{n-c-h}} c_1(i^* \mathcal{L}_{k,p}) \right).$$

We have

$$\int_{X_{p^0}} c_1(i^* \mathcal{L}_{1,p})^h \prod_{k=i_1}^{i_{n-c-h}} c_1(i^* \mathcal{L}_{k,p}) = I_{i_1, \dots, i_{n-c-h}}$$

where  $I_{i_1, \dots, i_{n-c-h}}$  is the following intersection number

$$I_{i_1, \dots, i_{n-c-h}} := \overbrace{(L_1 \cdots L_1)^{h+1}} \cdot L_2 \cdot L_3 \cdots L_c \cdot L_{i_1} \cdot L_{i_2} \cdots L_{i_{n-c-h}}.$$

We obtain that  $\deg_{\mathcal{H}}(\mathbb{P}(E_X))$  is

$$\sum_{h=0}^{n-c} \binom{2n-2c}{h} p^{n-c-h} \sum_{1 \leq i_1 < \dots < i_{n-c-h} \leq c} I_{i_1, \dots, i_{n-c-h}},$$

and therefore the number of irreducible components of  $\text{Crit}^1(\mathcal{X}, \mathcal{A})$  is bounded by

$$p^{2n} (L_1^n) \sum_{h=0}^{n-c} \binom{2n-2c}{h} p^{n-c-h} \sum_{1 \leq i_1 < \dots < i_{n-c-h} \leq c} I_{i_1, \dots, i_{n-c-h}}.$$

### Acknowledgements

This paper constitutes part of my PhD thesis. I would like to thank professor D. Rössler for having introduced me to this topic and for his invaluable mentoring and advice. I am also indebted to the anonymous referee for the useful remarks.

This work was partly supported by the SFB 1085 “Higher Invariants” at the University of Regensburg.

### References

- [Bosch et al. 1990] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)] **21**, Springer, 1990. [MR](#) [Zbl](#)
- [Buium 1996] A. Buium, “Geometry of  $p$ -jets”, *Duke Math. J.* **82**:2 (1996), 349–367. [MR](#) [Zbl](#)
- [Buium and Voloch 1996] A. Buium and J. F. Voloch, “Lang’s conjecture in characteristic  $p$ : an explicit bound”, *Compositio Math.* **103**:1 (1996), 1–6. [MR](#) [Zbl](#)
- [Deligne and Illusie 1987] P. Deligne and L. Illusie, “Relèvements modulo  $p^2$  et décomposition du complexe de de Rham”, *Invent. Math.* **89**:2 (1987), 247–270. [MR](#) [Zbl](#)
- [Fulton 1998] W. Fulton, *Intersection theory*, 2nd ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics] **2**, Springer, 1998. [MR](#) [Zbl](#)
- [Greenberg 1961] M. J. Greenberg, “Schemata over local rings”, *Ann. of Math. (2)* **73** (1961), 624–648. [MR](#) [Zbl](#)
- [Greenberg 1963] M. J. Greenberg, “Schemata over local rings II”, *Ann. of Math. (2)* **78** (1963), 256–266. [MR](#) [Zbl](#)
- [Huybrechts and Lehn 2010] D. Huybrechts and M. Lehn, *The geometry of moduli spaces of sheaves*, 2nd ed., Cambridge University Press, 2010. [MR](#) [Zbl](#)
- [Langer 2004] A. Langer, “Semistable sheaves in positive characteristic”, *Ann. of Math. (2)* **159**:1 (2004), 251–276. [MR](#) [Zbl](#)
- [Martin-Deschamps 1984] M. Martin-Deschamps, “Propriétés de descente des variétés à fibré cotangent ample”, *Ann. Inst. Fourier (Grenoble)* **34**:3 (1984), 39–64. [MR](#) [Zbl](#)

- [Mumford 1970] D. Mumford, *Abelian varieties*, Tata Institute of Fundamental Research Studies in Mathematics **5**, Oxford University Press, London, 1970. [MR](#) [Zbl](#)
- [Raynaud 1970] M. Raynaud, *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, Lecture Notes in Mathematics **119**, Springer, 1970. [MR](#) [Zbl](#)
- [Raynaud 1983a] M. Raynaud, “Around the Mordell conjecture for function fields and a conjecture of Serge Lang”, pp. 1–19 in *Algebraic geometry* (Tokyo/Kyoto, 1982), edited by M. Raynaud and T. Shioda, Lecture Notes in Math. **1016**, Springer, 1983. [MR](#) [Zbl](#)
- [Raynaud 1983b] M. Raynaud, “Courbes sur une variété abélienne et points de torsion”, *Invent. Math.* **71**:1 (1983), 207–233. [MR](#) [Zbl](#)
- [Raynaud 1983c] M. Raynaud, “Sous-variétés d’une variété abélienne et points de torsion”, pp. 327–352 in *Arithmetic and geometry, Vol I*, edited by M. Artin and J. Tate, Progr. Math. **35**, Birkhäuser, Boston, 1983. [MR](#) [Zbl](#)
- [Rössler 2013] D. Rössler, “On the Manin–Mumford and Mordell–Lang conjectures in positive characteristic”, *Algebra Number Theory* **7**:8 (2013), 2039–2057. [MR](#) [Zbl](#)
- [Rössler 2016] D. Rössler, “Strongly semistable sheaves and the Mordell–Lang conjecture over function fields”, 2016. [arXiv](#)
- [SGA 3<sub>II</sub> 1970] M. Demazure and A. Grothendieck, *Schémas en groupes, Tome II: Groupes de type multiplicatif, et structure des schémas en groupes généraux, Exposés VIII–XVIII* (Séminaire de Géométrie Algébrique du Bois Marie 1962–1964), Lecture Notes in Math. **152**, Springer, 1970. [MR](#) [Zbl](#)
- [Szpiro 1981] L. Szpiro, “Propriétés numériques du faisceau dualisant relatif [Numerical properties of the relative dualizing sheaf]”, pp. 44–78 in *Séminaire sur les Pinceaux de Courbes de Genre au Moins Deux*, Astérisque **86**, Société Mathématique de France, Paris, 1981. [MR](#) [Zbl](#)
- [Xin 2016] H. Xin, “On  $W_2$ -lifting of Frobenius of algebraic surfaces”, *Collect. Math.* **67**:1 (2016), 69–83. [MR](#) [Zbl](#)

Communicated by Jonathan Pila

Received 2017-03-09

Revised 2017-11-09

Accepted 2017-12-19

[danny.scarponi@mathematik.uni-regensburg.de](mailto:danny.scarponi@mathematik.uni-regensburg.de)

Fakultät für Mathematik, Universität Regensburg, Germany

# Algebra & Number Theory

[msp.org/ant](http://msp.org/ant)

## EDITORS

### MANAGING EDITOR

Bjorn Poonen  
Massachusetts Institute of Technology  
Cambridge, USA

### EDITORIAL BOARD CHAIR

David Eisenbud  
University of California  
Berkeley, USA

### BOARD OF EDITORS

Richard E. Borcherds	University of California, Berkeley, USA	Martin Olsson	University of California, Berkeley, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Raman Parimala	Emory University, USA
Brian D. Conrad	Stanford University, USA	Jonathan Pila	University of Oxford, UK
Samit Dasgupta	University of California, Santa Cruz, USA	Anand Pillay	University of Notre Dame, USA
Hélène Esnault	Freie Universität Berlin, Germany	Michael Rapoport	Universität Bonn, Germany
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Victor Reiner	University of Minnesota, USA
Hubert Flenner	Ruhr-Universität, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Christopher Skinner	Princeton University, USA
Joseph Gubeladze	San Francisco State University, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Roger Heath-Brown	Oxford University, UK	J. Toby Stafford	University of Michigan, USA
Craig Huneke	University of Virginia, USA	Pham Huu Tiep	University of Arizona, USA
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Ravi Vakil	Stanford University, USA
János Kollár	Princeton University, USA	Michel van den Bergh	Hasselt University, Belgium
Yuri Manin	Northwestern University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Kei-Ichi Watanabe	Nihon University, Japan
Susan Montgomery	University of Southern California, USA	Shou-Wu Zhang	Princeton University, USA
Shigefumi Mori	RIMS, Kyoto University, Japan		

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

---

The subscription price for 2018 is US \$340/year for the electronic version, and \$535/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

---

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

ANT peer review and production are managed by EditFLOW® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 12 No. 2 2018

---

Proper $G_a$ -actions on $\mathbb{C}^4$ preserving a coordinate SHULIM KALIMAN	227
Nonemptiness of Newton strata of Shimura varieties of Hodge type DONG UK LEE	259
Towards Boij–Söderberg theory for Grassmannians: the case of square matrices NICOLAS FORD, JAKE LEVINSON and STEVEN V SAM	285
Chebyshev’s bias for products of $k$ primes XIANCHANG MENG	305
$D$ -groups and the Dixmier–Moeglin equivalence JASON BELL, OMAR LEÓN SÁNCHEZ and RAHIM MOOSA	343
Closures in varieties of representations and irreducible components KENNETH R. GOODEARL and BIRGE HUISGEN-ZIMMERMANN	379
Sparsity of $p$ -divisible unramified liftings for subvarieties of abelian varieties with trivial stabilizer DANNY SCARPONI	411
On a conjecture of Kato and Kuzumaki DIEGO IZQUIERDO	429
Height bounds and the Siegel property MARTIN ORR	455
Quadric surface bundles over surfaces and stable rationality STEFAN SCHREIEDER	479
Correction to the article Finite generation of the cohomology of some skew group algebras VAN C. NGUYEN and SARAH WITHERSPOON	491